

Relations

Rosen Section 9.1, 9.3

Tom Michael

MNF130V2020 – Week 14

Binary relations

Definition

The **Cartesian product** $A \times B$ is the set of ordered pairs

$$A \times B = \{(a, b) \mid a \in A, b \in B\}.$$

Binary relations

Definition

The **Cartesian product** $A \times B$ is the set of ordered pairs

$$A \times B = \{(a, b) \mid a \in A, b \in B\}.$$

Definition

A (binary) **relation from A to B** is a subset R of $A \times B$.

Binary relations

Definition

The **Cartesian product** $A \times B$ is the set of ordered pairs

$$A \times B = \{(a, b) \mid a \in A, b \in B\}.$$

Definition

A (binary) **relation from A to B** is a subset R of $A \times B$.

Definition

A **relation on a set A** is a relation from A to A .

Binary relations

Definition

The **Cartesian product** $A \times B$ is the set of ordered pairs

$$A \times B = \{(a, b) \mid a \in A, b \in B\}.$$

Definition

A (binary) **relation from A to B** is a subset R of $A \times B$.

Definition

A **relation on a set A** is a relation from A to A .

Example

\leq is a relation on \mathbb{R} . Formally, we can write

$$R_{\leq} = \{(a, b) \mid a, b \in \mathbb{R}, a \leq b\}$$

Relations vs. functions

- ▶ A *function* f from A to B assigns *exactly* one element of B to each element of A . The *graph* of f is the set $\{(a, f(a)) \mid a \in A\} \subset A \times B$. Hence **every function defines a binary relation**.

Relations vs. functions

- ▶ A *function* f from A to B assigns *exactly* one element of B to each element of A . The *graph* of f is the set $\{(a, f(a)) \mid a \in A\} \subset A \times B$. Hence **every function defines a binary relation**.
- ▶ Relations are a generalization of graphs of functions: **not every binary relation defines a function**.

Relations vs. functions


- ▶ A *function* f from A to B assigns *exactly* one element of B to each element of A . The *graph* of f is the set $\{(a, f(a)) \mid a \in A\} \subset A \times B$. Hence **every function defines a binary relation**.
- ▶ Relations are a generalization of graphs of functions: **not every binary relation defines a function**.
- ▶ Main difference between relations and (graphs of) functions: relations can be **one-to-many** (assigning more than one element of B to an element of A .)

Example: $R_{\leq} = \{(a, b) \mid a, b \in \mathbb{R}, a \leq b\}$

Representing relations using directed graphs

Definition

A **directed graph**, or **digraph**, $\mathcal{G} = (V, E)$ consists of a set of *vertices* (or *nodes*) V together with a set E of ordered pairs of elements of V called *edges* (or *arcs*).



$$R = \{(a, b), (a, d), (b, b), (b, d), (c, a), (c, b), (d, b)\}$$


Figure 1: A directed graph and the relation it represents

Representing relations using directed graphs

Definition

A **directed graph**, or **digraph**, $\mathcal{G} = (V, E)$ consists of a set of *vertices* (or *nodes*) V together with a set E of ordered pairs of elements of V called *edges* (or *arcs*).

- E is a relation on V , that is, a subset of $V \times V$.



$$R = \{(a, b), (a, d), (b, b), (b, d), (c, a), (c, b), (d, b)\}$$


Figure 1: A directed graph and the relation it represents

Representing relations using directed graphs

Definition

A **directed graph**, or **digraph**, $\mathcal{G} = (V, E)$ consists of a set of *vertices* (or *nodes*) V together with a set E of ordered pairs of elements of V called *edges* (or *arcs*).

- ▶ E is a relation on V , that is, a subset of $V \times V$.
- ▶ The vertex a is called the *initial* vertex of the edge (a, b) and the vertex b is called the *terminal* vertex of this edge. An edge of the form (a, a) is called a *loop*.



$$R = \{(a, b), (a, d), (b, b), (b, d), (c, a), (c, b), (d, b)\}$$


Figure 1: A directed graph and the relation it represents

Representing relations using directed graphs

Definition

A **directed graph**, or **digraph**, $\mathcal{G} = (V, E)$ consists of a set of *vertices* (or *nodes*) V together with a set E of ordered pairs of elements of V called *edges* (or *arcs*).

- ▶ E is a relation on V , that is, a subset of $V \times V$.
- ▶ The vertex a is called the *initial* vertex of the edge (a, b) and the vertex b is called the *terminal* vertex of this edge. An edge of the form (a, a) is called a *loop*.
- ▶ A relation R on a set A can be represented by the digraph $\mathcal{G} = (A, R)$



$$R = \{(a, b), (a, d), (b, b), (b, d), (c, a), (c, b), (d, b)\}$$

Figure 1: A directed graph and the relation it represents

Relations on a set: reflexive

Definition

- ▶ A relation R on a set A is **reflexive** if $(a, a) \in R$ for all $a \in A$.




Figure 2: A reflexive relation

Relations on a set: reflexive

Definition

- ▶ A relation R on a set A is **reflexive** if $(a, a) \in R$ for all $a \in A$.
- ▶ A relation is reflexive if and only if its digraph representation has a loop at every vertex.




Figure 2: A reflexive relation

Relations on a set: symmetric

Definition

- A relation R on a set A is **symmetric** if $(a, b) \in R \rightarrow (b, a) \in R$, for all $a, b \in A$.




Figure 3: A symmetric relation

Relations on a set: symmetric

Definition

- ▶ A relation R on a set A is **symmetric** if $(a, b) \in R \rightarrow (b, a) \in R$, for all $a, b \in A$.
- ▶ A relation is symmetric if and only if for every edge between distinct vertices in its digraph there is also an edge in the opposite direction.




Figure 3: A symmetric relation

Relations on a set: antisymmetric

Definition

- ▶ A relation R on a set A is **antisymmetric** if $((a, b) \in R \wedge (b, a) \in R) \rightarrow a = b$, for all $a, b \in A$.




Figure 4: An antisymmetric relation

Relations on a set: antisymmetric

Definition

- ▶ A relation R on a set A is **antisymmetric** if $((a, b) \in R \wedge (b, a) \in R) \rightarrow a = b$, for all $a, b \in A$.
- ▶ A relation is antisymmetric if and only if there are never two edges in opposite direction between two distinct vertices.




Figure 4: An antisymmetric relation

Relations on a set: antisymmetric

Definition

- ▶ A relation R on a set A is **antisymmetric** if $((a, b) \in R \wedge (b, a) \in R) \rightarrow a = b$, for all $a, b \in A$.
- ▶ A relation is antisymmetric if and only if there are never two edges in opposite direction between two distinct vertices.
- ▶ Relations on a set can be both symmetric and antisymmetric.
Example: Let $R = \{(a, a) \mid a \in A\}$.




Figure 4: An antisymmetric relation

Relations on a set: transitive

Definition

- ▶ A relation R on a set A is **transitive** if $((a, b) \in R \wedge (b, c) \in R) \rightarrow (a, c) \in R$, for all $a, b, c \in A$.




Figure 5: A transitive relation

Relations on a set: transitive

Definition

- ▶ A relation R on a set A is **transitive** if $((a, b) \in R \wedge (b, c) \in R) \rightarrow (a, c) \in R$, for all $a, b, c \in A$.
- ▶ A relation is transitive if and only if whenever there is an edge from vertex x to y and an edge from vertex y to z , there is an edge from x to z (completing triangles).




Figure 5: A transitive relation

Examples

Relation	Refl	Symm	Antisymm	Trans
----------	------	------	----------	-------

Table 1: Example relations on \mathbb{R}

Examples

Relation	Refl	Symm	Antisymm	Trans
$R_1 = \{(a, b) \mid a \leq b\}$	✓	–	✓	✓

Table 1: Example relations on \mathbb{R}

Examples

Relation	Refl	Symm	Antisymm	Trans
$R_1 = \{(a, b) \mid a \leq b\}$	✓	–	✓	✓
$R_2 = \{(a, b) \mid a > b\}$	–	–	✓	✓

Table 1: Example relations on \mathbb{R}

Examples

Relation	Refl	Symm	Antisymm	Trans
$R_1 = \{(a, b) \mid a \leq b\}$	✓	–	✓	✓
$R_2 = \{(a, b) \mid a > b\}$	–	–	✓	✓
$R_3 = \{(a, b) \mid a = \pm b\}$	✓	✓	–	✓

Table 1: Example relations on \mathbb{R}

Examples

Relation	Refl	Symm	Antisymm	Trans
$R_1 = \{(a, b) \mid a \leq b\}$	✓	–	✓	✓
$R_2 = \{(a, b) \mid a > b\}$	–	–	✓	✓
$R_3 = \{(a, b) \mid a = \pm b\}$	✓	✓	–	✓
$R_4 = \{(a, b) \mid a = b\}$	✓	✓	✓	✓

Table 1: Example relations on \mathbb{R}

Examples

Relation	Refl	Symm	Antisymm	Trans
$R_1 = \{(a, b) \mid a \leq b\}$	✓	–	✓	✓
$R_2 = \{(a, b) \mid a > b\}$	–	–	✓	✓
$R_3 = \{(a, b) \mid a = \pm b\}$	✓	✓	–	✓
$R_4 = \{(a, b) \mid a = b\}$	✓	✓	✓	✓
$R_5 = \{(a, b) \mid b = a + 1\}$	–	–	✓	–

Table 1: Example relations on \mathbb{R}

Examples

Relation	Refl	Symm	Antisymm	Trans
$R_1 = \{(a, b) \mid a \leq b\}$	✓	–	✓	✓
$R_2 = \{(a, b) \mid a > b\}$	–	–	✓	✓
$R_3 = \{(a, b) \mid a = \pm b\}$	✓	✓	–	✓
$R_4 = \{(a, b) \mid a = b\}$	✓	✓	✓	✓
$R_5 = \{(a, b) \mid b = a + 1\}$	–	–	✓	–
$R_6 = \{(a, b) \mid a \equiv b \pmod{3}\}$	✓	✓	–	✓

Table 1: Example relations on \mathbb{R}

Composite relations

Definition

The **composite** $S \circ R$ of a relation R from A to B and a relation S from B to C is the relation consisting of ordered pairs (a, c) , where




Figure 6: A composite relation

Composite relations

Definition

The **composite** $S \circ R$ of a relation R from A to B and a relation S from B to C is the relation consisting of ordered pairs (a, c) , where

- $a \in A, c \in C$




Figure 6: A composite relation

Composite relations

Definition

The **composite** $S \circ R$ of a relation R from A to B and a relation S from B to C is the relation consisting of ordered pairs (a, c) , where

- ▶ $a \in A, c \in C$
- ▶ there exists $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$.




Figure 6: A composite relation

Composite relations

Definition

The **composite** $S \circ R$ of a relation R from A to B and a relation S from B to C is the relation consisting of ordered pairs (a, c) , where

- ▶ $a \in A, c \in C$
- ▶ there exists $b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$.
- ▶ In one line:

$$S \circ R = \{(a, c) \mid a \in A, c \in C, \exists b \in B((a, b) \in R \wedge (b, c) \in S)\}$$




Figure 6: A composite relation

Composing the parent relation with itself

Definition

The **powers** R^n , $n = 1, 2, 3, \dots$, of a relation R on a set A are defined recursively by

$$R^1 = R \quad \text{and} \quad R^{n+1} = R^n \circ R$$

Example

Composing the parent relation with itself

Definition

The **powers** R^n , $n = 1, 2, 3, \dots$, of a relation R on a set A are defined recursively by

$$R^1 = R \quad \text{and} \quad R^{n+1} = R^n \circ R$$

Example

- Let $R = \{(a, b) \mid a, b \in \mathbb{N}, b - a = 1\} = \{(a, a + 1) \mid a \in \mathbb{N}\}$.

Composing the parent relation with itself

Definition

The **powers** R^n , $n = 1, 2, 3, \dots$, of a relation R on a set A are defined recursively by

$$R^1 = R \quad \text{and} \quad R^{n+1} = R^n \circ R$$

Example

- ▶ Let $R = \{(a, b) \mid a, b \in \mathbb{N}, b - a = 1\} = \{(a, a + 1) \mid a \in \mathbb{N}\}$.
- ▶ Then $(a, c) \in R^2$ if and only if $\exists b$ such that $(a, b) \in R$ and $(b, c) \in R$, that is $b = a + 1$ and $c = b + 1$, or $c = a + 2$.

Composing the parent relation with itself

Definition

The **powers** R^n , $n = 1, 2, 3, \dots$, of a relation R on a set A are defined recursively by

$$R^1 = R \quad \text{and} \quad R^{n+1} = R^n \circ R$$

Example

- ▶ Let $R = \{(a, b) \mid a, b \in \mathbb{N}, b - a = 1\} = \{(a, a + 1) \mid a \in \mathbb{N}\}$.
- ▶ Then $(a, c) \in R^2$ if and only if $\exists b$ such that $(a, b) \in R$ and $(b, c) \in R$, that is $b = a + 1$ and $c = b + 1$, or $c = a + 2$.
- ▶ Hence $R^2 = \{(a, b) \mid a, b \in \mathbb{N}, b - a = 2\}$.

Composing the parent relation with itself

Definition

The **powers** R^n , $n = 1, 2, 3, \dots$, of a relation R on a set A are defined recursively by

$$R^1 = R \quad \text{and} \quad R^{n+1} = R^n \circ R$$

Example

- ▶ Let $R = \{(a, b) \mid a, b \in \mathbb{N}, b - a = 1\} = \{(a, a + 1) \mid a \in \mathbb{N}\}$.
- ▶ Then $(a, c) \in R^2$ if and only if $\exists b$ such that $(a, b) \in R$ and $(b, c) \in R$, that is $b = a + 1$ and $c = b + 1$, or $c = a + 2$.
- ▶ Hence $R^2 = \{(a, b) \mid a, b \in \mathbb{N}, b - a = 2\}$.
- ▶ More generally, $R^n = \{(a, b) \mid a, b \in \mathbb{N}, b - a = n\}$ for all n .

Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.
 - ▶ *Induction hypothesis:* Assume $Q(k)$ is true for arbitrary $k \geq 1$.



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.
 - ▶ *Induction hypothesis:* Assume $Q(k)$ is true for arbitrary $k \geq 1$.
 - ▶ *Induction step:*



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.
 - ▶ *Induction hypothesis:* Assume $Q(k)$ is true for arbitrary $k \geq 1$.
 - ▶ *Induction step:*
 - ▶ Take $(a, c) \in R^{k+1}$ arbitrarily.



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.
 - ▶ *Induction hypothesis:* Assume $Q(k)$ is true for arbitrary $k \geq 1$.
 - ▶ *Induction step:*
 - ▶ Take $(a, c) \in R^{k+1}$ arbitrarily.
 - ▶ Because $R^{k+1} = R^k \circ R$, there must exist $b \in R$ such that $(a, b) \in R \wedge (b, c) \in R^k$.



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.
 - ▶ *Induction hypothesis:* Assume $Q(k)$ is true for arbitrary $k \geq 1$.
 - ▶ *Induction step:*
 - ▶ Take $(a, c) \in R^{k+1}$ arbitrarily.
 - ▶ Because $R^{k+1} = R^k \circ R$, there must exist $b \in R$ such that $(a, b) \in R \wedge (b, c) \in R^k$.
 - ▶ By the inductive hypothesis $R^k \subseteq R$ and hence $(b, c) \in R$.



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.
 - ▶ *Induction hypothesis:* Assume $Q(k)$ is true for arbitrary $k \geq 1$.
 - ▶ *Induction step:*
 - ▶ Take $(a, c) \in R^{k+1}$ arbitrarily.
 - ▶ Because $R^{k+1} = R^k \circ R$, there must exist $b \in R$ such that $(a, b) \in R \wedge (b, c) \in R^k$.
 - ▶ By the inductive hypothesis $R^k \subseteq R$ and hence $(b, c) \in R$.
 - ▶ By the assumption that R is transitive it follows that $(a, c) \in R$.



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.
 - ▶ *Induction hypothesis:* Assume $Q(k)$ is true for arbitrary $k \geq 1$.
 - ▶ *Induction step:*
 - ▶ Take $(a, c) \in R^{k+1}$ arbitrarily.
 - ▶ Because $R^{k+1} = R^k \circ R$, there must exist $b \in R$ such that $(a, b) \in R \wedge (b, c) \in R^k$.
 - ▶ By the inductive hypothesis $R^k \subseteq R$ and hence $(b, c) \in R$.
 - ▶ By the assumption that R is transitive it follows that $(a, c) \in R$.
 - ▶ Hence $\forall (a, c) ((a, c) \in R^{k+1} \rightarrow (a, c) \in R)$, or $R^{k+1} \subseteq R$.



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.
 - ▶ *Induction hypothesis:* Assume $Q(k)$ is true for arbitrary $k \geq 1$.
 - ▶ *Induction step:*
 - ▶ Take $(a, c) \in R^{k+1}$ arbitrarily.
 - ▶ Because $R^{k+1} = R^k \circ R$, there must exist $b \in R$ such that $(a, b) \in R \wedge (b, c) \in R^k$.
 - ▶ By the inductive hypothesis $R^k \subseteq R$ and hence $(b, c) \in R$.
 - ▶ By the assumption that R is transitive it follows that $(a, c) \in R$.
 - ▶ Hence $\forall (a, c) ((a, c) \in R^{k+1} \rightarrow (a, c) \in R)$, or $R^{k+1} \subseteq R$.
 - ▶ Hence $Q(k) \rightarrow Q(k+1)$ is true for arbitrary k .



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.
 - ▶ *Induction hypothesis:* Assume $Q(k)$ is true for arbitrary $k \geq 1$.
 - ▶ *Induction step:*
 - ▶ Take $(a, c) \in R^{k+1}$ arbitrarily.
 - ▶ Because $R^{k+1} = R^k \circ R$, there must exist $b \in R$ such that $(a, b) \in R \wedge (b, c) \in R^k$.
 - ▶ By the inductive hypothesis $R^k \subseteq R$ and hence $(b, c) \in R$.
 - ▶ By the assumption that R is transitive it follows that $(a, c) \in R$.
 - ▶ Hence $\forall (a, c) ((a, c) \in R^{k+1} \rightarrow (a, c) \in R)$, or $R^{k+1} \subseteq R$.
 - ▶ Hence $Q(k) \rightarrow Q(k+1)$ is true for arbitrary k .
- ▶ By the principle of induction $\forall n Q(n)$ is true.



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.
 - ▶ *Induction hypothesis:* Assume $Q(k)$ is true for arbitrary $k \geq 1$.
 - ▶ *Induction step:*
 - ▶ Take $(a, c) \in R^{k+1}$ arbitrarily.
 - ▶ Because $R^{k+1} = R^k \circ R$, there must exist $b \in R$ such that $(a, b) \in R \wedge (b, c) \in R^k$.
 - ▶ By the inductive hypothesis $R^k \subseteq R$ and hence $(b, c) \in R$.
 - ▶ By the assumption that R is transitive it follows that $(a, c) \in R$.
 - ▶ Hence $\forall (a, c) ((a, c) \in R^{k+1} \rightarrow (a, c) \in R)$, or $R^{k+1} \subseteq R$.
 - ▶ Hence $Q(k) \rightarrow Q(k+1)$ is true for arbitrary k .
 - ▶ By the principle of induction $\forall n Q(n)$ is true.
- ▶ Next, assume $\forall n Q(n)$ is true:



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.
 - ▶ *Induction hypothesis:* Assume $Q(k)$ is true for arbitrary $k \geq 1$.
 - ▶ *Induction step:*
 - ▶ Take $(a, c) \in R^{k+1}$ arbitrarily.
 - ▶ Because $R^{k+1} = R^k \circ R$, there must exist $b \in R$ such that $(a, b) \in R \wedge (b, c) \in R^k$.
 - ▶ By the inductive hypothesis $R^k \subseteq R$ and hence $(b, c) \in R$.
 - ▶ By the assumption that R is transitive it follows that $(a, c) \in R$.
 - ▶ Hence $\forall (a, c) ((a, c) \in R^{k+1} \rightarrow (a, c) \in R)$, or $R^{k+1} \subseteq R$.
 - ▶ Hence $Q(k) \rightarrow Q(k+1)$ is true for arbitrary k .
 - ▶ By the principle of induction $\forall n Q(n)$ is true.
- ▶ Next, assume $\forall n Q(n)$ is true:
 - ▶ Take any $a, b, c \in A$ such that $(a, b) \in R$ and $(b, c) \in R$.



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.
 - ▶ *Induction hypothesis:* Assume $Q(k)$ is true for arbitrary $k \geq 1$.
 - ▶ *Induction step:*
 - ▶ Take $(a, c) \in R^{k+1}$ arbitrarily.
 - ▶ Because $R^{k+1} = R^k \circ R$, there must exist $b \in R$ such that $(a, b) \in R \wedge (b, c) \in R^k$.
 - ▶ By the inductive hypothesis $R^k \subseteq R$ and hence $(b, c) \in R$.
 - ▶ By the assumption that R is transitive it follows that $(a, c) \in R$.
 - ▶ Hence $\forall (a, c) ((a, c) \in R^{k+1} \rightarrow (a, c) \in R)$, or $R^{k+1} \subseteq R$.
 - ▶ Hence $Q(k) \rightarrow Q(k+1)$ is true for arbitrary k .
 - ▶ By the principle of induction $\forall n Q(n)$ is true.
- ▶ Next, assume $\forall n Q(n)$ is true:
 - ▶ Take any $a, b, c \in A$ such that $(a, b) \in R$ and $(b, c) \in R$.
 - ▶ Then $(a, c) \in R^2 \subseteq R$ and hence $(a, c) \in R$.



Theorem

A relation R on a set A is transitive if and only if $R^n \subseteq R$ for $n \geq 1$.

Proof.

- ▶ Let $p = "R \text{ is transitive}"$ and $Q(n) = R^n \subseteq R$. To prove the equivalence $p \leftrightarrow \forall n Q(n)$, we first prove $p \rightarrow \forall n Q(n)$ and then $\forall n Q(n) \rightarrow p$.
- ▶ Assume p is true. We will show that this implies $\forall n Q(n)$ using mathematical induction:
 - ▶ *Basis step:* $Q(1)$ is true because $R^1 = R \subseteq R$.
 - ▶ *Induction hypothesis:* Assume $Q(k)$ is true for arbitrary $k \geq 1$.
 - ▶ *Induction step:*
 - ▶ Take $(a, c) \in R^{k+1}$ arbitrarily.
 - ▶ Because $R^{k+1} = R^k \circ R$, there must exist $b \in R$ such that $(a, b) \in R \wedge (b, c) \in R^k$.
 - ▶ By the inductive hypothesis $R^k \subseteq R$ and hence $(b, c) \in R$.
 - ▶ By the assumption that R is transitive it follows that $(a, c) \in R$.
 - ▶ Hence $\forall (a, c) ((a, c) \in R^{k+1} \rightarrow (a, c) \in R)$, or $R^{k+1} \subseteq R$.
 - ▶ Hence $Q(k) \rightarrow Q(k+1)$ is true for arbitrary k .
 - ▶ By the principle of induction $\forall n Q(n)$ is true.
- ▶ Next, assume $\forall n Q(n)$ is true:
 - ▶ Take any $a, b, c \in A$ such that $(a, b) \in R$ and $(b, c) \in R$.
 - ▶ Then $(a, c) \in R^2 \subseteq R$ and hence $(a, c) \in R$.
 - ▶ Hence p is true.



Representing relations using matrices

A relation between finite sets $A = \{a_1, a_2, \dots, a_m\}$ and $B = \{b_1, b_2, \dots, b_n\}$ can also be represented using the $m \times n$ **zero-one** or **Boolean matrix** $\mathbf{M}_R = (M_{ij})$ where

$$M_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R \\ 0 & \text{otherwise} \end{cases}$$

Example

Representing relations using matrices

A relation between finite sets $A = \{a_1, a_2, \dots, a_m\}$ and $B = \{b_1, b_2, \dots, b_n\}$ can also be represented using the $m \times n$ **zero-one** or **Boolean matrix** $\mathbf{M}_R = (M_{ij})$ where

$$M_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R \\ 0 & \text{otherwise} \end{cases}$$

Example

- Suppose that $A = \{1, 2, 3\}$ and $B = \{1, 2\}$.

Representing relations using matrices

A relation between finite sets $A = \{a_1, a_2, \dots, a_m\}$ and $B = \{b_1, b_2, \dots, b_n\}$ can also be represented using the $m \times n$ **zero-one** or **Boolean matrix** $\mathbf{M}_R = (M_{ij})$ where

$$M_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R \\ 0 & \text{otherwise} \end{cases}$$

Example

- ▶ Suppose that $A = \{1, 2, 3\}$ and $B = \{1, 2\}$.
- ▶ Let R be the relation from A to B containing (a, b) if $a > b$.

Representing relations using matrices

A relation between finite sets $A = \{a_1, a_2, \dots, a_m\}$ and $B = \{b_1, b_2, \dots, b_n\}$ can also be represented using the $m \times n$ **zero-one** or **Boolean matrix** $\mathbf{M}_R = (M_{ij})$ where

$$M_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R \\ 0 & \text{otherwise} \end{cases}$$


Example

- ▶ Suppose that $A = \{1, 2, 3\}$ and $B = \{1, 2\}$.
- ▶ Let R be the relation from A to B containing (a, b) if $a > b$.
- ▶ Then


$$\mathbf{M}_R = \begin{pmatrix} 0 & 0 \\ 1 & 0 \\ 1 & 1 \end{pmatrix}$$

Matrix properties for reflexive, symmetric and antisymmetric relations

If R is a relation on a set A (that is, $A = B$), then R is:



(a) Symmetric




(b) Antisymmetric

Figure 7: Zero-one matrices for symmetric and antisymmetric relations


Matrix properties for reflexive, symmetric and antisymmetric relations

If R is a relation on a set A (that is, $A = B$), then R is:

► **reflexive** if and only if $\text{diag}(\mathbf{M}_R) = (1, 1, \dots, 1)$.



(a) Symmetric



(b) Antisymmetric

Figure 7: Zero-one matrices for symmetric and antisymmetric relations

Matrix properties for reflexive, symmetric and antisymmetric relations

If R is a relation on a set A (that is, $A = B$), then R is:

- ▶ **reflexive** if and only if $\text{diag}(\mathbf{M}_R) = (1, 1, \dots, 1)$.
- ▶ **symmetric** if and only if $(\mathbf{M}_R)^t = \mathbf{M}_R$.





Figure 7: Zero-one matrices for symmetric and antisymmetric relations


Matrix properties for reflexive, symmetric and antisymmetric relations

If R is a relation on a set A (that is, $A = B$), then R is:

- ▶ **reflexive** if and only if $\text{diag}(\mathbf{M}_R) = (1, 1, \dots, 1)$.
- ▶ **symmetric** if and only if $(\mathbf{M}_R)^t = \mathbf{M}_R$.
- ▶ **antisymmetric** if $M_{ij} = 0 \vee M_{ji} = 0$ for all $i \neq j$.



(a) Symmetric



(b) Antisymmetric

Figure 7: Zero-one matrices for symmetric and antisymmetric relations

What to do next?

What to do next?

- ▶ Read Section 9.1 and 9.3, especially all the extra examples.

What to do next?

- ▶ Read Section 9.1 and 9.3, especially all the extra examples.
- ▶ Solve exercises. Some recommended exercises are in the assignment for week 16:
<https://mitt.uib.no/courses/21678/assignments/26393>

What to do next?

- ▶ Read Section 9.1 and 9.3, especially all the extra examples.
- ▶ Solve exercises. Some recommended exercises are in the assignment for week 16:
<https://mitt.uib.no/courses/21678/assignments/26393>
- ▶ Post questions on the discussion forum and participate in the discussion:
https://mitt.uib.no/courses/21678/discussion_topics/159090