

# Permutations and Combinations

Rosen Section 6.3

Tom Michael

MNF130V2020 – Week 13

# Permutations

## Definition

- ▶ A **permutation** of a set of distinct objects is an *ordered* arrangement of these objects.

# Permutations

## Definition

- ▶ A **permutation** of a set of distinct objects is an *ordered* arrangement of these objects.
- ▶ If  $A$  is a *set* of  $n$  objects, a permutation of  $A$  is an *element* of the  $n$ -fold Cartesian product  $A^n = A \times A \times \cdots \times A$ , that is, an *ordered  $n$ -tuple*  $(a_1, a_2, \dots, a_n)$ , with  $a_i \in A$  for  $i = 1, 2, \dots, n$ .

# Permutations

## Definition

- ▶ A **permutation** of a set of distinct objects is an *ordered* arrangement of these objects.
- ▶ If  $A$  is a *set* of  $n$  objects, a permutation of  $A$  is an *element* of the  $n$ -fold Cartesian product  $A^n = A \times A \times \cdots \times A$ , that is, an *ordered  $n$ -tuple*  $(a_1, a_2, \dots, a_n)$ , with  $a_i \in A$  for  $i = 1, 2, \dots, n$ .

## Example

Let  $A = \{a, b, c\}$ . The permutations of  $A$  are:

$(a, b, c)$

$(b, a, c)$

$(c, a, b)$

$(a, c, b)$

$(b, c, a)$

$(c, b, a)$

## Theorem

*If  $n$  is a positive integer, then there are  $n!$  permutations of a set with  $n$  distinct elements.*

Proof (direct, see also slides Section 2.3).



## Theorem

*If  $n$  is a positive integer, then there are  $n!$  permutations of a set with  $n$  distinct elements.*

Proof (direct, see also slides Section 2.3).

- ▶ The number of permutations is the same as the number of ordered  $n$ -tuples that can be made with  $n$  distinct element.



## Theorem

*If  $n$  is a positive integer, then there are  $n!$  permutations of a set with  $n$  distinct elements.*

Proof (direct, see also slides Section 2.3).

- ▶ The number of permutations is the same as the number of ordered  $n$ -tuples that can be made with  $n$  distinct element.
- ▶ By the product rule for counting, to make an ordered  $n$ -tuple, we have:

$n$  choices for the 1st element  
 $\times (n - 1)$  choices for the 2nd element  
 $\times (n - 2)$  choices for the 3rd element  
 $\times \dots$   
 $\times 2$  choices for the  $(n - 1)$ th element  
 $\times 1$  choice for the  $n$ th element



## Theorem

*If  $n$  is a positive integer, then there are  $n!$  permutations of a set with  $n$  distinct elements.*

Proof (direct, see also slides Section 2.3).

- ▶ The number of permutations is the same as the number of ordered  $n$ -tuples that can be made with  $n$  distinct element.
- ▶ By the product rule for counting, to make an ordered  $n$ -tuple, we have:

$n$  choices for the 1st element

$\times (n - 1)$  choices for the 2nd element

$\times (n - 2)$  choices for the 3rd element

$\times \dots$

$\times 2$  choices for the  $(n - 1)$ th element

$\times 1$  choice for the  $n$ th element

- ▶ This makes  $n! = n \cdot (n - 1) \cdot (n - 2) \cdot \dots \cdot 2 \cdot 1$  ways of making an ordered  $n$ -tuple.





Proof (by mathematical induction).



## Proof (by mathematical induction).

- ▶ *Basis step:* there is  $1 = 1!$  permutation of a set with 1 element.



## Proof (by mathematical induction).

- ▶ *Basis step:* there is  $1 = 1!$  permutation of a set with 1 element.
- ▶ *Induction hypothesis:* Assume that for arbitrary  $m \geq 1$ , there are  $m!$  permutations of a set with  $m$  distinct elements.



## Proof (by mathematical induction).

- ▶ *Basis step:* there is  $1 = 1!$  permutation of a set with 1 element.
- ▶ *Induction hypothesis:* Assume that for arbitrary  $m \geq 1$ , there are  $m!$  permutations of a set with  $m$  distinct elements.
- ▶ *Induction step:*



## Proof (by mathematical induction).

- ▶ *Basis step:* there is  $1 = 1!$  permutation of a set with 1 element.
- ▶ *Induction hypothesis:* Assume that for arbitrary  $m \geq 1$ , there are  $m!$  permutations of a set with  $m$  distinct elements.
- ▶ *Induction step:*
  - ▶ Let  $A$  be an arbitrary set with  $m + 1$  elements, and let  $a$  be an arbitrary element of  $A$ .



## Proof (by mathematical induction).

- ▶ *Basis step:* there is  $1 = 1!$  permutation of a set with 1 element.
- ▶ *Induction hypothesis:* Assume that for arbitrary  $m \geq 1$ , there are  $m!$  permutations of a set with  $m$  distinct elements.
- ▶ *Induction step:*
  - ▶ Let  $A$  be an arbitrary set with  $m + 1$  elements, and let  $a$  be an arbitrary element of  $A$ .
  - ▶ By the induction hypothesis, there are  $m!$  permutations of the set  $A - \{a\}$ .



## Proof (by mathematical induction).

- ▶ *Basis step:* there is  $1 = 1!$  permutation of a set with 1 element.
- ▶ *Induction hypothesis:* Assume that for arbitrary  $m \geq 1$ , there are  $m!$  permutations of a set with  $m$  distinct elements.
- ▶ *Induction step:*
  - ▶ Let  $A$  be an arbitrary set with  $m + 1$  elements, and let  $a$  be an arbitrary element of  $A$ .
  - ▶ By the induction hypothesis, there are  $m!$  permutations of the set  $A - \{a\}$ .
  - ▶ Each permutation  $(a_1, a_2, \dots, a_m)$  of  $A - \{a\}$  can be combined with  $a$  to form a permutation  $(a, a_1, a_2, \dots, a_m)$  of  $A$ .



## Proof (by mathematical induction).

- ▶ *Basis step:* there is  $1 = 1!$  permutation of a set with 1 element.
- ▶ *Induction hypothesis:* Assume that for arbitrary  $m \geq 1$ , there are  $m!$  permutations of a set with  $m$  distinct elements.
- ▶ *Induction step:*
  - ▶ Let  $A$  be an arbitrary set with  $m + 1$  elements, and let  $a$  be an arbitrary element of  $A$ .
  - ▶ By the induction hypothesis, there are  $m!$  permutations of the set  $A - \{a\}$ .
  - ▶ Each permutation  $(a_1, a_2, \dots, a_m)$  of  $A - \{a\}$  can be combined with  $a$  to form a permutation  $(a, a_1, a_2, \dots, a_m)$  of  $A$ .
  - ▶ Hence for every element  $a \in A$ , there are  $m!$  permutations of  $A$  that have  $a$  in the first place.





## Proof (by mathematical induction).

- ▶ *Basis step:* there is  $1 = 1!$  permutation of a set with 1 element.
- ▶ *Induction hypothesis:* Assume that for arbitrary  $m \geq 1$ , there are  $m!$  permutations of a set with  $m$  distinct elements.
- ▶ *Induction step:*
  - ▶ Let  $A$  be an arbitrary set with  $m + 1$  elements, and let  $a$  be an arbitrary element of  $A$ .
  - ▶ By the induction hypothesis, there are  $m!$  permutations of the set  $A - \{a\}$ .
  - ▶ Each permutation  $(a_1, a_2, \dots, a_m)$  of  $A - \{a\}$  can be combined with  $a$  to form a permutation  $(a, a_1, a_2, \dots, a_m)$  of  $A$ .
  - ▶ Hence for every element  $a \in A$ , there are  $m!$  permutations of  $A$  that have  $a$  in the first place.
  - ▶ By the product rule, because there are  $m + 1$  choices for  $a$ , the total number of permutations of  $A$  is  $(m + 1) \cdot m! = (m + 1)!$



## Proof (by mathematical induction).

- ▶ *Basis step:* there is  $1 = 1!$  permutation of a set with 1 element.
- ▶ *Induction hypothesis:* Assume that for arbitrary  $m \geq 1$ , there are  $m!$  permutations of a set with  $m$  distinct elements.
- ▶ *Induction step:*
  - ▶ Let  $A$  be an arbitrary set with  $m + 1$  elements, and let  $a$  be an arbitrary element of  $A$ .
  - ▶ By the induction hypothesis, there are  $m!$  permutations of the set  $A - \{a\}$ .
  - ▶ Each permutation  $(a_1, a_2, \dots, a_m)$  of  $A - \{a\}$  can be combined with  $a$  to form a permutation  $(a, a_1, a_2, \dots, a_m)$  of  $A$ .
  - ▶ Hence for every element  $a \in A$ , there are  $m!$  permutations of  $A$  that have  $a$  in the first place.
  - ▶ By the product rule, because there are  $m + 1$  choices for  $a$ , the total number of permutations of  $A$  is  $(m + 1) \cdot m! = (m + 1)!$
- ▶ By the principle of mathematical induction, there are  $n!$  permutations of a set with  $n$  distinct elements, for all integers  $n \geq 1$ .



# $r$ -Permutations

## Definition

- ▶ An  **$r$ -permutation** is an *ordered* arrangement of a *subset* of  $r$  elements of a set.

# $r$ -Permutations

## Definition

- ▶ An  **$r$ -permutation** is an *ordered* arrangement of a *subset* of  $r$  elements of a set.
- ▶ If  $A$  is a set of  $n$  objects, an  $r$ -permutation of  $A$  is an *element* of the  $r$ -fold Cartesian product  $A^r = A \times A \times \cdots \times A$ , that is, an *ordered  $r$ -tuple*  $(a_1, a_2, \dots, a_r)$ , with  $a_i \in A$  for  $i = 1, 2, \dots, r$ .

# $r$ -Permutations

## Definition

- ▶ An  **$r$ -permutation** is an *ordered* arrangement of a *subset* of  $r$  elements of a set.
- ▶ If  $A$  is a set of  $n$  objects, an  $r$ -permutation of  $A$  is an *element* of the  $r$ -fold Cartesian product  $A^r = A \times A \times \cdots \times A$ , that is, an *ordered  $r$ -tuple*  $(a_1, a_2, \dots, a_r)$ , with  $a_i \in A$  for  $i = 1, 2, \dots, r$ .

## Example

Let  $A = \{a, b, c\}$ . The 2-permutations of  $A$  are:

$(a, b)$	$(b, a)$
$(a, c)$	$(c, a)$
$(b, c)$	$(c, b)$

## Theorem

If  $n, r$  are positive integers with  $1 \leq r \leq n$ , then there are

$$P(n, r) = n(n-1)(n-2) \dots (n-r+1) = \frac{n!}{(n-r)!}$$

$r$ -permutations of a set with  $n$  distinct elements.

Proof.



## Theorem

If  $n, r$  are positive integers with  $1 \leq r \leq n$ , then there are

$$P(n, r) = n(n-1)(n-2) \dots (n-r+1) = \frac{n!}{(n-r)!}$$

$r$ -permutations of a set with  $n$  distinct elements.

## Proof.

- The number of  $r$ -permutations is the same as the number of ordered  $r$ -tuples that can be made with  $n$  distinct element.



## Theorem

If  $n, r$  are positive integers with  $1 \leq r \leq n$ , then there are

$$P(n, r) = n(n-1)(n-2) \dots (n-r+1) = \frac{n!}{(n-r)!}$$

*$r$ -permutations of a set with  $n$  distinct elements.*

## Proof.

- ▶ The number of  $r$ -permutations is the same as the number of ordered  $r$ -tuples that can be made with  $n$  distinct element.
- ▶ To make an ordered  $r$ -tuple, we have  $n$  choices for the 1st element,  $n-1$  choices for the 2nd element,  $n-2$  choices for the 3rd element, and so on, until there are  $(n-(r-1)) = (n-r+1)$  choices for the  $r$ th element.





## Theorem

If  $n, r$  are positive integers with  $1 \leq r \leq n$ , then there are

$$P(n, r) = n(n-1)(n-2) \dots (n-r+1) = \frac{n!}{(n-r)!}$$

$r$ -permutations of a set with  $n$  distinct elements.

## Proof.

- ▶ The number of  $r$ -permutations is the same as the number of ordered  $r$ -tuples that can be made with  $n$  distinct element.
- ▶ To make an ordered  $r$ -tuple, we have  $n$  choices for the 1st element,  $n-1$  choices for the 2nd element,  $n-2$  choices for the 3rd element, and so on, until there are  $(n-(r-1)) = (n-r+1)$  choices for the  $r$ th element.
- ▶ By the product rule for counting, this makes

$$n \cdot (n-1) \cdot (n-2) \cdot \dots \cdot (n-r+1) = \frac{n!}{(n-r)!}$$

ways of making an ordered  $r$ -tuple.



## Example

Suppose there are 8 runners in a race. The winner receives a gold medal, the second a silver medal and the third a bronze medal. How many different ways are there to award these medals, if all possible outcomes can occur and there are no ties?

## Example

Suppose there are 8 runners in a race. The winner receives a gold medal, the second a silver medal and the third a bronze medal. How many different ways are there to award these medals, if all possible outcomes can occur and there are no ties?

- The number of ways to award the medals is the number of ordered 3-permutations of a set of 8 elements.

## Example

Suppose there are 8 runners in a race. The winner receives a gold medal, the second a silver medal and the third a bronze medal. How many different ways are there to award these medals, if all possible outcomes can occur and there are no ties?

- ▶ The number of ways to award the medals is the number of ordered 3-permutations of a set of 8 elements.
- ▶ Hence there are  $P(8, 3) = 8 \cdot 7 \cdot 6 = 336$  ways to award the medals.

# Combinations

## Definition

- ▶ An **r-combination** is an *unordered* selection of  $r$  elements of a set.

# Combinations

## Definition

- ▶ An **r-combination** is an *unordered* selection of  $r$  elements of a set.
- ▶ If  $A$  is a *set* of  $n$  objects, an  $r$ -combination of  $A$  is a *subset* of  $A$  of  $r$  elements.

# Combinations

## Definition

- ▶ An **r-combination** is an *unordered* selection of  $r$  elements of a set.
- ▶ If  $A$  is a *set* of  $n$  objects, an  $r$ -combination of  $A$  is a *subset* of  $A$  of  $r$  elements.
- ▶ The number of  $r$ -combinations of a set with  $n$  elements is denoted  $C(n, r)$  or  $\binom{n}{r}$ , and is called a **binomial coefficient**.

# Combinations

## Definition

- ▶ An **r-combination** is an *unordered* selection of  $r$  elements of a set.
- ▶ If  $A$  is a *set* of  $n$  objects, an  $r$ -combination of  $A$  is a *subset* of  $A$  of  $r$  elements.
- ▶ The number of  $r$ -combinations of a set with  $n$  elements is denoted  $C(n, r)$  or  $\binom{n}{r}$ , and is called a **binomial coefficient**.

## Example

Let  $A = \{a, b, c\}$ . The 2-combinations of  $A$  are:

$$\{a, b\} = \{b, a\}$$

$$\{a, c\} = \{c, a\}$$

$$\{b, c\} = \{c, b\}$$



## Theorem

If  $n, r$  are integers with  $0 \leq r \leq n$ , then there are

$$C(n, r) = \frac{n!}{(n-r)!r!}$$

$r$ -combinations of a set with  $n$  distinct elements.

Proof.



## Theorem

If  $n, r$  are integers with  $0 \leq r \leq n$ , then there are

$$C(n, r) = \frac{n!}{(n-r)!r!}$$

$r$ -combinations of a set with  $n$  distinct elements.

## Proof.

- ▶ Let  $A$  be an arbitrary set with  $n$  elements.
- ▶ There are  $P(n, r)$   $r$ -permutations of  $A$ .



## Theorem

If  $n, r$  are integers with  $0 \leq r \leq n$ , then there are

$$C(n, r) = \frac{n!}{(n-r)!r!}$$

$r$ -combinations of a set with  $n$  distinct elements.

## Proof.

- ▶ Let  $A$  be an arbitrary set with  $n$  elements.
- ▶ There are  $P(n, r)$   $r$ -permutations of  $A$ .
- ▶ To obtain the set of  $r$ -permutations of  $A$ , we can take first an  $r$ -combination (subset of  $r$  elements) of  $A$ , and then consider all  $r$ -permutations of this subset.



## Theorem

If  $n, r$  are integers with  $0 \leq r \leq n$ , then there are

$$C(n, r) = \frac{n!}{(n-r)!r!}$$

$r$ -combinations of a set with  $n$  distinct elements.

## Proof.

- ▶ Let  $A$  be an arbitrary set with  $n$  elements.
- ▶ There are  $P(n, r)$   $r$ -permutations of  $A$ .
- ▶ To obtain the set of  $r$ -permutations of  $A$ , we can take first an  $r$ -combination (subset of  $r$  elements) of  $A$ , and then consider all  $r$ -permutations of this subset.
- ▶ By the product rule,  $P(n, r) = C(n, r) \cdot P(r, r)$ .



## Theorem

If  $n, r$  are integers with  $0 \leq r \leq n$ , then there are

$$C(n, r) = \frac{n!}{(n-r)!r!}$$

$r$ -combinations of a set with  $n$  distinct elements.

## Proof.

- ▶ Let  $A$  be an arbitrary set with  $n$  elements.
- ▶ There are  $P(n, r)$   $r$ -permutations of  $A$ .
- ▶ To obtain the set of  $r$ -permutations of  $A$ , we can take first an  $r$ -combination (subset of  $r$  elements) of  $A$ , and then consider all  $r$ -permutations of this subset.
- ▶ By the product rule,  $P(n, r) = C(n, r) \cdot P(r, r)$ .
- ▶ Hence

$$C(n, r) = \frac{P(n, r)}{P(r, r)} = \frac{n!}{(n-r)!r!}$$



### Corollary

*If  $n, r$  are integers with  $0 \leq r \leq n$ , then  $C(n, r) = C(n, n - r)$ .*

Proof.



## Corollary

If  $n, r$  are integers with  $0 \leq r \leq n$ , then  $C(n, r) = C(n, n - r)$ .

## Proof.

- ▶ This follows immediately from the formula

$$C(n, r) = \frac{n!}{(n-r)!r!} = C(n, n-r)$$



## Example

How many poker hands of 5 cards can be dealt from a standard deck of 52 cards?



## Example

How many poker hands of 5 cards can be dealt from a standard deck of 52 cards?

- ▶ Because the order of the cards does not matter, each poker hand is a subset with 5 elements of a set with 52 elements.

## Example

How many poker hands of 5 cards can be dealt from a standard deck of 52 cards?

- ▶ Because the order of the cards does not matter, each poker hand is a subset with 5 elements of a set with 52 elements.
- ▶ Hence the total number of poker hands is

$$\begin{aligned}C(52, 5) &= \frac{52!}{5!47!} = \frac{52 \cdot 51 \cdot 50 \cdot 49 \cdot 48}{5 \cdot 4 \cdot 3 \cdot 2} = 13 \cdot 17 \cdot 10 \cdot 49 \cdot 24 \\&= 2,598,960\end{aligned}$$

## Example

How many ways are there to divide a standard deck of 52 cards over 4 players?

## Example

How many ways are there to divide a standard deck of 52 cards over 4 players?

- ▶ Each player gets 13 cards.

## Example

How many ways are there to divide a standard deck of 52 cards over 4 players?

- ▶ Each player gets 13 cards.
- ▶ For the first player, the number of possible hands is the number of ways to select 13 cards from 52, or  $C(52, 13)$ .

## Example

How many ways are there to divide a standard deck of 52 cards over 4 players?

- ▶ Each player gets 13 cards.
- ▶ For the first player, the number of possible hands is the number of ways to select 13 cards from 52, or  $C(52, 13)$ .
- ▶ For the second player, for each hand of the first player, 39 cards remain, and the number of possible hands is  $C(39, 13)$ .

## Example

How many ways are there to divide a standard deck of 52 cards over 4 players?

- ▶ Each player gets 13 cards.
- ▶ For the first player, the number of possible hands is the number of ways to select 13 cards from 52, or  $C(52, 13)$ .
- ▶ For the second player, for each hand of the first player, 39 cards remain, and the number of possible hands is  $C(39, 13)$ .
- ▶ For the third player, for each hand of the first and second players, 26 cards remain, and the number of possible hands is  $C(26, 13)$ .

## Example

How many ways are there to divide a standard deck of 52 cards over 4 players?

- ▶ Each player gets 13 cards.
- ▶ For the first player, the number of possible hands is the number of ways to select 13 cards from 52, or  $C(52, 13)$ .
- ▶ For the second player, for each hand of the first player, 39 cards remain, and the number of possible hands is  $C(39, 13)$ .
- ▶ For the third player, for each hand of the first and second players, 26 cards remain, and the number of possible hands is  $C(26, 13)$ .
- ▶ For the fourth player, for each hand of the first, second, and third players, 13 cards remain and that is their only possible hand.



## Example

How many ways are there to divide a standard deck of 52 cards over 4 players?

- ▶ Each player gets 13 cards.
- ▶ For the first player, the number of possible hands is the number of ways to select 13 cards from 52, or  $C(52, 13)$ .
- ▶ For the second player, for each hand of the first player, 39 cards remain, and the number of possible hands is  $C(39, 13)$ .
- ▶ For the third player, for each hand of the first and second players, 26 cards remain, and the number of possible hands is  $C(26, 13)$ .
- ▶ For the fourth player, for each hand of the first, second, and third players, 13 cards remain and that is their only possible hand.
- ▶ By the product rule, the total number of ways to divide a standard deck of 52 cards over 4 players is

$$\begin{aligned} C(52, 13) \cdot C(39, 13) \cdot C(26, 13) \cdot 1 &= \frac{52!}{13!39!} \frac{39!}{13!26!} \frac{26!}{13!13!} = \frac{52!}{(13!)^4} \\ &= 5.3645 \times 10^{28} \end{aligned}$$

What to do next?

## What to do next?

- ▶ Read Rosen Section 6.3, especially all the extra examples.

# What to do next?

- ▶ Read Rosen Section 6.3, especially all the extra examples.
- ▶ Solve exercises. Some recommended exercises are in the assignment for week 14:  
<https://mitt.uib.no/courses/21678/assignments/26392>

# What to do next?

- ▶ Read Rosen Section 6.3, especially all the extra examples.
- ▶ Solve exercises. Some recommended exercises are in the assignment for week 14:  
<https://mitt.uib.no/courses/21678/assignments/26392>
- ▶ Post questions on the discussion forum and participate in the discussion:  
[https://mitt.uib.no/courses/21678/discussion\\_topics/156554](https://mitt.uib.no/courses/21678/discussion_topics/156554)