

# Logic

Murat Osmanoglu

# Definitions

- Its origin : logos (Greek), 'the word' or 'what is spoken'

# Definitions

- Its origin : logos (Greek), 'the word' or 'what is spoken'
- 'thought' or 'reason'

# Definitions

- Its origin : logos (Greek), 'the word' or 'what is spoken'
- 'thought' or 'reason'
- 'art of reason', or 'science of reasoning'
- systematic study of the form of valid arguments

# Definitions

- Its origin : logos (Greek), 'the word' or 'what is spoken'
- 'thought' or 'reason'
- 'art of reason', or 'science of reasoning'
- systematic study of the form of valid arguments  
(study of the difference between valid arguments and invalid arguments)

# Definitions

- Its origin : logos (Greek), 'the word' or 'what is spoken'
- 'thought' or 'reason'
- 'art of reason', or 'science of reasoning'
- systematic study of the form of valid arguments  
(study of the difference between valid arguments and invalid arguments)  
(finding out what it is that makes an **argument** valid)

# Definitions

argument :

# Definitions

argument: sequence of sentences (**propositions**); premises at the beginning and conclusion at the end



# Definitions

argument: sequence of sentences (**propositions**); premises at the beginning and conclusion at the end

if the premises are all true, then the conclusion must be true

# Definitions

argument: sequence of sentences (**propositions**); premises at the beginning and conclusion at the end

if the premises are all true, then the conclusion must be true

- 1) All men are mortal  
Socrates is a man

# Definitions

argument: sequence of sentences (**propositions**); premises at the beginning and conclusion at the end

if the premises are all true, then the conclusion must be true

- 1) All men are mortal  
Socrates is a man

---

Socrates is mortal

# Definitions

argument: sequence of sentences (**propositions**); premises at the beginning and conclusion at the end

if the premises are all true, then the conclusion must be true

- |                       |                   |
|-----------------------|-------------------|
| 1) All men are mortal |                   |
| Socrates is a man     | <b>premises</b>   |
| <hr/>                 |                   |
| Socrates is mortal    | <b>conclusion</b> |

# Definitions

argument : sequence of sentences (**propositions**); premises at the beginning and conclusion at the end

if the premises are all true, then the conclusion must be true

- |                       |                   |
|-----------------------|-------------------|
| 1) All men are mortal |                   |
| Socrates is a man     | <b>premises</b>   |
| <hr/>                 |                   |
| Socrates is mortal    | <b>conclusion</b> |

- 2) John will come to the party, or Mary will come to the party  
John will not come to the party

# Definitions

argument : sequence of sentences (**propositions**); premises at the beginning and conclusion at the end

if the premises are all true, then the conclusion must be true

- 1) All men are mortal  
Socrates is a man      **premises**
- 
- Socrates is mortal      **conclusion**

- 2) John will come to the party, or Mary will come to the party  
John will not come to the party
- 
- Mary will come to the party

# Definitions

Proposition :

# Definitions

Proposition : a sentence that states a fact, true or false (not both)  
(the truthness of the sentence can be evaluated)



# Definitions

Proposition : a sentence that states a fact, true or false (not both)  
(the truthness of the sentence can be evaluated)

- Istanbul is the biggest city of Turkey

# Definitions

Proposition : a sentence that states a fact, true or false (not both)  
(the truthness of the sentence can be evaluated)

- Istanbul is the biggest city of Turkey
- $2 + 3 = 5$

# Definitions

Proposition : a sentence that states a fact, true or false (not both)  
(the truthness of the sentence can be evaluated)

- Istanbul is the biggest city of Turkey
- $2 + 3 = 5$
- $2 + 1 = 4$

# Definitions

Proposition : a sentence that states a fact, true or false (not both)  
(the truthness of the sentence can be evaluated)

- Istanbul is the biggest city of Turkey
- $2 + 3 = 5$
- $2 + 1 = 4$
- Antalya is the capital city of Turkey

# Definitions

Proposition : a sentence that states a fact, true or false (not both)  
(the truthness of the sentence can be evaluated)

- Istanbul is the biggest city of Turkey
- $2 + 3 = 5$
- $2 + 1 = 4$
- Antalya is the capital city of Turkey
- $2 + x = 8$

# Definitions

Proposition : a sentence that states a fact, true or false (not both)  
(the truthness of the sentence can be evaluated)

- Istanbul is the biggest city of Turkey
- $2 + 3 = 5$
- $2 + 1 = 4$
- Antalya is the capital city of Turkey
- $2 + x = 8$
- Ankara is the best place to live on Earth

# Definitions

Proposition : a sentence that states a fact, true or false (not both)  
(the truthness of the sentence can be evaluated)

- Istanbul is the biggest city of Turkey
- $2 + 3 = 5$
- $2 + 1 = 4$
- Antalya is the capital city of Turkey
- $2 + x = 8$
- Ankara is the best place to live on Earth

letters  $p, q, r, s$  are mostly used to represent propositional variables

# Definitions

Proposition : a sentence that states a fact, true or false (not both)  
(the truthness of the sentence can be evaluated)

- Istanbul is the biggest city of Turkey
- $2 + 3 = 5$
- $2 + 1 = 4$
- Antalya is the capital city of Turkey
- $2 + x = 8$
- Ankara is the best place to live on Earth

letters  $p, q, r, s$  are mostly used to represent propositional variables

most of the mathematical statements are constructed by combining one or more propositions using **logical operators**  
(connectives)



# Logical Operators

Negation ( $\sim p$ ) : "it's not the case that  $p$ " or "not  $p$ " .

# Logical Operators

Negation ( $\sim p$ ) : "it's not the case that  $p$ " or "not  $p$ ".

- $p : 2 + 3 = 5$ ,

# Logical Operators

Negation ( $\sim p$ ) : "it's not the case that  $p$ " or "not  $p$ ".

- $p : 2 + 3 = 5$ ,

$\sim p$  : it is not the case that  $2 + 3 = 5$

$\sim p : 2 + 3 \neq 5$

# Logical Operators

Negation ( $\sim p$ ) : "it's not the case that  $p$ " or "not  $p$ ".

- $p : 2 + 3 = 5$ ,

$\sim p$  : it is not the case that  $2 + 3 = 5$

$\sim p : 2 + 3 \neq 5$

p	$\sim p$

# Logical Operators

Negation ( $\sim p$ ) : "it's not the case that  $p$ " or "not  $p$ ".

- $p : 2 + 3 = 5$ ,

$\sim p$  : it is not the case that  $2 + 3 = 5$

$\sim p : 2 + 3 \neq 5$

p	$\sim p$
T	
F	

# Logical Operators

Negation ( $\sim p$ ) : "it's not the case that  $p$ " or "not  $p$ ".

- $p : 2 + 3 = 5$ ,

$\sim p$  : it is not the case that  $2 + 3 = 5$

$\sim p : 2 + 3 \neq 5$

$p$	$\sim p$
T	F
F	T

# Logical Operators

Conjunction ( $p \wedge q$ ) : "p and q" .

# Logical Operators

Conjunction ( $p \wedge q$ ) : "p and q" .

- p : Ali passed the course  
q : Hasan passed the course



# Logical Operators

Conjunction ( $p \wedge q$ ) : "p and q" .

- $p$  : Ali passed the course  
 $q$  : Hasan passed the course

$p \wedge q$  : Ali and Hasan both passed the course.

# Logical Operators

Conjunction ( $p \wedge q$ ) : "p and q" .

- $p$  : Ali passed the course  
 $q$  : Hasan passed the course

$p \wedge q$  : Ali and Hasan both passed the course.

$p$	$q$	$p \wedge q$

# Logical Operators

Conjunction ( $p \wedge q$ ) : "p and q" .

- $p$  : Ali passed the course  
 $q$  : Hasan passed the course

$p \wedge q$  : Ali and Hasan both passed the course.

$p$	$q$	$p \wedge q$
T	T	
T	F	
F	T	
F	F	

# Logical Operators

Conjunction ( $p \wedge q$ ) : "p and q" .

- $p$  : Ali passed the course  
 $q$  : Hasan passed the course

$p \wedge q$  : Ali and Hasan both passed the course.

$p$	$q$	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

# Logical Operators

Disjunction ( $p \vee q$ ) : "p or q" .

# Logical Operators

Disjunction ( $p \vee q$ ) : "p or q" .

- p : Ali passed the course  
q : Hasan passed the course

# Logical Operators

Disjunction ( $p \vee q$ ) : "p or q" .

- $p$  : Ali passed the course  
 $q$  : Hasan passed the course

$p \vee q$  : Ali or Hasan passed the course.

# Logical Operators

Disjunction ( $p \vee q$ ) : "p or q" .

- $p$  : Ali passed the course  
 $q$  : Hasan passed the course

$p \vee q$  : Ali or Hasan passed the course.

$p$	$q$	$p \vee q$



# Logical Operators

Disjunction ( $p \vee q$ ) : "p or q" .

- $p$  : Ali passed the course  
 $q$  : Hasan passed the course

$p \vee q$  : Ali or Hasan passed the course.

p	q	$p \vee q$
T	T	
T	F	
F	T	
F	F	

# Logical Operators

Disjunction ( $p \vee q$ ) : "p or q" .

- $p$  : Ali passed the course  
 $q$  : Hasan passed the course

$p \vee q$  : Ali or Hasan passed the course.

$p$	$q$	$p \vee q$
T	T	T
T	F	T
F	T	T
F	F	F

# Logical Operators

Exclusive or ( $p \oplus q$ ) : “p exclusive or q” .

# Logical Operators

Exclusive or ( $p \oplus q$ ) : "p exclusive or q" .

- p : Ali passed the course  
q : Hasan passed the course

# Logical Operators

Exclusive or ( $p \oplus q$ ) : “p exclusive or q” .

- $p$  : Ali passed the course  
 $q$  : Hasan passed the course

$p \oplus q$  : Ali or Hasan, but not both, passed the course.

# Logical Operators

Exclusive or ( $p \oplus q$ ) : "p exclusive or q" .

- $p$  : Ali passed the course  
 $q$  : Hasan passed the course

$p \oplus q$  : Ali or Hasan, but not both, passed the course.

$p$	$q$	$p \oplus q$

# Logical Operators

Exclusive or ( $p \oplus q$ ) : "p exclusive or q" .

- $p$  : Ali passed the course  
 $q$  : Hasan passed the course

$p \oplus q$  : Ali or Hasan, but not both, passed the course.

$p$	$q$	$p \oplus q$
T	T	
T	F	
F	T	
F	F	

# Logical Operators

Exclusive or ( $p \oplus q$ ) : "p exclusive or q" .

- $p$  : Ali passed the course  
 $q$  : Hasan passed the course

$p \oplus q$  : Ali or Hasan, but not both, passed the course.

$p$	$q$	$p \oplus q$
T	T	F
T	F	T
F	T	T
F	F	F



# Logical Operators

Conditional Statement ( $p \rightarrow q$ ) : "if p, then q" (p implies q).

# Logical Operators

Conditional Statement ( $p \rightarrow q$ ) : "if p, then q" (p implies q).

- p : it is raining  
q : the ground is wet

# Logical Operators

Conditional Statement ( $p \rightarrow q$ ) : "if p, then q" (p implies q).

- p : it is raining  
q : the ground is wet

$p \rightarrow q$  : If it is raining, then the ground is wet.

# Logical Operators

Conditional Statement ( $p \rightarrow q$ ) : "if p, then q" (p implies q).

- p : it is raining  
q : the ground is wet

$p \rightarrow q$  : If it is raining, then the ground is wet.

p	q	$p \rightarrow q$

# Logical Operators

Conditional Statement ( $p \rightarrow q$ ) : "if p, then q" (p implies q).

- p : it is raining  
q : the ground is wet

$p \rightarrow q$  : If it is raining, then the ground is wet.

p	q	$p \rightarrow q$
T	T	
T	F	
F	T	
F	F	

# Logical Operators

Conditional Statement ( $p \rightarrow q$ ) : "if p, then q" (p implies q).

- p : it is raining  
q : the ground is wet

$p \rightarrow q$  : If it is raining, then the ground is wet.

p	q	$p \rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

# Logical Operators

Conditional Statement ( $p \rightarrow q$ ) : "if p, then q" (p implies q).

- p : it is raining  
q : the ground is wet

$p \rightarrow q$  : If it is raining, then the ground is wet.

# Logical Operators

Conditional Statement ( $p \rightarrow q$ ) : "if p, then q" (p implies q).

- p : it is raining  
q : the ground is wet

$p \rightarrow q$  : If it is raining, then the ground is wet.

- the converse of  $p \rightarrow q$  :  $q \rightarrow p$   
if the ground is wet, then it is raining



# Logical Operators

Conditional Statement ( $p \rightarrow q$ ) : "if p, then q" (p implies q).

- p : it is raining  
q : the ground is wet

$p \rightarrow q$  : If it is raining, then the ground is wet.

- the converse of  $p \rightarrow q$  :  $q \rightarrow p$   
if the ground is wet, then it is raining
- the contrapositive of  $p \rightarrow q$  :  $\sim q \rightarrow \sim p$   
if the ground is not wet, then it is not raining

# Logical Operators

Conditional Statement ( $p \rightarrow q$ ) : "if p, then q" (p implies q).

- p : it is raining  
q : the ground is wet

$p \rightarrow q$  : If it is raining, then the ground is wet.

- the converse of  $p \rightarrow q$  :  $q \rightarrow p$   
if the ground is wet, then it is raining
- the contrapositive of  $p \rightarrow q$  :  $\sim q \rightarrow \sim p$   
if the ground is not wet, then it is not raining
- the inverse of  $p \rightarrow q$  :  $\sim p \rightarrow \sim q$   
if it is not raining, then the ground is not wet

# Logical Operators

Biconditional Statement ( $p \leftrightarrow q$ ) : "p if and only if q" (p implies q and q implies p).

# Logical Operators

Biconditional Statement ( $p \leftrightarrow q$ ) : "p if and only if q" (p implies q and q implies p).

- p : you can take the flight  
q : you have a ticket

# Logical Operators

Biconditional Statement ( $p \leftrightarrow q$ ) : "p if and only if q" (p implies q and q implies p).

- p : you can take the flight  
q : you have a ticket

$p \leftrightarrow q$  : you can take the flight if and only if you have a ticket.

# Logical Operators

Biconditional Statement ( $p \leftrightarrow q$ ) : "p if and only if q" (p implies q and q implies p).

- p : you can take the flight  
q : you have a ticket

$p \leftrightarrow q$  : you can take the flight if and only if you have a ticket.

p	q	$p \rightarrow q$	$q \rightarrow p$	$p \leftrightarrow q$

# Logical Operators

Biconditional Statement ( $p \leftrightarrow q$ ) : "p if and only if q" (p implies q and q implies p).

- p : you can take the flight  
q : you have a ticket

$p \leftrightarrow q$  : you can take the flight if and only if you have a ticket.

p	q	$p \rightarrow q$	$q \rightarrow p$	$p \leftrightarrow q$
T	T			
T	F			
F	T			
F	F			

# Logical Operators

Biconditional Statement ( $p \leftrightarrow q$ ) : "p if and only if q" (p implies q and q implies p).

- p : you can take the flight  
q : you have a ticket

$p \leftrightarrow q$  : you can take the flight if and only if you have a ticket.

p	q	$p \rightarrow q$	$q \rightarrow p$	$p \leftrightarrow q$
T	T	T	T	
T	F	T	F	
F	T	F	T	
F	F	T	T	



# Logical Operators

Biconditional Statement ( $p \leftrightarrow q$ ) : "p if and only if q" (p implies q and q implies p).

- p : you can take the flight  
q : you have a ticket

$p \leftrightarrow q$  : you can take the flight if and only if you have a ticket.

p	q	$p \rightarrow q$	$q \rightarrow p$	$p \leftrightarrow q$
T	T	T	T	T
T	F	T	F	F
F	T	F	T	F
F	F	T	T	T

# Logical Operators

## Truth Tables

$$(p \vee \sim q) \rightarrow (p \wedge q)$$

# Logical Operators

## Truth Tables

$$(p \vee \sim q) \rightarrow (p \wedge q)$$

p	q	$\sim q$	$p \wedge q$	$p \vee \sim q$	$(p \vee \sim q) \rightarrow (p \wedge q)$

# Logical Operators

## Truth Tables

$$(p \vee \sim q) \rightarrow (p \wedge q)$$

p	q	$\sim q$	$p \wedge q$	$p \vee \sim q$	$(p \vee \sim q) \rightarrow (p \wedge q)$
1	1				
1	0				
0	1				
0	0				

# Logical Operators

## Truth Tables

$$(p \vee \sim q) \rightarrow (p \wedge q)$$

p	q	$\sim q$	$p \wedge q$	$p \vee \sim q$	$(p \vee \sim q) \rightarrow (p \wedge q)$
1	1	0			
1	0	1			
0	1	0			
0	0	1			

# Logical Operators

## Truth Tables

$$(p \vee \sim q) \rightarrow (p \wedge q)$$

p	q	$\sim q$	$p \wedge q$	$p \vee \sim q$	$(p \vee \sim q) \rightarrow (p \wedge q)$
1	1	0	1		
1	0	1	0		
0	1	0	0		
0	0	1	0		

# Logical Operators

## Truth Tables

$$(p \vee \sim q) \rightarrow (p \wedge q)$$

p	q	$\sim q$	$p \wedge q$	$p \vee \sim q$	$(p \vee \sim q) \rightarrow (p \wedge q)$
1	1	0	1	1	
1	0	1	0	1	
0	1	0	0	0	
0	0	1	0	1	

# Logical Operators

## Truth Tables

$$(p \vee \sim q) \rightarrow (p \wedge q)$$

p	q	$\sim q$	$p \wedge q$	$p \vee \sim q$	$(p \vee \sim q) \rightarrow (p \wedge q)$
1	1	0	1	1	1
1	0	1	0	1	0
0	1	0	0	0	1
0	0	1	0	1	0



# Logical Operators

## Truth Tables

$$q \leftrightarrow (\sim p \vee \sim q)$$

# Logical Operators

## Truth Tables

$$q \leftrightarrow (\sim p \vee \sim q)$$

p	q	$\sim p$	$\sim q$	$\sim p \vee \sim q$	$q \leftrightarrow (\sim p \vee \sim q)$

# Logical Operators

## Truth Tables

$$q \leftrightarrow (\sim p \vee \sim q)$$

p	q	$\sim p$	$\sim q$	$\sim p \vee \sim q$	$q \leftrightarrow (\sim p \vee \sim q)$
1	1				
1	0				
0	1				
0	0				

# Logical Operators

## Truth Tables

$$q \leftrightarrow (\sim p \vee \sim q)$$

p	q	$\sim p$	$\sim q$	$\sim p \vee \sim q$	$q \leftrightarrow (\sim p \vee \sim q)$
1	1	0	0		
1	0	0	1		
0	1	1	0		
0	0	1	1		

# Logical Operators

## Truth Tables

$$q \leftrightarrow (\sim p \vee \sim q)$$

p	q	$\sim p$	$\sim q$	$\sim p \vee \sim q$	$q \leftrightarrow (\sim p \vee \sim q)$
1	1	0	0	0	
1	0	0	1	1	
0	1	1	0	1	
0	0	1	1	1	

# Logical Operators

## Truth Tables

$$q \leftrightarrow (\sim p \vee \sim q)$$

p	q	$\sim p$	$\sim q$	$\sim p \vee \sim q$	$q \leftrightarrow (\sim p \vee \sim q)$
1	1	0	0	0	0
1	0	0	1	1	0
0	1	1	0	1	1
0	0	1	1	1	0

# Logical Operators

## Truth Tables

$$p \rightarrow (p \vee q)$$

$$p \wedge (\sim p \wedge q)$$

# Logical Operators

## Truth Tables

$$p \rightarrow (p \vee q)$$

$$p \wedge (\sim p \wedge q)$$

p	q	$p \vee q$	$p \rightarrow (p \vee q)$	$\sim p$	$\sim p \wedge q$	$p \wedge (\sim p \wedge q)$



# Logical Operators

## Truth Tables

$$p \rightarrow (p \vee q)$$

$$p \wedge (\sim p \wedge q)$$

p	q	$p \vee q$	$p \rightarrow (p \vee q)$	$\sim p$	$\sim p \wedge q$	$p \wedge (\sim p \wedge q)$
1	1					
1	0					
0	1					
0	0					

# Logical Operators

## Truth Tables

$$p \rightarrow (p \vee q)$$

$$p \wedge (\sim p \wedge q)$$

p	q	$p \vee q$	$p \rightarrow (p \vee q)$	$\sim p$	$\sim p \wedge q$	$p \wedge (\sim p \wedge q)$
1	1	1		0	0	
1	0	1		0	0	
0	1	1		1	1	
0	0	0		1	0	

# Logical Operators

## Truth Tables

$$p \rightarrow (p \vee q)$$

$$p \wedge (\sim p \wedge q)$$

p	q	$p \vee q$	$p \rightarrow (p \vee q)$	$\sim p$	$\sim p \wedge q$	$p \wedge (\sim p \wedge q)$
1	1	1	1	0	0	
1	0	1	1	0	0	
0	1	1	1	1	1	
0	0	0	1	1	0	

# Logical Operators

## Truth Tables

$$p \rightarrow (p \vee q)$$

$$p \wedge (\sim p \wedge q)$$

p	q	$p \vee q$	$p \rightarrow (p \vee q)$	$\sim p$	$\sim p \wedge q$	$p \wedge (\sim p \wedge q)$
1	1	1	1	0	0	0
1	0	1	1	0	0	0
0	1	1	1	1	1	0
0	0	0	1	1	0	0

# Logical Operators

## Truth Tables

$$p \rightarrow (p \vee q)$$

$$p \wedge (\sim p \wedge q)$$

p	q	$p \vee q$	$p \rightarrow (p \vee q)$	$\sim p$	$\sim p \wedge q$	$p \wedge (\sim p \wedge q)$
1	1	1	1	0	0	0
1	0	1	1	0	0	0
0	1	1	1	1	1	0
0	0	0	1	1	0	0

- A compound proposition is called **tautology** if it's true for all the cases

# Logical Operators

## Truth Tables

$$p \rightarrow (p \vee q)$$

$$p \wedge (\sim p \wedge q)$$

p	q	$p \vee q$	$p \rightarrow (p \vee q)$	$\sim p$	$\sim p \wedge q$	$p \wedge (\sim p \wedge q)$
1	1	1	1	0	0	0
1	0	1	1	0	0	0
0	1	1	1	1	1	0
0	0	0	1	1	0	0

- A compound proposition is called **tautology** if it's true for all the cases
- A compound proposition is called **contradiction** if it's false for all the cases

# Logical Equivalences

# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**



# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**

$p$	$q$	$\sim p$	$\sim p \vee q$	$p \rightarrow q$

# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**

$p$	$q$	$\sim p$	$\sim p \vee q$	$p \rightarrow q$
1	1			
1	0			
0	1			
0	0			

# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**

$p$	$q$	$\sim p$	$\sim p \vee q$	$p \rightarrow q$
1	1	0		
1	0	0		
0	1	1		
0	0	1		

# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**

$p$	$q$	$\sim p$	$\sim p \vee q$	$p \rightarrow q$
1	1	0	1	
1	0	0	0	
0	1	1	1	
0	0	1	1	

# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**

$p$	$q$	$\sim p$	$\sim p \vee q$	$p \rightarrow q$
1	1	0	1	1
1	0	0	0	0
0	1	1	1	1
0	0	1	1	1

# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**

$p$	$q$	$\sim p$	$\sim p \vee q$	$p \rightarrow q$
1	1	0	1	1
1	0	0	0	0
0	1	1	1	1
0	0	1	1	1

$$\sim p \vee q \equiv p \rightarrow q$$

# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**

$p$	$q$	$\sim p$	$\sim p \vee q$	$p \rightarrow q$
1	1	0	1	1
1	0	0	0	0
0	1	1	1	1
0	0	1	1	1

$$\sim p \vee q \equiv p \rightarrow q$$

$p$	$q$	$\sim p$	$\sim q$	$\sim(p \wedge q)$	$\sim p \vee \sim q$

# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**

$p$	$q$	$\sim p$	$\sim p \vee q$	$p \rightarrow q$
1	1	0	1	1
1	0	0	0	0
0	1	1	1	1
0	0	1	1	1

$$\sim p \vee q \equiv p \rightarrow q$$

$p$	$q$	$\sim p$	$\sim q$	$\sim(p \wedge q)$	$\sim p \vee \sim q$
1	1				
1	0				
0	1				
0	0				



# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**

$p$	$q$	$\sim p$	$\sim p \vee q$	$p \rightarrow q$
1	1	0	1	1
1	0	0	0	0
0	1	1	1	1
0	0	1	1	1

$$\sim p \vee q \equiv p \rightarrow q$$

$p$	$q$	$\sim p$	$\sim q$	$\sim(p \wedge q)$	$\sim p \vee \sim q$
1	1	0	0		
1	0	0	1		
0	1	1	0		
0	0	0	1		

# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**

$p$	$q$	$\sim p$	$\sim p \vee q$	$p \rightarrow q$
1	1	0	1	1
1	0	0	0	0
0	1	1	1	1
0	0	1	1	1

$$\sim p \vee q \equiv p \rightarrow q$$

$p$	$q$	$\sim p$	$\sim q$	$\sim(p \wedge q)$	$\sim p \vee \sim q$
1	1	0	0	0	
1	0	0	1	1	
0	1	1	0	1	
0	0	0	1	1	

# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**

$p$	$q$	$\sim p$	$\sim p \vee q$	$p \rightarrow q$
1	1	0	1	1
1	0	0	0	0
0	1	1	1	1
0	0	1	1	1

$$\sim p \vee q \equiv p \rightarrow q$$

$p$	$q$	$\sim p$	$\sim q$	$\sim(p \wedge q)$	$\sim p \vee \sim q$
1	1	0	0	0	0
1	0	0	1	1	1
0	1	1	0	1	1
0	0	0	1	1	1

# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**

$p$	$q$	$\sim p$	$\sim p \vee q$	$p \rightarrow q$
1	1	0	1	1
1	0	0	0	0
0	1	1	1	1
0	0	1	1	1

$$\sim p \vee q \equiv p \rightarrow q$$

$$\sim(p \wedge q) \equiv \sim p \vee \sim q$$

$p$	$q$	$\sim p$	$\sim q$	$\sim(p \wedge q)$	$\sim p \vee \sim q$
1	1	0	0	0	0
1	0	0	1	1	1
0	1	1	0	1	1
0	0	0	1	1	1

# Logical Equivalences

- If the compound propositions  $p$  and  $q$  have same truth values for all the cases, they are called **logically equivalent**

$p$	$q$	$\sim p$	$\sim p \vee q$	$p \rightarrow q$
1	1	0	1	1
1	0	0	0	0
0	1	1	1	1
0	0	1	1	1

$$\sim p \vee q \equiv p \rightarrow q$$

## De Morgan's Law

$$\sim(p \vee q) \equiv \sim p \wedge \sim q$$

$$\sim(p \wedge q) \equiv \sim p \vee \sim q$$

$p$	$q$	$\sim p$	$\sim q$	$\sim(p \wedge q)$	$\sim p \vee \sim q$
1	1	0	0	0	0
1	0	0	1	1	1
0	1	1	0	1	1
0	0	0	1	1	1

# Logical Equivalences

- De Morgan's Law

$$\sim(p \vee q) \equiv \sim p \wedge \sim q$$

$$\sim(p \wedge q) \equiv \sim p \vee \sim q$$

- $\sim(\sim p) \equiv p$

- $p \wedge 1 \equiv p$

$$p \vee 0 \equiv p$$

- $p \wedge 0 \equiv 0$

$$p \vee 1 \equiv 1$$

- $p \wedge p \equiv p$

$$p \vee p \equiv p$$

- $(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$

$$(p \vee q) \vee r \equiv p \vee (q \vee r)$$

- $p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$

$$p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$$

- $p \wedge \sim p \equiv 0$

$$p \vee \sim p \equiv 1$$

- $p \rightarrow q \equiv \sim p \vee q$

$$p \rightarrow q \equiv \sim q \rightarrow \sim p$$

# Logical Equivalences

- $\sim(p \vee (\sim p \wedge q)) \equiv$

# Logical Equivalences

- $\sim(p \vee (\sim p \wedge q)) \equiv \sim p \wedge \sim(\sim p \wedge q)$



# Logical Equivalences

- $\sim(p \vee (\sim p \wedge q)) \equiv \sim p \wedge \sim(\sim p \wedge q)$   
 $\equiv \sim p \wedge (p \vee \sim q)$

# Logical Equivalences

- $\sim(p \vee (\sim p \wedge q)) \equiv \sim p \wedge \sim(\sim p \wedge q)$   
 $\equiv \sim p \wedge (p \vee \sim q)$   
 $\equiv (\sim p \wedge p) \vee (\sim p \wedge \sim q)$

# Logical Equivalences

- $\sim(p \vee (\sim p \wedge q)) \equiv \sim p \wedge \sim(\sim p \wedge q)$   
 $\equiv \sim p \wedge (p \vee \sim q)$   
 $\equiv (\sim p \wedge p) \vee (\sim p \wedge \sim q)$   
 $\equiv 0 \vee (\sim p \wedge \sim q)$

# Logical Equivalences

- $\sim(p \vee (\sim p \wedge q)) \equiv \sim p \wedge \sim(\sim p \wedge q)$   
 $\equiv \sim p \wedge (p \vee \sim q)$   
 $\equiv (\sim p \wedge p) \vee (\sim p \wedge \sim q)$   
 $\equiv 0 \vee (\sim p \wedge \sim q)$   
 $\equiv \sim p \wedge \sim q$

# Logical Equivalences

- $\sim(p \vee (\sim p \wedge q)) \equiv \sim p \wedge \sim(\sim p \wedge q)$   
 $\equiv \sim p \wedge (p \vee \sim q)$   
 $\equiv (\sim p \wedge p) \vee (\sim p \wedge \sim q)$   
 $\equiv 0 \vee (\sim p \wedge \sim q)$   
 $\equiv \sim p \wedge \sim q$
- $(p \rightarrow r) \wedge (p \rightarrow q) \equiv$

# Logical Equivalences

- $\sim(p \vee (\sim p \wedge q)) \equiv \sim p \wedge \sim(\sim p \wedge q)$   
 $\equiv \sim p \wedge (p \vee \sim q)$   
 $\equiv (\sim p \wedge p) \vee (\sim p \wedge \sim q)$   
 $\equiv 0 \vee (\sim p \wedge \sim q)$   
 $\equiv \sim p \wedge \sim q$
- $(p \rightarrow r) \wedge (p \rightarrow q) \equiv (\sim p \vee r) \wedge$

# Logical Equivalences

- $\sim(p \vee (\sim p \wedge q)) \equiv \sim p \wedge \sim(\sim p \wedge q)$   
 $\equiv \sim p \wedge (p \vee \sim q)$   
 $\equiv (\sim p \wedge p) \vee (\sim p \wedge \sim q)$   
 $\equiv 0 \vee (\sim p \wedge \sim q)$   
 $\equiv \sim p \wedge \sim q$
- $(p \rightarrow r) \wedge (p \rightarrow q) \equiv (\sim p \vee r) \wedge (\sim p \vee q)$

# Logical Equivalences

- $\sim(p \vee (\sim p \wedge q)) \equiv \sim p \wedge \sim(\sim p \wedge q)$   
 $\equiv \sim p \wedge (p \vee \sim q)$   
 $\equiv (\sim p \wedge p) \vee (\sim p \wedge \sim q)$   
 $\equiv 0 \vee (\sim p \wedge \sim q)$   
 $\equiv \sim p \wedge \sim q$
- $(p \rightarrow r) \wedge (p \rightarrow q) \equiv (\sim p \vee r) \wedge (\sim p \vee q)$   
 $\equiv \sim p \vee (r \wedge q)$



# Logical Equivalences

- $\sim(p \vee (\sim p \wedge q)) \equiv \sim p \wedge \sim(\sim p \wedge q)$   
 $\equiv \sim p \wedge (p \vee \sim q)$   
 $\equiv (\sim p \wedge p) \vee (\sim p \wedge \sim q)$   
 $\equiv 0 \vee (\sim p \wedge \sim q)$   
 $\equiv \sim p \wedge \sim q$
- $(p \rightarrow r) \wedge (p \rightarrow q) \equiv (\sim p \vee r) \wedge (\sim p \vee q)$   
 $\equiv \sim p \vee (r \wedge q)$   
 $\equiv p \rightarrow (r \wedge q)$

# Propositional Logic

- deals with propositions and logical relationship between propositions

(has only propositions and logical operators to represent, to interpret, to infer the knowledge)

(utilizes only propositions and operators to establish valid arguments)

- If Hasan does not die, then Zeynep will not get any money and Hasan's family will be happy

p : Hasan dies

q : Zenep will get money

r : Hasan's family will be happy

$$\sim p \rightarrow (\sim q \wedge r)$$

# Propositional Logic

- deals with propositions and logical relationship between propositions

(has only propositions and logical operators to represent, to interpret, to infer the knowledge)

(utilizes only propositions and operators to establish valid arguments)

- All men are mortal  
Socrates is a man

---

Socrates is mortal

How do we represent such argument ?

- propositional logic does not work for 'all' everyone'
- propositional logic does not have quantifiers

# Predicate Logic

# Predicate Logic

- $p : '2 + 3 = 5'$

$q : 'my\ computer\ is\ vulnerable\ to\ side\ channel\ attacks'$

# Predicate Logic

- $p : '2 + 3 = 5'$

$q : 'my\ computer\ is\ vulnerable\ to\ side\ channel\ attacks'$

- $'x + 3 = 5'$

$'computer\ x\ is\ vulnerable\ to\ side\ channel\ attacks'$

# Predicate Logic

- $p : '2 + 3 = 5'$

$q : 'my\ computer\ is\ vulnerable\ to\ side\ channel\ attacks'$

- $'x + 3 = 5'$

$'computer\ x\ is\ vulnerable\ to\ side\ channel\ attacks'$

Definition Propositions (or statements) that contains variables

# Predicate Logic

- $p : '2 + 3 = 5'$

$q : 'my\ computer\ is\ vulnerable\ to\ side\ channel\ attacks'$

- $P(x) : 'x + 3 = 5'$

$Q(x) : 'computer\ x\ is\ vulnerable\ to\ side\ channel\ attacks'$

Definition Propositions (or statements) that contains variables



# Predicate Logic

- $p : '2 + 3 = 5'$

$q : 'my\ computer\ is\ vulnerable\ to\ side\ channel\ attacks'$

- $P(x) : 'x + 3 = 5'$

$Q(x) : 'computer\ x\ is\ vulnerable\ to\ side\ channel\ attacks'$

Definition Propositions (or statements) that contains variables

- When a value is assigned to the variable  $x$ , then  $P(x)$  becomes a proposition and has a truth value.

# Predicate Logic

- $P(x) : 'x > 3'$
- $Q(x,y) : 'x + 3 = y'$
- $R(x,y,z) : 'x + y = z'$

# Predicate Logic

- $P(x) : 'x > 3'$   
 $P(4)$  is true
- $Q(x,y) : 'x + 3 = y'$
- $R(x,y,z) : 'x + y = z'$

# Predicate Logic

- $P(x) : 'x > 3'$   
 $P(4)$  is true, but  $P(2)$  is false
- $Q(x,y) : 'x + 3 = y'$
- $R(x,y,z) : 'x + y = z'$

# Predicate Logic

- $P(x) : 'x > 3'$   
 $P(4)$  is true, but  $P(2)$  is false
- $Q(x,y) : 'x + 3 = y'$   
 $Q(4,7)$  is true, but  $Q(4,2)$  is false
- $R(x,y,z) : 'x + y = z'$

# Predicate Logic

- $P(x) : 'x > 3'$   
 $P(4)$  is true, but  $P(2)$  is false
- $Q(x,y) : 'x + 3 = y'$   
 $Q(4,7)$  is true, but  $Q(4,2)$  is false
- $R(x,y,z) : 'x + y = z'$   
 $R(2,1,3)$  is true, but  $R(3,2,2)$  is false

# Quantifiers

- Another way of creating a proposition from a propositional function

# Quantifiers

- Another way of creating a proposition from a propositional function

## Universal Quantifier



# Quantifiers

- Another way of creating a proposition from a propositional function

## Universal Quantifier

$$Q : \forall x P(x)$$

# Quantifiers

- Another way of creating a proposition from a propositional function

## Universal Quantifier

$Q : \forall x P(x)$       If  $P(x)$  is true for all  $x$  in the domain,  
then  $Q$  is true

# Quantifiers

- Another way of creating a proposition from a propositional function

## Universal Quantifier

$Q : \forall x P(x)$

If  $P(x)$  is true **for all  $x$**  in the domain,  
then  $Q$  is true

If there is an  $x_0$  such that  $P(x_0)$  is not  
true, then  $Q$  is false

# Quantifiers

- Another way of creating a proposition from a propositional function

## Universal Quantifier

$Q : \forall x P(x)$       If  $P(x)$  is true **for all  $x$**  in the domain,  
then  $Q$  is true  
If there is an  $x_0$  such that  $P(x_0)$  is not  
true, then  $Q$  is false

## Existential Quantifier

# Quantifiers

- Another way of creating a proposition from a propositional function

## Universal Quantifier

$Q : \forall x P(x)$

If  $P(x)$  is true for all  $x$  in the domain,  
then  $Q$  is true

If there is an  $x_0$  such that  $P(x_0)$  is not  
true, then  $Q$  is false

## Existential Quantifier

$R : \exists x P(x)$

# Quantifiers

- Another way of creating a proposition from a propositional function

## Universal Quantifier

$Q : \forall x P(x)$       If  $P(x)$  is true **for all  $x$**  in the domain,  
then  $Q$  is true  
If there is an  $x_0$  such that  $P(x_0)$  is not  
true, then  $Q$  is false

## Existential Quantifier

$R : \exists x P(x)$       If there **exists an  $x_0$**  such that  $P(x_0)$  is true,  
then  $R$  is true

# Quantifiers

- Another way of creating a proposition from a propositional function

## Universal Quantifier

$Q : \forall x P(x)$       If  $P(x)$  is true **for all  $x$**  in the domain,  
then  $Q$  is true  
If there is an  $x_0$  such that  $P(x_0)$  is not  
true, then  $Q$  is false

## Existential Quantifier

$R : \exists x P(x)$       If there **exists an  $x_0$**  such that  $P(x_0)$  is true,  
then  $R$  is true  
If  $P(x)$  is false for all  $x$  in the domain,  
then  $R$  is false

# Quantifiers

- $P(x) : x^2 \geq x$

What is the truth value of  $\forall x P(x)$  if the domain is  $\mathbb{Z}^+$  ?



# Quantifiers

- $P(x) : x^2 \geq x$

What is the truth value of  $\forall x P(x)$  if the domain is  $Z^+$  ?

For all  $x \in Z^+$   $x^2 \geq x$  .

# Quantifiers

- $P(x) : x^2 \geq x$

What is the truth value of  $\forall x P(x)$  if the domain is  $Z^+$  ?

For all  $x \in Z^+$   $x^2 \geq x$  . So  $\forall x P(x)$  is true for  $Z^+$  .

# Quantifiers

- $P(x) : x^2 \geq x$

What is the truth value of  $\forall x P(x)$  if the domain is  $Z^+$  ?

For all  $x \in Z^+$   $x^2 \geq x$  . So  $\forall x P(x)$  is true for  $Z^+$  .

- $Q(x) : x = x + 1$

What is the truth value of  $\exists x Q(x)$  if the domain is  $R$ ?

# Quantifiers

- $P(x) : x^2 \geq x$

What is the truth value of  $\forall x P(x)$  if the domain is  $Z^+$  ?

For all  $x \in Z^+$   $x^2 \geq x$  . So  $\forall x P(x)$  is true for  $Z^+$  .

- $Q(x) : x = x + 1$

What is the truth value of  $\exists x Q(x)$  if the domain is  $R$ ?

There is no real number  $x$  such that  $x = x + 1$ .

# Quantifiers

- $P(x) : x^2 \geq x$

What is the truth value of  $\forall x P(x)$  if the domain is  $Z^+$  ?

For all  $x \in Z^+$   $x^2 \geq x$  . So  $\forall x P(x)$  is true for  $Z^+$  .

- $Q(x) : x = x + 1$

What is the truth value of  $\exists x Q(x)$  if the domain is  $R$ ?

There is no real number  $x$  such that  $x = x + 1$ . So  $\exists x Q(x)$  is false for  $R$ .

# Quantifiers

- $P(x) : x^2 + 1 < 10$  ,  $D = \{ 1, 2, 3 \}$

What is the truth value of  $\forall x P(x)$  if the domain is  $D$ ?

# Quantifiers

- $P(x) : x^2 + 1 < 10$  ,  $D = \{ 1, 2, 3 \}$

What is the truth value of  $\forall x P(x)$  if the domain is  $D$ ?

If the domain consists of  $n$  elements,  
then  $\forall x P(x) \equiv P(x_1) \wedge P(x_2) \wedge \dots \wedge P(x_n)$

# Quantifiers

- $P(x) : x^2 + 1 < 10$  ,  $D = \{ 1, 2, 3 \}$

What is the truth value of  $\forall x P(x)$  if the domain is  $D$ ?

If the domain consists of  $n$  elements,  
then  $\forall x P(x) \equiv P(x_1) \wedge P(x_2) \wedge \dots \wedge P(x_n)$

$P(1) : 2 < 10$ , true



# Quantifiers

- $P(x) : x^2 + 1 < 10$  ,  $D = \{ 1, 2, 3 \}$

What is the truth value of  $\forall x P(x)$  if the domain is  $D$ ?

If the domain consists of  $n$  elements,  
then  $\forall x P(x) \equiv P(x_1) \wedge P(x_2) \wedge \dots \wedge P(x_n)$

$P(1) : 2 < 10$ , true

$P(2) : 5 < 10$ , true

# Quantifiers

- $P(x) : x^2 + 1 < 10$  ,  $D = \{ 1, 2, 3 \}$

What is the truth value of  $\forall x P(x)$  if the domain is  $D$ ?

If the domain consists of  $n$  elements,  
then  $\forall x P(x) \equiv P(x_1) \wedge P(x_2) \wedge \dots \wedge P(x_n)$

$P(1) : 2 < 10$ , true

$P(2) : 5 < 10$ , true

$P(3) : 10 < 10$ , false

# Quantifiers

- $P(x) : x^2 + 1 < 10$  ,  $D = \{ 1, 2, 3 \}$

What is the truth value of  $\forall x P(x)$  if the domain is  $D$ ?

If the domain consists of  $n$  elements,  
then  $\forall x P(x) \equiv P(x_1) \wedge P(x_2) \wedge \dots \wedge P(x_n)$

$P(1) : 2 < 10$ , true

$P(2) : 5 < 10$ , true

$P(3) : 10 < 10$ , false

Since  $1 \wedge 1 \wedge 0 \equiv 0$ , then  $\forall x P(x)$  is false for  $D$ .

# Quantifiers

- $Q(x) : x^2 < 3$  ,  $D = \{1, 2, 3\}$

What is the truth value of  $\exists x Q(x)$  if the domain is  $D$ ?

# Quantifiers

- $Q(x) : x^2 < 3$  ,  $D = \{1, 2, 3\}$

What is the truth value of  $\exists x Q(x)$  if the domain is  $D$ ?

If the domain consists of  $n$  elements,  
then  $\exists x P(x) \equiv P(x_1) \vee P(x_2) \vee \dots \vee P(x_n)$

# Quantifiers

- $Q(x) : x^2 < 3$  ,  $D = \{1, 2, 3\}$

What is the truth value of  $\exists x Q(x)$  if the domain is  $D$ ?

If the domain consists of  $n$  elements,  
then  $\exists x P(x) \equiv P(x_1) \vee P(x_2) \vee \dots \vee P(x_n)$

$P(1) : 1 < 3$ , true

# Quantifiers

- $Q(x) : x^2 < 3$  ,  $D = \{1, 2, 3\}$

What is the truth value of  $\exists x Q(x)$  if the domain is  $D$ ?

If the domain consists of  $n$  elements,  
then  $\exists x P(x) \equiv P(x_1) \vee P(x_2) \vee \dots \vee P(x_n)$

$P(1) : 1 < 3$ , true

$P(2) : 4 < 3$ , false

# Quantifiers

- $Q(x) : x^2 < 3$  ,  $D = \{1, 2, 3\}$

What is the truth value of  $\exists x Q(x)$  if the domain is  $D$ ?

If the domain consists of  $n$  elements,  
then  $\exists x P(x) \equiv P(x_1) \vee P(x_2) \vee \dots \vee P(x_n)$

$P(1) : 1 < 3$ , true

$P(2) : 4 < 3$ , false

$P(3) : 9 < 3$ , false



# Quantifiers

- $Q(x) : x^2 < 3$  ,  $D = \{1, 2, 3\}$

What is the truth value of  $\exists x Q(x)$  if the domain is  $D$ ?

If the domain consists of  $n$  elements,  
then  $\exists x P(x) \equiv P(x_1) \vee P(x_2) \vee \dots \vee P(x_n)$

$P(1) : 1 < 3$ , true

$P(2) : 4 < 3$ , false

$P(3) : 9 < 3$ , false

Since  $1 \vee 0 \vee 0 \equiv 1$ , then  $\exists x P(x)$  is true for  $D$ .

# Quantifiers

Negation

# Quantifiers

- Every student in this class has entered the entrance exam

## Negation

# Quantifiers

- Every student in this class has entered the entrance exam

$\forall x P(x)$ , 'x has taken the entrance exam'

Negation

# Quantifiers

- Every student in this class has entered the entrance exam

$\forall x P(x)$ , 'x has taken the entrance exam'

## Negation

- It's not the case that every student in this class has entered the entrance exam.

# Quantifiers

- Every student in this class has entered the entrance exam

$\forall x P(x)$ , 'x has taken the entrance exam'

## Negation

- It's not the case that every student in this class has entered the entrance exam.

There is a student in this class who has not taken the entrance exam.

# Quantifiers

- Every student in this class has entered the entrance exam

$\forall x P(x)$ , 'x has taken the entrance exam'

## Negation

- It's not the case that every student in this class has entered the entrance exam.

There is a student in this class who has not taken the entrance exam.

$$\sim(\forall x P(x)) \equiv$$

# Quantifiers

- Every student in this class has entered the entrance exam

$\forall x P(x)$ , 'x has taken the entrance exam'

## Negation

- It's not the case that every student in this class has entered the entrance exam.

There is a student in this class who has not taken the entrance exam.

$$\sim(\forall x P(x)) \equiv \sim(P(x_1) \wedge P(x_2) \wedge \dots \wedge P(x_n))$$



# Quantifiers

- Every student in this class has entered the entrance exam

$\forall x P(x)$ , 'x has taken the entrance exam'

## Negation

- It's not the case that every student in this class has entered the entrance exam.

There is a student in this class who has not taken the entrance exam.

$$\begin{aligned}\sim(\forall x P(x)) &\equiv \sim(P(x_1) \wedge P(x_2) \wedge \dots \wedge P(x_n)) \\ &\equiv \sim P(x_1) \vee \sim P(x_2) \vee \dots \vee \sim P(x_n)\end{aligned}$$

# Quantifiers

- Every student in this class has entered the entrance exam

$\forall x P(x)$ , 'x has taken the entrance exam'

## Negation

- It's not the case that every student in this class has entered the entrance exam.

There is a student in this class who has not taken the entrance exam.

$$\begin{aligned}\sim(\forall x P(x)) &\equiv \sim(P(x_1) \wedge P(x_2) \wedge \dots \wedge P(x_n)) \\ &\equiv \sim P(x_1) \vee \sim P(x_2) \vee \dots \vee \sim P(x_n) \\ &\equiv \exists x \sim P(x)\end{aligned}$$

# Quantifiers

Negation

# Quantifiers

- There is a student in this class who has taken the entrance exam.

## Negation

# Quantifiers

- There is a student in this class who has taken the entrance exam.

$\exists x P(x)$ , 'x has taken the entrance exam'

## Negation

# Quantifiers

- There is a student in this class who has taken the entrance exam.

$\exists x P(x),$  'x has taken the entrance exam'

## Negation

- It's not the case that There is a student in this class who has taken the entrance exam

# Quantifiers

- There is a student in this class who has taken the entrance exam.

$\exists x P(x)$ , 'x has taken the entrance exam'

## Negation

- It's not the case that There is a student in this class who has taken the entrance exam

None of the students in this class has taken the entrance exam.

# Quantifiers

- There is a student in this class who has taken the entrance exam.

$\exists x P(x)$ , 'x has taken the entrance exam'

## Negation

- It's not the case that There is a student in this class who has taken the entrance exam

None of the students in this class has taken the entrance exam.

$$\sim(\exists x P(x)) \equiv$$



# Quantifiers

- There is a student in this class who has taken the entrance exam.

$\exists x P(x),$  'x has taken the entrance exam'

## Negation

- It's not the case that There is a student in this class who has taken the entrance exam

None of the students in this class has taken the entrance exam.

$$\sim(\exists x P(x)) \equiv \sim(P(x_1) \vee P(x_2) \vee \dots \vee P(x_n))$$

# Quantifiers

- There is a student in this class who has taken the entrance exam.

$\exists x P(x)$ , 'x has taken the entrance exam'

## Negation

- It's not the case that There is a student in this class who has taken the entrance exam

None of the students in this class has taken the entrance exam.

$$\begin{aligned}\sim(\exists x P(x)) &\equiv \sim(P(x_1) \vee P(x_2) \vee \dots \vee P(x_n)) \\ &\equiv \sim P(x_1) \wedge \sim P(x_2) \wedge \dots \wedge \sim P(x_n)\end{aligned}$$

# Quantifiers

- There is a student in this class who has taken the entrance exam.

$\exists x P(x)$ , 'x has taken the entrance exam'

## Negation

- It's not the case that There is a student in this class who has taken the entrance exam

None of the students in this class has taken the entrance exam.

$$\begin{aligned}\sim(\exists x P(x)) &\equiv \sim(P(x_1) \vee P(x_2) \vee \dots \vee P(x_n)) \\ &\equiv \sim P(x_1) \wedge \sim P(x_2) \wedge \dots \wedge \sim P(x_n) \\ &\equiv \forall x \sim P(x)\end{aligned}$$

# Quantifiers

- $\sim(\forall x(x^2 > x)) \equiv$

- $\sim(\exists x(x^2 = 7)) \equiv$

# Quantifiers

- $\sim(\forall x(x^2 > x)) \equiv \exists x \sim(x^2 > x)$

- $\sim(\exists x(x^2 = 7)) \equiv$

# Quantifiers

- $\sim(\forall x(x^2 > x)) \equiv \exists x \sim(x^2 > x)$   
 $\equiv \exists x x^2 \leq x$

- $\sim(\exists x(x^2 = 7)) \equiv$

# Quantifiers

- $\sim(\forall x(x^2 > x)) \equiv \exists x \sim(x^2 > x)$   
 $\equiv \exists x x^2 \leq x$
- $\sim(\exists x(x^2 = 7)) \equiv \forall x \sim(x^2 = 7)$

# Quantifiers

- $\sim(\forall x(x^2 > x)) \equiv \exists x \sim(x^2 > x)$   
 $\equiv \exists x x^2 \leq x$

- $\sim(\exists x(x^2 = 7)) \equiv \forall x \sim(x^2 = 7)$   
 $\equiv \forall x x^2 \neq 7$



# Quantifiers

- $\forall x \forall y ((x > 0) \wedge (y < 0) \rightarrow (xy < 0))$        $D = \mathbb{R}$

# Quantifiers

If  $x$  is positive and  $y$  is negative,  
then  $xy$  is negative

- $\forall x \forall y ((x > 0) \wedge (y < 0) \rightarrow (xy < 0))$   $D = \mathbb{R}$

# Quantifiers

If  $x$  is positive and  $y$  is negative,  
then  $xy$  is negative

- $\forall x \forall y ((x > 0) \wedge (y < 0) \rightarrow (xy < 0))$   $D = \mathbb{R}$

For every real numbers  $x$  and  $y$ , if  $x$  is positive and  $y$  is negative, then  $xy$  is negative

# Quantifiers

- For every two integers, if these integers are both positive, then the sum of these integers is also positive

# Quantifiers

- For every two integers, if these integers are both positive, then the sum of these integers is also positive
- For two integers  $x$  and  $y$ , if  $x > 0$  and  $y > 0$ , then  $x + y > 0$

# Quantifiers

- For every two integers, if these integers are both positive, then the sum of these integers is also positive
- For two integers  $x$  and  $y$ , if  $x > 0$  and  $y > 0$ , then  $x + y > 0$


$$(x > 0) \wedge (y > 0) \rightarrow (x + y > 0)$$

# Quantifiers

- For every two integers, if these integers are both positive, then the sum of these integers is also positive
- For two integers  $x$  and  $y$ , if  $x > 0$  and  $y > 0$ , then  $x + y > 0$


$$(x > 0) \wedge (y > 0) \rightarrow (x + y > 0)$$


$$\forall x \forall y ((x > 0) \wedge (y > 0) \rightarrow (x + y > 0))$$

# Quantifiers

- There exist integers  $x$  and  $y$  such that  $x + y = 6$



# Quantifiers

- There exist integers  $x$  and  $y$  such that  $x + y = 6$

$$\exists x \exists y (x + y = 6)$$

# Quantifiers

- There exist integers  $x$  and  $y$  such that  $x + y = 6$

$$\exists x \exists y (x + y = 6)$$

or

$$\exists y \exists x (x + y = 6)$$

# Quantifiers

- There exist integers  $x$  and  $y$  such that  $x + y = 6$

$$\exists x \exists y (x + y = 6)$$

or

$$\exists y \exists x (x + y = 6)$$

- $\forall x \exists y (x + y = 6)$

# Quantifiers

- There exist integers  $x$  and  $y$  such that  $x + y = 6$

$$\exists x \exists y (x + y = 6)$$

or

$$\exists y \exists x (x + y = 6)$$

- $\forall x \exists y (x + y = 6)$

For every integer  $x$ , there exists an integer  $y$  such that  
 $x + y = 6$

# Quantifiers

- There exist integers  $x$  and  $y$  such that  $x + y = 6$

$$\exists x \exists y (x + y = 6)$$

or

$$\exists y \exists x (x + y = 6)$$

- $\forall x \exists y (x + y = 6)$

For every integer  $x$ , there exists an integer  $y$  such that  
 $x + y = 6$  (It's true)

# Quantifiers

- There exist integers  $x$  and  $y$  such that  $x + y = 6$

$$\exists x \exists y (x + y = 6)$$

or

$$\exists y \exists x (x + y = 6)$$

- $\forall x \exists y (x + y = 6)$

For every integer  $x$ , there exists an integer  $y$  such that  
 $x + y = 6$  (It's true)

- $\exists y \forall x (x + y = 6)$

# Quantifiers

- There exist integers  $x$  and  $y$  such that  $x + y = 6$

$$\exists x \exists y (x + y = 6)$$

or

$$\exists y \exists x (x + y = 6)$$

- $\forall x \exists y (x + y = 6)$

For every integer  $x$ , there exists an integer  $y$  such that  
 $x + y = 6$  (It's true)

- $\exists y \forall x (x + y = 6)$

There exists an integer  $y$  so that for all integers  $x$ ,  
 $x + y = 6$

# Quantifiers

- There exist integers  $x$  and  $y$  such that  $x + y = 6$

$$\exists x \exists y (x + y = 6)$$

or

$$\exists y \exists x (x + y = 6)$$

- $\forall x \exists y (x + y = 6)$

For every integer  $x$ , there exists an integer  $y$  such that  
 $x + y = 6$  (It's true)

- $\exists y \forall x (x + y = 6)$

There exists an integer  $y$  so that for all integers  $x$ ,  
 $x + y = 6$  (It's false)