

Mathematical Induction

Murat Osmanoglu

Definition

- To prove $P(n)$ is true for all positive integers n ,

Definition

- To prove $P(n)$ is true for all positive integers n ,
 - verify that $P(1)$ is true (Basic Step)

Definition

- To prove $P(n)$ is true for all positive integers n ,
 - verify that $P(1)$ is true (Basic Step)
 - prove that the implication $P(k) \rightarrow P(k + 1)$ for all $k \in \mathbb{Z}^+$ (Inductive Step)

Definition


- To prove $P(n)$ is true for all positive integers n ,
 - verify that $P(1)$ is true (Basic Step)
 - prove that the implication $P(k) \rightarrow P(k + 1)$ for all $k \in \mathbb{Z}^+$ (Inductive Step)

$$[P(1) \wedge \forall k P(k) \rightarrow P(k + 1)] \rightarrow \forall n P(n)$$

Definition

- To prove $P(n)$ is true for all positive integers n ,
 - verify that $P(1)$ is true (Basic Step)
 - prove that the implication $P(k) \rightarrow P(k + 1)$ for all $k \in \mathbb{Z}^+$ (Inductive Step)

$$[P(1) \wedge \forall k P(k) \rightarrow P(k + 1)] \rightarrow \forall n P(n)$$



Proofs

- Prove that $\forall x \in \mathbb{Z}^+, x^3 - x$ is divisible by 3

Proofs

- Prove that $\forall x \in \mathbb{Z}^+, x^3 - x$ is divisible by 3

Basic Step $P(1) : 1^3 - 1 = 0$ is divisible by 3

Proofs

- Prove that $\forall x \in \mathbb{Z}^+, x^3 - x$ is divisible by 3

Basic Step $P(1) : 1^3 - 1 = 0$ is divisible by 3

Inductive Step $P(k) \rightarrow P(k + 1)$

Proofs

- Prove that $\forall x \in \mathbb{Z}^+, x^3 - x$ is divisible by 3

Basic Step $P(1) : 1^3 - 1 = 0$ is divisible by 3

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $k^3 - k$ is divisible by 3

Proofs

- Prove that $\forall x \in \mathbb{Z}^+, x^3 - x$ is divisible by 3

Basic Step $P(1) : 1^3 - 1 = 0$ is divisible by 3

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $k^3 - k$ is divisible by 3

$$[k^3 - k = 3a, \exists a \in \mathbb{Z}]$$

Proofs

- Prove that $\forall x \in \mathbb{Z}^+, x^3 - x$ is divisible by 3

Basic Step $P(1) : 1^3 - 1 = 0$ is divisible by 3

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $k^3 - k$ is divisible by 3

$$[k^3 - k = 3a, \exists a \in \mathbb{Z}] \rightarrow (k + 1)^3 - (k + 1)$$

Proofs

- Prove that $\forall x \in \mathbb{Z}^+, x^3 - x$ is divisible by 3

Basic Step $P(1) : 1^3 - 1 = 0$ is divisible by 3

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $k^3 - k$ is divisible by 3

$$[k^3 - k = 3a, \exists a \in \mathbb{Z}] \rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k + 1 - k - 1$$

Proofs

- Prove that $\forall x \in \mathbb{Z}^+, x^3 - x$ is divisible by 3

Basic Step $P(1) : 1^3 - 1 = 0$ is divisible by 3

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $k^3 - k$ is divisible by 3

$$\begin{aligned} [k^3 - k = 3a, \exists a \in \mathbb{Z}] &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k + 1 - k - 1 \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k - k \end{aligned}$$

Proofs

- Prove that $\forall x \in \mathbb{Z}^+, x^3 - x$ is divisible by 3

Basic Step $P(1) : 1^3 - 1 = 0$ is divisible by 3

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $k^3 - k$ is divisible by 3

$$\begin{aligned} [k^3 - k = 3a, \exists a \in \mathbb{Z}] &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k + 1 - k - 1 \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k - k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3k^2 + 3k \end{aligned}$$

Proofs

- Prove that $\forall x \in \mathbb{Z}^+, x^3 - x$ is divisible by 3

Basic Step $P(1) : 1^3 - 1 = 0$ is divisible by 3

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $k^3 - k$ is divisible by 3

$$\begin{aligned}[k^3 - k = 3a, \exists a \in \mathbb{Z}] &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k + 1 - k - 1 \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k - k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3k^2 + 3k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3(k^2 + k)\end{aligned}$$

Proofs

- Prove that $\forall x \in \mathbb{Z}^+, x^3 - x$ is divisible by 3

Basic Step $P(1) : 1^3 - 1 = 0$ is divisible by 3

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $k^3 - k$ is divisible by 3

$$\begin{aligned} [k^3 - k = 3a, \exists a \in \mathbb{Z}] &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k + 1 - k - 1 \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k - k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3k^2 + 3k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3(k^2 + k) \\ &\rightarrow (k + 1)^3 - (k + 1) = 3a + 3b, \exists a, b \in \mathbb{Z} \end{aligned}$$

Proofs

- Prove that $\forall x \in \mathbb{Z}^+, x^3 - x$ is divisible by 3

Basic Step $P(1) : 1^3 - 1 = 0$ is divisible by 3

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $k^3 - k$ is divisible by 3

$$\begin{aligned} [k^3 - k = 3a, \exists a \in \mathbb{Z}] &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k + 1 - k - 1 \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 + 3k^2 + 3k - k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3k^2 + 3k \\ &\rightarrow (k + 1)^3 - (k + 1) = k^3 - k + 3(k^2 + k) \\ &\rightarrow (k + 1)^3 - (k + 1) = 3a + 3b, \exists a, b \in \mathbb{Z} \\ &\rightarrow (k + 1)^3 - (k + 1) \text{ is divisible by 3} \end{aligned}$$

Proofs

- Prove that $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$ is divisible by 57

Proofs

- Prove that $\forall n \in N, 7^{n+2} + 8^{2n+1}$ is divisible by 57

Basic Step $P(0) : 7^2 + 8 = 57$ is divisible by 57

Proofs

- Prove that $\forall n \in N, 7^{n+2} + 8^{2n+1}$ is divisible by 57

Basic Step $P(0) : 7^2 + 8 = 57$ is divisible by 57

Inductive Step $P(k) \rightarrow P(k + 1)$

Proofs

- Prove that $\forall n \in N, 7^{n+2} + 8^{2n+1}$ is divisible by 57

Basic Step $P(0) : 7^2 + 8 = 57$ is divisible by 57

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $7^{k+2} + 8^{2k+1}$ is divisible by 57

Proofs

- Prove that $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$ is divisible by 57

Basic Step $P(0) : 7^2 + 8 = 57$ is divisible by 57

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $7^{k+2} + 8^{2k+1}$ is divisible by 57

$$[7^{k+2} + 8^{2k+1} = 57a, \exists a \in \mathbb{Z}]$$

Proofs

- Prove that $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$ is divisible by 57

Basic Step $P(0) : 7^2 + 8 = 57$ is divisible by 57

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $7^{k+2} + 8^{2k+1}$ is divisible by 57

$$[7^{k+2} + 8^{2k+1} = 57a, \exists a \in \mathbb{Z}] \rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 64 \cdot 8^{2k+1}$$

Proofs

- Prove that $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$ is divisible by 57

Basic Step $P(0) : 7^2 + 8 = 57$ is divisible by 57

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $7^{k+2} + 8^{2k+1}$ is divisible by 57

$$\begin{aligned} [7^{k+2} + 8^{2k+1} = 57a, \exists a \in \mathbb{Z}] &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 64 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 7 \cdot 8^{2k+1} + 57 \cdot 8^{2k+1} \end{aligned}$$

Proofs

- Prove that $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$ is divisible by 57

Basic Step $P(0) : 7^2 + 8 = 57$ is divisible by 57

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $7^{k+2} + 8^{2k+1}$ is divisible by 57

$$\begin{aligned} [7^{k+2} + 8^{2k+1} = 57a, \exists a \in \mathbb{Z}] &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 64 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 7 \cdot 8^{2k+1} + 57 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7(7^{k+2} + 8^{2k+1}) + 57 \cdot 8^{2k+1} \end{aligned}$$

Proofs

- Prove that $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$ is divisible by 57

Basic Step $P(0) : 7^2 + 8 = 57$ is divisible by 57

Inductive Step $P(k) \rightarrow P(k+1)$

assume that $P(k)$ is true, i.e $7^{k+2} + 8^{2k+1}$ is divisible by 57

$$\begin{aligned} [7^{k+2} + 8^{2k+1} = 57a, \exists a \in \mathbb{Z}] &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 64 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 7 \cdot 8^{2k+1} + 57 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7(7^{k+2} + 8^{2k+1}) + 57 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 57a + 57b, \exists a, b \in \mathbb{Z} \end{aligned}$$

Proofs

- Prove that $\forall n \in \mathbb{N}, 7^{n+2} + 8^{2n+1}$ is divisible by 57

Basic Step $P(0) : 7^2 + 8 = 57$ is divisible by 57

Inductive Step $P(k) \rightarrow P(k+1)$

assume that $P(k)$ is true, i.e $7^{k+2} + 8^{2k+1}$ is divisible by 57

$$\begin{aligned} [7^{k+2} + 8^{2k+1} = 57a, \exists a \in \mathbb{Z}] &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 64 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7 \cdot 7^{k+2} + 7 \cdot 8^{2k+1} + 57 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 7(7^{k+2} + 8^{2k+1}) + 57 \cdot 8^{2k+1} \\ &\rightarrow 7^{k+3} + 8^{2k+3} = 57a + 57b, \exists a, b \in \mathbb{Z} \\ &\rightarrow 7^{k+3} + 8^{2k+3} \text{ is divisible by } 57 \end{aligned}$$

Proofs

- Prove that if $\forall n \in \mathbb{Z}^+$, then $1 + 2 + \dots + n = n \cdot (n + 1) / 2$

Proofs

- Prove that if $\forall n \in \mathbb{Z}^+$, then $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step $P(1) : 1 = 1 \cdot 2/2$

Proofs

- Prove that if $\forall n \in \mathbb{Z}^+$, then $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step $P(1) : 1 = 1 \cdot 2/2$

Inductive Step $P(k) \rightarrow P(k + 1)$

Proofs

- Prove that if $\forall n \in \mathbb{Z}^+$, then $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step $P(1) : 1 = 1 \cdot 2/2$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $1 + 2 + \dots + k = k \cdot (k + 1)/2$

Proofs

- Prove that if $\forall n \in \mathbb{Z}^+$, then $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step $P(1) : 1 = 1 \cdot 2/2$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $1 + 2 + \dots + k = k \cdot (k + 1)/2$

$[1 + 2 + \dots + k = k \cdot (k + 1)/2]$

Proofs

- Prove that if $\forall n \in \mathbb{Z}^+$, then $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step $P(1) : 1 = 1 \cdot 2/2$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $1 + 2 + \dots + k = k \cdot (k + 1)/2$

$$[1 + 2 + \dots + k = k \cdot (k + 1)/2] \rightarrow \left[1 + 2 + \dots + (k + 1) = k \cdot \frac{k+1}{2} + k + 1 \right]$$

Proofs

- Prove that if $\forall n \in \mathbb{Z}^+$, then $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step $P(1) : 1 = 1 \cdot 2/2$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $1 + 2 + \dots + k = k \cdot (k + 1)/2$

$$\begin{aligned} [1 + 2 + \dots + k = k \cdot (k + 1)/2] &\rightarrow [1 + 2 + \dots + (k + 1) = k \cdot \frac{k+1}{2} + k + 1] \\ &\rightarrow [1 + 2 + \dots + (k + 1) = \frac{k(k+1) + 2(k+1)}{2}] \end{aligned}$$

Proofs

- Prove that if $\forall n \in \mathbb{Z}^+$, then $1 + 2 + \dots + n = n \cdot (n + 1)/2$

Basic Step $P(1) : 1 = 1 \cdot 2/2$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $1 + 2 + \dots + k = k \cdot (k + 1)/2$

$$[1 + 2 + \dots + k = k \cdot (k + 1)/2] \rightarrow [1 + 2 + \dots + (k + 1) = k \cdot \frac{k+1}{2} + k + 1]$$

$$\rightarrow [1 + 2 + \dots + (k + 1) = \frac{k(k+1) + 2(k+1)}{2}]$$

$$\rightarrow [1 + 2 + \dots + (k + 1) = \frac{(k+1)(k+2)}{2}]$$

Proofs

Conjecture a formula for the sum of the first n positive odd integers, then prove your conjecture using mathematical induction

Proofs

Conjecture a formula for the sum of the first n positive odd integers, then prove your conjecture using mathematical induction

- $1 = 1$ $1 + 3 = 4$ $1 + 3 + 5 = 9$ $1 + 3 + 5 + 7 = 16$

Proofs

Conjecture a formula for the sum of the first n positive odd integers, then prove your conjecture using mathematical induction

- $$\begin{array}{cccc} 1 = 1 & 1 + 3 = 4 & 1 + 3 + 5 = 9 & 1 + 3 + 5 + 7 = 16 \\ 1^2 & 2^2 & 3^2 & 4^2 \end{array}$$

Proofs

Conjecture a formula for the sum of the first n positive odd integers, then prove your conjecture using mathematical induction

- $1 = 1$ $1 + 3 = 4$ $1 + 3 + 5 = 9$ $1 + 3 + 5 + 7 = 16$
 1^2 2^2 3^2 4^2
 $1 + 3 + \dots + (2n - 1) = n^2$

Proofs

Conjecture a formula for the sum of the first n positive odd integers, then prove your conjecture using mathematical induction

$$\begin{array}{cccc} \bullet & 1 = 1 & 1 + 3 = 4 & 1 + 3 + 5 = 9 & 1 + 3 + 5 + 7 = 16 \\ & 1^2 & 2^2 & 3^2 & 4^2 \end{array}$$

$$1 + 3 + \dots + (2n - 1) = n^2$$

Basic Step $P(1) : 1 = 1^2$

Proofs

Conjecture a formula for the sum of the first n positive odd integers, then prove your conjecture using mathematical induction

- $1 = 1$ $1 + 3 = 4$ $1 + 3 + 5 = 9$ $1 + 3 + 5 + 7 = 16$
 1^2 2^2 3^2 4^2
 $1 + 3 + \dots + (2n - 1) = n^2$

Basic Step $P(1) : 1 = 1^2$

Inductive Step $P(k) \rightarrow P(k + 1)$

Proofs

Conjecture a formula for the sum of the first n positive odd integers, then prove your conjecture using mathematical induction

- $$\begin{array}{ccccccc} 1 = 1 & 1 + 3 = 4 & 1 + 3 + 5 = 9 & 1 + 3 + 5 + 7 = 16 \\ 1^2 & 2^2 & 3^2 & 4^2 \\ 1 + 3 + \dots + (2n - 1) = n^2 \end{array}$$

Basic Step $P(1) : 1 = 1^2$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $1 + 3 + \dots + (2k - 1) = k^2$

Proofs

Conjecture a formula for the sum of the first n positive odd integers, then prove your conjecture using mathematical induction

- $$\begin{array}{ccccccc} 1 = 1 & 1 + 3 = 4 & 1 + 3 + 5 = 9 & 1 + 3 + 5 + 7 = 16 \\ 1^2 & 2^2 & 3^2 & 4^2 \\ 1 + 3 + \dots + (2n - 1) = n^2 \end{array}$$

Basic Step $P(1) : 1 = 1^2$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $1 + 2 + \dots + (2k - 1) = k^2$

$$[1 + 2 + \dots + (2k - 1) = k^2]$$

Proofs

Conjecture a formula for the sum of the first n positive odd integers, then prove your conjecture using mathematical induction

- $$\begin{array}{ccccccc} 1 = 1 & 1 + 3 = 4 & 1 + 3 + 5 = 9 & 1 + 3 + 5 + 7 = 16 \\ 1^2 & 2^2 & 3^2 & 4^2 \\ 1 + 3 + \dots + (2n - 1) = n^2 \end{array}$$

Basic Step $P(1) : 1 = 1^2$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $1 + 2 + \dots + (2k - 1) = k^2$

$$[1 + 2 + \dots + (2k - 1) = k^2] \rightarrow [1 + 2 + \dots + (2k - 1) + (2k + 1) = k^2 + 2k + 1]$$

Proofs

Conjecture a formula for the sum of the first n positive odd integers, then prove your conjecture using mathematical induction

- $$\begin{array}{ccccccc} 1 = 1 & 1 + 3 = 4 & 1 + 3 + 5 = 9 & 1 + 3 + 5 + 7 = 16 \\ 1^2 & 2^2 & 3^2 & 4^2 \\ 1 + 3 + \dots + (2n - 1) = n^2 \end{array}$$

Basic Step $P(1) : 1 = 1^2$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $1 + 2 + \dots + (2k - 1) = k^2$

$$\begin{aligned} [1 + 2 + \dots + (2k - 1) = k^2] &\rightarrow [1 + 2 + \dots + (2k - 1) + (2k + 1) = k^2 + 2k + 1] \\ &\rightarrow [1 + 2 + \dots + (2k - 1) + (2k + 1) = (k + 1)^2] \end{aligned}$$

Proofs

- Prove that if $\forall n \in \mathbb{N}$, then $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Proofs

- Prove that if $\forall n \in \mathbb{N}$, then $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step $P(1) : 1 = 2^{0+1} - 1$

Proofs

- Prove that if $\forall n \in \mathbb{N}$, then $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step $P(1) : 1 = 2^{0+1} - 1$

Inductive Step $P(k) \rightarrow P(k + 1)$

Proofs

- Prove that if $\forall n \in \mathbb{N}$, then $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step $P(1) : 1 = 2^{0+1} - 1$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1$

Proofs

- Prove that if $\forall n \in \mathbb{N}$, then $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step $P(1) : 1 = 2^{0+1} - 1$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1$

$$[1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1]$$

Proofs

- Prove that if $\forall n \in \mathbb{N}$, then $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step $P(1) : 1 = 2^{0+1} - 1$

Inductive Step $P(k) \rightarrow P(k+1)$

assume that $P(k)$ is true, i.e $1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1$

$$[1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1] \rightarrow [1 + 2 + 2^2 + \dots + 2^k + 2^{k+1} = 2^{k+1} - 1 + 2^{k+1}]$$

Proofs

- Prove that if $\forall n \in \mathbb{N}$, then $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step $P(1) : 1 = 2^{0+1} - 1$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1$

$$[1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1] \rightarrow [1 + 2 + 2^2 + \dots + 2^k + 2^{k+1} = 2^{k+1} - 1 + 2^{k+1}]$$

$$\rightarrow [1 + 2 + 2^2 + \dots + 2^{k+1} = 2 \cdot 2^{k+1} - 1]$$

Proofs

- Prove that if $\forall n \in \mathbb{N}$, then $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basic Step $P(1) : 1 = 2^{0+1} - 1$

Inductive Step $P(k) \rightarrow P(k+1)$

assume that $P(k)$ is true, i.e $1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1$

$$[1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1] \rightarrow [1 + 2 + 2^2 + \dots + 2^k + 2^{k+1} = 2^{k+1} - 1 + 2^{k+1}]$$

$$\rightarrow [1 + 2 + 2^2 + \dots + 2^{k+1} = 2 \cdot 2^{k+1} - 1]$$

$$\rightarrow [1 + 2 + 2^2 + \dots + 2^{k+1} = 2^{k+1+1} - 1]$$

Proofs

- Prove that for every integer $n \geq 4$, $2^n < n!$

Proofs

- Prove that for every integer $n \geq 4$, $2^n < n!$

Basic Step $P(4)$: $2^4 = 16 < 4! = 24$

Proofs

- Prove that for every integer $n \geq 4$, $2^n < n!$

Basic Step $P(4) : 2^4 = 16 < 4! = 24$

Inductive Step $P(k) \rightarrow P(k + 1)$

Proofs

- Prove that for every integer $n \geq 4$, $2^n < n!$

Basic Step $P(4) : 2^4 = 16 < 4! = 24$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $2^k < k!$

Proofs

- Prove that for every integer $n \geq 4$, $2^n < n!$

Basic Step $P(4) : 2^4 = 16 < 4! = 24$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $2^k < k!$

$[2^k < k!]$

Proofs

- Prove that for every integer $n \geq 4$, $2^n < n!$

Basic Step $P(4)$: $2^4 = 16 < 4! = 24$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $2^k < k!$

$$[2^k < k!] \rightarrow [2^{k+1} = 2 \cdot 2^k < 2 \cdot k!]$$

Proofs

- Prove that for every integer $n \geq 4$, $2^n < n!$

Basic Step $P(4)$: $2^4 = 16 < 4! = 24$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $2^k < k!$

$$[2^k < k!] \rightarrow [2^{k+1} = 2 \cdot 2^k < 2 \cdot k!] \rightarrow [2^{k+1} < 2 \cdot k!]$$

Proofs

- Prove that for every integer $n \geq 4$, $2^n < n!$

Basic Step $P(4)$: $2^4 = 16 < 4! = 24$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $2^k < k!$

$$[2^k < k!] \rightarrow [2^{k+1} = 2 \cdot 2^k < 2 \cdot k!] \rightarrow [2^{k+1} < 2 \cdot k! < (k+1) \cdot k!]$$

Proofs

- Prove that for every integer $n \geq 4$, $2^n < n!$

Basic Step $P(4)$: $2^4 = 16 < 4! = 24$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e $2^k < k!$

$$\begin{aligned} [2^k < k!] &\rightarrow [2^{k+1} = 2 \cdot 2^k < 2 \cdot k!] \rightarrow [2^{k+1} < 2 \cdot k! < (k+1) \cdot k!] \\ &\rightarrow [2^{k+1} < (k+1)!] \end{aligned}$$

Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1]$

Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step $P(1) : [H_1 \stackrel{?}{=} 2.H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step $P(k) \rightarrow P(k + 1)$

Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $H_1 + \dots + H_k = (k + 1)H_k - k$

Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $H_1 + \dots + H_k = (k + 1)H_k - k$

$$[H_1 + \dots + H_k = (k + 1)H_k - k]$$

Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $H_1 + \dots + H_k = (k + 1)H_k - k$

$$[H_1 + \dots + H_k = (k + 1)H_k - k] \rightarrow [H_1 + \dots + H_k + H_{k+1} = (k + 1)H_k - k + H_{k+1}]$$

Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $H_1 + \dots + H_k = (k + 1)H_k - k$

$$\begin{aligned} [H_1 + \dots + H_k = (k + 1)H_k - k] &\rightarrow [H_1 + \dots + H_k + H_{k+1} = (k + 1)H_k - k + H_{k+1}] \\ &\rightarrow \left[H_1 + \dots + H_{k+1} = (k + 1)\left(H_k - \frac{1}{k+1} + \frac{1}{k+1}\right) - k + H_{k+1} \right] \end{aligned}$$

Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $H_1 + \dots + H_k = (k + 1)H_k - k$

$$\begin{aligned} [H_1 + \dots + H_k = (k + 1)H_k - k] &\rightarrow [H_1 + \dots + H_k + H_{k+1} = (k + 1)H_k - k + H_{k+1}] \\ &\rightarrow \left[H_1 + \dots + H_{k+1} = (k + 1)\left(H_k - \frac{1}{k+1} + \frac{1}{k+1}\right) - k + H_{k+1} \right] \\ &\rightarrow \left[H_1 + \dots + H_{k+1} = (k + 1)\left(H_{k+1} - \frac{1}{k+1}\right) - k + H_{k+1} \right] \end{aligned}$$

Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $H_1 + \dots + H_k = (k + 1)H_k - k$

$$\begin{aligned} [H_1 + \dots + H_k = (k + 1)H_k - k] &\rightarrow [H_1 + \dots + H_k + H_{k+1} = (k + 1)H_k - k + H_{k+1}] \\ &\rightarrow \left[H_1 + \dots + H_{k+1} = (k + 1)\left(H_k - \frac{1}{k+1} + \frac{1}{k+1}\right) - k + H_{k+1} \right] \\ &\rightarrow \left[H_1 + \dots + H_{k+1} = (k + 1)\left(H_{k+1} - \frac{1}{k+1}\right) - k + H_{k+1} \right] \\ &\rightarrow [H_1 + \dots + H_{k+1} = (k + 1)H_{k+1} - 1 - k + H_{k+1}] \end{aligned}$$

Proofs

$$H_j = 1 + \frac{1}{2} + \dots + \frac{1}{j}$$

- Prove that $H_1 + H_2 + \dots + H_n = (n + 1)H_n - n$

Basic Step $P(1) : [H_1 \stackrel{?}{=} 2 \cdot H_1 - 1] \rightarrow [1 = 2 - 1]$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $H_1 + \dots + H_k = (k + 1)H_k - k$

$$\begin{aligned} [H_1 + \dots + H_k = (k + 1)H_k - k] &\rightarrow [H_1 + \dots + H_k + H_{k+1} = (k + 1)H_k - k + H_{k+1}] \\ &\rightarrow \left[H_1 + \dots + H_{k+1} = (k + 1)\left(H_k - \frac{1}{k+1} + \frac{1}{k+1}\right) - k + H_{k+1} \right] \\ &\rightarrow \left[H_1 + \dots + H_{k+1} = (k + 1)\left(H_{k+1} - \frac{1}{k+1}\right) - k + H_{k+1} \right] \\ &\rightarrow [H_1 + \dots + H_{k+1} = (k + 1)H_{k+1} - 1 - k + H_{k+1}] \\ &\rightarrow [H_1 + \dots + H_{k+1} = (k + 2)H_{k+1} - (k + 1)] \end{aligned}$$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step $P(4) : 14 = 2.3 + 1.8$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step $P(4) : 14 = 2.3 + 1.8$

Inductive Step $P(k) \rightarrow P(k + 1)$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step $P(4) : 14 = 2.3 + 1.8$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $k = a.3 + b.8, \exists a, b \in \mathbb{N}$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step $P(4) : 14 = 2.3 + 1.8$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $k = a.3 + b.8, \exists a, b \in N$



if $b > 0$,

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step $P(4) : 14 = 2.3 + 1.8$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $k = a.3 + b.8, \exists a, b \in N$



if $b > 0$, $k + 1 = a.3 + b.8 + 1$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step $P(4) : 14 = 2.3 + 1.8$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $k = a.3 + b.8, \exists a, b \in \mathbb{N}$



if $b > 0$, $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step $P(4) : 14 = 2.3 + 1.8$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $k = a.3 + b.8, \exists a, b \in \mathbb{N}$



if $b > 0$, $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step $P(4) : 14 = 2.3 + 1.8$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $k = a.3 + b.8, \exists a, b \in N$



if $b > 0$, $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$

if $b = 0$, $k + 1 = a.3 + 1$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step $P(4) : 14 = 2.3 + 1.8$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $k = a.3 + b.8, \exists a, b \in N$



if $b > 0$, $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$

if $b = 0$, $k + 1 = a.3 + 1$

$$k + 1 = (a - 5).3 + 15 + 1$$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step $P(4) : 14 = 2.3 + 1.8$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $k = a.3 + b.8, \exists a, b \in N$



if $b > 0$, $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$

if $b = 0$, $k + 1 = a.3 + 1$

$$k + 1 = (a - 5).3 + 15 + 1$$

$$k + 1 = (a - 5).3 + 2.8$$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step $P(4) : 14 = 2.3 + 1.8$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $k = a.3 + b.8, \exists a, b \in N$



if $b > 0$, $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$

if $b = 0$, $k + 1 = a.3 + 1$

$$k + 1 = (a - 5).3 + 15 + 1$$

$$k + 1 = (a - 5).3 + 2.8$$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step $P(4) : 14 = 2.3 + 1.8$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $k = a.3 + b.8, \exists a, b \in N$



if $b > 0$, $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$

$P(k - 8)$

if $b = 0$, $k + 1 = a.3 + 1$

$$k + 1 = (a - 5).3 + 15 + 1$$

$$k + 1 = (a - 5).3 + 2.8$$

$P(k - 15)$

Proofs

- For every integer $n \geq 14$, n can be written as a sum of 3's and 8's

$$19 = 3 + 8 + 8 = 1.3 + 2.8$$

$$20 = 3 + 3 + 3 + 3 + 8 = 4.3 + 1.8$$

Basic Step $P(4) : 14 = 2.3 + 1.8$

Inductive Step $P(k) \rightarrow P(k + 1)$

assume that $P(k)$ is true, i.e. $k = a.3 + b.8, \exists a, b \in \mathbb{N}$

if $b > 0$, $k + 1 = a.3 + b.8 + 1$

$$k + 1 = a.3 + (b - 1).8 + 8 + 1$$

$$k + 1 = (a + 3).3 + (b - 1).8$$

$P(k - 8)$

if $b = 0$, $k + 1 = a.3 + 1$

$$k + 1 = (a - 5).3 + 15 + 1$$

$$k + 1 = (a - 5).3 + 2.8$$

$P(k - 15)$

$$[P(k - 8) \wedge P(k - 15)] \rightarrow P(k + 1)$$

Strong Induction

- To prove $P(n)$ is true for all positive integers n ,
 - verify that $P(1)$ is true (Basic Step)

Strong Induction

- To prove $P(n)$ is true for all positive integers n ,
 - verify that $P(1)$ is true (Basic Step)
 - prove that the implication

$$[P(1) \wedge P(2) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$$

for all $k \in \mathbb{Z}^+$ (Inductive Step)

Strong Induction

- Prove that for every integer $n \geq 2$, n can be written as the product of primes

Strong Induction

- Prove that for every integer $n \geq 2$, n can be written as the product of primes

Basic Step $P(2)$ is true, i.e. 2 can be written as the product of primes

Strong Induction

- Prove that for every integer $n \geq 2$, n can be written as the product of primes

Basic Step $P(2)$ is true, i.e. 2 can be written as the product of primes

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

Strong Induction

- Prove that for every integer $n \geq 2$, n can be written as the product of primes

Basic Step $P(2)$ is true, i.e. 2 can be written as the product of primes

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e i can be written as the product of primes, then

Strong Induction

- Prove that for every integer $n \geq 2$, n can be written as the product of primes

Basic Step $P(2)$ is true, i.e. 2 can be written as the product of primes

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e i can be written as the product of primes, then

if $(k + 1)$ is prime, then $P(k + 1)$ is true

Strong Induction

- Prove that for every integer $n \geq 2$, n can be written as the product of primes

Basic Step $P(2)$ is true, i.e. 2 can be written as the product of primes

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e i can be written as the product of primes, then

if $(k + 1)$ is prime, then $P(k + 1)$ is true

if $(k + 1)$ is composite, then $k + 1 = a.b$, where $2 \leq a \leq b < k + 1$.

Strong Induction

- Prove that for every integer $n \geq 2$, n can be written as the product of primes

Basic Step $P(2)$ is true, i.e. 2 can be written as the product of primes

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e i can be written as the product of primes, then

if $(k + 1)$ is prime, then $P(k + 1)$ is true

if $(k + 1)$ is composite, then $k + 1 = a.b$, where $2 \leq a \leq b < k + 1$.
Since $a, b < k + 1$, $P(a)$ and $P(b)$ are true from the assumption,

Strong Induction

- Prove that for every integer $n \geq 2$, n can be written as the product of primes

Basic Step $P(2)$ is true, i.e. 2 can be written as the product of primes

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e i can be written as the product of primes, then

if $(k + 1)$ is prime, then $P(k + 1)$ is true

if $(k + 1)$ is composite, then $k + 1 = a.b$, where $2 \leq a \leq b < k + 1$.

Since $a, b < k + 1$, $P(a)$ and $P(b)$ are true from the assumption, i.e. a and b can be written as the product of primes.

Strong Induction

- Prove that for every integer $n \geq 2$, n can be written as the product of primes

Basic Step $P(2)$ is true, i.e. 2 can be written as the product of primes

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e i can be written as the product of primes, then

if $(k + 1)$ is prime, then $P(k + 1)$ is true

if $(k + 1)$ is composite, then $k + 1 = a.b$, where $2 \leq a \leq b < k + 1$.

Since $a, b < k + 1$, $P(a)$ and $P(b)$ are true from the assumption, i.e. a and b can be written as the product of primes.


Thus, $k + 1 = a.b$ can also be written as the product of primes.

Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?



Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?





Strong Induction

- Consider a puzzle. How do we assemble a puzzle?



Strong Induction



- Consider a puzzle. How do we assemble a puzzle ?



- Show that no matter which move we make, $n-1$ moves required to assemble a puzzle with n pieces.

Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?





- Show that no matter which move we make, $n-1$ moves required to assemble a puzzle with n pieces.

Basic Step $P(1)$ is true, i.e. no move required for just 1 piece

Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?





- Show that no matter which move we make, $n-1$ moves required to assemble a puzzle with n pieces.

Basic Step $P(1)$ is true, i.e. no move required for just 1 piece

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k + 1)$

Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?



- Show that no matter which move we make, $n-1$ moves required to assemble a puzzle with n pieces.



Basic Step $P(1)$ is true, i.e. no move required for just 1 piece

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e a puzzle with i pieces can be assembled with $i-1$ moves

Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?



- Show that no matter which move we make, $n-1$ moves required to assemble a puzzle with n pieces.

Basic Step $P(1)$ is true, i.e. no move required for just 1 piece



Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e a puzzle with i pieces can be assembled with $i-1$ moves

$k+1$ pieces

Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?




- Show that no matter which move we make, $n-1$ moves required to assemble a puzzle with n pieces.

Basic Step $P(1)$ is true, i.e. no move required for just 1 piece

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e a puzzle with i pieces can be assembled with $i-1$ moves

$k+1$ pieces



Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?




- Show that no matter which move we make, $n-1$ moves required to assemble a puzzle with n pieces.

Basic Step $P(1)$ is true, i.e. no move required for just 1 piece

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e a puzzle with i pieces can be assembled with $i-1$ moves



Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?




- Show that no matter which move we make, $n-1$ moves required to assemble a puzzle with n pieces.

Basic Step $P(1)$ is true, i.e. no move required for just 1 piece

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e a puzzle with i pieces can be assembled with $i-1$ moves



Strong Induction

- Consider a puzzle. How do we assemble a puzzle ?




- Show that no matter which move we make, $n-1$ moves required to assemble a puzzle with n pieces.

Basic Step $P(1)$ is true, i.e. no move required for just 1 piece

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e a puzzle with i pieces can be assembled with $i-1$ moves



Strong Induction

- Prove that for every integer $n \geq 3$, $F(n) > \alpha^{n-2}$ where $\alpha = (1 + \sqrt{5})/2$

Strong Induction

- Prove that for every integer $n \geq 3$, $F(n) > \alpha^{n-2}$ where $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence : $F(1) = 1, F(2) = 1$, and $F(n) = F(n-1) + F(n-2)$

Strong Induction

- Prove that for every integer $n \geq 3$, $F(n) > \alpha^{n-2}$ where $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence : $F(1) = 1, F(2) = 1$, and $F(n) = F(n-1) + F(n-2)$

Basic Step $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Strong Induction

- Prove that for every integer $n \geq 3$, $F(n) > \alpha^{n-2}$ where $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence : $F(1) = 1, F(2) = 1$, and $F(n) = F(n-1) + F(n-2)$

Basic Step $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

Strong Induction

- Prove that for every integer $n \geq 3$, $F(n) > \alpha^{n-2}$ where $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence : $F(1) = 1, F(2) = 1$, and $F(n) = F(n-1) + F(n-2)$

Basic Step $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e $F(i) > \alpha^{i-2}$

Strong Induction

- Prove that for every integer $n \geq 3$, $F(n) > \alpha^{n-2}$ where $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence : $F(1) = 1, F(2) = 1$, and $F(n) = F(n-1) + F(n-2)$

Basic Step $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e $F(i) > \alpha^{i-2}$

for $P(k+1) : F(k+1) = F(k) + F(k-1)$

Strong Induction

- Prove that for every integer $n \geq 3$, $F(n) > \alpha^{n-2}$ where $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence : $F(1) = 1, F(2) = 1$, and $F(n) = F(n-1) + F(n-2)$

Basic Step $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e $F(i) > \alpha^{i-2}$

for $P(k+1) : F(k+1) = F(k) + F(k-1) > \alpha^{k-2} + \alpha^{k-3}$

Strong Induction

- Prove that for every integer $n \geq 3$, $F(n) > \alpha^{n-2}$ where $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence : $F(1) = 1, F(2) = 1$, and $F(n) = F(n-1) + F(n-2)$

Basic Step $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e $F(i) > \alpha^{i-2}$

for $P(k+1) : F(k+1) = F(k) + F(k-1) > \alpha^{k-2} + \alpha^{k-3}$
 $= \alpha \cdot \alpha^{k-3} + \alpha^{k-3}$

Strong Induction

- Prove that for every integer $n \geq 3$, $F(n) > \alpha^{n-2}$ where $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence : $F(1) = 1, F(2) = 1$, and $F(n) = F(n-1) + F(n-2)$

Basic Step $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e $F(i) > \alpha^{i-2}$

for $P(k+1) : F(k+1) = F(k) + F(k-1) > \alpha^{k-2} + \alpha^{k-3}$
 $= \alpha \cdot \alpha^{k-3} + \alpha^{k-3}$
 $= (\alpha + 1) \cdot \alpha^{k-3}$

Strong Induction

- Prove that for every integer $n \geq 3$, $F(n) > \alpha^{n-2}$ where $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence : $F(1) = 1, F(2) = 1$, and $F(n) = F(n-1) + F(n-2)$

Basic Step $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e $F(i) > \alpha^{i-2}$

$$\begin{aligned} \text{for } P(k+1) : F(k+1) &= F(k) + F(k-1) > \alpha^{k-2} + \alpha^{k-3} \\ &= \alpha \cdot \alpha^{k-3} + \alpha^{k-3} \\ &= (\alpha + 1) \cdot \alpha^{k-3} \end{aligned}$$

$\alpha = \frac{1+\sqrt{5}}{2}$ is a solution of the equation $\alpha^2 - \alpha - 1 = 0$. Thus, $\alpha^2 = \alpha + 1$

Strong Induction

- Prove that for every integer $n \geq 3$, $F(n) > \alpha^{n-2}$ where $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence : $F(1) = 1, F(2) = 1$, and $F(n) = F(n-1) + F(n-2)$

Basic Step $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e $F(i) > \alpha^{i-2}$

$$\begin{aligned} \text{for } P(k+1) : F(k+1) &= F(k) + F(k-1) > \alpha^{k-2} + \alpha^{k-3} \\ &= \alpha \cdot \alpha^{k-3} + \alpha^{k-3} \\ &= (\alpha + 1) \cdot \alpha^{k-3} = \alpha^2 \cdot \alpha^{k-3} \end{aligned}$$

$\alpha = \frac{1+\sqrt{5}}{2}$ is a solution of the equation $\alpha^2 - \alpha - 1 = 0$. Thus, $\alpha^2 = \alpha + 1$

Strong Induction

- Prove that for every integer $n \geq 3$, $F(n) > \alpha^{n-2}$ where $\alpha = (1 + \sqrt{5})/2$

Fibonacci sequence : $F(1) = 1, F(2) = 1$, and $F(n) = F(n-1) + F(n-2)$

Basic Step $P(3) : F(3) = 2 > \alpha^{3-2} = (1 + \sqrt{5})/2$

Inductive Step $[P(1) \wedge \dots \wedge P(k)] \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e $F(i) > \alpha^{i-2}$

$$\begin{aligned} \text{for } P(k+1) : F(k+1) &= F(k) + F(k-1) > \alpha^{k-2} + \alpha^{k-3} \\ &= \alpha \cdot \alpha^{k-3} + \alpha^{k-3} \\ &= (\alpha + 1) \cdot \alpha^{k-3} = \alpha^2 \cdot \alpha^{k-3} \end{aligned}$$

$$F(k+1) > \alpha^{k-1}$$

$\alpha = \frac{1+\sqrt{5}}{2}$ is a solution of the equation $\alpha^2 - \alpha - 1 = 0$. Thus, $\alpha^2 = \alpha + 1$

Proofs

Conjecture a formula for the sum of the squares of the first n terms in Fibonacci sequence, then prove your conjecture using mathematical induction

Proofs

Conjecture a formula for the sum of the squares of the first n terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1, F(1)^2 + F(2)^2 = 2, F(1)^2 + F(2)^2 + F(3)^2 = 6,$

Proofs

Conjecture a formula for the sum of the squares of the first n terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1, F(1)^2 + F(2)^2 = 2, F(1)^2 + F(2)^2 + F(3)^2 = 6,$
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15,$

Proofs

Conjecture a formula for the sum of the squares of the first n terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1, F(1)^2 + F(2)^2 = 2, F(1)^2 + F(2)^2 + F(3)^2 = 6,$
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15,$
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40,$

Proofs

Conjecture a formula for the sum of the squares of the first n terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$, $F(1)^2 + F(2)^2 = 2$, $F(1)^2 + F(2)^2 + F(3)^2 = 6$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$,
1.1 1.2 2.3 3.5 5.8

Proofs

Conjecture a formula for the sum of the squares of the first n terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$, $F(1)^2 + F(2)^2 = 2$, $F(1)^2 + F(2)^2 + F(3)^2 = 6$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$,
1.1 1.2 2.3 3.5 5.8
- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$, where $n \geq 2$

Proofs

Conjecture a formula for the sum of the squares of the first n terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$, $F(1)^2 + F(2)^2 = 2$, $F(1)^2 + F(2)^2 + F(3)^2 = 6$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$,
1.1 1.2 2.3 3.5 5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$, where $n \geq 2$

Basic Step $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$

Proofs

Conjecture a formula for the sum of the squares of the first n terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$, $F(1)^2 + F(2)^2 = 2$, $F(1)^2 + F(2)^2 + F(3)^2 = 6$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$,
1.1 1.2 2.3 3.5 5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$, where $n \geq 2$

Basic Step $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$

Inductive Step $P(k) \rightarrow P(k+1)$

Proofs

Conjecture a formula for the sum of the squares of the first n terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$, $F(1)^2 + F(2)^2 = 2$, $F(1)^2 + F(2)^2 + F(3)^2 = 6$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$,
1.1 1.2 2.3 3.5 5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$, where $n \geq 2$

Basic Step $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$

Inductive Step $P(k) \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e. $\sum_{j=1}^i F(j)^2 = F(i) \cdot F(i+1)$

Proofs

Conjecture a formula for the sum of the squares of the first n terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$, $F(1)^2 + F(2)^2 = 2$, $F(1)^2 + F(2)^2 + F(3)^2 = 6$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$,
1.1 1.2 2.3 3.5 5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$, where $n \geq 2$

Basic Step $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$

Inductive Step $P(k) \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e. $\sum_{j=1}^i F(j)^2 = F(i) \cdot F(i+1)$

for $P(k+1) : F(1)^2 + \dots + F(k)^2 + F(k+1)^2$

Proofs

Conjecture a formula for the sum of the squares of the first n terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$, $F(1)^2 + F(2)^2 = 2$, $F(1)^2 + F(2)^2 + F(3)^2 = 6$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$,
1.1 1.2 2.3 3.5 5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$, where $n \geq 2$

Basic Step $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$

Inductive Step $P(k) \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e. $\sum_{j=1}^i F(j)^2 = F(i) \cdot F(i+1)$

for $P(k+1) : F(1)^2 + \dots + F(k)^2 + F(k+1)^2 = F(k) \cdot F(k+1) + F(k+1)^2$

Proofs

Conjecture a formula for the sum of the squares of the first n terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$, $F(1)^2 + F(2)^2 = 2$, $F(1)^2 + F(2)^2 + F(3)^2 = 6$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$,
1.1 1.2 2.3 3.5 5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$, where $n \geq 2$

Basic Step $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$

Inductive Step $P(k) \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e. $\sum_{j=1}^i F(j)^2 = F(i) \cdot F(i+1)$

for $P(k+1) : F(1)^2 + \dots + F(k)^2 + F(k+1)^2 = F(k) \cdot F(k+1) + F(k+1)^2$
 $= F(k+1)(F(k) + F(k+1))$

Proofs

Conjecture a formula for the sum of the squares of the first n terms in Fibonacci sequence, then prove your conjecture using mathematical induction

- $F(1)^2 = 1$, $F(1)^2 + F(2)^2 = 2$, $F(1)^2 + F(2)^2 + F(3)^2 = 6$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 = 15$,
 $F(1)^2 + F(2)^2 + F(3)^2 + F(4)^2 + F(5)^2 = 40$,
1.1 1.2 2.3 3.5 5.8

- $\sum_{j=1}^n F(j)^2 = F(n) \cdot F(n+1)$, where $n \geq 2$

Basic Step $P(2) : F(1)^2 + F(2)^2 = 2 = F(2) \cdot F(3)$

Inductive Step $P(k) \rightarrow P(k+1)$

assume that $P(i)$ is true for all i such that $2 \leq i \leq k$, i.e. $\sum_{j=1}^i F(j)^2 = F(i) \cdot F(i+1)$

$$\begin{aligned} \text{for } P(k+1) : F(1)^2 + \dots + F(k)^2 + F(k+1)^2 &= F(k) \cdot F(k+1) + F(k+1)^2 \\ &= F(k+1)(F(k) + F(k+1)) \\ &= F(k+1)F(k+2) \end{aligned}$$