

Sets

Murat Osmanoglu

Definitions

- A set is an unordered (well-defined) collection of objects.

Definitions

- A set is an unordered (well-defined) collection of objects.
- These objects are called elements (or members) of the set.

Definitions

- A set is an unordered (well-defined) collection of objects.
- These objects are called elements (or members) of the set.
- $x \in A$, x is an element of the set A
- $x \notin A$, x is not an element of the set A

Definitions


- $Z = \{2, -22, 12, 0, 43, -1287, \dots\}$
 $Z = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$

Definitions

- $Z = \{2, -22, 12, 0, 43, -1287, \dots\}$
 $Z = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$
- $A = \{x \in Z^+ | x < 10\}$ (set builder or common property)
 $B = \{x \in Z | x^2 < 10\}$

Definitions

- $Z = \{2, -22, 12, 0, 43, -1287, \dots\}$
 $Z = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$
- $A = \{x \in Z^+ | x < 10\}$ (set builder or common property)
 $B = \{x \in Z | x^2 < 10\}$
- Venn Diagram



Definitions

- Two sets are equal if and only if they have same elements. A and B are equal if and only if $\forall x(x \in A \leftrightarrow x \in B)$.

Definitions

- Two sets are equal if and only if they have same elements. A and B are equal if and only if $\forall x(x \in A \leftrightarrow x \in B)$.

$$A = \{x \in \mathbb{Z}^+ | x < 6\} \text{ and } B = \{1, 2, 3, 4, 5\}, A = B$$

Definitions

- Two sets are equal if and only if they have same elements. A and B are equal if and only if $\forall x(x \in A \leftrightarrow x \in B)$.

$$A = \{x \in \mathbb{Z}^+ | x < 6\} \text{ and } B = \{1, 2, 3, 4, 5\}, A = B$$

- The universal set, denoted by U , contains all possible elements under the consideration

Definitions

- Two sets are equal if and only if they have same elements. A and B are equal if and only if $\forall x(x \in A \leftrightarrow x \in B)$.

$$A = \{x \in \mathbb{Z}^+ | x < 6\} \text{ and } B = \{1, 2, 3, 4, 5\}, A = B$$

- The universal set, denoted by U , contains all possible elements under the consideration
- The empty set, denoted by \emptyset , has no element

Definitions

- A set A is a subset of a set B if and only if every element of A is also an element of B .

Definitions

- A set A is a subset of a set B if and only if every element of A is also an element of B .

$$A \subseteq B \leftrightarrow \forall x [x \in A \rightarrow x \in B]$$

Definitions

- A set A is a subset of a set B if and only if every element of A is also an element of B .

$$A \subseteq B \leftrightarrow \forall x [x \in A \rightarrow x \in B]$$

$$A \not\subseteq B$$

Definitions

- A set A is a subset of a set B if and only if every element of A is also an element of B .

$$A \subseteq B \leftrightarrow \forall x [x \in A \rightarrow x \in B]$$

$$A \not\subseteq B \leftrightarrow \sim \forall x [x \in A \rightarrow x \in B]$$

Definitions

- A set A is a subset of a set B if and only if every element of A is also an element of B .

$$A \subseteq B \leftrightarrow \forall x [x \in A \rightarrow x \in B]$$

$$\begin{aligned} A \not\subseteq B &\leftrightarrow \sim \forall x [x \in A \rightarrow x \in B] \\ &\leftrightarrow \exists x \sim [x \in A \rightarrow x \in B] \end{aligned}$$

Definitions

- A set A is a subset of a set B if and only if every element of A is also an element of B .

$$A \subseteq B \leftrightarrow \forall x [x \in A \rightarrow x \in B]$$

$$A \not\subseteq B \leftrightarrow \sim \forall x [x \in A \rightarrow x \in B]$$

$$\leftrightarrow \exists x \sim [x \in A \rightarrow x \in B]$$

$$\leftrightarrow \exists x \sim [\sim x \in A \vee x \in B]$$

$$(p \rightarrow q \equiv \sim p \vee q)$$

Definitions

- A set A is a subset of a set B if and only if every element of A is also an element of B .

$$A \subseteq B \leftrightarrow \forall x [x \in A \rightarrow x \in B]$$

$$A \not\subseteq B \leftrightarrow \sim \forall x [x \in A \rightarrow x \in B]$$

$$\leftrightarrow \exists x \sim [x \in A \rightarrow x \in B]$$

$$\leftrightarrow \exists x \sim [\sim x \in A \vee x \in B]$$

$$\leftrightarrow \exists x [x \in A \wedge \sim x \in B]$$

$$(p \rightarrow q \equiv \sim p \vee q)$$

Definitions

- A set A is a subset of a set B if and only if every element of A is also an element of B .

$$A \subseteq B \leftrightarrow \forall x [x \in A \rightarrow x \in B]$$

$$\begin{aligned} A \not\subseteq B &\leftrightarrow \sim \forall x [x \in A \rightarrow x \in B] \\ &\leftrightarrow \exists x \sim [x \in A \rightarrow x \in B] \\ &\leftrightarrow \exists x \sim [\sim x \in A \vee x \in B] & (p \rightarrow q \equiv \sim p \vee q) \\ &\leftrightarrow \exists x [x \in A \wedge \sim x \in B] \\ &\leftrightarrow \exists x [x \in A \wedge x \notin B] \end{aligned}$$

Definitions

- A set A is a subset of a set B if and only if every element of A is also an element of B .

$$A \subseteq B \leftrightarrow \forall x [x \in A \rightarrow x \in B]$$

- $\emptyset \subseteq A$ and $A \subseteq A$.

Definitions

- A set A is a subset of a set B if and only if every element of A is also an element of B .

$$A \subseteq B \leftrightarrow \forall x [x \in A \rightarrow x \in B]$$

- $\emptyset \subseteq A$ and $A \subseteq A$.
- $A = B$ if and only if $A \subseteq B$ and $B \subseteq A$

Definitions

- $A = \{x | x = 4k + 1 \text{ for some } k \in \mathbb{Z}\},$

- $B = \{x | x = 4k - 3 \text{ for some } k \in \mathbb{Z}\}$

Show that whether the sets A and B are equal or not.

of A is

Definitions

- $A = \{x | x = 4k + 1 \text{ for some } k \in \mathbb{Z}\},$

- $B = \{x | x = 4k - 3 \text{ for some } k \in \mathbb{Z}\}$

Show that whether the sets A and B are equal or not.

- $(A \subseteq B)$

of A is

Definitions

- $A = \{x | x = 4k + 1 \text{ for some } k \in \mathbb{Z}\}$,
 $B = \{x | x = 4k - 3 \text{ for some } k \in \mathbb{Z}\}$
Show that whether the sets A and B are equal or not.
- $(A \subseteq B)$ For any $x \in A$, $x = 4k + 1$ for some $k \in \mathbb{Z}$
-
-

of A is

Definitions

- $A = \{x | x = 4k + 1 \text{ for some } k \in \mathbb{Z}\}$,
 $B = \{x | x = 4k - 3 \text{ for some } k \in \mathbb{Z}\}$
Show that whether the sets A and B are equal or not.

($A \subseteq B$) For any $x \in A$, $x = 4k + 1$ for some $k \in \mathbb{Z}$
 $x = 4k + 1 + 3 - 3$

Definitions

- $A = \{x | x = 4k + 1 \text{ for some } k \in \mathbb{Z}\},$

- $B = \{x | x = 4k - 3 \text{ for some } k \in \mathbb{Z}\}$

Show that whether the sets A and B are equal or not.

- $(A \subseteq B)$ For any $x \in A$, $x = 4k + 1$ for some $k \in \mathbb{Z}$

- $$x = 4k + 1 + 3 - 3$$

- $$x = 4(k + 1) - 3$$

of A is

Definitions

- $A = \{x | x = 4k + 1 \text{ for some } k \in \mathbb{Z}\},$

- $B = \{x | x = 4k - 3 \text{ for some } k \in \mathbb{Z}\}$

Show that whether the sets A and B are equal or not.

- $(A \subseteq B)$ For any $x \in A$, $x = 4k + 1$ for some $k \in \mathbb{Z}$

- $x = 4k + 1 + 3 - 3$

- $x = 4(k + 1) - 3$

- $x = 4m - 3$ for some $m \in \mathbb{Z}$, so $x \in B$

of A is

Definitions

- $A = \{x | x = 4k + 1 \text{ for some } k \in \mathbb{Z}\},$

- $B = \{x | x = 4k - 3 \text{ for some } k \in \mathbb{Z}\}$

Show that whether the sets A and B are equal or not.

- $(A \subseteq B)$ For any $x \in A$, $x = 4k + 1$ for some $k \in \mathbb{Z}$

- $x = 4k + 1 + 3 - 3$

- $x = 4(k + 1) - 3$

- $x = 4m - 3$ for some $m \in \mathbb{Z}$, so $x \in B$

- $(B \subseteq A)$

of A is

Definitions

- $A = \{x | x = 4k + 1 \text{ for some } k \in \mathbb{Z}\},$

- $B = \{x | x = 4k - 3 \text{ for some } k \in \mathbb{Z}\}$

Show that whether the sets A and B are equal or not.

- $(A \subseteq B)$ For any $x \in A$, $x = 4k + 1$ for some $k \in \mathbb{Z}$

- $x = 4k + 1 + 3 - 3$

- $x = 4(k + 1) - 3$

- $x = 4m - 3$ for some $m \in \mathbb{Z}$, so $x \in B$

- $(B \subseteq A)$ For any $x \in B$, $x = 4k - 3$ for some $k \in \mathbb{Z}$

of A is

Definitions

- $A = \{x | x = 4k + 1 \text{ for some } k \in \mathbb{Z}\},$

- $B = \{x | x = 4k - 3 \text{ for some } k \in \mathbb{Z}\}$

Show that whether the sets A and B are equal or not.

- $(A \subseteq B)$ For any $x \in A$, $x = 4k + 1$ for some $k \in \mathbb{Z}$

- $x = 4k + 1 + 3 - 3$

- $x = 4(k + 1) - 3$

- $x = 4m - 3$ for some $m \in \mathbb{Z}$, so $x \in B$

- $(B \subseteq A)$ For any $x \in B$, $x = 4k - 3$ for some $k \in \mathbb{Z}$

- $x = 4k - 3 + 1 - 1$

of A is

Definitions

- $A = \{x \mid x = 4k + 1 \text{ for some } k \in \mathbb{Z}\},$

- $B = \{x \mid x = 4k - 3 \text{ for some } k \in \mathbb{Z}\}$

Show that whether the sets A and B are equal or not.

- $(A \subseteq B)$ For any $x \in A$, $x = 4k + 1$ for some $k \in \mathbb{Z}$

- $x = 4k + 1 + 3 - 3$

- $x = 4(k + 1) - 3$

- $x = 4m - 3$ for some $m \in \mathbb{Z}$, so $x \in B$

- $(B \subseteq A)$ For any $x \in B$, $x = 4k - 3$ for some $k \in \mathbb{Z}$

- $x = 4k - 3 + 1 - 1$

- $x = 4(k - 1) + 1$

of A is

Definitions

- $A = \{x \mid x = 4k + 1 \text{ for some } k \in \mathbb{Z}\},$

- $B = \{x \mid x = 4k - 3 \text{ for some } k \in \mathbb{Z}\}$

Show that whether the sets A and B are equal or not.

- $(A \subseteq B)$ For any $x \in A$, $x = 4k + 1$ for some $k \in \mathbb{Z}$

- $x = 4k + 1 + 3 - 3$

- $x = 4(k + 1) - 3$

- $x = 4m - 3$ for some $m \in \mathbb{Z}$, so $x \in B$

- $(B \subseteq A)$ For any $x \in B$, $x = 4k - 3$ for some $k \in \mathbb{Z}$

- $x = 4k - 3 + 1 - 1$

- $x = 4(k - 1) + 1$

- $x = 4m + 1$ for some $m \in \mathbb{Z}$, so $x \in A$

of A is

Definitions

- $A = \{x | x = 4k + 1 \text{ for some } k \in \mathbb{Z}\},$

- $B = \{x | x = 4k - 3 \text{ for some } k \in \mathbb{Z}\}$

Show that whether the sets A and B are equal or not.

- $(A \subseteq B)$ For any $x \in A$, $x = 4k + 1$ for some $k \in \mathbb{Z}$

- $x = 4k + 1 + 3 - 3$

- $x = 4(k + 1) - 3$

- $x = 4m - 3$ for some $m \in \mathbb{Z}$, so $x \in B$

- $(B \subseteq A)$ For any $x \in B$, $x = 4k - 3$ for some $k \in \mathbb{Z}$

- $x = 4k - 3 + 1 - 1$

- $x = 4(k - 1) + 1$

- $x = 4m + 1$ for some $m \in \mathbb{Z}$, so $x \in A$

Thus, $A=B$.

of A is

Definitions

- A set A is a subset of a set B if and only if every element of A is also an element of B .

$$A \subseteq B \leftrightarrow \forall x [x \in A \rightarrow x \in B]$$

- $\emptyset \subseteq A$ and $A \subseteq A$.
- $A = B$ if and only if $A \subseteq B$ and $B \subseteq A$
- A set A is a proper subset of a set B if and only if $A \subseteq B$ and $A \neq B$

Definitions

- A set A is a subset of a set B if and only if every element of A is also an element of B .

$$A \subseteq B \leftrightarrow \forall x [x \in A \rightarrow x \in B]$$

- $\emptyset \subseteq A$ and $A \subseteq A$.
- $A = B$ if and only if $A \subseteq B$ and $B \subseteq A$
- A set A is a proper subset of a set B if and only if $A \subseteq B$ and $A \neq B$

$$B = \{x \in \mathbb{Z}^+ | x < 10\} \text{ and } A = \{1, 2, 3, 4, 5\}$$

Definitions

- A set A is a subset of a set B if and only if every element of A is also an element of B .

$$A \subseteq B \leftrightarrow \forall x [x \in A \rightarrow x \in B]$$

- $\emptyset \subseteq A$ and $A \subseteq A$.
- $A = B$ if and only if $A \subseteq B$ and $B \subseteq A$
- A set A is a proper subset of a set B if and only if $A \subseteq B$ and $A \neq B$

$$B = \{x \in \mathbb{Z}^+ | x < 10\} \text{ and } A = \{1, 2, 3, 4, 5\}, A \subseteq B$$

Definitions

- The **cardinality** of a set A is defined as the size of A . It's denoted by $|A|$. (only for finite set)

Definitions

- The **cardinality** of a set A is defined as the size of A . It's denoted by $|A|$. (only for finite set)

For the set $A = \{x \in \mathbb{Z}^+ | x < 10\}$, $|A| = 9$

Definitions

- The **cardinality** of a set A is defined as the size of A . It's denoted by $|A|$. (only for finite set)

For the set $A = \{x \in \mathbb{Z}^+ | x < 10\}$, $|A| = 9$

- The **power set** of a given set is the set of all possible subsets.

Definitions

- The **cardinality** of a set A is defined as the size of A . It's denoted by $|A|$. (only for finite set)

For the set $A = \{x \in \mathbb{Z}^+ | x < 10\}$, $|A| = 9$

- The **power set** of a given set is the set of all possible subsets.

$$S = \{1\}$$

$$P(S) = \{\emptyset, \{1\}\}$$

Definitions

- The **cardinality** of a set A is defined as the size of A . It's denoted by $|A|$. (only for finite set)

For the set $A = \{x \in \mathbb{Z}^+ | x < 10\}$, $|A| = 9$

- The **power set** of a given set is the set of all possible subsets.

$$S = \{1\}$$

$$P(S) = \{\emptyset, \{1\}\}$$

$$S = \{a, b\}$$

$$P(S) = \{\emptyset, \{a\}, \{b\}, \{a, b\}\}$$

Definitions

- The **cardinality** of a set A is defined as the size of A . It's denoted by $|A|$. (only for finite set)

For the set $A = \{x \in \mathbb{Z}^+ | x < 10\}$, $|A| = 9$

- The **power set** of a given set is the set of all possible subsets.

$$S = \{1\}$$

$$P(S) = \{\emptyset, \{1\}\}$$

$$|P(S)| = 2$$

$$S = \{a, b\}$$

$$P(S) = \{\emptyset, \{a\}, \{b\}, \{a, b\}\}$$

$$|P(S)| = 4$$

Definitions

- The **cardinality** of a set A is defined as the size of A . It's denoted by $|A|$. (only for finite set)

For the set $A = \{x \in \mathbb{Z}^+ | x < 10\}$, $|A| = 9$

- The **power set** of a given set is the set of all possible subsets.

$$S = \{1\} \qquad P(S) = \{\emptyset, \{1\}\} \qquad |P(S)| = 2$$

$$S = \{a, b\} \qquad P(S) = \{\emptyset, \{a\}, \{b\}, \{a, b\}\} \qquad |P(S)| = 4$$

- If $|S| = n$, then $|P(S)| = 2^n$

Set Operations

Set Operations

- The union of A and B , denoted by $A \cup B$, contains elements that are either in A or B .

$$A \cup B = \{x | x \in A \vee x \in B\}$$

Set Operations

- The union of A and B , denoted by $A \cup B$, contains elements that are either in A or B .

$$A \cup B = \{x | x \in A \vee x \in B\}$$

- The intersection of A and B , denoted by $A \cap B$, contains elements that are in both A or B .

$$A \cap B = \{x | x \in A \wedge x \in B\}$$

Set Operations

- The union of A and B , denoted by $A \cup B$, contains elements that are either in A or B .

$$A \cup B = \{x | x \in A \vee x \in B\}$$

- The intersection of A and B , denoted by $A \cap B$, contains elements that are in both A or B .

$$A \cap B = \{x | x \in A \wedge x \in B\}$$

The sets A and B are called disjoint sets if $A \cap B = \emptyset$

Set Operations

- The union of A and B , denoted by $A \cup B$, contains elements that are either in A or B .

$$A \cup B = \{x | x \in A \vee x \in B\}$$

- The intersection of A and B , denoted by $A \cap B$, contains elements that are in both A or B .

$$A \cap B = \{x | x \in A \wedge x \in B\}$$

The sets A and B are called disjoint sets if $A \cap B = \emptyset$

- The difference of A and B , denoted by $A - B$, contains elements that are in A but not in B

$$A - B = \{x | x \in A \wedge x \notin B\}$$

Set Operations

- The union of A and B , denoted by $A \cup B$, contains elements that are either in A or B .

$$A \cup B = \{x | x \in A \vee x \in B\}$$

- The intersection of A and B , denoted by $A \cap B$, contains elements that are in both A or B .

$$A \cap B = \{x | x \in A \wedge x \in B\}$$

The sets A and B are called disjoint sets if $A \cap B = \emptyset$

- The difference of A and B , denoted by $A - B$, contains elements that are in A but not in B

$$A - B = \{x | x \in A \wedge x \notin B\}$$

- The complement of A , denoted by \bar{A} , contains elements that are in U but not in A .

$$\bar{A} = \{x \in U | x \notin A\}$$

Set Operations

- $A \cup \emptyset = A$
 $A \cap U = A$

Set Operations

- $A \cup \emptyset = A$
 $A \cap U = A$
- $A \cup U = U$
 $A \cap \emptyset = \emptyset$

Set Operations

- $A \cup \emptyset = A$
 $A \cap U = A$
- $A \cup U = U$
 $A \cap \emptyset = \emptyset$
- $A \cup A = A$
 $A \cap A = A$

Set Operations

- $A \cup \emptyset = A$
 $A \cap U = A$
- $A \cup U = U$
 $A \cap \emptyset = \emptyset$
- $A \cup A = A$
 $A \cap A = A$
- $A \cup B = B \cup A$
 $A \cap B = B \cap A$

Set Operations

- $A \cup \emptyset = A$
 $A \cap U = A$
- $A \cup U = U$
 $A \cap \emptyset = \emptyset$
- $A \cup A = A$
 $A \cap A = A$
- $A \cup B = B \cup A$
 $A \cap B = B \cap A$
- $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
 $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

Set Operations

- $A \cup \emptyset = A$
 $A \cap U = A$
- $A \cup U = U$
 $A \cap \emptyset = \emptyset$
- $A \cup A = A$
 $A \cap A = A$
- $A \cup B = B \cup A$
 $A \cap B = B \cap A$
- $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
 $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
- $A \cap \bar{A} = \emptyset$
 $A \cup \bar{A} = U$

Set Operations

- $A \cup \emptyset = A$
 $A \cap U = A$
- $A \cup U = U$
 $A \cap \emptyset = \emptyset$
- $A \cup A = A$
 $A \cap A = A$
- $A \cup B = B \cup A$
 $A \cap B = B \cap A$
- $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
 $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
- $A \cap \bar{A} = \emptyset$
 $A \cup \bar{A} = U$
- $\overline{(\bar{A})} = A$

Set Operations

- $A \cup \emptyset = A$
 $A \cap U = A$
- $A \cup U = U$
 $A \cap \emptyset = \emptyset$
- $A \cup A = A$
 $A \cap A = A$
- $A \cup B = B \cup A$
 $A \cap B = B \cap A$
- $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
 $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
- $A \cap \bar{A} = \emptyset$
 $A \cup \bar{A} = U$
- $\overline{(\bar{A})} = A$
- $\overline{A \cup B} = \bar{A} \cap \bar{B}$ (De Morgan)
 $\overline{A \cap B} = \bar{A} \cup \bar{B}$

Set Operations

- $A \cup \emptyset = A$
 $A \cap U = A$
 $p \vee 0 \equiv p$
 $p \wedge 1 \equiv p$
- $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
 $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
- $A \cup U = U$
 $A \cap \emptyset = \emptyset$
 $p \vee 1 \equiv 1$
 $p \wedge 0 \equiv 0$
- $A \cap \bar{A} = \emptyset$
 $A \cup \bar{A} = U$
 $p \wedge \sim p \equiv 0$
 $p \vee \sim p \equiv 1$
- $A \cup A = A$
 $A \cap A = A$
 $p \vee p \equiv p$
 $p \wedge p \equiv p$
- $\overline{(\bar{A})} = A$
 $\sim(\sim p) \equiv p$
- $\overline{A \cup B} = \bar{A} \cap \bar{B}$ (De Morgan)
 $\overline{A \cap B} = \bar{A} \cup \bar{B}$
 $\sim(p \vee q) \equiv \sim p \wedge \sim q$
 $\sim(p \wedge q) \equiv \sim p \vee \sim q$

Set Operations

- $A \cup \emptyset = A$
 $A \cap U = A$
- $A \cup U = U$
 $A \cap \emptyset = \emptyset$
- $A \cup A = A$
 $A \cap A = A$
- $A \cup B = B \cup A$
 $A \cap B = B \cap A$
- $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
 $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
- $A \cap \bar{A} = \emptyset$
 $A \cup \bar{A} = U$
- $\overline{(\bar{A})} = A$
- $\overline{A \cup B} = \bar{A} \cap \bar{B}$ (De Morgan)
 $\overline{A \cap B} = \bar{A} \cup \bar{B}$

$$|A \cup B| = |A| + |B| - |A \cap B|$$

Definitions

Show that $\overline{A \cup B} = \bar{A} \cap \bar{B}$.

A is

Definitions

Show that $\overline{A \cup B} = \overline{A} \cap \overline{B}$.

• $(\overline{A \cup B} \subseteq \overline{A} \cap \overline{B})$

A is

Definitions

Show that $\overline{A \cup B} = \overline{A} \cap \overline{B}$.

- $(\overline{A \cup B} \subseteq \overline{A} \cap \overline{B})$ assume $x \in \overline{A \cup B}$, then $(x \notin A \cup B)$
- $(\overline{A} \cap \overline{B} \subseteq \overline{A \cup B})$ assume $x \in \overline{A} \cap \overline{B}$, then $(x \notin A \cup B)$
- $\overline{A \cup B} = \overline{A} \cap \overline{B}$

A is

Definitions

Show that $\overline{A \cup B} = \bar{A} \cap \bar{B}$.

- $\overline{A \cup B} \subseteq \bar{A} \cap \bar{B}$ assume $x \in \overline{A \cup B}$, then $(x \notin A \cup B)$
 $(x \notin A \cup B) \leftrightarrow \sim((x \in A) \vee (x \in B))$
- $\bar{A} \cap \bar{B} \subseteq \overline{A \cup B}$ assume $x \in \bar{A} \cap \bar{B}$, then $(x \notin A) \wedge (x \notin B)$
 $(x \notin A) \wedge (x \notin B) \leftrightarrow \sim((x \in A) \vee (x \in B))$
- $\overline{A \cup B} = \bar{A} \cap \bar{B}$

Definitions

Show that $\overline{A \cup B} = \overline{A} \cap \overline{B}$.

- $\overline{A \cup B} \subseteq \overline{A} \cap \overline{B}$ assume $x \in \overline{A \cup B}$, then $(x \notin A \cup B)$
 $(x \notin A \cup B) \leftrightarrow \sim((x \in A) \vee (x \in B))$
 $\leftrightarrow (x \notin A) \wedge (x \notin B)$
- $\overline{A} \cap \overline{B} \subseteq \overline{A \cup B}$ assume $x \in \overline{A} \cap \overline{B}$, then $(x \notin A) \wedge (x \notin B)$
 $\leftrightarrow \sim((x \in A) \vee (x \in B))$
 $\leftrightarrow x \notin A \cup B$
- $\overline{A \cup B} = \overline{A} \cap \overline{B}$

Definitions

Show that $\overline{A \cup B} = \overline{A} \cap \overline{B}$.

- $\overline{A \cup B} \subseteq \overline{A} \cap \overline{B}$ assume $x \in \overline{A \cup B}$, then $(x \notin A \cup B)$
 $(x \notin A \cup B) \leftrightarrow \sim((x \in A) \vee (x \in B))$
 $\leftrightarrow (x \notin A) \wedge (x \notin B)$
 $\leftrightarrow (x \in \overline{A}) \wedge (x \in \overline{B})$
- $\overline{A} \cap \overline{B} \subseteq \overline{A \cup B}$ assume $x \in \overline{A} \cap \overline{B}$, then $(x \in \overline{A}) \wedge (x \in \overline{B})$
 $(x \in \overline{A}) \wedge (x \in \overline{B}) \leftrightarrow (x \notin A) \wedge (x \notin B)$
 $\leftrightarrow \sim((x \in A) \vee (x \in B))$
 $\leftrightarrow x \notin A \cup B$
 $\leftrightarrow x \in \overline{A \cup B}$
- $\overline{A \cup B} = \overline{A} \cap \overline{B}$

Definitions

Show that $\overline{A \cup B} = \overline{A} \cap \overline{B}$.

- $\overline{A \cup B} \subseteq \overline{A} \cap \overline{B}$ assume $x \in \overline{A \cup B}$, then $(x \notin A \cup B)$
 $(x \notin A \cup B) \leftrightarrow \sim((x \in A) \vee (x \in B))$
 $\leftrightarrow (x \notin A) \wedge (x \notin B)$
 $\leftrightarrow (x \in \overline{A}) \wedge (x \in \overline{B})$
 $\leftrightarrow x \in \overline{A} \cap \overline{B}$
- $\overline{A} \cap \overline{B} \subseteq \overline{A \cup B}$
- $\overline{A \cup B} = \overline{A} \cap \overline{B}$

A is

Definitions

Show that $\overline{A \cup B} = \overline{A} \cap \overline{B}$.

- $\overline{A \cup B} \subseteq \overline{A} \cap \overline{B}$ assume $x \in \overline{A \cup B}$, then $(x \notin A \cup B)$
 $(x \notin A \cup B) \leftrightarrow \sim((x \in A) \vee (x \in B))$
 $\leftrightarrow (x \notin A) \wedge (x \notin B)$
 $\leftrightarrow (x \in \overline{A}) \wedge (x \in \overline{B})$
 $\leftrightarrow x \in \overline{A} \cap \overline{B}$
- $\overline{A} \cap \overline{B} \subseteq \overline{A \cup B}$

Definitions

Show that $\overline{A \cup B} = \bar{A} \cap \bar{B}$.

- $(\overline{A \cup B} \subseteq \bar{A} \cap \bar{B})$ assume $x \in \overline{A \cup B}$, then $(x \notin A \cup B)$
 $(x \notin A \cup B) \leftrightarrow \sim((x \in A) \vee (x \in B))$
 $\leftrightarrow (x \notin A) \wedge (x \notin B)$
 $\leftrightarrow (x \in \bar{A}) \wedge (x \in \bar{B})$
 $\leftrightarrow x \in \bar{A} \cap \bar{B}$
- $(\bar{A} \cap \bar{B} \subseteq \overline{A \cup B})$ assume $x \in \bar{A} \cap \bar{B}$, then $(x \in \bar{A}) \wedge (x \in \bar{B})$

Definitions

Show that $\overline{A \cup B} = \bar{A} \cap \bar{B}$.

- $(\overline{A \cup B} \subseteq \bar{A} \cap \bar{B})$ assume $x \in \overline{A \cup B}$, then $(x \notin A \cup B)$
 $(x \notin A \cup B) \leftrightarrow \sim((x \in A) \vee (x \in B))$
 $\leftrightarrow (x \notin A) \wedge (x \notin B)$
 $\leftrightarrow (x \in \bar{A}) \wedge (x \in \bar{B})$
 $\leftrightarrow x \in \bar{A} \cap \bar{B}$
- $(\bar{A} \cap \bar{B} \subseteq \overline{A \cup B})$ assume $x \in \bar{A} \cap \bar{B}$, then $(x \in \bar{A}) \wedge (x \in \bar{B})$
 $(x \in \bar{A}) \wedge (x \in \bar{B}) \leftrightarrow (x \notin A) \wedge (x \notin B)$

Definitions

Show that $\overline{A \cup B} = \overline{A} \cap \overline{B}$.

- $(\overline{A \cup B} \subseteq \overline{A} \cap \overline{B})$ assume $x \in \overline{A \cup B}$, then $(x \notin A \cup B)$
 $(x \notin A \cup B) \leftrightarrow \sim((x \in A) \vee (x \in B))$
 $\leftrightarrow (x \notin A) \wedge (x \notin B)$
 $\leftrightarrow (x \in \overline{A}) \wedge (x \in \overline{B})$
 $\leftrightarrow x \in \overline{A} \cap \overline{B}$
- $(\overline{A} \cap \overline{B} \subseteq \overline{A \cup B})$ assume $x \in \overline{A} \cap \overline{B}$, then $(x \in \overline{A}) \wedge (x \in \overline{B})$
 $(x \in \overline{A}) \wedge (x \in \overline{B}) \leftrightarrow (x \notin A) \wedge (x \notin B)$
 $\leftrightarrow \sim(x \in A) \wedge \sim(x \in B)$

Definitions

Show that $\overline{A \cup B} = \bar{A} \cap \bar{B}$.

- $(\overline{A \cup B} \subseteq \bar{A} \cap \bar{B})$ assume $x \in \overline{A \cup B}$, then $(x \notin A \cup B)$
 $(x \notin A \cup B) \leftrightarrow \sim((x \in A) \vee (x \in B))$
 $\leftrightarrow (x \notin A) \wedge (x \notin B)$
 $\leftrightarrow (x \in \bar{A}) \wedge (x \in \bar{B})$
 $\leftrightarrow x \in \bar{A} \cap \bar{B}$
- $(\bar{A} \cap \bar{B} \subseteq \overline{A \cup B})$ assume $x \in \bar{A} \cap \bar{B}$, then $(x \in \bar{A}) \wedge (x \in \bar{B})$
 $(x \in \bar{A}) \wedge (x \in \bar{B}) \leftrightarrow (x \notin A) \wedge (x \notin B)$
 $\leftrightarrow \sim(x \in A) \wedge \sim(x \in B)$
 $\leftrightarrow \sim((x \in A) \vee (x \in B))$

Definitions

Show that $\overline{A \cup B} = \overline{A} \cap \overline{B}$.

- $(\overline{A \cup B} \subseteq \overline{A} \cap \overline{B})$ assume $x \in \overline{A \cup B}$, then $(x \notin A \cup B)$
 $(x \notin A \cup B) \leftrightarrow \sim((x \in A) \vee (x \in B))$
 $\leftrightarrow (x \notin A) \wedge (x \notin B)$
 $\leftrightarrow (x \in \overline{A}) \wedge (x \in \overline{B})$
 $\leftrightarrow x \in \overline{A} \cap \overline{B}$
- $(\overline{A} \cap \overline{B} \subseteq \overline{A \cup B})$ assume $x \in \overline{A} \cap \overline{B}$, then $(x \in \overline{A}) \wedge (x \in \overline{B})$
 $(x \in \overline{A}) \wedge (x \in \overline{B}) \leftrightarrow (x \notin A) \wedge (x \notin B)$
 $\leftrightarrow \sim(x \in A) \wedge \sim(x \in B)$
 $\leftrightarrow \sim((x \in A) \vee (x \in B))$
 $\leftrightarrow (x \notin A \cup B) \leftrightarrow (x \in \overline{A \cup B})$

Cartesian Products

- The cartesian product of A and B , denoted by $A \times B$, is the set of all pairs (x,y) where $x \in A$ and $y \in B$

Cartesian Products

- The cartesian product of A and B , denoted by $A \times B$, is the set of all pairs (x,y) where $x \in A$ and $y \in B$

$$A \times B = \{(x,y) | x \in A \wedge y \in B\}$$

Cartesian Products

- The cartesian product of A and B , denoted by $A \times B$, is the set of all pairs (x,y) where $x \in A$ and $y \in B$

$$A \times B = \{(x,y) | x \in A \wedge y \in B\}$$

- $A=\{a,b\}$, $B=\{1, 2, 3\}$

Cartesian Products

- The cartesian product of A and B , denoted by $A \times B$, is the set of all pairs (x,y) where $x \in A$ and $y \in B$

$$A \times B = \{(x,y) | x \in A \wedge y \in B\}$$

- $A=\{a,b\}$, $B=\{1, 2, 3\}$

$$A \times B = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$$

Cartesian Products

- The cartesian product of A and B , denoted by $A \times B$, is the set of all pairs (x,y) where $x \in A$ and $y \in B$

$$A \times B = \{(x,y) | x \in A \wedge y \in B\}$$

- $A=\{a,b\}$, $B=\{1, 2, 3\}$

$$A \times B = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$$

$$|A \times B| = |A| \cdot |B|$$

Cartesian Products

- The cartesian product of A and B , denoted by $A \times B$, is the set of all pairs (x, y) where $x \in A$ and $y \in B$

$$A \times B = \{(x, y) | x \in A \wedge y \in B\}$$

- $A = \{a, b\}$, $B = \{1, 2, 3\}$

$$A \times B = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$$

$$|A \times B| = |A| \cdot |B|$$

- *The Cartesian products of the sets A_1, A_2, \dots, A_n is the set of ordered n -tuples (a_1, a_2, \dots, a_n) where $a_1 \in A_1, a_2 \in A_2, \dots, a_n \in A_n$.*

Cartesian Products

- The cartesian product of A and B , denoted by $A \times B$, is the set of all pairs (x, y) where $x \in A$ and $y \in B$

$$A \times B = \{(x, y) \mid x \in A \wedge y \in B\}$$

- $A = \{a, b\}$, $B = \{1, 2, 3\}$

$$A \times B = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$$

$$|A \times B| = |A| \cdot |B|$$

- The Cartesian products of the sets A_1, A_2, \dots, A_n is the set of ordered n -tuples (a_1, a_2, \dots, a_n) where $a_1 \in A_1, a_2 \in A_2, \dots, a_n \in A_n$.*

$$A_1 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) \mid a_i \in A_i, i = 1..n\}$$