

# More on Dynamic Programming

Murat Osmanoglu

# SSSP

- given a weighted graph  $G=(V,E)$  and a source vertex  $s$  in  $V$ , find the shortest path from  $s$  to every other vertex in  $V$
- the weight of each edge fixed as 1


--BFS--

- the weight of each edge non-negative

--Dijkstra—


- the weight of each edge can be negative

# Dijkstra's Algorithm



- find the best paths from tether to all other cryptocurrencies

# Dijkstra's Algorithm




- find the best paths from tether to all other cryptocurrencies

tether - cardano - bitcoin - litecoin

$$1 \text{ tether} = 4,255 * 0,000028 * 63,694 = 0,0075 \text{ LTC}$$

# Dijkstra's Algorithm



- find the best paths from tether to all other cryptocurrencies


tether - cardano - bitcoin - litecoin

$$1 \text{ tether} = 4,255 * 0,000028 * 63,694 = 0,0075 \text{ LTC}$$

tether - ripple - etherbase - litecoin

$$1 \text{ tether} = 1,515 * 0,0013 * 3,98 = 0,0078 \text{ LTC}$$


# Dijkstra's Algorithm



- find the best paths from tether to all other cryptocurrencies

$$\max_{P \in \{s \rightarrow u\}} (\prod_{e \in P} w(e))$$


# Dijkstra's Algorithm



- find the best paths from tether to all other cryptocurrencies

$$K = \max_{P \in \{s \rightarrow u\}} (\prod_{e \in P} w(e))$$

# Dijkstra's Algorithm




- find the best paths from tether to all other cryptocurrencies

$$K = \max_{P \in \{s \rightarrow u\}} (\prod_{e \in P} w(e))$$

$$\log K = \log (\max_{P \in \{s \rightarrow u\}} \sum_{e \in P} w(e))$$



# Dijkstra's Algorithm




- find the best paths from tether to all other cryptocurrencies

$$K = \max_{P \in \{s \rightarrow u\}} (\prod_{e \in P} w(e))$$

$$\log K = \log (\max_{P \in \{s \rightarrow u\}} \sum_{e \in P} w(e))$$

$$\log K = \max_{P \in \{s \rightarrow u\}} \log (\sum_{e \in P} w(e))$$

# Dijkstra's Algorithm



- find the best paths from tether to all other cryptocurrencies

$$K = \max_{P \in \{s \rightarrow u\}} (\prod_{e \in P} w(e))$$

$$\log K = \log (\max_{P \in \{s \rightarrow u\}} \sum_{e \in P} w(e))$$


$$\log K = \max_{P \in \{s \rightarrow u\}} \log (\sum_{e \in P} w(e))$$

$$\log (0,00011) = - 3,95$$


$$\log (0,0629) = - 1,201$$

$$\log (0,658) = - 0,181$$


# Dijkstra's Algorithm




# Dijkstra's Algorithm




# Dijkstra's Algorithm



# Dijkstra's Algorithm




# Dijkstra's Algorithm



$C$   
 $S = \{A \ B \ D\}$

# Dijkstra's Algorithm



The algorithm outputs 4 as the weight of the shortest path from A to D. However, the weight of the shortest path is 2.

Dijkstra's Algorithm fails for this type of problem!



# Dijkstra's Algorithm


## A Naive Approach

- find the lightest edge of the graph
- add the weight of that edge to all edges of the graph in order to make them non-negative
- apply Dijkstra to find the shortest paths

# Dijkstra's Algorithm

## A Naive Approach


- find the lightest edge of the graph
- add the weight of that edge to all edges of the graph in order to make them non-negative
- apply Dijkstra to find the shortest paths



# Dijkstra's Algorithm

## A Naive Approach


- find the lightest edge of the graph
- add the weight of that edge to all edges of the graph in order to make them non-negative
- apply Dijkstra to find the shortest paths




# Dijkstra's Algorithm

## A Naive Approach


- find the lightest edge of the graph
- add the weight of that edge to all edges of the graph in order to make them non-negative
- apply Dijkstra to find the shortest paths




# Dijkstra's Algorithm




# Dijkstra's Algorithm



# Dijkstra's Algorithm



# Dijkstra's Algorithm





# Bellman-Ford

- define a subproblem

# Bellman-Ford

- define a subproblem


$PATH(i,v)$  : the weight of the shortest path to  $v$  that contains  $\leq i$  edges

# Bellman-Ford

- define a subproblem

$\text{PATH}(i,v)$  : the weight of the shortest path to  $v$  that contains  $\leq i$  edges

- construct recurrence relation



## Two Cases


1)

# Bellman-Ford

- define a subproblem

$\text{PATH}(i,v)$  : the weight of the shortest path to  $v$  that contains  $\leq i$  edges

- construct recurrence relation



## Two Cases


1)  $\text{PATH}(i-1, v)$

# Bellman-Ford

- define a subproblem

$\text{PATH}(i,v)$  : the weight of the shortest path to  $v$  that contains  $\leq i$  edges

- construct recurrence relation



## Two Cases

1)  $\text{PATH}(i-1, v)$

2)  $\text{PATH}(i-1, x) + w(x, v)$

relaxation




# Bellman-Ford

- define a subproblem

$PATH(i,v)$  : the weight of the shortest path to  $v$  that contains  $\leq i$  edges

- construct recurrence relation



## Two Cases

1)  $PATH(i-1, v)$

2)  $PATH(i-1, x) + w(x, v)$

relaxation

$$PATH(i, v) = \begin{cases} \infty & , \quad i = 0 \\ 0 & , \quad s = v \\ \min \{PATH(i-1, v), PATH(i-1, x) + w(x, v)\} & , \text{ otherwise} \end{cases}$$

# Bellman-Ford

## Bellman-Ford( $G,s$ )

for each  $u$  of  $V$

$$\text{PATH}(0, v) = \infty$$
$$\text{PATH}(0,s) = 0$$

```
for i = 1 to |V| - 1
```

for each edge  $(u,v)$  in  $E$

$$\text{PATH}(i,v) = \min \{ \text{PATH}(i-1,v), \text{PATH}(i-1,u) + w(u,v) \}$$

# Bellman-Ford

Bellman-Ford( $G,s$ )

for each  $u$  of  $V$

$PATH(0,v) = \infty$

$PATH(0,s) = 0$

for  $i = 1$  to  $|V| - 1$

    for each edge  $(u,v)$  in  $E$

$PATH(i,v) = \min \{PATH(i-1,v),$   
                                 $PATH(i-1,u) + w(u,v)\}$

} Initialize( $G,s$ )  
     $O(|V|)$

}  $O(|E| \cdot |V|)$



# Bellman-Ford

Bellman-Ford( $G,s$ )

for each  $u$  of  $V$


$PATH(0,v) = \infty$

$PATH(0,s) = 0$

for  $i = 1$  to  $|V| - 1$

    for each edge  $(u,v)$  in  $E$

$PATH(i,v) = \min \{PATH(i-1,v),$   
                             $PATH(i-1,u) + w(u,v)\}$



# Bellman-Ford

Bellman-Ford( $G,s$ )

for each  $u$  of  $V$


$PATH(0,v) = \infty$

$PATH(0,s) = 0$

for  $i = 1$  to  $|V| - 1$

for each edge  $(u,v)$  in  $E$

$PATH(i,v) = \min \{PATH(i-1,v),$   
 $PATH(i-1,u) + w(u,v)\}$



	0	1	2
A	$\infty$		
B	$\infty$		
C	$\infty$		
D	0		
E	$\infty$		
F	$\infty$		
H	$\infty$		

# Bellman-Ford

Bellman-Ford( $G,s$ )

for each  $u$  of  $V$


$$\text{PATH}(0,v) = \infty$$

$$\text{PATH}(0,s) = 0$$

for  $i = 1$  to  $|V| - 1$

for each edge  $(u,v)$  in  $E$

$$\text{PATH}(i,v) = \min \{ \text{PATH}(i-1,v), \\ \text{PATH}(i-1,u) + w(u,v) \}$$



	0	1	2
A	$\infty$	$\infty$	
B	$\infty$		
C	$\infty$		
D	0		
E	$\infty$		
F	$\infty$		
H	$\infty$		

$$\text{PATH}(1,A) = \min \begin{cases} \text{PATH}(0,A) \\ \text{PATH}(0,B) + w(B,A) \\ \text{PATH}(0,C) + w(C,A) \end{cases}$$

# Bellman-Ford

Bellman-Ford( $G,s$ )

for each  $u$  of  $V$


$$\text{PATH}(0,v) = \infty$$

$$\text{PATH}(0,s) = 0$$

for  $i = 1$  to  $|V| - 1$

for each edge  $(u,v)$  in  $E$

$$\text{PATH}(i,v) = \min \{ \text{PATH}(i-1,v), \text{PATH}(i-1,u) + w(u,v) \}$$



	0	1	2
A	$\infty$	$\infty$	
B	$\infty$	8	
C	$\infty$		
D	0		
E	$\infty$		
F	$\infty$		
H	$\infty$		

$$\text{PATH}(1,A) = \min \begin{cases} \text{PATH}(0,A) \\ \text{PATH}(0,B) + w(B,A) \\ \text{PATH}(0,C) + w(C,A) \end{cases}$$

$$\text{PATH}(1,B) = \min \begin{cases} \text{PATH}(0,B) \\ \text{PATH}(0,D) + w(D,B) \end{cases}$$

# Bellman-Ford

Bellman-Ford( $G,s$ )

for each  $u$  of  $V$


$$\text{PATH}(0,v) = \infty$$

$$\text{PATH}(0,s) = 0$$

for  $i = 1$  to  $|V| - 1$

for each edge  $(u,v)$  in  $E$

$$\text{PATH}(i,v) = \min \{ \text{PATH}(i-1,v), \\ \text{PATH}(i-1,u) + w(u,v) \}$$



	0	1	2
A	$\infty$	$\infty$	18
B	$\infty$	8	
C	$\infty$	$\infty$	
D	0	0	
E	$\infty$	7	
F	$\infty$	$\infty$	
H	$\infty$	5	

$$\text{PATH}(2,A) = \min \begin{cases} \text{PATH}(1,A) \\ \text{PATH}(1,B) + w(B,A) \\ \text{PATH}(1,C) + w(C,A) \end{cases}$$

# Bellman-Ford

Bellman-Ford( $G,s$ )

for each  $u$  of  $V$


$$\text{PATH}(0,v) = \infty$$

$$\text{PATH}(0,s) = 0$$

for  $i = 1$  to  $|V| - 1$

for each edge  $(u,v)$  in  $E$

$$\text{PATH}(i,v) = \min \{ \text{PATH}(i-1,v), \text{PATH}(i-1,u) + w(u,v) \}$$



	0	1	2
A	$\infty$	$\infty$	18
B	$\infty$	8	
C	$\infty$	$\infty$	
D	0	0	
E	$\infty$	7	
F	$\infty$	$\infty$	4
H	$\infty$	5	

$$\text{PATH}(2,A) = \min \begin{cases} \text{PATH}(1,A) \\ \text{PATH}(1,B) + w(B,A) \\ \text{PATH}(1,C) + w(C,A) \end{cases}$$

$$\text{PATH}(2,F) = \min \begin{cases} \text{PATH}(1,F) \\ \text{PATH}(1,E) + w(E,F) \\ \text{PATH}(1,H) + w(H,F) \end{cases}$$

# Bellman-Ford

Bellman-Ford( $G,s$ )

for each  $u$  of  $V$


$PATH(0,v) = \infty$

$PATH(0,s) = 0$

for  $i = 1$  to  $|V| - 1$

for each edge  $(u,v)$  in  $E$

$PATH(i,v) = \min \{PATH(i-1,v),$   
 $PATH(i-1,u) + w(u,v)\}$



	0	1	2
A	$\infty$	$\infty$	18
B	$\infty$	8	8
C	$\infty$	$\infty$	4
D	0	0	0
E	$\infty$	7	7
F	$\infty$	$\infty$	4
H	$\infty$	5	5

# Bellman-Ford

Bellman-Ford( $G,s$ )

for each  $u$  of  $V$


$PATH(0,v) = \infty$

$PATH(0,s) = 0$

for  $i = 1$  to  $|V| - 1$

    for each edge  $(u,v)$  in  $E$

$PATH(i,v) = \min \{PATH(i-1,v),$   
                                 $PATH(i-1,u) + w(u,v)\}$





# Bellman-Ford

Bellman-Ford( $G, s$ )

for each  $u$  of  $V$

$PATH(0, v) = \infty$

$PATH(0, s) = 0$

for  $i = 1$  to  $|V| - 1$

    for each edge  $(u, v)$  in  $E$

$PATH(i, v) = \min \{PATH(i-1, v),$   
                                 $PATH(i-1, u) + w(u, v)\}$

- the shortest path contains at most  $|V| - 1$  edges.
- more than this creates a cycle  
(use this fact to detect a negative cycle)

# Bellman-Ford

Bellman-Ford( $G,s$ )

for each  $u$  of  $V$


$PATH(0,v) = \infty$

$PATH(0,s) = 0$

for  $i = 1$  to  $|V| - 1$

    for each edge  $(u,v)$  in  $E$

$PATH(i,v) = \min \{PATH(i-1,v),$   
                                 $PATH(i-1,u) + w(u,v)\}$



# Bellman-Ford

Bellman-Ford( $G,s$ )

for each  $u$  of  $V$

$PATH(0,v) = \infty$


$PATH(0,s) = 0$

for  $i = 1$  to  $|V| - 1$

    for each edge  $(u,v)$  in  $E$

$PATH(i,v) = \min \{PATH(i-1,v),$   
                                 $PATH(i-1,u) + w(u,v)\}$

	0	1	2	3
A	0	0	0	0
B	$\infty$	2	2	2
C	$\infty$	$\infty$	5	5
D	$\infty$	$\infty$	$\infty$	6



# Bellman-Ford

Bellman-Ford( $G, s$ )

for each  $u$  of  $V$

$PATH(0, v) = \infty$


$PATH(0, s) = 0$

for  $i = 1$  to  $|V| - 1$

    for each edge  $(u, v)$  in  $E$

$PATH(i, v) = \min \{PATH(i-1, v),$   
                                 $PATH(i-1, u) + w(u, v)\}$

	0	1	2	3		
A	0	0	0	0	0	0
B	$\infty$	2	2	2	1	1
C	$\infty$	$\infty$	5	5	5	4
D	$\infty$	$\infty$	$\infty$	6	6	6



# Bellman-Ford

Bellman-Ford( $G, s$ )

for each  $u$  of  $V$

$PATH(0, v) = \infty$


$PATH(0, s) = 0$

for  $i = 1$  to  $|V| - 1$

    for each edge  $(u, v)$  in  $E$

$PATH(i, v) = \min \{PATH(i-1, v),$   
                                 $PATH(i-1, u) + w(u, v)\}$

	0	1	2	3		
A	0	0	0	0	0	0
B	$\infty$	2	2	2	1	1
C	$\infty$	$\infty$	5	5	5	4
D	$\infty$	$\infty$	$\infty$	6	6	6



negative cycle!

# APSP

- given a weighted graph  $G=(V,E)$  and a source vertex  $s$  in  $V$ , find the shortest paths from each vertex to every other vertex in  $V$

# APSP

- given a weighted graph  $G=(V,E)$  and a source vertex  $s$  in  $V$ , find the shortest paths from each vertex to every other vertex in  $V$
- If it does not contain negative weight edge, run Dijkstra's Algorithm on each vertex
- If it contains negative weight edge, run Bellman-Ford on each vertex

# APSP

- given a weighted graph  $G=(V,E)$  and a source vertex  $s$  in  $V$ , find the shortest paths from each vertex to every other vertex in  $V$
- If it does not contain negative weight edge, run Dijkstra's Algorithm on each vertex  
 $O(|V|.|E|. \log|V|)$   
(for dense graph  $\approx O(|V|^3 \log|V|)$  )
- If it contains negative weight edge, run Bellman-Ford on each vertex  
 $O(|V|^2|E|)$   
(for dense graph  $\approx O(|V|^4)$  )



# APSP

- define a subproblem

Suppose the graph given with adjacency matrix of weight :

$$W = (w_{ij}) \quad \text{such that} \quad w_{ij} = \begin{cases} 0, & \text{if } i = j \\ \infty, & \text{if } i \neq j \text{ and } (i,j) \text{ not in } E \\ w(i,j), & \text{if } i \neq j \text{ and } (i,j) \text{ in } E \end{cases}$$

# APSP

- define a subproblem

$PATH_{ij}^{(m)}$ : the weight of the shortest path from  $i$  to  $j$  that contains  $\leq m$  edges

# APSP

- define a subproblem

$PATH_{ij}^{(m)}$ : the weight of the shortest path from  $i$  to  $j$  that contains  $\leq m$  edges


- construct recurrence relation

# APSP

- define a subproblem

$PATH_{ij}^{(m)}$ : the weight of the shortest path from  $i$  to  $j$  that contains  $\leq m$  edges

- construct recurrence relation




Two Cases

# APSP

- define a subproblem

$PATH_{ij}^{(m)}$ : the weight of the shortest path from  $i$  to  $j$  that contains  $\leq m$  edges

- construct recurrence relation



## Two Cases


1)  $PATH_{ij}^{(m-1)}$

# APSP

- define a subproblem

$PATH_{ij}^{(m)}$ : the weight of the shortest path from  $i$  to  $j$  that contains  $\leq m$  edges

- construct recurrence relation



## Two Cases

1)  $PATH_{ij}^{(m-1)}$


2)  $PATH_{ik}^{(m-1)} + w_{kj}$

# APSP

- define a subproblem

$PATH_{ij}^{(m)}$ : the weight of the shortest path from  $i$  to  $j$  that contains  $\leq m$  edges

- construct recurrence relation



## Two Cases

1)  $PATH_{ij}^{(m-1)}$

2)  $PATH_{ik}^{(m-1)} + w_{kj}$


$$PATH_{ij}^{(m)} = \min \left\{ PATH_{ij}^{(m-1)}, \min_{1 \leq k \leq n} \left\{ PATH_{ik}^{(m-1)} + w_{kj} \right\} \right\}$$

# APSP

- define a subproblem

$PATH_{ij}^{(m)}$ : the weight of the shortest path from  $i$  to  $j$  that contains  $\leq m$  edges

- construct recurrence relation



## Two Cases

1)  $PATH_{ij}^{(m-1)}$

2)  $PATH_{ik}^{(m-1)} + w_{kj}$

$$PATH_{ij}^{(m)} = \min \left\{ PATH_{ij}^{(m-1)}, \min_{1 \leq k \leq n} \left\{ PATH_{ik}^{(m-1)} + w_{kj} \right\} \right\}$$

it's contained in this part




# APSP

- define a subproblem

$PATH_{ij}^{(m)}$ : the weight of the shortest path from  $i$  to  $j$  that contains  $\leq m$  edges

- construct recurrence relation



## Two Cases

1)  $PATH_{ij}^{(m-1)}$

2)  $PATH_{ik}^{(m-1)} + w_{kj}$


$$PATH_{ij}^{(m)} = \min_{1 \leq k \leq n} \{ PATH_{ik}^{(m-1)} + w_{kj} \}$$

# APSP

- define a subproblem

$PATH_{ij}^{(m)}$ : the weight of the shortest path from  $i$  to  $j$  that contains  $\leq m$  edges

- construct recurrence relation



## Two Cases

1)  $PATH_{ij}^{(m-1)}$

2)  $PATH_{ik}^{(m-1)} + w_{kj}$

$$PATH_{ij}^{(m)} = \min_{1 \leq k \leq n} \{ PATH_{ik}^{(m-1)} + w_{kj} \}$$

$$PATH_{ii}^{(0)} = 0$$

$$PATH_{ij}^{(0)} = \infty$$

# APSP

- Let's define a matrix

$$P^{(m)} = PATH_{ij}^{(m)}$$

# APSP

- Let's define a matrix

$$P^{(m)} = PATH_{ij}^{(m)}$$

- $P^{(1)} = W$     since     $w_{ij} = PATH_{ij}^{(1)}$

# APSP

- Let's define a matrix

$$P^{(m)} = PATH_{ij}^{(m)}$$

- $P^{(1)} = W$  since  $w_{ij} = PATH_{ij}^{(1)}$
- Then, we are computing the matrix  $P^{(n-1)}$  since  $PATH_{ij}^{(n-1)} = \delta(i, j)$

# APSP

- Let's define a matrix

$$P^{(m)} = PATH_{ij}^{(m)}$$

- $P^{(1)} = W$  since  $w_{ij} = PATH_{ij}^{(1)}$

- Then, we are computing the matrix  $P^{(n-1)}$  since  $PATH_{ij}^{(n-1)} = \delta(i, j)$

$$\begin{bmatrix} PATH_{11}^{(m-1)} & \cdots & PATH_{1n}^{(m-1)} \\ \vdots & \ddots & \vdots \\ PATH_{n1}^{(m-1)} & \cdots & PATH_{nn}^{(m-1)} \end{bmatrix} \cdot \begin{bmatrix} w_{11} & \cdots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{n1} & \cdots & w_{nn} \end{bmatrix} = \begin{bmatrix} PATH_{11}^{(m)} & \cdots & PATH_{1n}^{(m)} \\ \vdots & \ddots & \vdots \\ PATH_{n1}^{(m)} & \cdots & PATH_{nn}^{(m)} \end{bmatrix}$$

$$PATH_{ij}^{(m)} = PATH_{i1}^{(m-1)} \cdot w_{1j} + PATH_{i2}^{(m-1)} \cdot w_{2j} + \cdots + PATH_{in}^{(m-1)} \cdot w_{nj}$$

# APSP

- Let's define a matrix


$$P^{(m)} = PATH_{ij}^{(m)}$$

- $P^{(1)} = W$  since  $w_{ij} = PATH_{ij}^{(1)}$

- Then, we are computing the matrix  $P^{(n-1)}$  since  $PATH_{ij}^{(n-1)} = \delta(i, j)$

$$\begin{bmatrix} PATH_{11}^{(m-1)} & \cdots & PATH_{1n}^{(m-1)} \\ \vdots & \ddots & \vdots \\ PATH_{n1}^{(m-1)} & \cdots & PATH_{nn}^{(m-1)} \end{bmatrix} \cdot \begin{bmatrix} w_{11} & \cdots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{n1} & \cdots & w_{nn} \end{bmatrix} = \begin{bmatrix} PATH_{11}^{(m)} & \cdots & PATH_{1n}^{(m)} \\ \vdots & \ddots & \vdots \\ PATH_{n1}^{(m)} & \cdots & PATH_{nn}^{(m)} \end{bmatrix}$$

$$PATH_{ij}^{(m)} = PATH_{i1}^{(m-1)} \cdot w_{1j} + PATH_{i2}^{(m-1)} \cdot w_{2j} + \cdots + PATH_{in}^{(m-1)} \cdot w_{nj}$$

$$PATH_{ij}^{(m)} = \min_{1 \leq k \leq n} \{ PATH_{ik}^{(m-1)} + w_{kj} \}$$


# APSP

- The idea is to extend  $P^{(1)}$  using the weight matrix  $W$



# APSP

- The idea is to extend  $P^{(1)}$  using the weight matrix  $W$

$$P^{(1)} = W$$

for  $m = 2$  to  $n-1$

$$P^{(m)} = \text{Extend}(P^{(m-1)}, W, n)$$

return  $P^{(n-1)}$

# APSP


- The idea is to extend  $P^{(1)}$  using the weight matrix  $W$

$$P^{(1)} = W$$

for  $m = 2$  to  $n-1$

$$P^{(m)} = \text{Extend}(P^{(m-1)}, W, n)$$

return  $P^{(n-1)}$



Extend( $P, W, n$ )

initialize  $P'$  as  $n \times n$  matrix

for  $i = 1$  to  $n$

for  $j = 1$  to  $n$

$$PATH'_{ij} = \infty$$

for  $k = 1$  to  $n$

$$PATH'_{ij} = \min\{PATH'_{ij}, PATH_{ij} + w_{ij}\}$$

# APSP

- The idea is to extend  $P^{(1)}$  using the weight matrix  $W$

$$P^{(1)} = W$$

for  $m = 2$  to  $n-1$

$$P^{(m)} = \text{Extend}(P^{(m-1)}, W, n)$$

return  $P^{(n-1)}$

Extend(P,W,n)

initialize  $P'$  as  $n \times n$  matrix

for  $i = 1$  to  $n$

for  $j = 1$  to  $n$

$$PATH'_{ij} = \infty$$

for  $k = 1$  to  $n$

$$PATH'_{ij} = \min\{PATH'_{ij}, PATH_{ij} + w_{ij}\}$$

$O(n^3)$

# APSP

- The idea is to extend  $P^{(1)}$  using the weight matrix  $W$

$$P^{(1)} = W$$

for  $m = 2$  to  $n-1$

$$O(n^4)$$

$$P^{(m)} = \text{Extend}(P^{(m-1)}, W, n)$$

return  $P^{(n-1)}$

Extend(P,W,n)

$$O(n^3)$$

initialize  $P'$  as  $n \times n$  matrix

for  $i = 1$  to  $n$

for  $j = 1$  to  $n$

$$PATH'_{ij} = \infty$$

for  $k = 1$  to  $n$

$$PATH'_{ij} = \min\{PATH'_{ij}, PATH_{ij} + w_{ij}\}$$

# APSP

- The idea is to extend  $P^{(1)}$  using the weight matrix  $W$

$$P^{(1)} = W$$

for  $m = 2$  to  $n-1$

$$P^{(m)} = \text{Extend}(P^{(m-1)}, W, n)$$

return  $P^{(n-1)}$

Extend( $P, W, n$ )

initialize  $P'$  as  $n \times n$  matrix


for  $i = 1$  to  $n$

for  $j = 1$  to  $n$

$$PATH_{ij}' = \infty$$

for  $k = 1$  to  $n$

$$PATH_{ij}' = \min\{PATH_{ij}', PATH_{ik} + w_{kj}\}$$



# APSP

- The idea is to extend  $P^{(1)}$  using the weight matrix  $W$


$$P^{(1)} = W$$

for  $m = 2$  to  $n-1$

$$P^{(m)} = \text{Extend}(P^{(m-1)}, W, n)$$

return  $P^{(n-1)}$

$$P^{(2)} = \text{Extend}(P^{(1)}, W, n)$$



# APSP

- The idea is to extend  $P^{(1)}$  using the weight matrix  $W$


$$P^{(1)} = W$$

for  $m = 2$  to  $n-1$

$$P^{(m)} = \text{Extend}(P^{(m-1)}, W, n)$$

return  $P^{(n-1)}$

$$\begin{aligned} P^{(2)} &= \text{Extend}(P^{(1)}, W, n) \\ &= P^{(1)} * W \end{aligned}$$



# APSP

- The idea is to extend  $P^{(1)}$  using the weight matrix  $W$

$$P^{(1)} = W$$

for  $m = 2$  to  $n-1$


$$P^{(m)} = \text{Extend}(P^{(m-1)}, W, n)$$

return  $P^{(n-1)}$

$$P^{(2)} = \text{Extend}(P^{(1)}, W, n)$$

$$= P^{(1)} * W$$

$$P^{(2)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix}$$





# APSP


- The idea is to extend  $P^{(1)}$  using the weight matrix  $W$

$$P^{(1)} = W$$

for  $m = 2$  to  $n-1$

$$P^{(m)} = \text{Extend}(P^{(m-1)}, W, n)$$

return  $P^{(n-1)}$



$$P^{(2)} = \text{Extend}(P^{(1)}, W, n)$$

$$= P^{(1)} * W$$

$$PATH'_{ij} = \min\{PATH'_{ij}, PATH_{ij} + w_{ij}\}$$

$$P^{(2)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix} = \begin{pmatrix} 0 & & & \end{pmatrix}$$

$$\min\{0+0, 3+\infty, \infty+1, -4+\infty\}$$

# APSP


- The idea is to extend  $P^{(1)}$  using the weight matrix  $W$

$$P^{(1)} = W$$

for  $m = 2$  to  $n-1$

$$P^{(m)} = \text{Extend}(P^{(m-1)}, W, n)$$

return  $P^{(n-1)}$



$$P^{(2)} = \text{Extend}(P^{(1)}, W, n)$$

$$= P^{(1)} * W$$

$$PATH'_{ij} = \min\{PATH'_{ij}, PATH_{ij} + w_{ij}\}$$

$$P^{(2)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 & & \end{pmatrix}$$

$$\min\{0+3, 3+0, \infty+\infty, -4+5\}$$

# APSP

- The idea is to extend  $P^{(1)}$  using the weight matrix  $W$


$$P^{(1)} = W$$

for  $m = 2$  to  $n-1$

$$P^{(m)} = \text{Extend}(P^{(m-1)}, W, n)$$

return  $P^{(n-1)}$

$$P^{(2)} = \text{Extend}(P^{(1)}, W, n) \\ = P^{(1)} * W$$



$$PATH'_{ij} = \min\{PATH'_{ij}, PATH_{ij} + w_{ij}\}$$

$$P^{(2)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 & -1 \\ & & \uparrow \\ & & \end{pmatrix}$$

$$\min\{0+\infty, 3-2, \infty+0, -4+3\}$$

# APSP


- The idea is to extend  $P^{(1)}$  using the weight matrix  $W$

$$P^{(1)} = W$$

for  $m = 2$  to  $n-1$

$$P^{(m)} = \text{Extend}(P^{(m-1)}, W, n)$$

return  $P^{(n-1)}$



$$P^{(2)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix} \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 & -1 & -4 \\ -1 & 0 & -2 & \infty \\ 1 & 4 & 0 & -3 \\ 4 & 5 & 3 & 0 \end{pmatrix}$$

# Floyd-Warshall


- For a given graph  $G$  with  $n$  vertices, all vertices are labeled as  $\{1, 2, 3, \dots, n\}$

# Floyd-Warshall

- For a given graph  $G$  with  $n$  vertices, all vertices are labeled as  $\{1, 2, 3, \dots, n\}$
- For a path  $p = \langle v_1, v_2, \dots, v_m \rangle$ ,  $\{v_2, \dots, v_{m-1}\}$  are called intermediate vertices
- Among all the paths from  $u$  to  $v$ , we consider only the ones in which all intermediate vertices are drawn from a subset  $\{1, 2, \dots, k\}$

# Floyd-Warshall

- For a given graph  $G$  with  $n$  vertices, all vertices are labeled as  $\{1, 2, 3, \dots, n\}$
- For a path  $p = \langle v_1, v_2, \dots, v_m \rangle$ ,  $\{v_2, \dots, v_{m-1}\}$  are called intermediate vertices
- Among all the paths from  $u$  to  $v$ , we consider only the ones in which all intermediate vertices are drawn from a subset  $\{1, 2, \dots, k\}$




1  $\rightsquigarrow$  5

intermediate vertices  
are drawn from  $\{1, 2, 3\}$

# Floyd-Warshall

- For a given graph  $G$  with  $n$  vertices, all vertices are labeled as  $\{1, 2, 3, \dots, n\}$
- For a path  $p = \langle v_1, v_2, \dots, v_m \rangle$ ,  $\{v_2, \dots, v_{m-1}\}$  are called intermediate vertices
- Among all the paths from  $u$  to  $v$ , we consider only the ones in which all intermediate vertices are drawn from a subset  $\{1, 2, \dots, k\}$




1  $\rightsquigarrow$  5

intermediate vertices  
are drawn from  $\{1, 2, 3\}$



# Floyd-Warshall

- For a given graph  $G$  with  $n$  vertices, all vertices are labeled as  $\{1, 2, 3, \dots, n\}$
- For a path  $p = \langle v_1, v_2, \dots, v_m \rangle$ ,  $\{v_2, \dots, v_{m-1}\}$  are called intermediate vertices
- Among all the paths from  $u$  to  $v$ , we consider only the ones in which all intermediate vertices are drawn from a subset  $\{1, 2, \dots, k\}$




1  $\rightsquigarrow$  5

intermediate vertices  
are drawn from  $\{1, 2, 3\}$

# Floyd-Warshall

- For a given graph  $G$  with  $n$  vertices, all vertices are labeled as  $\{1, 2, 3, \dots, n\}$
- For a path  $p = \langle v_1, v_2, \dots, v_m \rangle$ ,  $\{v_2, \dots, v_{m-1}\}$  are called intermediate vertices
- Among all the paths from  $u$  to  $v$ , we consider only the ones in which all intermediate vertices are drawn from a subset  $\{1, 2, \dots, k\}$




1  $\rightsquigarrow$  5

intermediate vertices  
are drawn from  $\{1, 2, 3\}$

# Floyd-Warshall

- For a given graph  $G$  with  $n$  vertices, all vertices are labeled as  $\{1, 2, 3, \dots, n\}$
- For a path  $p = \langle v_1, v_2, \dots, v_m \rangle$ ,  $\{v_2, \dots, v_{m-1}\}$  are called intermediate vertices
- Among all the paths from  $u$  to  $v$ , we consider only the ones in which all intermediate vertices are drawn from a subset  $\{1, 2, \dots, k\}$



1  $\rightsquigarrow$  5

intermediate vertices  
are drawn from  $\{1, 2, 3\}$

negative-weight  
cycle not allowed

# Floyd-Warshall


- define a subproblem

$d_{ij}^{(k)}$  : the weight of a shortest path from vertex  $i$  to vertex  $j$   
such that all intermediate vertices drawn from  $\{1, 2, \dots, k\}$

# Floyd-Warshall

- define a subproblem


$d_{ij}^{(k)}$  : the weight of a shortest path from vertex  $i$  to vertex  $j$   
such that all intermediate vertices drawn from  $\{1, 2, \dots, k\}$



# Floyd-Warshall

- define a subproblem

$d_{ij}^{(k)}$  : the weight of a shortest path from vertex  $i$  to vertex  $j$   
such that all intermediate vertices drawn from  $\{1, 2, \dots, k\}$




$$d_{56}^{(0)} = \infty$$

# Floyd-Warshall

- define a subproblem

$d_{ij}^{(k)}$  : the weight of a shortest path from vertex  $i$  to vertex  $j$   
such that all intermediate vertices drawn from  $\{1, 2, \dots, k\}$




$$d_{56}^{(0)} = \infty$$

$$d_{56}^{(1)} = 13$$

# Floyd-Warshall

- define a subproblem

$d_{ij}^{(k)}$  : the weight of a shortest path from vertex  $i$  to vertex  $j$   
such that all intermediate vertices drawn from  $\{1, 2, \dots, k\}$



$$d_{56}^{(0)} = \infty$$

$$d_{56}^{(1)} = 13$$


$$d_{56}^{(2)} = 9$$



# Floyd-Warshall

- define a subproblem

$d_{ij}^{(k)}$  : the weight of a shortest path from vertex  $i$  to vertex  $j$   
such that all intermediate vertices drawn from  $\{1, 2, \dots, k\}$



$$d_{56}^{(0)} = \infty$$

$$d_{56}^{(1)} = 13$$


$$d_{56}^{(2)} = 9$$

$$d_{56}^{(3)} = 8$$

# Floyd-Warshall

- define a subproblem

$d_{ij}^{(k)}$  : the weight of a shortest path from vertex  $i$  to vertex  $j$   
such that all intermediate vertices drawn from  $\{1, 2, \dots, k\}$




$$\begin{aligned}d_{56}^{(0)} &= \infty \\d_{56}^{(1)} &= 13 \\d_{56}^{(2)} &= 9 \\d_{56}^{(3)} &= 8 \\d_{56}^{(4)} &= 6\end{aligned}$$

# Floyd-Warshall

- define a subproblem

$d_{ij}^{(k)}$  : the weight of a shortest path from vertex  $i$  to vertex  $j$   
such that all intermediate vertices drawn from  $\{1, 2, \dots, k\}$

- construct recurrence relation




Two Cases

# Floyd-Warshall

- define a subproblem

$d_{ij}^{(k)}$  : the weight of a shortest path from vertex  $i$  to vertex  $j$   
such that all intermediate vertices drawn from  $\{1, 2, \dots, k\}$

- construct recurrence relation



## Two Cases

$$1) d_{ij}^{(k)} = d_{ij}^{(k-1)}$$


- $k$  is not intermediate vertex in the path  
(the intermediate vertices are drawn  
from  $\{1, 2, \dots, k-1\}$  )

# Floyd-Warshall

- define a subproblem

$d_{ij}^{(k)}$  : the weight of a shortest path from vertex  $i$  to vertex  $j$   
such that all intermediate vertices drawn from  $\{1, 2, \dots, k\}$

- construct recurrence relation



## Two Cases

$$1) d_{ij}^{(k)} = d_{ij}^{(k-1)}$$

$$2) d_{ij}^{(k)} = d_{ik}^{(k-1)} + d_{kj}^{(k-1)}$$


- $k$  is an intermediate vertex in the path  
(the intermediate vertices in the smaller  
paths are drawn from  $\{1, 2, \dots, k-1\}$  )

# Floyd-Warshall

- define a subproblem

$d_{ij}^{(k)}$  : the weight of a shortest path from vertex  $i$  to vertex  $j$   
such that all intermediate vertices drawn from  $\{1, 2, \dots, k\}$

- construct recurrence relation



## Two Cases

$$1) d_{ij}^{(k)} = d_{ij}^{(k-1)}$$

$$2) d_{ij}^{(k)} = d_{ik}^{(k-1)} + d_{kj}^{(k-1)}$$

- $k$  is an intermediate vertex in the path  
(the intermediate vertices in the smaller paths are drawn from  $\{1, 2, \dots, k-1\}$ )

$$d_{ij}^{(k)} = \begin{cases} w_{ij} & , \quad \text{if } k = 0 \\ \min \{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \} & , \quad \text{if } k \geq 1 \end{cases}$$

# Floyd-Warshall

## Floyd-Warshall

$D^{(0)} = W$

for  $k = 1$  to  $n$

  let  $D^{(k)}$  be a new  $n \times n$  matrix

  for  $i = 1$  to  $n$

    for  $j = 1$  to  $n$

$$d_{ij}^{(k)} = \min \left\{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \right\}$$

# Floyd-Warshall

## Floyd-Warshall

$D^{(0)} = W$


for  $k = 1$  to  $n$

  let  $D^{(k)}$  be a new  $n \times n$  matrix

  for  $i = 1$  to  $n$

    for  $j = 1$  to  $n$

$$d_{ij}^{(k)} = \min \left\{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \right\}$$





# Floyd-Warshall

## Floyd-Warshall

$$D^{(0)} = W$$


for k = 1 to n

  let  $D^{(k)}$  be a new  $n \times n$  matrix

  for i = 1 to n

    for j = 1 to n

$$d_{ij}^{(k)} = \min \{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \}$$



$$D^{(0)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix}$$

# Floyd-Warshall

## Floyd-Warshall

$$D^{(0)} = W$$


for k = 1 to n

  let  $D^{(k)}$  be a new  $n \times n$  matrix

  for i = 1 to n

    for j = 1 to n

$$d_{ij}^{(k)} = \min \{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \}$$



$$D^{(0)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix} \quad \longrightarrow \quad D^{(1)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & & & \end{pmatrix}$$

# Floyd-Warshall

## Floyd-Warshall

$$D^{(0)} = W$$


for k = 1 to n

  let  $D^{(k)}$  be a new  $n \times n$  matrix

  for i = 1 to n

    for j = 1 to n

$$d_{ij}^{(k)} = \min \{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \}$$



$$D^{(0)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix}$$



$$D^{(1)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & 0 & -2 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix}$$



$$d_{32}^{(1)} = \min \{ d_{32}^{(0)}, d_{31}^{(0)} + d_{12}^{(0)} \}$$

# Floyd-Warshall

## Floyd-Warshall

$$D^{(0)} = W$$


for k = 1 to n

let  $D^{(k)}$  be a new  $n \times n$  matrix

for i = 1 to n

for j = 1 to n

$$d_{ij}^{(k)} = \min \{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \}$$



$$D^{(0)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix}$$

$$D^{(1)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & 4 & & \end{pmatrix}$$

$$d_{32}^{(1)} = \min \{ d_{32}^{(0)}, d_{31}^{(0)} + d_{12}^{(0)} \}$$

$$d_{32}^{(1)} = \min \{ \infty, 1 + 3 \} = 4$$

# Floyd-Warshall

## Floyd-Warshall

$$D^{(0)} = W$$


for k = 1 to n

  let  $D^{(k)}$  be a new  $n \times n$  matrix

  for i = 1 to n

    for j = 1 to n

$$d_{ij}^{(k)} = \min \{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \}$$



$$D^{(0)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix} \quad \longrightarrow \quad D^{(1)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & 4 & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix}$$

# Floyd-Warshall

## Floyd-Warshall

$$D^{(0)} = W$$


for k = 1 to n

let  $D^{(k)}$  be a new  $n \times n$  matrix

for i = 1 to n

for j = 1 to n

$$d_{ij}^{(k)} = \min \{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \}$$



$$D^{(0)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix}$$



$$D^{(1)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & 4 & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix}$$



$$d_{34}^{(1)} = \min \{ d_{34}^{(0)}, d_{31}^{(0)} + d_{14}^{(0)} \}$$

# Floyd-Warshall

## Floyd-Warshall

$$D^{(0)} = W$$


for k = 1 to n

let  $D^{(k)}$  be a new  $n \times n$  matrix

for i = 1 to n

for j = 1 to n

$$d_{ij}^{(k)} = \min \{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \}$$



$$D^{(0)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix}$$



$$D^{(1)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & 4 & 0 & -3 \\ \infty & 5 & 3 & 0 \end{pmatrix}$$



$$d_{34}^{(1)} = \min \{ d_{34}^{(0)}, d_{31}^{(0)} + d_{14}^{(0)} \}$$

$$d_{34}^{(1)} = \min \{ \infty, 1 - 4 \} = -3$$

# Floyd-Warshall

## Floyd-Warshall

$$D^{(0)} = W$$

for k = 1 to n

let  $D^{(k)}$  be a new  $n \times n$  matrix


for i = 1 to n

for j = 1 to n

$$d_{ij}^{(k)} = \min \{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \}$$

$$D^{(0)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix}$$

$$D^{(1)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & 4 & 0 & -3 \\ \infty & 5 & 3 & 0 \end{pmatrix}$$





# Floyd-Warshall

## Floyd-Warshall

$$D^{(0)} = W$$


for k = 1 to n

let  $D^{(k)}$  be a new  $n \times n$  matrix

for i = 1 to n

for j = 1 to n

$$d_{ij}^{(k)} = \min \{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \}$$



$$D^{(0)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix} \rightarrow D^{(1)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & 4 & 0 & -3 \\ \infty & 5 & 3 & 0 \end{pmatrix} \rightarrow D^{(2)} = \begin{pmatrix} 0 & 3 & 1 & -4 \\ \infty & 0 & -2 & \infty \\ 1 & 4 & 0 & -3 \\ \infty & 5 & 3 & 0 \end{pmatrix}$$

# Floyd-Warshall

## Floyd-Warshall

$$D^{(0)} = W$$


for k = 1 to n

let  $D^{(k)}$  be a new  $n \times n$  matrix

for i = 1 to n

for j = 1 to n

$$d_{ij}^{(k)} = \min \{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \}$$



$$D^{(0)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix}$$

$$D^{(1)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & 4 & 0 & -3 \\ \infty & 5 & 3 & 0 \end{pmatrix}$$

$$D^{(2)} = \begin{pmatrix} 0 & 3 & 1 & -4 \\ \infty & 0 & -2 & \infty \\ 1 & 4 & 0 & -3 \\ \infty & 5 & 3 & 0 \end{pmatrix}$$

$$D^{(3)} = \begin{pmatrix} 0 & 3 & 1 & -4 \\ -1 & 0 & -2 & -5 \\ 1 & 4 & 0 & -3 \\ 4 & 5 & 3 & 0 \end{pmatrix}$$

# Floyd-Warshall

## Floyd-Warshall

$$D^{(0)} = W$$


for  $k = 1$  to  $n$

let  $D^{(k)}$  be a new  $n \times n$  matrix

for  $i = 1$  to  $n$

for  $j = 1$  to  $n$

$$d_{ij}^{(k)} = \min \{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \}$$



$$D^{(0)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & \infty & 0 & \infty \\ \infty & 5 & 3 & 0 \end{pmatrix}$$

$$D^{(1)} = \begin{pmatrix} 0 & 3 & \infty & -4 \\ \infty & 0 & -2 & \infty \\ 1 & 4 & 0 & -3 \\ \infty & 5 & 3 & 0 \end{pmatrix}$$

$$D^{(2)} = \begin{pmatrix} 0 & 3 & 1 & -4 \\ \infty & 0 & -2 & \infty \\ 1 & 4 & 0 & -3 \\ \infty & 5 & 3 & 0 \end{pmatrix}$$

$$D^{(4)} = \begin{pmatrix} 0 & 1 & -1 & -4 \\ -1 & 0 & -2 & -5 \\ 1 & 2 & 0 & -3 \\ 4 & 5 & 3 & 0 \end{pmatrix}$$

$$D^{(3)} = \begin{pmatrix} 0 & 3 & 1 & -4 \\ -1 & 0 & -2 & -5 \\ 1 & 4 & 0 & -3 \\ 4 & 5 & 3 & 0 \end{pmatrix}$$