

## Level 1 BLAS

	dim	scalar	vector	vector	scalars	5-element array		prefixes
SUBROUTINE xROTG (					A, B, C, S )		Generate plane rotation	S, D
SUBROUTINE xROTMG(					D1, D2, A, B,	PARAM )	Generate modified plane rotation	S, D
SUBROUTINE xROT ( N,			X, INCX, Y, INCY,		C, S )		Apply plane rotation	S, D
SUBROUTINE xROTM ( N,			X, INCX, Y, INCY,			PARAM )	Apply modified plane rotation	S, D
SUBROUTINE xSWAP ( N,			X, INCX, Y, INCY )				$x \leftrightarrow y$	S, D, C, Z
SUBROUTINE xSCAL ( N,	ALPHA,		X, INCX )				$x \leftarrow \alpha x$	S, D, C, Z, CS, ZD
SUBROUTINE xCOPY ( N,			X, INCX, Y, INCY )				$y \leftarrow x$	S, D, C, Z
SUBROUTINE xAXPY ( N,	ALPHA,		X, INCX, Y, INCY )				$y \leftarrow \alpha x + y$	S, D, C, Z
FUNCTION xDOT ( N,			X, INCX, Y, INCY )				$dot \leftarrow x^T y$	S, D, DS
FUNCTION xDOTU ( N,			X, INCX, Y, INCY )				$dot \leftarrow x^T y$	C, Z
FUNCTION xDOTC ( N,			X, INCX, Y, INCY )				$dot \leftarrow x^H y$	C, Z
FUNCTION xxDOT ( N,			X, INCX, Y, INCY )				$dot \leftarrow \alpha + x^T y$	SDS
FUNCTION xNRM2 ( N,			X, INCX )				$nrm2 \leftarrow   x  _2$	S, D, SC, DZ
FUNCTION xASUM ( N,			X, INCX )				$asum \leftarrow   re(x)  _1 +   im(x)  _1$	S, D, SC, DZ
FUNCTION IxAMAX( N,			X, INCX )				$amax \leftarrow 1^{st} k \ni  re(x_k)  +  im(x_k) $ $= \max( re(x_i)  +  im(x_i) )$	S, D, C, Z

## Level 2 BLAS

	options	dim	b-width	scalar	matrix	vector	scalar	vector	
xGEMV (	TRANS,	M, N,		ALPHA, A, LDA,	X, INCX,	BETA, Y, INCY )			S, D, C, Z
xGBMV (	TRANS,	M, N, KL, KU,		ALPHA, A, LDA,	X, INCX,	BETA, Y, INCY )			S, D, C, Z
xHEMV (	UPL0,	N,		ALPHA, A, LDA,	X, INCX,	BETA, Y, INCY )			C, Z
xHBMV (	UPL0,	N, K,		ALPHA, A, LDA,	X, INCX,	BETA, Y, INCY )			C, Z
xHPMV (	UPL0,	N,		ALPHA, AP,	X, INCX,	BETA, Y, INCY )			C, Z
xSYMV (	UPL0,	N,		ALPHA, A, LDA,	X, INCX,	BETA, Y, INCY )			S, D
xSBMV (	UPL0,	N, K,		ALPHA, A, LDA,	X, INCX,	BETA, Y, INCY )			S, D
xSPMV (	UPL0,	N,		ALPHA, AP,	X, INCX,	BETA, Y, INCY )			S, D
xTRMV (	UPL0, TRANS, DIAG,	N,		A, LDA,	X, INCX )				S, D, C, Z
xTBMV (	UPL0, TRANS, DIAG,	N, K,		A, LDA,	X, INCX )				S, D, C, Z
xTPMV (	UPL0, TRANS, DIAG,	N,		AP,	X, INCX )				S, D, C, Z
xTRSV (	UPL0, TRANS, DIAG,	N,		A, LDA,	X, INCX )				S, D, C, Z
xTBSV (	UPL0, TRANS, DIAG,	N, K,		A, LDA,	X, INCX )				S, D, C, Z
xTPSV (	UPL0, TRANS, DIAG,	N,		AP,	X, INCX )				S, D, C, Z
	options	dim	scalar	vector	vector	matrix			
xGER (		M, N,	ALPHA, X, INCX, Y, INCY, A, LDA )						S, D
xGERU (		M, N,	ALPHA, X, INCX, Y, INCY, A, LDA )						C, Z
xGERC (		M, N,	ALPHA, X, INCX, Y, INCY, A, LDA )						C, Z
xHER (	UPL0,	N,	ALPHA, X, INCX,		A, LDA )				C, Z
xHPR (	UPL0,	N,	ALPHA, X, INCX,		AP )				C, Z
xHER2 (	UPL0,	N,	ALPHA, X, INCX, Y, INCY, A, LDA )						C, Z
xHPR2 (	UPL0,	N,	ALPHA, X, INCX, Y, INCY, AP )						C, Z
xSYR (	UPL0,	N,	ALPHA, X, INCX,		A, LDA )				S, D
xSPR (	UPL0,	N,	ALPHA, X, INCX,		AP )				S, D
xSYR2 (	UPL0,	N,	ALPHA, X, INCX, Y, INCY, A, LDA )						S, D
xSPR2 (	UPL0,	N,	ALPHA, X, INCX, Y, INCY, AP )						S, D

## Level 3 BLAS

	options		dim	scalar	matrix	matrix	scalar	matrix		
xGEMM (	TRANSA, TRANSB,		M, N, K,	ALPHA, A, LDA,	B, LDB,	BETA, C, LDC )	$C \leftarrow \alpha op(A)op(B) + \beta C, op(X) = X, X^T, X^H, C - m \times n$			S, D, C, Z
xSYMM (	SIDE, UPLO,		M, N,	ALPHA, A, LDA,	B, LDB,	BETA, C, LDC )	$C \leftarrow \alpha AB + \beta C, C \leftarrow \alpha BA + \beta C, C - m \times n, A = A^T$			S, D, C, Z
xHEMM (	SIDE, UPLO,		M, N,	ALPHA, A, LDA,	B, LDB,	BETA, C, LDC )	$C \leftarrow \alpha AB + \beta C, C \leftarrow \alpha BA + \beta C, C - m \times n, A = A^H$			C, Z
xSYRK (	UPLO, TRANS,		N, K,	ALPHA, A, LDA,		BETA, C, LDC )	$C \leftarrow \alpha AA^T + \beta C, C \leftarrow \alpha A^T A + \beta C, C - n \times n$			S, D, C, Z
xHERK (	UPLO, TRANS,		N, K,	ALPHA, A, LDA,		BETA, C, LDC )	$C \leftarrow \alpha AA^H + \beta C, C \leftarrow \alpha A^H A + \beta C, C - n \times n$			C, Z
xSYR2K(	UPLO, TRANS,		N, K,	ALPHA, A, LDA,	B, LDB,	BETA, C, LDC )	$C \leftarrow \alpha AB^T + \bar{\alpha} BA^T + \beta C, C \leftarrow \alpha A^T B + \bar{\alpha} B^T A + \beta C, C - n \times n$			S, D, C, Z
xHER2K(	UPLO, TRANS,		N, K,	ALPHA, A, LDA,	B, LDB,	BETA, C, LDC )	$C \leftarrow \alpha AB^H + \bar{\alpha} BA^H + \beta C, C \leftarrow \alpha A^H B + \bar{\alpha} B^H A + \beta C, C - n \times n$			C, Z
xTRMM (	SIDE, UPLO, TRANSA,	DIAG, M, N,		ALPHA, A, LDA,	B, LDB )		$B \leftarrow \alpha op(A)B, B \leftarrow \alpha Bop(A), op(A) = A, A^T, A^H, B - m \times n$			S, D, C, Z
xTRSM (	SIDE, UPLO, TRANSA,	DIAG, M, N,		ALPHA, A, LDA,	B, LDB )		$B \leftarrow \alpha op(A^{-1})B, B \leftarrow \alpha Bop(A^{-1}), op(A) = A, A^T, A^H, B - m \times n$			S, D, C, Z

### Meaning of prefixes

S - REAL	C - COMPLEX
D - DOUBLE PRECISION	Z - COMPLEX*16

(this may not be supported by all machines)

For the Level 2 BLAS a set of extended-precision routines with the prefixes ES, ED, EC, EZ may also be available.

### Level 1 BLAS

In addition to the listed routines there are two further extended-precision dot product routines DQDOTI and DQDOTA.

### Level 2 and Level 3 BLAS

Matrix types:

GE - GEneral	GB - General Band	
SY - SYmmetric	SB - Sym. Band	SP - Sum. Packed
HE - HErmitian	HB - Herm. Band	HP - Herm. Packed
TR - TRiangular	TB - Triang. Band	TP - Triang. Packed

### Level 2 and Level 3 BLAS Options

Dummy options arguments are declared as CHARACTER\*1 and may be passed as character strings.

TRANx	= 'No transpose', 'Transpose', 'Conjugate transpose' ( $X, X^T, X^H$ )
UPLO	= 'Upper triangular', 'Lower triangular'
DIAG	= 'Non-unit triangular', 'Unit triangular'
SIDE	= 'Left', 'Right' (A or op(A) on the left, or A or op(A) on the right)

For real matrices, TRANSx = 'T' and TRANSx = 'C' have the same meaning.

For Hermitian matrices, TRANSx = 'T' is not allowed.

For complex symmetric matrices, TRANSx = 'H' is not allowed.

### References

C. Lawson, R. Hanson, D. Kincaid, and F. Krogh, "Basic Linear Algebra Subprograms for Fortran Usage," *ACM Trans. on Math. Soft.* 5 (1979) 308-325

J.J. Dongarra, J. DuCroz, S. Hammarling, and R. Hanson, "An Extended Set of Fortran Basic Linear Algebra Subprograms," *ACM Trans. on Math. Soft.* 14,1 (1988) 1-32

J.J. Dongarra, I. Duff, J. DuCroz, and S. Hammarling, "A Set of Level 3 Basic Linear Algebra Subprograms," *ACM Trans. on Math. Soft.* (1989)

### Obtaining the Software via [netlib@ornl.gov](mailto:netlib@ornl.gov)

To receive a copy of the single-precision software, type in a mail message:

```
send sblas from blas
send sblas2 from blas
send sblas3 from blas
```

To receive a copy of the double-precision software, type in a mail message:

```
send dblas from blas
send dblas2 from blas
send dblas3 from blas
```

To receive a copy of the complex single-precision software, type in a mail message:

```
send cblas from blas
send cblas2 from blas
send cblas3 from blas
```

To receive a copy of the complex double-precision software, type in a mail message:

```
send zblas from blas
send zblas2 from blas
send zblas3 from blas
```

Send comments and questions to [lapack@cs.utk.edu](mailto:lapack@cs.utk.edu).

# Basic

# Linear

# Algebra

# Subprograms

# A Quick Reference Guide

University of Tennessee  
Oak Ridge National Laboratory  
Numerical Algorithms Group Ltd.

May 11, 1997