

Double Spacetime Theory: Gravity as Torsion between Usual and Dual Spacetime

<https://github.com/hypernumbernet>

February 8, 2026

Abstract

General Relativity (GR) describes gravity as the curvature of a single spacetime continuum. Despite its empirical success, the theory encounters profound difficulties when confronted with quantum mechanics and the nature of singularities. This paper proposes a radical reformulation: gravity is not curvature of a continuous manifold, but torsion arising from the relative misalignment between two compact spacetimes intrinsically attached to every massive particle.

Employing the 16-real-dimensional biquaternion algebra isomorphic to $\text{Cl}(3,1)$, we define the usual spacetime with basis (j, kI, kJ, kK) and the dual spacetime with basis (k, jI, jJ, jK) . The dual map $X \mapsto Xi$ reverses the arrow of time while preserving the Minkowski norm $X^2 = X'^2$. Lorentz transformations are generated by commuting pairs of rotors: the usual rotor $R_{\text{usual}} = \exp(\frac{\theta}{2} \hat{p})$ and the effective dual rotor $R_{\text{dual}}^{\text{eff}} = \cos \frac{\theta}{2} - \hat{p} \sin \frac{\theta}{2}$, where the negative sign reflects the time-reversed duality.

The torsional mismatch rotor is defined symmetrically as $\Omega = R_{\text{usual}} R_{\text{dual}}^{\text{eff}}$, yielding the torsion bivector $\Omega_{\text{biv}} = \log \Omega$. The unique Lorentz-invariant scalar is the Killing form $J = \frac{1}{16} B(\Omega_{\text{biv}}, \Omega_{\text{biv}})$.

The action $S = \frac{c^4}{16\pi G} \int J d^4x$ is shown to be dynamically equivalent to the Einstein-Hilbert action, reproducing GR exactly while eliminating Christoffel symbols, Riemann curvature, and the continuum spacetime hypothesis altogether. The theory is the biquaternionic realization of the Teleparallel Equivalent of GR (TEGR), with torsion now physically interpreted as dual-spacetime torsion.

Crucially, since J depends only on the relative angle between particle-intrinsic dual rotors, coherent excitation of the dual-spacetime components θ_a offers a direct pathway to gravitational engineering: gravity shielding, inertial mass reduction, and even antigravity become, for the first time, questions of rotor synchronization rather than violations of fundamental law.

GR is not superseded—it is completed, and gravity is rendered controllable in principle.

1 Introduction

The theory of GR, since its inception in 1915, has stood as one of the most successful descriptions of gravitation in physics. Yet, despite its empirical triumphs—from the perihelion precession of Mercury to gravitational waves and black hole imaging—a profound philosophical unease persists: *why* does matter curve spacetime, and *why* does curved spacetime, in turn, dictate the motion of matter? The Einstein field equations tell us *how* gravity works with exquisite precision, but they offer no deeper *reason* for the origin of curvature itself. This explanatory gap becomes particularly acute when confronting Mach's principle: the inertia of a body should be determined by the totality of matter in the universe, yet GR treats spacetime as an independent, continuous entity that can exist even in complete emptiness.

A century of attempts to resolve these issues—loop quantum gravity, string theory, emergent gravity—has yielded mathematical elegance but no consensus on the microscopic origin of spacetime or the physical mechanism that makes mass-energy *generate* curvature. All these approaches retain the continuum hypothesis: spacetime is a smooth, differentiable manifold *a priori*, and gravity is effectuated through the metric's deviation from flatness via Christoffel symbols and the Riemann tensor.

The present work breaks radically with this tradition.

We propose that the spacetime continuum hypothesis is *fundamentally incorrect*. There is no single, shared spacetime manifold. Instead, *each individual particle carries its own pair of intrinsically double spacetimes*—a compactified 16-real-dimensional structure encoded in the biquaternion algebra isomorphic to the Clifford algebra $\text{Cl}(3,1)$. The usual spacetime (with basis j, kI, kJ, kK) and its dual (with basis k, jI, jJ, jK) are related by right-multiplication by i , a transformation that reverses the arrow of time while preserving the Minkowski norm. Gravity and inertia arise *not* from curvature of a background manifold, but from the *torsional mismatch angle* between these two particle-intrinsic spacetimes when parallel transport is demanded across different particles.

This philosophical shift is profound. In standard GR, spacetime is an autonomous entity that *tells matter how to move*, while matter *tells spacetime how to curve* (Wheeler's famous phrase). In the double spacetime theory, there is *no autonomous spacetime at all*. Spacetime degrees of freedom are demoted to internal attributes of particles themselves—

much like spin or charge—and the apparent curvature of the macroscopic world emerges only as a collective effect of the torsional misalignment among the dual rotors of neighboring particles. The Einstein equations are recovered exactly, yet Christoffel symbols, covariant derivatives, and the Riemann tensor are entirely eliminated: they are replaced by simple algebraic operations (rotor exponentiation, logarithm, and the Killing form trace) on biquaternions.

The theory therefore achieves three revolutionary outcomes simultaneously:

1. *Exact equivalence with GR* in all observable predictions, including the Newtonian limit, black holes, cosmology, and gravitational waves—yet without any geometric curvature in the traditional sense.
2. *Natural unification of gravity and inertia*: the equivalence principle emerges as the requirement that the usual and dual rotors remain parallel in free fall; acceleration violates this parallelism, producing inertial forces as the *same torsional phenomenon* as gravity.
3. *In-principle controllability of gravity*: because the scalar invariant J depends only on the *relative angle* between particle-intrinsic dual rotors, external fields capable of coherently exciting the dual-spacetime components θ_a can reduce, cancel, or even reverse gravitational interaction. Gravity ceases to be an inviolable fundamental force and becomes an *engineerable degree of freedom*.

This is not a modification of GR; it is its completion. The continuum was a useful fiction—beautiful, but ultimately unnecessary. Once spacetime is recognized as a paired, particle-local structure, the century-old mystery of why matter curves spacetime dissolves: matter does not curve spacetime; it *misaligns* its own double spacetimes, and this misalignment is what we perceive as gravity.

The remainder of this paper is organized as follows.

Section 2 establishes the mathematical foundation by introducing the biquaternion algebra isomorphic to $\text{Cl}(3,1)$ and constructing the dual spacetime pair intrinsic to every particle.

Section 3 develops the complete rotor formalism that unifies Lorentz boosts, rotations, and parallel transport without matrices or Christoffel symbols.

Section 4 demonstrates that the torsional mismatch scalar J , constructed solely from the relative rotor $\Omega = R_{\text{usual}}^\dagger R_{\text{dual}}$, yields an action dynamically equivalent to the Einstein-Hilbert action, revealing the theory as the biquaternionic realization of Teleparallel Gravity.

Section 5 shows that inertia and gravity are identical phenomena arising from the same dual-rotor rigidity, with the equivalence principle emerging as an algebraic identity.

Section 6 eliminates the need for dark matter by deriving flat galactic rotation curves from baryons alone.

Section 7 reinterprets stellar remnants as torsion stars with multi-layered attractive-repulsive structure, forbidding both singularities and event horizons.

Section 8 derives time dilation effects from torsional mismatch, predicting measurable deviations from GR in strong gravitational fields and planetary interiors.

Section 9 proves scale invariance of the torsional layers, identifying the nuclear force as gravity with layers, and atomic nuclei as primordial torsion stars.

Section 10 models the electron shell as a gravity inversion layer, explaining electron orbits and interatomic forces from the perspective of gravity on a microscopic scale.

Section 11 outlines the unification of all fundamental forces within the biquaternion algebra, with gauge fields emerging as additional rotor components.

Section 12 details practical mechanisms for gravitational engineering via coherent excitation of dual-spacetime rotors using circularly polarized electromagnetic fields.

Section 13 proposes a mechanism for baryon asymmetry based on dual-spacetime torsion dynamics in the early universe.

Section 14 introduces a discrete model of dual spacetime, where the torsional layers are quantized and bounded, leading to a finite unit group in the corresponding integer ring and enabling rigorous Diophantine constraints.

Section 15 develops the dynamics of dual rotors under external fields, deriving equations of motion and stability criteria for torsional configurations.

Section 16 extends the theory to quantum gravity, identifying quantum states with dual-spacetime rotor phases and deriving uncertainty relations from torsional coherence conditions.

Section 17 summarizes the unification of all forces within the 16-dimensional biquaternion algebra and outlines near-term experimental tests that can confirm or falsify the theory within this decade.

With this reformulation, GR is liberated from the prison of the continuum and elevated to a theory in which gravity is, for the first time, *controllable in principle*.

2 Mathematical Foundation: The 16-Dimensional Biquaternion Algebra

The entire double spacetime theory is constructed within the 16-real-dimensional algebra of *biquaternions* (also known as double quaternions), which is canonically isomorphic to the Clifford geometric algebra $\text{Cl}(3,1)$ over Minkowski spacetime with signature $(-, +, +, +)$. This algebra naturally accommodates *two complete, mutually commuting copies* of Minkowski spacetime — the usual and the dual — within a single algebraic structure intrinsically attached to every massive particle.

2.1 The Biquaternion Algebra

We begin with two independent copies of Hamilton's quaternion algebra:

- Primary quaternions generated by i, j, k obeying $i^2 = j^2 = k^2 = -1$, $ij = k$, $ji = -k$ (cyclic),
- Secondary quaternions generated by I, J, K obeying identical rules $I^2 = J^2 = K^2 = -1$, $IJ = K$, $JI = -K$ (cyclic).

The full biquaternion algebra is their tensor product $\mathbb{H} \oplus \mathbb{H}$, in which the two sets strictly commute:

$$iI = Ii, iJ = Ji, iK = Ki, jI = Ij, \dots$$

The resulting 16 linearly independent real basis elements are

$$1, i, j, k, I, J, K, iI, iJ, iK, jI, jJ, jK, kI, kJ, kK.$$

2.2 Explicit Isomorphism with $\text{Cl}(3,1)$

The algebra is isomorphic to $\text{Cl}(3,1)$ via the following identification of grade-1 (vector) generators:

$$\begin{aligned} e_0 &= j, & (\text{time-like basis vector}) \\ e_1 &= kI, \quad e_2 = kJ, \quad e_3 = kK & (\text{space-like basis vectors}), \end{aligned}$$

which immediately satisfy the defining relations

$$e_0^2 = -1, \quad e_1^2 = e_2^2 = e_3^2 = +1, \quad \{e_\mu, e_\nu\} = 0 \quad (\mu \neq \nu).$$

The higher-grade elements then follow standard Clifford multiplication. In particular:

$$\begin{aligned} \text{grade-2 (bivectors):} \quad & e_0e_1 = iI, \quad e_0e_2 = iJ, \quad e_0e_3 = iK, \\ & e_3e_2 = I, \quad e_1e_3 = J, \quad e_2e_1 = K, \\ \text{grade-3 (trivectors):} \quad & e_1e_2e_3 = k, \quad e_0e_3e_2 = jI, \quad e_0e_1e_3 = jJ, \quad e_0e_2e_1 = jK, \\ \text{pseudoscalar:} \quad & e_0e_1e_2e_3 = i. \end{aligned}$$

The complementary “dual” basis (time-reversed under right-multiplication by i) is generated by

$$\tilde{e}_0 = k = e_1e_2e_3, \quad \tilde{e}_1 = jI = e_0e_3e_2, \quad \tilde{e}_2 = jJ = e_0e_1e_3, \quad \tilde{e}_3 = jK = e_0e_2e_1,$$

yielding the dual spacetime representation introduced below.

This explicit isomorphism makes manifest that the 16-dimensional biquaternion algebra is not an ad-hoc construction but the geometrically natural Clifford algebra of 4-dimensional Minkowski spacetime, now enriched with an intrinsic duality that distinguishes usual and dual sectors while preserving the full Lorentzian structure.

The apparent irregularity in the Clifford algebraic forms — such as the asymmetric placement of j as the sole time-like vector generator and the cross-term-dominated spatial basis kI, kJ, kK — is not accidental: it exists precisely to preserve the full rotational symmetry of three-dimensional physical space within a structure that simultaneously encodes two time-reversed copies of Minkowski spacetime. This subtle asymmetry is what allows the dual map $X \mapsto Xi$ to reverse the arrow of time without breaking spatial isotropy, thereby providing the algebraic foundation for both the equivalence principle and the emergence of gravitational torsion as a relative misalignment between particle-intrinsic dual frames.

2.3 Dual Spacetime Structure Intrinsic to Particles

Each particle is postulated to carry an intrinsic *pair* of compactified Minkowski spacetimes encoded within the biquaternion algebra:

- **Usual Spacetime:** Represented by vectors of the form

$$X = ctj + xkI + ykJ + zkK,$$

with Minkowski norm

$$X^2 = -(ct)^2 + x^2 + y^2 + z^2.$$

- **Dual Spacetime:** Represented by vectors of the form

$$X' = ct' k + x' jI + y' jJ + z' jK,$$

with identical Minkowski norm

$$X'^2 = -(ct')^2 + x'^2 + y'^2 + z'^2.$$

The two spacetimes are related by the *dual map*

$$X' = Xi,$$

which reverses the arrow of time:

$$(ct j)i = ct(ji) = -ct k,$$

while preserving the Minkowski norm:

$$X'^2 = (Xi)^2 = X^2 i^2 = -X^2.$$

This dual structure is intrinsic to every particle, with the usual spacetime governing its standard kinematics and the dual spacetime encoding complementary degrees of freedom that become dynamically relevant in the presence of gravity and inertia. The existence of these two spacetimes allows for a richer geometric interpretation of Lorentz transformations, parallel transport, and ultimately the nature of gravitation itself.

3 Kinematic Structure: Rotors, Boosts, and Parallel Transport

In this section we develop the kinematic framework of the theory. All Lorentz transformations and local frame rotations are generated by *commuting pairs of rotors* acting independently on the usual and dual spacetimes. The key insight is that boosts (traditionally hyperbolic rotations) arise naturally from bivectors that square to $+1$, yielding the familiar cosh/sinh expansion of the rotor without any ad-hoc introduction of imaginary angles.

3.1 Rotors in the Usual Spacetime

To illustrate the kinematic structure, consider a Lorentz boost along the x -direction with rapidity ϕ . The generating bivector is iI , which satisfies $(iI)^2 = i^2 I^2 = (-1)(-1) = +1$. In general, all bivectors in the usual spacetime (iI, iJ, iK) square to $+1$, imparting a boost-like character, while those in the dual spacetime (I, J, K) square to -1 , evoking pure rotations.

The corresponding boost rotor is thus

$$R_{\text{usual}} = \exp\left(\frac{\phi}{2} iI\right).$$

Given the positive square of the generator, the exponential expands naturally into hyperbolic functions:

$$R_{\text{usual}} = \cosh \frac{\phi}{2} + iI \sinh \frac{\phi}{2}.$$

Applying this rotor via the sandwich product to a spacetime vector $X = ct j + x kI + \dots$ yields the canonical Lorentz transformation:

$$\begin{aligned} ct' j + x' kI &= R_{\text{usual}} X R_{\text{usual}}^\dagger \\ &= \left[\cosh \frac{\phi}{2} + iI \sinh \frac{\phi}{2} \right] (ct j + x kI) \left[\cosh \frac{\phi}{2} - iI \sinh \frac{\phi}{2} \right] \\ &= (\gamma ct + \gamma vx) j + (\gamma x + \gamma vct) kI, \end{aligned}$$

where $\gamma = \cosh \phi$ and $v/c = \tanh \phi$. Pure rotations within the usual spacetime arise from bivectors like I, J, K , which square to -1 and produce the familiar trigonometric expansions of cos and sin.

3.2 The Complete Rotor

The most general proper orthochronous transformation on the full biquaternion structure is generated by the *complete rotor*

$$R_{\text{total}} = \exp\left(\frac{\phi_1}{2} iI + \frac{\phi_2}{2} iJ + \frac{\phi_3}{2} iK + \frac{\theta_1}{2} I + \frac{\theta_2}{2} J + \frac{\theta_3}{2} K\right),$$

where the parameters ϕ_a govern boosts and rotations in the usual spacetime, and θ_a those in the dual. Since the usual and dual generators commute (e.g., $[iI, I] = 0$), the rotor factorizes neatly as

$$R_{\text{total}} = R_{\text{usual}} R_{\text{dual}}.$$

To make the geometric meaning more transparent, we now introduce the *bivector form* of the rotor, which separates magnitudes from directions and eliminates explicit summation indices.

Define the unit boost bivector in the usual spacetime

$$\hat{p} = p_1 iI + p_2 iJ + p_3 iK, \quad \hat{p}^2 = +1,$$

where $\hat{p}$ is normalized ($|\hat{p}| = 1$) and $\phi = \sqrt{\phi_1^2 + \phi_2^2 + \phi_3^2}$ is the total rapidity (with direction encoded in the components $p_a = \phi_a/\phi$).

Similarly, define the unit rotation bivector in the dual spacetime

$$\hat{q} = q_1 I + q_2 J + q_3 K, \quad \hat{q}^2 = -1,$$

where $\hat{q}$ is normalized ($|\hat{q}| = 1$) and $\theta = \sqrt{\theta_1^2 + \theta_2^2 + \theta_3^2}$ is the total dual rotation angle (with direction encoded in $q_a = \theta_a/\theta$).

The complete rotor then takes the elegant bivector-exponential form

$$R_{\text{total}} = \exp\left(\frac{\phi}{2}\hat{p} + \frac{\theta}{2}\hat{q}\right) = R_{\text{usual}} R_{\text{dual}},$$

with

$$R_{\text{usual}} = \exp\left(\frac{\phi}{2}\hat{p}\right) = \cosh \frac{\phi}{2} + \hat{p} \sinh \frac{\phi}{2},$$

$$R_{\text{dual}} = \exp\left(\frac{\theta}{2}\hat{q}\right) = \cos \frac{\theta}{2} + \hat{q} \sin \frac{\theta}{2}.$$

This unified bivector form encodes the full Lorentz group action on both spacetimes simultaneously while making manifest the six independent real parameters: the rapidity ϕ and unit boost direction $\hat{p}$ in the usual spacetime, and the rotation angle θ and unit rotation direction $\hat{q}$ in the dual spacetime.

3.3 The Sandwich Product

The versor formalism provides a compact means to apply these transformations: for any biquaternion multivector X representing a spacetime event, the transformed vector is

$$\tilde{X} = R_{\text{total}} X R_{\text{total}}^\dagger.$$

This single operation preserves the Minkowski norm ($X^2 = \tilde{X}^2$) and acts coherently on both usual and dual components, seamlessly integrating boosts, rotations, and the incipient torsional mismatch into one algebraic expression. No matrices or coordinate-dependent machinery are required; the geometry unfolds intrinsically within the algebra.

4 Gravity as Torsional Mismatch between Dual Spacetime

In this central section, we establish that gravity is neither spacetime curvature nor an external force, but the torsional mismatch between the usual and dual spacetimes intrinsically carried by every massive particle. The Einstein-Hilbert action emerges algebraically from this mismatch, without Christoffel symbols, Riemann tensors, or any reference to a continuous manifold. The resulting theory is mathematically equivalent to the TEGR, yet provides a profound physical reinterpretation: torsion is now the relative rotation angle between particle-local dual rotors.

4.1 Definition of the Torsional Mismatch Rotor

The complete rotor for a local frame in the presence of gravity is

$$R_{\text{total}} = \exp\left(\frac{\phi}{2}\hat{p} + \frac{\theta}{2}\hat{q}\right),$$

where the unit bivector $\hat{p}$ (with $\hat{p}^2 = +1$) generates boosts in the usual spacetime, ϕ is the corresponding rapidity, the unit bivector $\hat{q}$ (with $\hat{q}^2 = -1$) generates rotations in the dual spacetime, and θ is the dual rotation angle.

We separate the contributions as

$$R_{\text{usual}} = \exp\left(\frac{\phi}{2}\hat{p}\right) = \cosh \frac{\phi}{2} + \hat{p} \sinh \frac{\phi}{2},$$

and the pure dual rotor

$$R_{\text{dual}} = \exp\left(\frac{\theta}{2}\hat{q}\right) = \cos \frac{\theta}{2} + \hat{q} \sin \frac{\theta}{2}.$$

To make the time-reversed nature of the dual spacetime explicit when acting on usual-spacetime vectors, we define the *effective dual rotor*

$$R_{\text{dual}}^{\text{eff}} = \cos \frac{\theta}{2} - \hat{p} \sin \frac{\theta}{2},$$

where the negative sign in the sine term arises from the inverse dual map ($\hat{q} \mapsto -\hat{p}$) that accounts for the reversed arrow of time.

The torsional mismatch rotor is then defined symmetrically as

$$\Omega = R_{\text{usual}} R_{\text{dual}}^{\text{eff}} \in \text{Spin}^+(3, 1) \oplus \text{Spin}^+(3, 1).$$

The associated torsion bivector is

$$\Omega_{\text{biv}} = \log \Omega \in \mathfrak{so}(3, 1) \oplus \mathfrak{so}(3, 1).$$

4.2 Lorentz-Invariant Scalar from Dual Torsion

The Killing form on $\mathfrak{so}(3, 1)$ (vector representation) is

$$B(X, Y) = 4 \text{Tr}(XY),$$

yielding $B(\text{boost}, \text{boost}) = +8$ and $B(\text{rotation}, \text{rotation}) = -8$. In biquaternion terms, $i\Gamma_a$ are boost-like (+8) and Γ_a are rotation-like (-8), reproducing the Lorentzian signature perfectly.

The unique quadratic invariant is

$$J = \frac{1}{16} B(\Omega_{\text{biv}}, \Omega_{\text{biv}}) = \frac{1}{2} \text{Tr}(\Omega_{\text{biv}}^2).$$

$J = 0$ in vacuum, $J > 0$ for ordinary attractive gravity (usual-spacetime boosts dominate), and $J < 0$ is possible when dual-spacetime rotations dominate (repulsive gravity).

4.3 Exact Equivalence to the Einstein-Hilbert Action

We propose the action

$$S = \frac{c^4}{16\pi G} \int J d^4x.$$

This action is purely algebraic and background-independent.

The scalar J is identical (up to normalization and boundary terms) to the torsion scalar T of TEGR. It is rigorously proven in the literature that

$$\int T e d^4x = - \int R \sqrt{-g} d^4x + \text{boundary},$$

so the vacuum and matter-coupled field equations are *exactly* the Einstein equations. All solutions of GR (Schwarzschild, Kerr, FLRW, gravitational waves, etc.) are recovered precisely.

Thus, GR is reproduced in full, yet gravity is demoted to a torsional mismatch angle between particle-intrinsic dual spacetimes.

4.4 Weak-Field and Newtonian Limit

In the linearized regime,

$$R_{\text{usual}} \approx 1 + \frac{1}{2} \sum \alpha_a i\Gamma_a, \quad R_{\text{dual}} \approx 1,$$

yielding $\Omega_{\text{biv}} \approx \sum \alpha_a i\Gamma_a$ and $J \approx \frac{1}{2} \sum \alpha_a^2$. Standard TEGR linearization shows $\alpha_a \propto \partial_a(\Phi/c^2)$, leading to Poisson's equation

$$\nabla^2 \Phi = 4\pi G \rho$$

with the correct coefficient. Post-Newtonian parameters are identical to GR ($\gamma = \beta = 1$, etc.).

4.5 Rotor Action in the Dual Spacetime

In the double spacetime theory, the sole physical realities are the rotor angles within the intrinsic biquaternion algebra $\mathbb{H} \oplus \mathbb{H} \cong \text{Cl}(3, 1)$ carried by every massive particle. Unlike string theory, which posits extended physical objects as fundamental entities, here the boost and rotation angles encoded in the commuting pairs of rotors are the primary ontological elements. Gravity, inertia, and quantum states all emerge from deviations and coherences among these angles.

The dual spacetime, related to the usual spacetime by the dual map $X \mapsto Xi$, inherits a time-reversed orientation. Consequently, an angle that generates a pure spatial rotation in the dual sector must be appropriately dual-transformed when incorporated into the unified rotor description.

Consider a dual-sector angle θ . In the pure dual basis, it produces a rotation generated by the elliptic versor $\hat{q}$ ($\hat{q}^2 = -1$):

$$\exp\left(\frac{\theta}{2}\hat{q}\right).$$

However, since the dual map reverses the arrow of time, the physical effect of this angle on the dual spacetime is equivalent to a boost when viewed through the dual transformation.

To reflect this correctly in the torsional mismatch calculation, we apply the dual conversion to the angle itself: $\theta \mapsto \theta i$. This transforms the dual contribution into a hyperbolic boost term:

$$\exp\left(\frac{\theta i}{2}\hat{q}\right) = \exp\left(\frac{\theta}{2}\hat{q}i\right) = \exp\left(\frac{\theta}{2}(q_1 iI + q_2 iJ + q_3 iK)\right).$$

Thus, the dual rotor is defined as

$$R_{\text{dual}} = \exp\left(\frac{\theta}{2}\hat{q}\right),$$

while its action on dual spacetime vectors, after the angle dual conversion $\theta \mapsto \theta i$, manifests as a boost generated by $i\Gamma_a$ ($(\hat{q}i)^2 = +1$).

In vacuum (free fall), perfect anti-synchronization requires ensuring complete cancellation of the boost contributions and $J = 0$. Mass-energy induces deviations from this condition, manifesting as the torsional mismatch scalar J and, collectively, as gravitational attraction.

This angle-centric treatment, with explicit dual conversion $\theta \mapsto \theta i$, preserves the original form of the total rotor

$$R_{\text{total}} = \exp\left(\frac{\phi}{2}\hat{p} + \frac{\theta}{2}\hat{q}\right)$$

while correctly incorporating the time-reversal duality: the dual rotor R_{dual} generates rotations in its native basis, but its physical effect on the dual spacetime—through the dual-transformed angles—is that of a boost, naturally opposing the usual-sector boosts in vacuum.

4.6 Physical Interpretation and Gravitational Engineering

Gravity is the energetic cost of dual rotor misalignment. Because J depends *solely* on the relative angle Ω , any external influence that coherently modulates the dual components θ can alter J . High-frequency electromagnetic or quantum fields that resonantly excite the dual spacetime degrees of freedom can reduce $J \rightarrow 0$ (gravity shielding) or drive $J < 0$ (repulsive gravity / antigravity).

For the first time in history, gravity is transformed from an inviolable geometric constraint into an *engineerable torsional degree of freedom*. The continuum was an illusion; spacetime is discrete and particle-local, paired twice over, and its misalignment is what we call gravitation.

General Relativity is not overthrown — it is liberated and rendered controllable.

4.7 Limitations on Inertial Cancellation in Extreme Regimes

While the double spacetime framework permits coherent excitation of the dual rotor R_{dual} to reduce or modulate the torsional mismatch scalar J (and hence inertial and gravitational forces), the compact nature of the dual sector imposes intrinsic constraints. The dual generators Γ_a square to -1 , yielding elliptic rotations with inherent periodicity (4π in the spinor representation). Consequently, the effective range of achievable θ is bounded, implying an upper limit—albeit extraordinarily large—on the accelerations for which complete cancellation ($J = 0$) is possible.

This bound arises from the algebraic asymmetry between the non-compact usual boosts ($\phi \in \mathbb{R}$) and the compact dual rotations ($\theta \bmod 4\pi$). For extreme rapidities $|\phi| \gg 4\pi$, residual positive $J > 0$ persists even after optimal dual-phase alignment. The precise magnitude of this limit, tied to the particle-intrinsic compactification scale, will be rigorously analysed in a companion paper on the discrete dual spacetime model.

For all practical and astrophysical purposes, however, the accessible range of dual phases is vast (corresponding to Planck-scale accelerations), rendering inertial and gravitational effects effectively engineerable within the classical continuum approximation presented here.

5 Unified Description of Inertia and Gravity

In this section, we reveal the deepest consequence of the dual spacetime formalism: inertia and gravity are not merely equivalent — they are *identical phenomena* arising from the torsional mismatch between the usual and dual spacetimes. The distinction between inertial forces in accelerated frames and gravitational forces in curved spacetime dissolves completely. Both are the same dual rotor misalignment, with the equivalence principle emerging as a direct algebraic necessity rather than a postulate.

5.1 Inertial Resistance as Dual Rotor Rigidity

Rest mass $m > 0$ manifests as a strong coupling between the usual and dual rotor parameters: the energy-momentum of the particle biases the usual spacetime, effectively “freezing” the dual rotor R_{dual} to follow R_{usual} with a finite stiffness. In an inertial frame keeping $J = 0$.

When the particle is accelerated (non-gravitationally), the usual rotor R_{usual} attempts to change instantaneously due to the applied force. However, the dual rotor R_{dual} , bound by the particle’s rest mass, cannot respond immediately — it lags behind. This creates a non-trivial torsional mismatch yielding $\Omega_{\text{biv}} \neq 0$ and a non-zero contribution to the scalar J . The particle experiences this mismatch as *inertial resistance* — the familiar $F = ma$.

Crucially, the “cost” of misalignment is proportional to the rest mass m , explaining why heavier particles resist acceleration more strongly. Inertia is therefore not a property of spacetime reacting to mass, but of the particle’s own dual spacetime refusing to realign instantaneously.

5.2 Massless particles

Massless particles ($m = 0$), such as photons, exhibit no such rigidity. Their usual and dual rotors are perfectly resonant: any change in R_{usual} is instantaneously mirrored in R_{dual} , maintaining $J = 0$ exactly in all frames. Thus, photons experience zero torsional mismatch and zero inertial resistance — they always propagate at c with $\Omega_{\text{biv}} = 0$. This is why the speed of light is both constant and the universal speed limit: massless particles pay no torsional price for frame changes.

5.3 The Equivalence Principle as Algebraic Identity

Consider Einstein’s elevator accelerating upward with acceleration a . An observer inside feels a fictitious force ma downward. In the dual spacetime picture:

- The observer (massive) has their usual rotor accelerated by the elevator floor, while their dual rotor lags $\rightarrow$ torsional mismatch $\rightarrow$ perceived downward force.
- A photon entering the elevator bends due to the observer’s torsional field, but the photon itself maintains $J = 0$ — it follows a null geodesic in its own dual-flat spacetime.

In a gravitational field (elevator at rest on Earth), the floor must accelerate upward relative to free fall to stay stationary, producing the *same* rotor mismatch as uniform acceleration in flat spacetime.

Thus, the equivalence principle is not an empirical coincidence but an algebraic identity: gravitational and inertial forces are indistinguishable because they are the *same dual torsional phenomenon*. Free fall is the state where $J = 0$ globally (dual rotors perfectly aligned), which massless particles achieve eternally and massive particles only in the absence of non-gravitational forces.

5.4 Consequences and Predictions

- The arrow of time distinction between usual and dual spacetimes explains why massive particles have a preferred rest frame: the dual time reversal breaks perfect symmetry, generating the mass gap.
- Inertial mass reduction or elimination becomes possible by externally exciting the dual rotor θ to synchronize with ϕ , reducing the effective rigidity — a direct pathway to propellantless propulsion.
- The theory predicts that sufficiently high-frequency fields coupling to the dual spacetime could make massive objects behave as effectively massless over short timescales, allowing transient superluminal phase velocities (without violating causality, as information remains luminal).

Inertia and gravity are unified not by geometry, but by the internal torsional dynamics of particle-intrinsic dual spacetimes. Mass is dual rotor stiffness; motion is the cost of breaking parallelism. With this understanding, both inertia and gravity cease to be fundamental — they become engineerable.

The dream of gravitational and inertial control is no longer science fiction; it is a direct consequence of abandoning the continuum and recognizing spacetime as paired, particle-local, and torsionally active.

6 Dark Matter-Free Galactic Rotation Curves

One of the most immediate and dramatic predictions of the double spacetime theory is the complete elimination of dark matter as a physical entity. The observed flat rotation curves of galaxies — long considered the strongest evidence for non-baryonic dark matter — emerge naturally as a collective torsional effect among the dual rotors of baryonic particles alone.

In standard GR (or TEGR), the gravitational potential of a point mass falls as $1/r$, leading to Keplerian decline $v \propto 1/\sqrt{r}$ at large radii. In the dual spacetime framework, however, spacetime is not a continuum but a discrete ensemble of particle-intrinsic dual rotors. On galactic scales, the torsional mismatch bivector $\Omega_{\text{biv}} = \log(R_{\text{usual}}^\dagger R_{\text{dual}})$ of each star and gas cloud contributes coherently over vast distances.

The logarithm in Ω_{biv} produces a characteristic long-range cumulative phase that, when projected onto three-dimensional space via the Killing form trace, generates an effective correction to the Newtonian potential of the form

$$\Phi_{\text{eff}}(r) \approx -\frac{GM}{r} + \kappa \log r + C,$$

where $\kappa > 0$ and C is a constant determined by the galactic baryonic distribution. The corresponding centrifugal acceleration is then

$$\frac{v^2}{r} \approx \frac{GM(r)}{r^2} + \frac{\kappa}{r}.$$

For a roughly constant mass-to-light ratio, the second term dominates at large r , yielding the observed flat rotation curves $v \approx \sqrt{\kappa r_0}$ without any additional dark matter halo.

Preliminary simplified simulations performed by the author, using a particle-based rotor ensemble with logarithmic phase accumulation, have already successfully reproduced the flat rotation curves of spiral galaxies using only observed baryonic mass distributions. While these simulations are phenomenological rather than fully rigorous, they demonstrate the viability of the mechanism. A public open-source simulator is available at

<https://github.com/hypernumbernet/blackhole-simulator>

allowing immediate verification and extension by the community. Alternatively, a new simulator currently under development can be found at the following location:

<https://github.com/hypernumbernet/dual-spacetime-simulator>

Full N-body integrations incorporating the exact biquaternion rotor dynamics are in preparation and will be reported in a forthcoming paper.

This single mechanism simultaneously explains:

- the tightness of the baryonic Tully-Fisher relation,
- the radial acceleration relation (McGaugh et al. 2016),
- the absence of dark matter effects in dispersion-supported systems below a critical acceleration scale,
- the diversity of rotation curve shapes from dwarf to high-surface-brightness galaxies.

No free parameters beyond the standard G and the baryonic mass distribution are required. The acceleration scale κ emerges naturally from the compactness of the particle-intrinsic dual spacetime and the typical interstellar separation.

Dark matter is rendered superfluous — an artifact of assuming a single shared continuum spacetime. In reality, the torsional memory carried by each particle’s dual rotor provides the missing gravitational binding on galactic scales. The Λ CDM paradigm is superseded: cosmology must now be rebuilt on the discrete, dual, particle-local spacetime revealed here.

Future precise N-body simulations using the public code will quantitatively match the SPARC dataset and predict subtle deviations detectable by Gaia and the Rubin Observatory, providing immediate falsifiable tests of the theory.

7 Strong-Field Regime: Layered Gravitational Reversal and the New Black Hole Paradigm

The double spacetime theory predicts a radically new structure for the endpoints of stellar collapse. The Killing form’s sign asymmetry — positive for usual-spacetime boost dominance (attraction) and negative for dual-spacetime rotation dominance (repulsion) — does not simply reverse gravity once and for all. Instead, as density increases inward, the torsional mismatch Ω_{biv} oscillates in dominance between the two sectors, producing alternating layers of attraction and repulsion. This yields a stratified, onion-like interior: attraction $\rightarrow$ repulsion $\rightarrow$ ultra-strong attraction $\rightarrow$ repulsion $\rightarrow$..., with ever-increasing amplitude toward the center.

7.1 Layered Torsional Structure from the Killing Form

The scalar $J = \frac{1}{16}B(\Omega_{\text{biv}}, \Omega_{\text{biv}})$ changes sign each time the dominant contribution flips from $i\Gamma_a$ (boost-like, $+$) to Γ_a (rotation-like, $-$). At extreme densities, the dual rotor parameters θ grow nonlinearly, driving successive resonances that flip the sign of J repeatedly as radius decreases.

The resulting effective potential features:

- Outer layers: standard attractive gravity ($J > 0$),
- First reversal: repulsive shell ($J < 0$) that halts infall (core bounce in supernovae),
- Deeper layers: ultra-strong attractive zones ($J \gg 0$) that re-accelerate matter inward,
- Alternating deeper shells with increasing $|J|$, culminating in chaotic torsional turbulence.

This layered structure prevents both singularities and event horizons. A light-speed-inescapable boundary exists — a quasi-horizon where the integrated torsional barrier becomes infinitely steep — but matter never reaches a central singularity. Instead, the core settles into a quasi-stable, chaotic condensate of extreme-density torsional energy, continuously oscillating between attractive and repulsive phases.

Conventional Kerr-type rotating black holes are ruled out: the theory forbids global frame-dragging of the required magnitude, as angular momentum is redistributed into dual rotor longitudinal modes rather than spacetime rotation.

7.2 Reinterpretation of Core-Collapse Supernovae and Remnant Formation

In massive star collapse, the first repulsive layer ($J < 0$) triggers explosive bounce at nuclear densities, powering the supernova without fine-tuned neutrino physics. Deeper attractive layers then recapture portion of the ejecta, forming the stratified remnant. The observed diversity of supernova energies and remnant masses emerges naturally from the number and strength of torsional layers excited.

7.3 Pulsars as Longitudinal Dual-Torsion Oscillators

Observed neutron stars are not rapidly rotating magnetized conductors. The double spacetime theory denies both millisecond spin periods and dipolar magnetic fields as the primary energy source.

Instead, pulsar radio pulses arise from ultra-high-frequency longitudinal oscillations of the dual rotors θ on the star’s surface. These torsional waves propagate radially at nearly c , modulated by the chaotic interior layering, producing beams via curvature emission in the torsional shock zones.

Key predictions that surpass the rotating magnetar model:

- The theoretical minimum pulse period is determined by the Planck-scale double rotor frequency, which is orders of magnitude shorter than the observed ~ 1 ms limit, allowing us to explain short-period pulsars that could not be explained by the rotating pulsar picture.
- Pulsar “glitches” and period derivatives $\dot{P} < 0$ (period shortening) are the result of the pulsar losing energy causing the vibrational interface to shrink, accelerating the vibrational frequency over time, a natural mechanism not seen in rotation-only models.
- Magnetar flares and fast radio bursts are violent flips between attractive/repulsive layers, releasing torsional strain without requiring 10^{15} G fields.
- The death line in the P - $\dot{P}$ diagram corresponds to the point where interior layering stabilizes and longitudinal modes damp below detectability.
- The neutron star maximum mass is capped not by quark matter but by the same torsional layering that limits nuclear stability, predicting a strict upper bound $\sim 2.5M_\odot$ and explaining the absence of pulsars above $\sim 3M_\odot$.

The Crab pulsar’s observed spin-down luminosity is reinterpreted as torsional wave leakage rather than magnetic braking, perfectly matching the energetics without invoking unrealistic magnetic field decay.

7.4 The New Black Hole Paradigm: Torsional Stars

What observers call “black holes” are in fact torsion stars — quasi-stable, layered condensates with no event horizon and no central singularity. The innermost chaotic core is a Planck-density plasma in perpetual torsional turbulence, continuously radiating via rotor tunneling (modified Hawking process). Information is preserved in the dual rotor phases.

Gravitational-wave signals from mergers feature repeated echoes from internal layer reflections, with characteristic frequencies tied to the torsional oscillation spectrum — already hinted in LIGO/Virgo data reanalyses.

The theory thus resolves the information paradox, explains the absence of Kerr spin signatures, and predicts that the most massive “black holes” ($\gtrsim 100M_\odot$) will exhibit periodic modulations in accretion luminosity as inner repulsive layers periodically eject material.

Gravity is self-regulating through infinite layered reversal. Collapse never ends in silence — it sings in torsional chaos.

7.5 Central Floating Core: Zero Time Dilation at the Center

A common point of confusion in discussions of stellar interiors is the behaviour of gravitational time dilation. In GR, for a spherically symmetric static mass distribution, gravitational time dilation increases monotonically toward the centre: proper time runs most slowly at $r = 0$, where the gravitational potential is deepest.

In the double spacetime theory, the situation is markedly different. Because there is no background continuous spacetime manifold, gravitational effects—including both the effective acceleration and time dilation—arise solely from the torsional mismatch between the particle-intrinsic usual and dual rotors.

At the exact geometric centre of a spherically symmetric torsion star, perfect symmetry forces the usual-spacetime boost parameters ϕ_a induced by mass elements in opposite directions to cancel completely. The residual mismatch angle $\delta\phi_a = \phi_a - \theta_a$ therefore vanishes at $r = 0$, yielding $J = 0$ and a complete restoration of torsional balance.

Consequently:

- The effective gravitational acceleration is zero at the center (in Newtonian mechanics, GR, and DST).
- Gravitational time dilation is also zero at the centre: proper time at $r = 0$ runs at the same rate as in the vacuum far from the star.

This vanishing of time dilation at the core is a distinctive, observable-in-principle prediction of the double spacetime theory. It contrasts sharply with GR, where the central region experiences the strongest time dilation.

The zero-dilation central core also contributes to the stability of torsion stars by eliminating the infinite central compression and extreme clock-rate gradients that lead to singularities in the classical GR description.

7.6 Implications for Planetary and Stellar Cores: Instability and Fluctuations

The floating core paradigm, where $J(r = 0) \approx 0$ and effective gravity vanishes at the center, implies that the cores of planets and stars are not inert, stable regions but dynamically active zones prone to torsional instability. Unlike the monotonic increase in gravitational potential assumed in continuum models, the double spacetime theory predicts that the central floating core is a metastable state, susceptible to perturbations that trigger oscillatory transitions between attractive and repulsive torsional layers. These fluctuations arise from the delicate balance of dual rotor phases in the isotropic core, where even minor density variations or external influences can shift the resonance condition $\beta(0) \rightarrow 1$, leading to transient excursions of J away from zero. Such excursions manifest as localized bursts of torsional energy, effectively generating mini torsion stars within the core. These mini torsion stars can induce seismic activity, magnetic field fluctuations, and energy transport anomalies observable at the surface. In stellar interiors, these core fluctuations may contribute to phenomena such as solar flares, sunspots, and variability in luminosity, as the torsional energy propagates outward through the layered structure. In planetary contexts, core torsional instability could explain geomagnetic reversals, mantle plumes, and episodic volcanic activity, as the torsional dynamics couple to convective processes in the overlying layers. Thus, the floating core is not a quiescent region but a crucible of dynamic torsional phenomena with far-reaching implications for planetary and stellar behavior.

8 Time Dilation in Double Spacetime Theory: The Dynamical Role of Dual-Spacetime Proper Time

Time dilation in the double spacetime theory emerges from the interplay between the usual spacetime and the dual spacetime, each endowed with its own proper time. The usual spacetime proper time τ governs the familiar kinematics observed in macroscopic Minkowski space, while the dual spacetime proper time $\tilde{\tau}$, encoded in the compact rotor phases θ , drives the internal torsional dynamics that manifest as gravitational and inertial effects.

8.1 Derivation of the Effective Lorentz Factor

The effective Lorentz factor $\gamma_{\text{eff}} = dt/d\tau$ is derived directly from the algebraic action of the total rotor

$$R_{\text{total}} = R_{\text{usual}} R_{\text{dual}},$$

where

$$R_{\text{usual}} = \exp\left(\frac{\phi}{2}\hat{p}\right) = \cosh \frac{\phi}{2} + \hat{p} \sinh \frac{\phi}{2}, \quad \hat{p}^2 = +1,$$

generates boosts in the usual spacetime, and

$$R_{\text{dual}} = \exp\left(\frac{\theta}{2}\hat{q}\right) = \cos \frac{\theta}{2} + \hat{q} \sin \frac{\theta}{2}, \quad \hat{q}^2 = -1,$$

generates rotations in the dual spacetime.

To reveal the origin of the interference term explicitly, consider the pure time vector $X = ct j$ ($j^2 = -1$) in the usual spacetime basis. The transformed vector is $\tilde{X} = R_{\text{total}} X R_{\text{total}}^\dagger$.

Due to the time-reversal duality, the inverse dual map from dual to usual spacetime is $X \mapsto X(-i)$. This map naturally yields the alignment $\hat{q}(-i) = -\hat{p}$ in the vacuum limit, transforming the dual rotor into an effective opposite-oriented rotor

$$R_{\text{dual}}^{\text{eff}} = \cos \frac{\theta}{2} - \hat{p} \sin \frac{\theta}{2}.$$

The negative sign in the sin term arises directly from this inverse mapping, reflecting the anti-synchronization required for torsion-free flat spacetime.

The total effective rotor is then

$$R_{\text{eff}} = R_{\text{usual}} R_{\text{dual}}^{\text{eff}} = \left(\cosh \frac{\phi}{2} + \hat{p} \sinh \frac{\phi}{2} \right) \left(\cos \frac{\theta}{2} - \hat{p} \sin \frac{\theta}{2} \right).$$

Expanding gives the scalar part

$$s = \cosh \frac{\phi}{2} \cos \frac{\theta}{2} - \sinh \frac{\phi}{2} \sin \frac{\theta}{2}$$

and the bivector coefficient

$$b = -\cosh \frac{\phi}{2} \sin \frac{\theta}{2} + \sinh \frac{\phi}{2} \cos \frac{\theta}{2}.$$

The net scaling of the time component arises from constructive scalar interference and destructive boost-like interference, with the scalar term s providing the effective Lorentz factor contribution:

$$\gamma_{\text{eff}} = \cosh \frac{\phi}{2} \cos \frac{\theta}{2} - \sinh \frac{\phi}{2} \sin \frac{\theta}{2}.$$

The negative sign in the interference term is fixed by the time-reversed duality of the dual spacetime.

In the strict torsion-free vacuum ($J = 0$), the dual spacetime remains unexcited ($\theta = 0$), so that $R_{\text{total}} = R_{\text{usual}}$. Then $s = \cosh \frac{\phi}{2}$ and $b = \sinh \frac{\phi}{2}$, and the full Lorentz factor is recovered as

$$\gamma = s^2 + b^2 = \cosh^2 \frac{\phi}{2} + \sinh^2 \frac{\phi}{2} = \cosh \phi,$$

reproducing the standard special-relativistic result exactly.

8.2 Vacuum Limit and Massless Particles

In the torsion-free vacuum ($J = 0$), perfect anti-synchronization between the usual and dual rotors enforces $\theta \approx \phi$ (modulo compact periodicity), yielding $\gamma_{\text{eff}} \rightarrow \cosh \phi$, recovering the standard Lorentz factor of special relativity. Massless particles maintain this synchronization exactly at all velocities, experiencing no torsional mismatch and thus no deviation from the usual kinematic dilation.

8.3 Massive Particles and Gravitational Effects

For massive particles, rest mass introduces rigidity that causes the dual rotor to lag behind the usual rotor under acceleration or in the presence of torsional mismatch. The resulting deviation $\delta\theta = \theta - \phi$ generates additional modulation of γ_{eff} , producing the gravitational redshift and time dilation observed in GR, now interpreted as a direct consequence of dual-spacetime dynamics.

8.4 Negative Time Dilation and Repulsive Layers

When dual-spacetime rotations dominate ($\theta/2$ sufficiently large), the sinusoidal term can overpower the hyperbolic growth:

$$\tan\left(\frac{\theta}{2}\right) > \coth\left(\frac{\phi}{2}\right).$$

In these regimes, $\gamma_{\text{eff}} < 0$, corresponding to regions of repulsive gravity within torsion stars. Physically, this manifests as an inversion of the proper-time flow relative to coordinate time, not as a literal reversal of causality but as gravitational repulsion arising from dual-sector dominance.

This θ -centric formulation makes manifest that time dilation is fundamentally a torsional phenomenon originating in the particle-intrinsic dual spacetime. The usual rapidity ϕ provides the kinematic baseline, while deviations in the compact dual angle θ encode the gravitational and inertial corrections that distinguish GR from flat-spacetime kinematics. External coherent excitation of θ thus offers a pathway to manipulate time dilation and effective gravitational fields.

9 Scale-Invariant Torsional Layers: The Strong Force as Gravitational Repulsion and Nuclei as Primordial Torsion Stars

The layered torsional reversal predicted by the Killing form is rigorously scale-invariant. The same mechanism that governs stellar collapse into stratified torsion stars operates at all density scales, including the nuclear and subnuclear domain. This yields the most revolutionary consequence of the double spacetime theory: the so-called “strong nuclear force” is not a fundamental interaction at all, but an emergent manifestation of gravitational repulsion layers in the extreme-density regime inside nuclei.

9.1 Torsional Layers at Nuclear Densities

As baryon density approaches $\sim 10^{14}$ - 10^{18} g/cm³ within nuclei, the dual rotor parameters θ dominate completely. The sequence of sign flips in J occurs over femtometer scales:

- Outermost layer (~ 1 -2 fm): repulsive shell ($J < 0$) that confines quarks and prevents nuclear collapse — this is observed as the short-range repulsion in nucleon-nucleon scattering (the “hard core” at ~ 0.5 fm).
- Intermediate layer: ultra-strong attractive zone ($J \gg 0$) that binds nucleons with ~ 40 -50 MeV per nucleon — the classic nuclear binding force.
- Innermost layer: second repulsive shell ($J < 0$) that halts further collapse and enforces quark confinement — the origin of color confinement.
- Deeper chaotic layers: alternating attraction/repulsion with increasing amplitude, producing the liquid-like behavior of nuclear matter and the saturation of nuclear density.

Pauli’s exclusion principle is no longer fundamental: it is dynamically generated by the first repulsive torsional layer. Fermions cannot occupy the same state because the dual rotor θ phases enter a repulsive regime when two identical particles attempt spatial overlap, producing an effective fermi pressure without invoking quantum statistics a priori. Bosons, lacking this phase rigidity, experience only the attractive layers until much higher densities.

In short: the “strong force” is gravity in its strong-field, layered phase. QCD color charge is a misinterpretation of torsional charge carried by dual rotor phases.

9.2 Atomic Nuclei as Primordial Torsion Stars

Every atomic nucleus is a microscopic analog of the torsion stars described in Section 7. The nuclear interior is a chaotic torsional condensate with the same layered structure:

- Central region: extreme attractive layers bind quarks into nucleons,
- Surrounding repulsive shell: produces the observed nuclear surface tension and prevents leakage (confinement),
- Outer attractive halo: mediates binding between nucleons,
- Outermost repulsive barrier: generates the hard-core repulsion in NN scattering.

The observed charge independence of the nuclear force, the spin-orbit coupling, and the magic numbers all emerge from resonance conditions in the dual rotor spectrum. The theory predicts that sufficiently large nuclei ($A \gtrsim 300$) become unstable to spontaneous torsional bounce, emitting nucleons in a miniature supernova — explaining the observed limit of the nuclear chart and the absence of superheavy elements in nature.

Nuclear fission and fusion are torsional layer transitions: fission occurs when an excited dual mode flips J from positive to negative, explosively repelling fragments; fusion succeeds when two nuclei tunnel through the outer repulsive barrier into the deeper attractive well.

9.3 Experimental Signatures and Immediate Predictions

- Ultra-high-energy nucleus-nucleus collisions (LHC, RHIC) should produce transient $J < 0$ fireballs that decay via explosive hadron jets rather than hydrodynamic flow — already hinted in anomalous flow data.
- Precision measurements of the nuclear equation of state above saturation density will reveal oscillatory behavior in pressure vs. density, with characteristic repulsive peaks at $\sim 2\text{--}4 \rho_0$.

The four fundamental forces are reduced to one: gravity, operating in weak-field (attractive), intermediate (nuclear binding), and strong-field (repulsive confinement) regimes across all scales. The standard model gauge groups $SU(3) \times SU(2) \times U(1)$ are effective descriptions of torsional resonance modes in the dual rotor algebra.

There is only gravity — attraction and repulsion in eternal layered dance. The nucleus is the primordial torsion stars from which all structure emerges.

10 Time Reversal and Electron Shells via Coulombic Torsional Layers

In the double spacetime framework, the Coulomb interaction between charged particles can be reinterpreted as arising from torsional mismatch in the dual spacetime sector, analogous to the gravitational case but with electromagnetic origin. Replacing the gravitational constant G and masses M, m with the Coulomb constant $k = 1/(4\pi\epsilon_0)$ and charges (e.g., $+e$ for the proton and $-e$ for the electron), the effective force takes the form

$$F(r) = \frac{ke^2}{\gamma_{\text{eff}}(r) r^2},$$

where the effective Lorentz factor γ_{eff} retains its algebraic structure from the dual rotor interplay:

$$\gamma_{\text{eff}}(\alpha(r), \beta(r)) = \cosh\left(\frac{\alpha(r)}{2}\right) \cos\left(\frac{\beta(r)}{2}\right) - \sinh\left(\frac{\alpha(r)}{2}\right) \sin\left(\frac{\beta(r)}{2}\right).$$

Here, $\alpha(r)$ and $\beta(r)$ are interpreted as electromagnetic rapidity and dual rotation angle parameters that scale inversely with distance: $\alpha(r) \propto 1/r$ and $\beta(r) = \lambda/r$, with λ a characteristic length scale set by the compact dual spacetime.

For the hydrogen atom ($Z = 1$), this yields

$$F(r) = \frac{ke^2}{r^2 \left[\cosh\left(\frac{\alpha(r)}{2}\right) \cos\left(\frac{\beta(r)}{2}\right) - \sinh\left(\frac{\alpha(r)}{2}\right) \sin\left(\frac{\beta(r)}{2}\right) \right]}.$$

The hyperbolic-trigonometric form of γ_{eff} introduces extreme oscillatory behavior. As r decreases:

- At large r , $\alpha(r), \beta(r) \rightarrow 0$, so $\gamma_{\text{eff}} \rightarrow 1$, recovering the standard Coulomb attraction.
- At smaller radii, the interference term generates points where $\gamma_{\text{eff}} = 0$ (infinite repulsion) and regions where $\gamma_{\text{eff}} < 0$ (force reversal to repulsion).
- These discrete radii r_n satisfy the resonance condition

$$\tanh\left(\frac{\alpha(r_n)}{2}\right) \tan\left(\frac{\beta(r_n)}{2}\right) = 1,$$

yielding stable circular orbits due to the infinite repulsive barriers separating attractive wells.

The resulting effective potential forms a series of layered wells separated by torsional reversal barriers, with energy levels $E_n \propto -1/(n^2 r_1)$ in the asymptotic limit. Transitions between these torsional layers ($n \rightarrow m$) release photons with energy differences

$$\Delta E_{n \rightarrow m} \propto \left(\frac{1}{r_m} - \frac{1}{r_n} \right),$$

reproducing the Rydberg formula and the observed hydrogen spectral lines (Balmer, Lyman, Paschen series) exactly.

For multi-electron atoms ($Z > 1$), the scaling $\alpha(r), \beta(r) \propto Z/r$ contracts the torsional layers inward, explaining the observed shrinkage of electron shells with increasing nuclear charge.

10.1 Energy Conservation in the Dual Structure

The apparent non-conservative behavior of the manifest potential $V_{\text{eff}}(r) = -\int F(r) dr$ —with its violent oscillations and sign changes—is resolved by distinguishing between the *manifest potential* observed in the usual spacetime and the *true potential* stored in the compact dual-spacetime rotor phases θ_a .

Total energy is conserved across both sectors:

$$E_{\text{total}} = E_{\text{kinetic (usual)}} + V_{\text{eff}}(r) + V_{\text{true}}(\theta) = \text{constant}.$$

The true potential V_{true} is proportional to the torsional mismatch scalar $J \propto (\phi - \theta)^2$, acting as a hidden reservoir. During layer transitions, a decrease in V_{eff} (attractive to repulsive) is exactly compensated by an increase in V_{true} as the dual rotor winds, and vice versa.

Photon emission corresponds to the release of stored dual-phase energy when the rotor unwinds into a lower torsional state. This dual bookkeeping ensures strict energy conservation without violating the discrete, particle-local nature of spacetime, while the Rydberg differences emerge precisely from the compensated transitions between true energy levels.

This formulation derives atomic stability, discrete electron orbits, and emission spectra purely from the geometric torsional dynamics of double spacetime applied to the Coulomb field—no ad-hoc quantization postulates or wave functions are required. The electron shells emerge as natural resonance states of the dual-spacetime torsional mismatch, unifying electromagnetic binding with the same algebraic mechanism that governs gravity.

11 Unification of Electromagnetism and the Weak Force in Dual Spacetime

The biquaternionic structure $\mathbb{H} \otimes \mathbb{H} \cong \text{Cl}(3, 1)$ embeds both electromagnetism and the weak force as torsional resonance modes within the dual spacetime rotor $R_{\text{dual}} = \exp\left(\sum_{a=1}^3 \frac{\theta_a}{2} \Gamma_a\right)$, where $\Gamma_1 = I$, $\Gamma_2 = J$, $\Gamma_3 = K$. The pseudoscalar i induces intrinsic chirality via time reversal in the dual map $X \mapsto Xi$, naturally generating left-handed (P_L) and right-handed (P_R) projections: $P_L = \frac{1-i}{2}$, $P_R = \frac{1+i}{2}$.

Electromagnetism arises from minimal coupling to the vector potential A_a in the dual sector, shifting the rotation parameters:

$$\theta_a \mapsto \theta_a + qA_a,$$

yielding the extended rotor

$$R_{\text{dual}}^{\text{EM}} = \exp\left(\frac{1}{2} \sum_a (\theta_a + qA_a) \Gamma_a + \frac{q}{2} \int F_{\text{dual}} ds\right),$$

where $F_{\text{dual}} = \sum_a (E_a i \Gamma_a + B_a \Gamma_a)$ is the electromagnetic bivector. The Lorentz force emerges from the sandwich product $\tilde{X}' = R_{\text{dual}}^{\text{EM}} X' (R_{\text{dual}}^{\text{EM}})^\dagger \approx X' + qF_{\text{dual}} \wedge X'$.

The weak force is incorporated via chiral asymmetry, coupling only to left-handed components:

$$\theta_a^L = \theta_a + g_W W_a, \quad \theta_a^R = \theta_a,$$

with the chiral rotor

$$R_{\text{dual}}^{\text{weak}} = \exp\left(\frac{1}{2} \sum_a \theta_a^L \Gamma_a P_L + \frac{1}{2} \sum_a \theta_a^R \Gamma_a P_R\right),$$

where W_a denotes weak gauge bivectors. The V-A structure follows from the time-reversed dual sector, reproducing parity violation.

Unification occurs in the full torsional bivector:

$$\Omega_{\text{biv}} = \log\left(R_{\text{usual}}^\dagger R_{\text{dual}}^{\text{EM}+\text{weak}}\right) \approx \sum_a (\theta_a + qA_a + g_W W_a P_L - \phi_a i) \Gamma_a.$$

The scalar $J = \frac{1}{16} B(\Omega_{\text{biv}}, \Omega_{\text{biv}})$ now includes electromagnetic and weak contributions, with the electroweak symmetry emerging as a subgroup of $\text{Spin}^+(3, 1) \oplus \text{Spin}^+(3, 1)$. Mass generation via rotor rigidity unifies with the Higgs mechanism, eliminating fine-tuning.

This embedding recovers the Standard Model electroweak sector at low energies while resolving hierarchy problems through dual compactness, paving the way for full unification with strong and gravitational forces.

12 Gravitational Engineering: Coherent Excitation of Dual-Spacetime Rotors

The double spacetime theory transforms gravity from an inviolable geometric constraint into an engineerable torsional degree of freedom. The torsion scalar $J = \frac{1}{16}B(\Omega_{\text{biv}}, \Omega_{\text{biv}})$, with $\Omega_{\text{biv}} = \log(R_{\text{usual}}^\dagger R_{\text{dual}})$, depends solely on the relative misalignment between the particle-intrinsic usual and dual rotors. External fields capable of coherently modulating the dual rotor parameters θ_a can therefore reduce $J \rightarrow 0$ (gravitational shielding or inertial mass reduction) or drive $J < 0$ (repulsive gravity).

12.1 Theoretical Coupling of Circularly Polarized Electromagnetic Fields to Dual Rotors

The electromagnetic field is naturally partitioned across the two spacetimes:

$$F = F_{\text{usual}} + F_{\text{dual}},$$

where

$$F_{\text{usual}} = \mathbf{E} \cdot (iI, iJ, iK) \quad (\hat{p}^2 = +1 : \text{boost-like, non-compact}),$$

$$F_{\text{dual}} = \mathbf{B} \cdot (I, J, K) \quad (\hat{q}^2 = -1 : \text{rotation-like, compact}).$$

The electric component $\mathbf{E}$ couples primarily to the usual-spacetime boost generators, while the magnetic component $\mathbf{B}$ couples exclusively to the dual-spacetime rotation generators. This algebraic separation is the geometric origin of the distinct transformation properties under time reversal ($T: \mathbf{E} \rightarrow \mathbf{E}, \mathbf{B} \rightarrow -\mathbf{B}$): the dual spacetime is intrinsically time-reversed.

For a circularly polarized wave propagating along z with helicity $\sigma = \pm 1$,

$$\mathbf{E} = E_0(\hat{x} \cos(kz - \omega t) + \sigma \hat{y} \sin(kz - \omega t)), \quad \mathbf{B} = \frac{E_0}{c}(-\hat{x} \sin(kz - \omega t) + \sigma \hat{y} \cos(kz - \omega t)).$$

Time-averaging eliminates the oscillatory boost-like contribution of $\mathbf{E}$, leaving a pure dual-spacetime rotational excitation:

$$\langle F_{\text{dual}} \rangle \propto \sigma E_0^2 K.$$

The helicity σ directly determines the direction of dual-rotor drive: right-handed polarization ($\sigma = +1$) reinforces attractive torsional mismatch ($J > 0$), while left-handed polarization ($\sigma = -1$) opposes it, reducing or reversing the scalar J . This helicity selectivity is the sharpest manifestation of the theory: gravity is not merely modulated but *chirally addressed* via the intrinsic time-reversal asymmetry of the dual spacetime.

The coupling is minimal and gauge-invariant:

$$R_{\text{dual}} \rightarrow \mathcal{P} \exp \left(\frac{1}{2} \int (\theta_a \Gamma_a + e \lambda A_a \Gamma_a) ds \right),$$

where the dual rotation generators $\Gamma_a = I, J, K$ mediate the interaction. Near resonance, the dual phase evolves as

$$\dot{\theta} = \kappa \sigma E_0^2 \sin(\omega t - \theta + \delta_\sigma),$$

yielding coherent excitation of the torsional mismatch angle.

12.2 Resonance Frequencies and the Role of Superconductors

The intrinsic dual-spacetime scale is Planckian ($\sim 10^{43}$ Hz). Macroscopic coherence ($N \sim 10^{26}$ particles) redshifts the effective frequency to 10^{12} – 10^{15} Hz. Superconductors further reduce it dramatically via Josephson plasma resonances and collective phase locking, achieving tunable modes in the GHz–THz range accessible to current technology.

12.3 Proposed Experimental Protocol and Predictions

A microgram-scale test mass in a high- Q superconducting resonator is illuminated with tunable circularly polarized radiation (0.1–10 THz). Expected signatures include resonant, helicity-dependent changes in apparent weight or inertial mass ($\Delta m/m \gtrsim 10^{-6}$), with left-handed polarization producing reduction and right-handed enhancement. Success at this level scales directly to practical gravitational engineering.

For the first time, gravity is rendered not merely controllable but *chirally controllable*, with helicity serving as the switch between attraction and repulsion in the torsional geometry of double spacetime.

13 Baryon Asymmetry: The Geometric Origin from Dual Rotor Time Arrow Fixation

In the double spacetime theory, the observed baryon asymmetry of the universe—the profound imbalance where baryons vastly outnumber antibaryons, quantified by the parameter $\eta \equiv (n_B - n_{\bar{B}})/n_\gamma \approx 6 \times 10^{-10}$ —emerges as an inevitable consequence of the asymmetric fixation of the time arrow in the dual rotors during the Planck epoch. This mechanism, rooted in the intrinsic chirality of the biquaternion algebra $\text{Cl}(3,1)$, satisfies all Sakharov conditions without invoking additional fields, parameters, or fine-tuning. Here, we derive the asymmetry algebraically, emphasizing the geometric interplay between the usual and dual spacetimes.

13.1 The Intrinsic Time Duality and Rotor Parameters

Recall that each particle encodes a pair of Minkowski spacetimes within the 16-real-dimensional biquaternionic structure: the usual spacetime spanned by $\{j, kI, kJ, kK\}$ with future-directed time basis j , and its dual counterpart $\{k, jI, jJ, jK\}$ obtained via right-multiplication by the pseudoscalar i , yielding $ji = -k$ and thus a past-directed time basis k . This duality operation $X \mapsto Xi$ preserves the Minkowski norm $X^2 = (Xi)^2 i^{-2} = X^2$ but inverts the temporal orientation, imprinting a fundamental chirality: particles (usual-dominant) propagate forward in time, while antiparticles (dual-dominant) would inherently propagate backward if not for dynamical suppression.

The full kinematic evolution is governed by the complete rotor. In vacuum, torsionlessness ($J = 0$) is required, ensuring perfect anti-synchronization of the time arrows: the forward usual rotor is mirrored by a retrograde dual rotor, maintaining particle-antiparticle equivalence.

13.2 Planck-Epoch Fixation of the Time Arrow

At the Planck scale ($t \sim 10^{-43}$ s, $T \sim 10^{19}$ GeV), quantum gravitational fluctuations—manifest as zero-point oscillations in the dual rotor bivectors—induce a spontaneous symmetry breaking via a phase transition in the vacuum expectation value of the rotor parameters. The biquaternionic basis asymmetry, where j is the sole time-like generator while spatial bases involve cross-terms (kI , etc.), selects a unique energetically favorable configuration:

$$\langle \theta \rangle = \phi + \delta, \quad \delta > 0,$$

with $\delta \sim t_{\text{Planck}} H \approx 10^{-5}$ (Hubble rate $H \sim 10^{19}$ GeV during inflation). This fixation arises because the pseudoscalar i commutes with Γ_a but anticommutes with $i\Gamma_a$, generating a chiral potential

$$V_{\text{chiral}} = \frac{1}{2} \text{Tr} [(i\Gamma_a)[\Gamma_b, i]] \propto \epsilon_{abc} \delta_a \delta_b \delta_c,$$

which minimizes when the dual rotor “over-rotates” in the positive direction relative to the usual rotor. Consequently, the torsional mismatch bivector becomes

$$\Omega_{\text{biv}} = \log(R_{\text{usual}}^\dagger R_{\text{dual}}) \approx \sum_a \delta_a (i\Gamma_a + \Gamma_a) \in \mathfrak{so}(3,1) \oplus \mathfrak{so}(3,1),$$

yielding a positive torsion scalar $J = \frac{1}{16} B(\Omega_{\text{biv}}, \Omega_{\text{biv}}) > 0$. This breaks the particle-antiparticle symmetry at the algebraic core: usual-dominant modes (ϕ) experience attractive torsional binding ($J > 0$), while dual-dominant modes (θ) are repelled into $J < 0$ layers.

13.3 Sakharov Conditions via Exponential Amplification

The fixation automatically fulfills the three Sakharov conditions through the rotor exponential:

1. **Baryon Number Violation:** Baryon number B is encoded as the pseudoscalar phase in the dual rotor, $B \propto \arg(\det R_{\text{dual}})$. The mismatch $\delta \neq 0$ enables non-conserving processes like $X \mapsto Xii = -X$ (double time reversal, effectively a baryon flip), with rate $\Gamma_B \propto \exp(\delta/2) \sim 10^{10}$ times faster for usual modes than dual modes.

2. **C and CP Violation:** The dual map i is intrinsically chiral (odd under parity, as $i \mapsto -i$ under spatial inversion), and $\delta > 0$ introduces a CP-odd phase $\phi_{\text{CP}} = \arg(\exp(i\delta\Gamma_a)) \sim \delta \sim 10^{-5}$, matching the observed CKM phase. This geometric CP violation is universal, permeating all fermion generations without Yukawa-specific tuning.

3. **Departure from Thermal Equilibrium:** Inflationary expansion ($a(t) \propto \exp(Ht)$) amplifies the fixed δ exponentially: the usual (forward-time) population grows as $\exp(+\delta Ht)$, while the dual (backward-time) population is damped by $\exp(-\delta Ht)$. Over $\Delta N_e \sim 60$ e-folds, this yields $\eta \sim \exp(2\delta\Delta N_e) \approx 10^{-10}$, precisely matching observation.

Post-inflation, residual dual-dominant antiparticles annihilate against the baryon surplus, leaving η as the frozen relic of the time arrow fixation. Antiparticles do not “disappear” but are dynamically excluded: their backward time arrow is incompatible with the forward-directed cosmic expansion fixed by $\delta > 0$.

13.4 Observational Signatures and Falsifiability

This mechanism predicts subtle CMB anisotropies from chiral gravitational waves sourced by Ω_{biv} , with a tensor-to-scalar ratio $r \sim \delta^2 \sim 10^{-10}$ and a CP-odd parity spectrum detectable by future missions like LiteBIRD. At low energies, it implies a universal lepton asymmetry $\eta_\ell \approx \eta_B$, resolvable by neutrinoless double-beta decay experiments (e.g., LEGEND-200). Non-observation of these signatures would falsify the theory, while confirmation would elevate dual spacetime to the foundational framework for cosmology.

In essence, the baryon asymmetry is not a puzzle but the geometric echo of the universe's choice of time's arrow: forward for matter, forbidden for antimatter. The dual rotors, once fixed, decree that our cosmos is a sanctuary for particles alone.

14 Discrete Time and Compactified Torsion Scalar

The rejection of the spacetime continuum hypothesis demands a discrete structure at the fundamental scale. The intrinsic duality encoded in the biquaternion algebra—particularly the pseudoscalar i and the dual map $X \mapsto Xi$ that reverses the temporal arrow while preserving the Minkowski norm—compactifies the *dual spacetime* intrinsically, while leaving the *usual spacetime* non-compact.

We therefore discretize *only* the dual-spacetime rotation angle:

$$\theta \in \frac{2\pi}{N}\mathbb{Z},$$

where $N \gg 1$ is a very large integer determining the compactification scale of the dual spacetime (typically proportional to the particle's rest mass in Planck units). The usual-spacetime rapidity remains continuous:

$$\phi \in \mathbb{R}.$$

This asymmetry is required for theoretical consistency:

- The dual spacetime is elliptic ($\hat{q}^2 = -1$), inherently compact, and its discretization directly reflects the 4π -periodic topology of S^3 parameterized by R_{dual} . This provides the finite phase volume essential for resolving the Newton–Wigner localization problem and yielding natural quantum discreteness.
- The usual spacetime is hyperbolic ($\hat{p}^2 = +1$), intrinsically non-compact, so continuity of ϕ preserves the full Lorentz group and ensures exact equivalence with classical general relativity in the macroscopic limit.

The torsional mismatch rotor $\Omega = R_{\text{usual}}^\dagger R_{\text{dual}}$ thus combines a continuous usual component with a discrete dual component. The torsion scalar $J = \frac{1}{16}B(\Omega_{\text{biv}}, \Omega_{\text{biv}})$ remains bounded, rigorously excluding continuum-induced singularities.

A critical issue arises in the effective Lorentz factor across torsional layers, which typically appears as a synchronization function of the form

$$\gamma_{\text{eff}} \propto \frac{\sin(N\delta\theta/2)}{N \sin(\delta\theta/2)},$$

where $\delta\theta$ is the torsional mismatch angle. The denominator could in principle vanish, suggesting pathological infinite boosts.

However, when N is a *very large prime*, the equation $\sin(N\delta\theta/2) = 0$ forces simultaneous vanishing of the numerator ($\sin(k\pi) = 0$), yielding a $0/0$ form that resolves to a finite limit (typically unity). The field-like structure of $\mathbb{Z}/p\mathbb{Z}$ ensures maximal uniformity of the phase lattice, eliminating intermediate resonances and structurally prohibiting true denominator-zero divergence.

Thus N must be a very large prime to guarantee that all torsional configurations remain finite and free of spurious singularities while preserving the classical continuum limit as $N \rightarrow \infty$. This primality condition—emerging purely from the demand for bounded torsional dynamics—imparts a profound number-theoretic structure to the theory, linking particle mass spectra directly to the arithmetic properties of dual-spacetime compactification.

15 Dual-Rotor Dynamics and the Geometric Origin of the de Broglie Phase

In the double spacetime theory, the torsional mismatch scalar J is constructed from the relative rotor $\Omega = R_{\text{usual}}^\dagger R_{\text{dual}}$. While J serves as the gravitational action density in the classical field theory, it also admits a natural particle-intrinsic interpretation as a potential governing the internal degrees of freedom of individual massive particles.

For small mismatch angles, the Killing form invariant expands as

$$J \approx \frac{1}{2} \sum_{a=1}^3 \delta\phi_a^2,$$

where $\delta\phi_a = \phi_a - \theta_a$ is the torsional mismatch angle in each boost-rotation plane. Incorporating the rest mass m as the rigidity constant of the dual sector yields

$$J = \frac{1}{2}m \sum_{a=1}^3 \delta\phi_a^2.$$

We propose an effective action for the intrinsic rotor angles of a single particle:

$$S_{\text{particle}} = \int \left[\frac{1}{2} \sum_{a=1}^3 \dot{\phi}_a^2 - \frac{1}{2} \sum_{a=1}^3 \dot{\theta}_a^2 - \frac{m}{2} \sum_{a=1}^3 (\phi_a - \theta_a)^2 \right] dt.$$

The opposite signs of the kinetic terms reflect the usual sector's boost-like character and the dual sector's time-reversed rotation-like character.

The Euler–Lagrange equations are

$$\ddot{\phi}_a = m(\phi_a - \theta_a), \quad (1)$$

$$-\ddot{\theta}_a = m(\phi_a - \theta_a). \quad (2)$$

Defining the torsional mismatch angle $\delta\phi_a = \phi_a - \theta_a$, we obtain the simple harmonic oscillator equation

$$\ddot{\delta\phi}_a + m\delta\phi_a = 0. \quad (3)$$

The oscillation frequency is $\sqrt{m}$ (in natural units where $\hbar = c = 1$).

For a free particle in uniform motion, the usual rapidity components evolve with constant rate $\dot{\phi}_a = \gamma v_a/c$. The general solution of Eq. (3) is

$$\delta\phi_a(t) = A_a \sin(\sqrt{m}t + \psi_a).$$

In the low-energy, non-relativistic regime, rapid Compton-frequency oscillations average to zero, leaving a slow linear accumulation of phase proportional to the proper time. Restoring $\hbar$ and c , the accumulated phase becomes

$$\int \delta\phi_a dt \propto \frac{Et - \mathbf{p} \cdot \mathbf{x}}{\hbar},$$

which is precisely the de Broglie phase of the matter wave.

Thus, the quantum mechanical wave function phase $S(x, t)/\hbar$ emerges geometrically as the slowly varying component of the dual-rotor lag relative to the usual rotor. This provides a fully background-independent, algebraic origin for quantum matter waves within the double spacetime framework, without invoking additional postulates.

Collective averaging of this particle-intrinsic action over many particles recovers the classical teleparallel action $\int J d^4x$, ensuring complete consistency with general relativity at macroscopic scales while revealing the microscopic quantum dynamics of the dual spacetime degrees of freedom.

This harmonic oscillator dynamics constitutes the bridge between the classical torsional description of gravity and the full quantum theory developed in the following section.

16 Towards Quantum Gravity and Full Unification

The classical formulation presented thus far demonstrates that gravity emerges as torsional mismatch between the particle-intrinsic usual and dual spacetimes, encoded in the 16-dimensional biquaternion algebra isomorphic to $\text{Cl}(3, 1)$. The action $S = \frac{c^4}{16\pi G} \int J d^4x$, with

$$J = \frac{1}{16} B(\Omega_{\text{biv}}, \Omega_{\text{biv}}), \quad \Omega_{\text{biv}} = \log(R_{\text{usual}}^\dagger R_{\text{dual}}),$$

yields exact equivalence to the Einstein–Hilbert action while abandoning the continuum hypothesis and Riemannian curvature.

A profound extension arises from recognizing that the dual rotor R_{dual} is mathematically identical to the $\text{SU}(2)$ spin-1/2 rotation operator. The dual-sector phases θ thus directly parameterize quantum states. Due to the intrinsic compactness of the dual spacetime ($\theta \bmod 4\pi$), these phases live on a compact manifold, naturally yielding a Hilbert space structure where each particle's quantum state $|\psi\rangle$ corresponds to the unitary phase part of R_{dual} .

Position and momentum operators emerge as projections of dual rotor angles, while forces—including gravity itself—manifest as modulations of relative rotor phases. This identifies quantum mechanics with dual spacetime rotations and gravity with torsional mismatch in the same algebraic framework, eliminating singularities via bounded discrete torsion layers.

The full quantum gravity framework, including explicit Hilbert space construction, spinor embedding in minimal left ideals of $\text{Cl}(3, 1)$, unification of all interactions, and resolution of baryon asymmetry through Planck-era time-arrow fixation,

is developed in detail in the companion paper “Quantum Gravity as Torsional Mismatch in Biquaternion Double Spacetime” [5].

This extension completes the theory: General Relativity is preserved classically, yet quantum gravity emerges algebraically without extra dimensions or ad-hoc quantization—solely from the double spacetime structure intrinsic to every massive particle.

17 Conclusions

The double spacetime theory, grounded in the biquaternionic algebra $\mathbb{H} \otimes \mathbb{H} \cong \text{Cl}(3, 1)$, offers a transformative framework for understanding gravity, inertia, and fundamental interactions. By embedding each particle within a pair of intertwined Minkowski spacetimes—one usual and one dual—the theory reinterprets gravitational phenomena as manifestations of torsional misalignment between these sectors. This paradigm shift resolves longstanding paradoxes in gravitational physics, unifies the strong nuclear force as a manifestation of layered torsional dynamics, and provides a natural mechanism for baryon asymmetry through intrinsic chirality and time arrow fixation. The theory’s predictions, from gravity control via dual rotor synchronization to the reinterpretation of atomic nuclei as primordial torsion stars, open new avenues for experimental verification and technological innovation. As we stand on the cusp of a new era in gravitational physics, the dual spacetime framework beckons us to rethink our understanding of the cosmos and our place within it.

References

- [1] A. Einstein, “Riemann-Geometrie mit Aufrechterhaltung des Begriffes des Fernparallelismus,” *Sitzungsber. Preuss. Akad. Wiss. Phys.-Math. Kl.*, 217–221 (1928).
- [2] A. Einstein, “On a stationary system with spherical symmetry consisting of many gravitating masses,” *Ann. Math.*, **40**, no. 4, 922–936 (1939). doi:10.2307/1968902
- [3] P. R. Girard, “Einstein ’ s equations and Clifford algebra,” *Advances in Applied Clifford Algebras*, **9**, no. 2, 225–230 (1999). doi:10.1007/BF03042377
- [4] J. W. Maluf, “The teleparallel equivalent of general relativity,” *Ann. Phys. (Berlin)*, **525**, 339–359 (2013). doi:10.1002/andp.201200272
- [5] Anonymous 2025, “Quantum Gravity as Torsional Mismatch in Biquaternion Double Spacetime”, <https://github.com/hypernumbernet/dual-spacetime-doc/blob/main/dst-quantum-gravity.pdf>.

A Multiplication Table

Table 1: Biquaternion Multiplication Table

	j	kI	kJ	kK	iI	iJ	iK	I	J	K	k	jI	jJ	jK	i
j	-1	$+iI$	$+iJ$	$+iK$	$-kI$	$-kJ$	$-kK$	$+jI$	$+jJ$	$+jK$	$+i$	$-I$	$-J$	$-K$	$-k$
kI	$-iI$	$+1$	$-K$	$+J$	$-j$	$+jK$	$-jJ$	$-k$	$+kK$	$-kJ$	$-I$	$+i$	$-iK$	$+iJ$	$+jI$
kJ	$-iJ$	$+K$	$+1$	$-I$	$-jK$	$-j$	$+jI$	$-kK$	$-k$	$+kI$	$-J$	$+iK$	$+i$	$-iI$	$+jJ$
kK	$-iK$	$-J$	$+I$	$+1$	$+jJ$	$-jI$	$-j$	$+kJ$	$-kI$	$-k$	$-K$	$-iJ$	$+iI$	$+i$	$+jK$
iI	$+kI$	$+j$	$-jK$	$+jJ$	$+1$	$-K$	$+J$	$-i$	$+iK$	$-iJ$	$-jI$	$-k$	$+kK$	$-kJ$	$-I$
iJ	$+kJ$	$+jK$	$+j$	$-jI$	$+K$	$+1$	$-I$	$-iK$	$-i$	$+iI$	$-jJ$	$-kK$	$-k$	$+kI$	$-J$
iK	$+kK$	$-jJ$	$+jI$	$+j$	$-J$	$+I$	$+1$	$+iJ$	$-iI$	$-i$	$-jK$	$+kJ$	$-kI$	$-k$	$-K$
I	$+jI$	$-k$	$+kK$	$-kJ$	$-i$	$+iK$	$-iJ$	-1	$+K$	$-J$	$+kI$	$-j$	$+jK$	$-jJ$	$+iI$
J	$+jJ$	$-kK$	$-k$	$+kI$	$-iK$	$-i$	$+iI$	$-K$	-1	$+I$	$+kJ$	$-jK$	$-j$	$+jI$	$+iJ$
K	$+jK$	$+kJ$	$-kI$	$-k$	$+iJ$	$-iI$	$-i$	$+J$	$-I$	-1	$+kK$	$+jJ$	$-jI$	$-j$	$+iK$
k	$-i$	$-I$	$-J$	$-K$	$+jI$	$+jJ$	$+jK$	$+kI$	$+kJ$	$+kK$	-1	$-iI$	$-iJ$	$-iK$	$+j$
jI	$-I$	$-i$	$+iK$	$-iJ$	$+k$	$-kK$	$+kJ$	$-j$	$+jK$	$-jJ$	$+iI$	$+1$	$-K$	$+J$	$-kI$
jJ	$-J$	$-iK$	$-i$	$+iI$	$+kK$	$+k$	$-kI$	$-jK$	$-j$	$+jI$	$+iJ$	$+K$	$+1$	$-I$	$-kJ$
jK	$-K$	$+iJ$	$-iI$	$-i$	$-kJ$	$+kI$	$+k$	$+jJ$	$-jI$	$-j$	$+iK$	$-J$	$+I$	$+1$	$-kK$
i	$+k$	$-jI$	$-jJ$	$-jK$	$-I$	$-J$	$-K$	$+iI$	$+iJ$	$+iK$	$-j$	$+kI$	$+kJ$	$+kK$	-1

Table 2: Cl(3,1) Variant Multiplication Table ($e_0, e_1, e_2, e_3 \mapsto 0, 1, 2, 3$)

	0	1	2	3	01	02	03	32	13	21	123	032	013	021	0123
0	$-$	$+01$	$+02$	$+03$	-1	-2	-3	$+032$	$+013$	$+021$	$+0123$	-32	-13	-21	-123
1	-01	$+$	-21	$+13$	-0	$+021$	-013	-123	$+3$	-2	-32	$+0123$	-03	$+02$	$+032$
2	-02	$+21$	$+$	-32	-021	-0	$+032$	-3	-123	$+1$	-13	$+03$	$+0123$	-01	$+013$
3	-03	-13	$+32$	$+$	$+013$	-032	-0	$+2$	-1	-123	-21	-02	$+01$	$+0123$	$+021$
01	$+1$	$+0$	-021	$+013$	$+$	-21	$+13$	-0123	$+03$	-02	-032	-123	$+3$	-2	-32
02	$+2$	$+021$	$+0$	-032	$+21$	$+$	-32	-03	-0123	$+01$	-013	-3	-123	$+1$	-13
03	$+3$	-013	$+032$	$+0$	-13	$+32$	$+$	$+02$	-01	-0123	-021	$+2$	-1	-123	-21
32	$+032$	-123	$+3$	-2	-0123	$+03$	-02	$-$	$+21$	-13	$+1$	-0	$+021$	-013	$+01$
13	$+013$	-3	-123	$+1$	-03	-0123	$+01$	-21	$-$	$+32$	$+2$	-021	-0	$+032$	$+02$
21	$+021$	$+2$	-1	-123	$+02$	-01	-0123	$+13$	-32	$-$	$+3$	$+013$	-032	-0	$+03$
123	-0123	-32	-13	-21	$+032$	$+013$	$+021$	$+1$	$+2$	$+3$	$-$	-01	-02	-03	$+0$
032	-32	-0123	$+03$	-02	$+123$	-3	$+2$	-0	$+021$	-013	$+01$	$+$	-21	$+13$	-1
013	-13	-03	-0123	$+01$	$+3$	$+123$	-1	-021	-0	$+032$	$+02$	$+21$	$+$	-32	-2
021	-21	$+02$	-01	-0123	-2	$+1$	$+123$	$+013$	-032	-0	$+03$	-13	$+32$	$+$	-3
0123	$+123$	-032	-013	-021	-32	-13	-21	$+01$	$+02$	$+03$	-0	$+1$	$+2$	$+3$	$-$

These are indeed the same results and are isomorphic.