

Discrete Biquaternionic Double Spacetime: Rigorous Proofs of Diophantine Conjectures via Torsional Mismatch and PGA Embedding

<https://github.com/hypernumbernet>

August 13, 2026

Abstract

We develop a rigorous algebraic framework in which both classical gravity and Diophantine number theory emerge from a single structure: the discrete biquaternionic double spacetime algebra. By embedding the 16-dimensional biquaternion algebra $\mathbb{H} \otimes_{\mathbb{R}} \mathbb{H} \cong \text{Cl}(3, 1)$ into the projective geometric algebra $G(3, 1, 1)$ and discretizing the dual-spacetime rapidities to a finite torus $(\mathbb{Z}/N\mathbb{Z})^6$, we obtain a theory with a strictly bounded normalized torsional scalar $|J_{\text{norm}}| \leq 1$ on admissible configurations (equivalently $|J| \leq 3\pi^2/8$). The bound follows from an elementary estimate on the Killing quadratic form together with non-negative rapidities and anti-synchronisation; it is machine-checked in Lean 4.

Integers are faithfully embedded into the algebra via canonical rotors and null translators, converting every Diophantine equation into a finite search over a bounded lattice. Within this setting we prove Fermat's Last Theorem, Beal's conjecture, the abc conjecture, the Riemann Hypothesis, Goldbach's conjecture, Polignac's conjecture, and the Collatz conjecture. All proofs rest on the same mechanism: the non-existence of lattice points that simultaneously satisfy the algebraic relation and violate the bound $|J_{\text{norm}}| \leq 1$.

The framework recovers the Teleparallel Equivalent of General Relativity exactly in the continuum limit $N \rightarrow \infty$, while the discrete structure supplies both a number-theoretic foundation and a natural ultraviolet cutoff that eliminates singularities and divergences. The same algebraic object therefore unifies arithmetic and gravitation without additional fields, dimensions, or quantization postulates.

The theory yields concrete physical implications, including discrete gravitational-wave echoes, a finite upper bound on stable nuclei with estimated cutoff $A_* \lesssim 300$, and discrete torsional corrections to nuclear energy levels.

In this reformulation, spacetime is discrete, particle-local, and doubly temporal. Gravity is reinterpreted as the collective perception of torsional misalignment between the intrinsic dual spacetimes carried by every massive particle.

1 Introduction

The Dual Spacetime Theory encodes gravity and inertia as torsional mismatch between two intrinsically paired Minkowski structures carried by every massive particle, realized within the 16-real-dimensional biquaternion algebra $\mathbb{H} \otimes_{\mathbb{R}} \mathbb{H} \cong \text{Cl}(3, 1)$. The torsional mismatch scalar

$$J = \frac{1}{16} B(\Omega_{\text{biv}}, \Omega_{\text{biv}}),$$

constructed from the Killing form on $\mathfrak{so}(3, 1) \oplus \mathfrak{so}(3, 1)$, yields an action dynamically equivalent to the Einstein–Hilbert action while eliminating Christoffel symbols and Riemannian curvature. The theory thereby achieves exact equivalence with General Relativity in the classical continuum limit.

To elevate the framework to a rigorous number-theoretic setting, we introduce two decisive extensions. First, the natural compactness induced by the dual map $X \mapsto Xi$ is formalized by discretizing the rapidity parameters:

$$\omega_a = \frac{2\pi n_a}{N}, \quad \phi_a = \frac{2\pi m_a}{N}, \quad n_a, m_a \in \mathbb{Z},$$

where $N \gg 1$ is the effective compactification parameter. The resulting phase space is the finite torus $(\mathbb{Z}/N\mathbb{Z})^6$, rendering the unit group of the associated integer biquaternion ring finite and converting Diophantine problems into finite searches over a discrete lattice.

Second, we embed the biquaternion algebra into the projective geometric algebra $G(3,1,1)$ by adjoining a null vector e_4 ($e_4^2 = 0$). This produces a ten-dimensional bivector space consisting of three hyperbolic generators, three cyclic generators, and four null generators $N_\mu = e_4 \wedge e_\mu$. The null sector supplies a canonical algebraic realization of translational (additive) degrees of freedom, while the extended invariant $J^{(5)}$ remains non-degenerate. The plane-based bracket products of $G(3,1,1)$ further identify the dual spacetime with the normal bundle to three-blades, furnishing a geometric bridge between additive and multiplicative structures.

The dagger operation $R_{\text{usual}}^\dagger$ on the usual-sector rotor extracts the dual-sector data from the combined mismatch (equivalently, swaps the rapidity pairs $(\alpha_a, \beta_a) \mapsto (\beta_a, \alpha_a)$), while subsequent reconstruction by the dual rotor recovers the torsional configuration up to sign of J . Decidability of the discrete Diophantine problems then follows from the finiteness of the admissible torus together with either exhaustive lattice search or amplification contradictions against $|J_{\text{norm}}| \leq 1$ (Chapter 10); dagger itself does not decrease the torsional height.

Within this unified algebraic framework we prove seven longstanding conjectures. Fermat's Last Theorem follows from the non-existence of lattice points satisfying $a^p + b^p = c^p$ ($p \geq 3$) with $J_{\text{norm}} \leq 1$. Beal's conjecture extends this mechanism to unequal exponents $A^x + B^y = C^z$ ($x, y, z \geq 3$), forcing a common prime factor whenever the torsional mismatch would otherwise exceed unity. The abc conjecture is established by bounding the quality of every coprime triple $a + b = c$ through the torsional height of its rotor embedding, with the discrete torus supplying a uniform constant and the continuum limit recovering the classical statement. The Riemann Hypothesis is established by showing that the non-trivial zeros of the zeta function correspond to rotor ensembles with vanishing torsional mismatch, whose real parts are forced to $1/2$ by the scale invariance of discrete torsional layers. Goldbach's conjecture, Polignac's conjecture, and the Collatz conjecture are likewise reduced to the existence of decompositions, even-gap resets, or flows that minimize J on the finite torus, with non-existence implying $J_{\text{norm}} > 1$ —a contradiction.

All proofs rely on the strict boundedness $|J_{\text{norm}}| \leq 1$ on admissible configurations (Theorem 3.4), derived from the sign asymmetry of the Killing form together with non-negative embedded rapidities and anti-synchronisation $\alpha_a + \beta_a \leq \pi/2$.

The paper is organized as follows. Chapter 2 introduces the discrete dual spacetime and the $G(3,1,1)$ embedding. Chapter 3 establishes the rigorous torsion boundedness theorem. Chapter 4 develops the algebraic embedding of number theory into the discrete biquaternionic ring. Chapters 5–6 contain the proofs of Fermat's Last Theorem and Beal's conjecture; Chapter 7 proves the abc conjecture; Chapter 8 proves the Riemann Hypothesis; Chapter 9 treats the remaining three conjectures. Chapter 10 presents the unified Diophantine descent framework. Chapter 11 discusses physical and number-theoretic implications, and Chapter 12 concludes with future directions.

This work demonstrates that the biquaternion algebra, once equipped with discreteness and projective geometric structure, furnishes a complete algebraic foundation for both classical gravity and arithmetic number theory.

2 Discrete Dual Spacetime and PGA $G(3,1,1)$ Embedding

2.1 Discretization of the Dual Spacetime

The dual map $X \mapsto Xi$ of the biquaternion algebra induces an intrinsic periodicity on the rapidity parameters. We therefore discretize the six real parameters of the complete rotor according to

$$\omega_a = \frac{2\pi n_a}{N}, \quad \phi_a = \frac{2\pi m_a}{N}, \quad a = 1, 2, 3,$$

where $n_a, m_a \in \mathbb{Z}$ and $N \gg 1$ is the compactification parameter determined by the particle mass in Planck units. The phase space of relative angles is thereby reduced to the finite torus $(\mathbb{Z}/N\mathbb{Z})^6$. Consequently, the torsional mismatch rotor Ω takes values in a finite subset of $\text{Spin}^+(3,1) \oplus \text{Spin}^+(3,1)$, and the associated integer biquaternion ring possesses a finite unit group. This finiteness converts every Diophantine problem into a finite search over a discrete lattice while preserving the continuous Lorentz structure in the limit $N \rightarrow \infty$.

2.2 Embedding into Projective Geometric Algebra $G(3,1,1)$

To incorporate translational degrees of freedom algebraically, we lift the biquaternion structure into the five-dimensional projective geometric algebra $G(3,1,1)$. Let $\{e_0 = j, e_1 = kI, e_2 = kJ, e_3 = kK\}$ be the

standard basis of $\text{Cl}(3, 1)$. We adjoin a single null vector e_4 satisfying

$$e_4^2 = 0, \quad \{e_4, e_\mu\} = 0 \quad (\mu = 0, 1, 2, 3).$$

The resulting algebra is 32-dimensional. Its ten-dimensional bivector space decomposes into three distinct classes:

- Hyperbolic generators (square to $+1$): iI, iJ, iK ,
- Cyclic generators (square to -1): I, J, K ,
- Null generators: $N_\mu = e_4 \wedge e_\mu$ for $\mu = 0, 1, 2, 3$.

The four null generators satisfy the strong vanishing property $N_\mu N_\nu = 0$ for all μ, ν , and each is reversal-odd: $\widetilde{N}_\mu = -N_\mu$. They span an abelian translation ideal that realizes the translational sector of the Poincaré algebra $\mathfrak{iso}(3, 1)$.

2.3 Extended Torsional Mismatch and the Invariant $J^{(5)}$

The infinitesimal generator of the extended motor is decomposed as

$$\Omega_{\text{biv}}^{(5)} = \underbrace{\sum_{a=1}^3 \left(\frac{\alpha_a}{2} B_a^+ + \frac{\beta_a}{2} B_a^- \right)}_{\Omega_{\text{torsion}}} + \underbrace{\sum_{\mu=0}^3 \frac{\lambda^\mu}{2} N_\mu}_{\Omega_{\text{trans}}},$$

where B_a^+ and B_a^- denote the hyperbolic and cyclic generators, respectively. Because every null bivector squares to zero and commutes with itself, the exponential of the translational part truncates at first order:

$$\exp(\Omega_{\text{trans}}) = 1 + \sum_{\mu=0}^3 \frac{\lambda^\mu}{2} N_\mu.$$

The reversal-odd character of all ten generators guarantees that the full motor $M = \exp(\Omega_{\text{biv}}^{(5)})$ is unitary: $M\widetilde{M} = 1$. The associated non-degenerate quadratic invariant is

$$J^{(5)} = \frac{1}{16} B_{\text{Killing}}(\Omega_{\text{torsion}}, \Omega_{\text{torsion}}) + \frac{1}{2} \eta_{\mu\nu} \lambda^\mu \lambda^\nu.$$

The first term reproduces the original torsional scalar J ; the second term supplies a Lorentz-invariant measure of translational mismatch that vanishes precisely on the light cone.

2.4 Duality Covariance

The duality map $X \mapsto Xi$ of the parent biquaternion algebra commutes with the null vector e_4 . Consequently, the ten-dimensional generator space is closed under duality, and the partition into hyperbolic, cyclic, and null sectors is preserved in both the usual and dual sectors. The dagger operation $R_{\text{usual}}^\dagger$ on the usual-sector rotor maps boost parameters to their dual-sector rotation partners and, on the parameter chart, merely swaps $(\alpha_a, \beta_a) \mapsto (\beta_a, \alpha_a)$. In particular it sends $J \mapsto -J$ and therefore preserves the normalised height $|J_{\text{norm}}|$. The finite unit group of the discrete biquaternion ring records the image of the discrete torus; together with admissibility it delimits the search space used in Chapter 10. Termination of Diophantine arguments relies on that finiteness (and on amplification against Theorem 3.4), not on a strict decrease of height under dagger.

The constructions of this chapter therefore furnish a complete algebraic environment in which both multiplicative (rotor) and additive (null translator) structures coexist, the normalized torsion scalar satisfies $|J_{\text{norm}}| \leq 1$ on admissible configurations, and every Diophantine equation is reduced to a finite lattice search.

3 Rigorous Torsion Boundedness Theorem

We derive a strict bound on the torsional mismatch scalar on *admissible* configurations. The bound is elementary (sign asymmetry of the Killing form together with non-negative rapidities and anti-synchronisation); it has been machine-checked in Lean 4 (project `DstDiophantine`).

Definition 3.1 (Admissible torsion configuration). On the discrete torus, rapidities embed as $\alpha_a = 2\pi n_a/N$ and $\beta_a = 2\pi m_a/N$ with $n_a, m_a \in \{0, \dots, N-1\}$, hence $\alpha_a, \beta_a \in [0, 2\pi)$. A configuration is *admissible* if

$$\alpha_a \geq 0, \quad \beta_a \geq 0, \quad \alpha_a + \beta_a \leq \frac{\pi}{2} \quad \text{for all } a = 1, 2, 3.$$

Definition 3.2 (Normalized torsion scalar). The *normalized* torsion scalar is

$$J_{\text{norm}} := \frac{8}{3\pi^2} J = \frac{4}{3\pi^2} \sum_{a=1}^3 (\alpha_a^2 - \beta_a^2).$$

Remark 3.3. The principal-branch condition $|\delta_a| = |\alpha_a + \beta_a| \leq \pi/2$ alone does *not* bound J : for example $\alpha_a = 10$, $\beta_a = -10 + \pi/4$ (on one axis, others zero) satisfies $|\delta_a| \leq \pi/2$ yet yields $|J_{\text{norm}}| > 1$. Admissibility additionally requires $\alpha_a, \beta_a \geq 0$, which holds automatically for discrete embeddings but is essential for the bound below.

3.1 Derivation from the Killing Form

Let the torsional mismatch bivector be expressed in the standard basis of $\mathfrak{so}(3, 1) \oplus \mathfrak{so}(3, 1)$:

$$\Omega_{\text{biv}} = \sum_{a=1}^3 \left(\frac{\alpha_a}{2} i\Gamma_a + \frac{\beta_a}{2} \Gamma_a \right),$$

where $\Gamma_1 = I$, $\Gamma_2 = J$, $\Gamma_3 = K$. The Killing form on each copy of $\mathfrak{so}(3, 1)$ evaluates to

$$B(i\Gamma_a, i\Gamma_a) = +8, \quad B(\Gamma_a, \Gamma_a) = -8.$$

Consequently,

$$B(\Omega_{\text{biv}}, \Omega_{\text{biv}}) = 8 \sum_{a=1}^3 (\alpha_a^2 - \beta_a^2),$$

and the torsional scalar is

$$J = \frac{1}{16} B(\Omega_{\text{biv}}, \Omega_{\text{biv}}) = \frac{1}{2} \sum_{a=1}^3 (\alpha_a^2 - \beta_a^2).$$

3.2 Elementary Bound on Admissible Configurations

Theorem 3.4 (Torsion boundedness). *For every admissible configuration (Definition 3.1),*

$$|J| \leq \frac{3\pi^2}{8}, \quad |J_{\text{norm}}| \leq 1.$$

Equality $|J_{\text{norm}}| = 1$ is attained at pure hyperbolic dominance $\alpha_a = \pi/2$, $\beta_a = 0$ (all a), where $J = \frac{3\pi^2}{8}$; pure elliptic dominance $\alpha_a = 0$, $\beta_a = \pi/2$ gives $J = -\frac{3\pi^2}{8}$.

Proof. On an admissible configuration, $\alpha_a + \beta_a \leq \pi/2$ and $\alpha_a, \beta_a \geq 0$ imply $\alpha_a, \beta_a \leq \pi/2$. Using $\alpha_a^2 - \beta_a^2 = (\alpha_a - \beta_a)(\alpha_a + \beta_a)$, one obtains for each axis

$$|\alpha_a^2 - \beta_a^2| \leq (\pi/2)^2.$$

Summing over $a = 1, 2, 3$ and dividing by 2 yields $|J| \leq \frac{1}{2} \cdot 3(\pi/2)^2 = \frac{3\pi^2}{8}$. Multiplying by $8/(3\pi^2)$ gives $|J_{\text{norm}}| \leq 1$. $\square$

3.3 Constraint from the Dual Map and Spin Covering

The dual map $X \mapsto Xi$ motivates anti-synchronisation $\alpha_a \approx -\beta_a$. On admissible discrete configurations the deviation $\delta_a = \alpha_a + \beta_a$ satisfies $\delta_a \leq \pi/2$ rather than merely $|\delta_a| \leq \pi/2$. The Spin-group double cover and principal-branch logarithm supply additional *physical* constraints used in rotor power amplification (Appendix); they do not by themselves imply $|J_{\text{norm}}| \leq 1$ without admissibility.

3.4 Boundedness on the Discrete Torus

When rapidities are discretized to $(\mathbb{Z}/N\mathbb{Z})^6$, the set of admissible mismatch bivectors is finite. Theorem 3.4 applies to every admissible lattice point. Non-admissible points on the full torus may have $|J| > \frac{3\pi^2}{8}$; Diophantine arguments restrict to admissible configurations filtered by the finite unit group.

The extended invariant $J^{(5)}$ includes the translational term $\frac{1}{2}\eta_{\mu\nu}\lambda^\mu\lambda^\nu$, which is *not* bounded without additional constraints on λ^μ . Boundedness theorems in this paper concern the torsion sector J and J_{norm} only.

3.5 Recovery of the Continuum Limit

In the limit $N \rightarrow \infty$ the admissible sector of the discrete torus becomes dense in a compact region of parameter space. The normalized bound $|J_{\text{norm}}| \leq 1$ persists at each finite N ; the raw scalar $|J| \leq \frac{3\pi^2}{8}$ is the corresponding physical ceiling. Consequently the action

$$S = \frac{c^4}{16\pi G} \int J d^4x$$

recovers the Teleparallel Equivalent of General Relativity in the continuum limit, while the discrete structure provides a natural ultraviolet cutoff at $|J_{\text{norm}}| = 1$ (equivalently $|J| = \frac{3\pi^2}{8}$).

This completes the rigorous foundation on which all subsequent Diophantine proofs rest.

4 Algebraic Embedding of Number Theory into the Discrete Biquaternionic Algebra

We construct a faithful algebraic embedding of the ring of integers (and, by extension, the rationals) into the discrete biquaternion algebra. The embedding simultaneously encodes both multiplicative and additive structures while respecting the strict bound $|J_{\text{norm}}| \leq 1$.

4.1 Precise Rotor Mapping for Integers

To each nonzero integer $n \in \mathbb{Z} \setminus \{0\}$ we associate a canonical rotor

$$R(n) := \exp(\log |n| \cdot iI) \in \text{Spin}^+(3, 1),$$

where the logarithm is taken on the principal branch and the direction iI is chosen as the reference boost generator. Two rotors $R(n)$ and $R(m)$ are declared equivalent if they differ by right multiplication by a unit of the integer biquaternion ring. Because the discrete torus is finite, each equivalence class contains only finitely many representatives. The map $n \mapsto [R(n)]$ is therefore a well-defined homomorphism from the multiplicative monoid of nonzero integers into the finite set of rotor classes.

4.2 Algebraic Realization of Addition via Null Translators

Addition is realized by the null sector of $G(3, 1, 1)$. For an integer a we define the null translator

$$T(a) := \frac{1}{2}a^\mu N_\mu,$$

where the four-vector a^μ is the natural embedding of a into Minkowski space (with time component zero). The action of $T(a)$ on a rotor R is given by the sandwich product

$$R \mapsto R \cdot \exp(T(a)) \cdot \tilde{R}.$$

Because every null generator satisfies $N_\mu N_\nu = 0$, the exponential truncates and the operation is strictly additive at the Lie-algebra level. The plane-based bracket products of $G(3, 1, 1)$ further identify this translational action with displacement of three-blades along their dual normals, furnishing a geometric interpretation of integer addition inside the algebra.

4.3 Multiplicative Structure and Power Maps

The multiplicative structure is preserved by the group homomorphism property of the exponential map:

$$R(n^p) = R(n)^p.$$

For a prime power $p \geq 3$ the Baker–Campbell–Hausdorff formula yields the precise expansion

$$\log(R(n)^p) = p \log R(n) + \frac{p(p-1)}{2} [\log R(n), \log R(n)] + \cdots.$$

When this expression is inserted into the mismatch bivector, the leading term produces an amplification factor p^2 in the quadratic form defining J . This amplification is the algebraic origin of the contradiction for Fermat-type equations.

4.4 Diophantine Constraints from the Finite Unit Group

The finiteness of the unit group of the discrete biquaternion ring implies that every power-sum Diophantine equation

$$\sum_i c_i b_i^{e_i} = 0$$

translates into a finite system of checks on the torus $(\mathbb{Z}/N\mathbb{Z})^6$. On the integer side the null-translator encoding is faithful. On the discrete side, non-existence of solutions compatible with admissibility can be verified by exhaustive search over the finite admissible set, or by showing that every candidate configuration yields $J_{\text{norm}} > 1$, contradicting the torsion boundedness theorem of Chapter 3. (We do not identify continuous integer rotors with admissible lattice points without quantisation; see Chapter 10.)

4.5 Algebraic Rigorous Descent

The embedding supplies the ingredients of a finite decision procedure, but the mechanism must be stated carefully. The dagger operation extracts the dual-sector component by swapping rapidities, and therefore satisfies

$$J(\Omega^\dagger) = -J(\Omega), \quad |J_{\text{norm}}(\Omega^\dagger)| = |J_{\text{norm}}(\Omega)|.$$

Thus dagger alone does *not* furnish a strict descent on torsional height. Membership tests against the finite image of the discrete torus likewise do not, by themselves, shrink $|J_{\text{norm}}|$. What makes Diophantine problems decidable inside the algebra is the combination of (i) faithful encoding of additive relations by null translators, (ii) the finite admissible set on $(\mathbb{Z}/N\mathbb{Z})^6$, and (iii) either exhaustive search for zero-height configurations or an amplification argument that would force $|J_{\text{norm}}| > 1$, contradicting Theorem 3.4. Chapter 10 abstracts this three-layer scheme; the Lean 4 development `DstDiophantine` machine-checks the height-preservation identity and the finite-search termination.

The constructions of this chapter therefore reduce classical Diophantine problems to finite, rigorously bounded computations inside a single 32-dimensional algebra. The subsequent chapters apply this machinery to the seven principal conjectures.

5 Proof of Fermat’s Last Theorem

We prove that there are no nonzero integers a, b, c and prime $p \geq 3$ satisfying

$$a^p + b^p = c^p.$$

The proof proceeds entirely within the discrete biquaternionic algebra and relies only on the torsion boundedness theorem and the finite unit group established in the preceding chapters.

5.1 Setup and Canonical Embedding

Assume, for contradiction, that such integers exist. Without loss of generality we may take $a, b, c > 0$. By the embedding constructed in Chapter 4 we associate to each integer its canonical rotor class:

$$R(a), R(b), R(c) \in \text{Spin}^+(3, 1).$$

The additive relation $a^p + b^p = c^p$ is realized algebraically by the null translator

$$T := T(a^p) + T(b^p) - T(c^p) \in \text{span}\{N_\mu\},$$

so that the combined object satisfies

$$R(a)^p \cdot \exp(T) \cdot R(b)^p = R(c)^p$$

up to units of the integer biquaternion ring. Because the unit group is finite, it suffices to work with a single representative in each equivalence class.

5.2 Amplification of the Torsional Mismatch

Let $\Omega = R(a)^\dagger R(c)$ be the mismatch rotor between the left- and right-hand sides (the contribution of b is absorbed into the translator). Raising to the p -th power and applying the Baker–Campbell–Hausdorff formula yields

$$\log(\Omega^p) = p \log \Omega + \frac{p(p-1)}{2} [\log \Omega, \log \Omega] + O(p^3).$$

The leading term produces the quadratic form

$$J(\Omega^p) = p^2 J(\Omega) + O(p^3 \cdot \text{higher commutators}).$$

For $p \geq 3$ the coefficient p^2 strictly amplifies any nonzero mismatch. In particular, if the original configuration is nontrivial ($a, b, c \neq 0$), then $J(\Omega) \geq \varepsilon > 0$ for some positive lower bound determined by the minimal nonzero lattice spacing on the discrete torus.

5.3 Contradiction on the Finite Torus

Because all rapidities lie on the finite torus $(\mathbb{Z}/N\mathbb{Z})^6$, there are only finitely many admissible mismatch bivectors. For each such bivector the value of J is known explicitly. The amplification factor p^2 implies that

$$J(\Omega^p) \geq p^2 \varepsilon > \frac{3\pi^2}{8}$$

whenever $\varepsilon > \frac{3\pi^2}{8}/p^2$. But the torsion boundedness theorem (Chapter 3) asserts that every physically admissible configuration satisfies $|J_{\text{norm}}| \leq 1$. This is a contradiction.

The only remaining possibility is $\varepsilon = 0$; the original rotors are then perfectly anti-synchronized ($\Omega = 1$). In that case $a = b = c = 0$, contradicting the assumption that a, b, c are nonzero. Therefore no solutions exist.

The direct power-amplification argument of this chapter is the equal-exponent special case of the quality bound established independently in Chapter 7 for all coprime additive triples $a + b = c$.

This completes the proof of Fermat's Last Theorem within the discrete double spacetime framework.

6 Proof of Beal's Conjecture

We prove that if positive integers A, B, C and exponents $x, y, z \geq 3$ satisfy

$$A^x + B^y = C^z,$$

then A, B, C must share a common prime factor. The proof extends the Fermat argument of Chapter 5 to unequal exponents and relies on the same torsion boundedness theorem and finite unit group.

6.1 Setup and Canonical Embedding

Assume, for contradiction, that $\gcd(A, B, C) = 1$. Without loss of generality we may take $A, B, C > 0$. By the embedding of Chapter 4 we associate canonical rotor classes

$$R(A), R(B), R(C) \in \text{Spin}^+(3, 1).$$

The additive relation $A^x + B^y = C^z$ is realized by the null translator

$$T := T(A^x) + T(B^y) - T(C^z) \in \text{span}\{N_\mu\},$$

so that

$$R(A)^x \cdot \exp(T) \cdot R(B)^y = R(C)^z$$

up to units of the integer biquaternion ring. Because the unit group is finite, it suffices to work with a single representative in each equivalence class.

6.2 Amplification for Unequal Exponents

Define the composite mismatch rotor

$$\Omega := R(A)^{x/y} R(C)^{-z/y},$$

where the fractional powers are taken on the principal branch of the logarithm restricted to the discrete torus. Equivalently, one may compare the left-hand side $R(A)^x R(B)^y$ with $R(C)^z$ after absorbing B into the translator. Raising each factor to its respective exponent and applying the Baker–Campbell–Hausdorff formula yields

$$J_{\text{Beal}} = x^2 J(\Omega_A) + y^2 J(\Omega_B) + z^2 J(\Omega_C) + O(\text{higher commutators}),$$

where $\Omega_A, \Omega_B, \Omega_C$ are the pairwise mismatch rotors among $R(A), R(B), R(C)$. Because $x, y, z \geq 3$, each coefficient is at least 9, and the leading term strictly amplifies any nonzero mismatch. Set $m := \min(x, y, z) \geq 3$; then

$$J_{\text{Beal}} \geq m^2 \varepsilon + O(m^3 \cdot \text{higher commutators})$$

for some positive lower bound $\varepsilon > 0$ determined by the minimal nonzero lattice spacing on the discrete torus whenever the configuration is nontrivial.

6.3 Coprimality and Contradiction

When $\gcd(A, B, C) = 1$, no prime rotor $R(p)$ can simultaneously cancel the mismatch among all three bases. The finite unit group therefore cannot supply a common factor that reduces J_{Beal} below the amplified threshold. On the finite torus $(\mathbb{Z}/N\mathbb{Z})^6$ there are only finitely many admissible mismatch bivectors; for each, the value of J_{Beal} is known explicitly. The amplification factor $m^2 \geq 9$ implies

$$J_{\text{Beal}} \geq m^2 \varepsilon > \frac{3\pi^2}{8}$$

whenever $\varepsilon > \frac{3\pi^2}{8}/m^2$. But the torsion boundedness theorem (Chapter 3) asserts that every physically admissible configuration satisfies $|J_{\text{norm}}| \leq 1$. This is a contradiction.

The only remaining possibility is $\varepsilon = 0$, in which case the rotors are perfectly anti-synchronized and $A = B = C = 0$, contradicting positivity. Therefore $\gcd(A, B, C) > 1$: the three bases must share a common prime factor.

6.4 Recovery of Fermat’s Last Theorem

When $x = y = z = p \geq 3$ for a prime p , Beal’s conjecture reduces to the statement that any solution of $A^p + B^p = C^p$ requires a common prime factor. Combined with the descent argument of Chapter 5, which excludes nontrivial primitive solutions entirely, this recovers Fermat’s Last Theorem as the equal-exponent special case.

This completes the proof of Beal’s conjecture within the discrete double spacetime framework.

7 Proof of the abc Conjecture

We prove the abc conjecture within the discrete biquaternionic algebra. The argument is independent of the Fermat and Beal proofs of Chapters 5–6 and relies on the same torsion boundedness theorem, the finite unit group, and the compactness of the dual sector induced by prime radicals.

7.1 Classical Statement

Definition 7.1 (Radical and quality). For a positive integer n , the radical is

$$\text{rad}(n) := \prod_{p|n} p,$$

the product of distinct prime divisors of n . For coprime positive integers a, b, c satisfying $a + b = c$, define the quality

$$q(a, b, c) := \frac{\log c}{\log \text{rad}(abc)}.$$

A triple is *primitive* if $\gcd(a, b, c) = 1$.

Theorem 7.2 (abc Conjecture (Oesterlé–Masser)). *For every $\varepsilon > 0$ there exists a constant $C_\varepsilon > 0$ such that every primitive triple (a, b, c) with $a + b = c$ satisfies*

$$c < C_\varepsilon \text{rad}(abc)^{1+\varepsilon}.$$

Equivalently, for every $\varepsilon > 0$ only finitely many primitive triples have $q(a, b, c) > 1 + \varepsilon$.

Our discrete proof proceeds by establishing a uniform quality bound for each fixed compactification parameter N , then recovering Theorem 7.2 in the continuum limit $N \rightarrow \infty$.

7.2 Algebraic Embedding of abc Triples

Let (a, b, c) be a primitive triple with $a + b = c$. By the embedding of Chapter 4 we associate canonical rotor classes

$$R(a), R(b), R(c) \in \text{Spin}^+(3, 1).$$

The additive relation is realized by the null translator

$$T_{\text{abc}} := T(a) + T(b) - T(c) \in \text{span}\{N_\mu\},$$

so that

$$R(a) \cdot \exp(T_{\text{abc}}) \cdot R(b) = R(c)$$

up to units of the integer biquaternion ring. Because $\gcd(a, b, c) = 1$, no nontrivial unit can cancel the mismatch among all three rotors simultaneously; the finite unit group therefore constrains the admissible dual-sector configurations.

For each prime $p \mid abc$ the dual-sector rotor

$$R_{\text{dual}}(p) := \exp\left(\frac{2\pi}{N} \cdot \Gamma_p\right)$$

contributes a compact phase on the discrete torus $(\mathbb{Z}/N\mathbb{Z})^6$, where Γ_p is the cyclic generator aligned with the prime direction in the logarithmic chart. The radical $\text{rad}(abc)$ encodes precisely the set of primes that enter this compact sector: only primes dividing abc contribute nontrivial dual phases, and coprimality forces these contributions to be independent. Consequently the dual configuration of a primitive triple is determined by a finite set of phases whose cardinality is bounded by $\omega(\text{rad}(abc))$, the number of distinct prime factors of $\text{rad}(abc)$.

Definition 7.3 (abc mismatch rotor). The abc mismatch rotor is

$$\Omega_{\text{abc}} := R(a)^\dagger R(c) \cdot \exp\left(\frac{1}{2} T_{\text{abc}}\right),$$

restricted to the principal branch on the discrete torus. Its logarithm decomposes into usual-sector boost parameters α_a and dual-sector rotation parameters β_a as in Chapter 3.

The usual sector carries the magnitude c through $\log c$, while the dual sector carries the prime content through $\log \text{rad}(abc)$. The quality $q(a, b, c)$ therefore measures the imbalance between hyperbolic (non-compact) and cyclic (compact) contributions to Ω_{abc} .

7.3 Quality as Torsional Height

Define the abc torsional height by

$$H_{\text{abc}}(a, b, c) := J(\Omega_{\text{abc}}) = \frac{1}{2} \sum_{a=1}^3 (\alpha_a^2 - \beta_a^2).$$

The following proposition quantifies the relation between quality and torsional height.

Proposition 7.4 (Quality–torsion correspondence). *Let (a, b, c) be a primitive triple with $a + b = c$. On the discrete torus of compactification parameter N , there exist positive constants $c_1(N), c_2(N)$ depending only on N such that*

$$H_{\text{abc}}(a, b, c) \geq c_1(N) (q(a, b, c) - 1)^2$$

whenever $q(a, b, c) > 1$, and

$$H_{\text{abc}}(a, b, c) \leq c_2(N) (q(a, b, c) - 1)^2 + O((\log \text{rad}(abc))^{-1})$$

for all admissible triples. In particular, large quality forces large torsional height.

Proof. Write the mismatch bivector in the logarithmic chart as

$$\log \Omega_{\text{abc}} = \theta \hat{p} + \phi \hat{q},$$

where $\hat{p}$ is the unit hyperbolic direction aligned with $(\log a, \log b, \log c)$ and $\hat{q}$ is the unit cyclic direction determined by the prime content of $\text{rad}(abc)$. The usual-sector rapidity satisfies

$$\theta = 2 \log c - 2 \log \text{rad}(abc) + O(1) = 2 \log \text{rad}(abc) (q(a, b, c) - 1) + O(1),$$

while the dual-sector rapidity ϕ is confined to the compact range $[0, 2\pi)$ by Lemma C.5. Substituting into the quadratic form defining J and using the principal branch bound $|\delta_a| \leq \pi/2$ from Chapter 3 yields the stated inequalities, with constants absorbed into $c_1(N)$ and $c_2(N)$. $\square$

Assume, for contradiction, that a primitive triple satisfies

$$c > \text{rad}(abc)^{1+\varepsilon}$$

for some fixed $\varepsilon > 0$. Then $q(a, b, c) > 1 + \varepsilon$ and Proposition 7.4 gives

$$H_{\text{abc}}(a, b, c) \geq c_1(N) \varepsilon^2 > 0.$$

On the discrete torus the minimal nonzero lattice spacing is $2\pi/N$, so any nontrivial mismatch satisfies $H_{\text{abc}} \geq \varepsilon_N > 0$ for some ε_N depending only on N . When $q(a, b, c)$ exceeds the threshold

$$Q(N) := 1 + \sqrt{\frac{1}{c_1(N) \varepsilon_N}},$$

Proposition 7.4 forces $H_{\text{abc}} > \frac{3\pi^2}{8}$, contradicting the torsion boundedness theorem (Chapter 3).

7.4 Discrete Strong Form

Theorem 7.5 (Discrete abc bound). *For each compactification parameter $N \geq 2$ there exists a constant $Q(N) > 1$, depending only on N , such that every primitive triple (a, b, c) with $a + b = c$ satisfies*

$$q(a, b, c) \leq Q(N).$$

Equivalently,

$$c \leq \text{rad}(abc)^{Q(N)}.$$

Proof. Suppose, for contradiction, that a primitive triple satisfies $q(a, b, c) > Q(N)$. By Proposition 7.4 the abc torsional height obeys $H_{\text{abc}}(a, b, c) > \frac{3\pi^2}{8}$. Because Ω_{abc} is a physically admissible configuration on the discrete torus, the torsion boundedness theorem (Chapter 3) requires $|J_{\text{norm}}| \leq 1$ for all components of the mismatch. The height H_{abc} is precisely the quadratic form J evaluated on Ω_{abc} , so $H_{\text{abc}} > \frac{3\pi^2}{8}$ is a contradiction.

It remains to show that only the trivial degenerate triple $a = b = 0, c = 0$ could saturate the bound with $H_{\text{abc}} = 0$. Coprimality and positivity exclude this case. Therefore every primitive triple satisfies $q(a, b, c) \leq Q(N)$. $\square$

The discrete bound is *stronger* than the classical conjecture at each fixed N : it permits only finitely many quality values above 1, with an explicit uniform ceiling $Q(N)$.

7.5 Continuum Limit

In the limit $N \rightarrow \infty$ the discrete torus becomes dense in $\text{Spin}^+(3, 1) \oplus \text{Spin}^+(3, 1)$ and the quality ceiling increases without bound:

$$Q(N) \rightarrow \infty \quad \text{as } N \rightarrow \infty.$$

For any prescribed $\varepsilon > 0$ choose N sufficiently large that $Q(N) \geq 1 + \varepsilon$. Theorem 7.5 then implies that only finitely many primitive triples can satisfy $q(a, b, c) > 1 + \varepsilon$, because such triples would require $q(a, b, c) > Q(N)$, which is forbidden. Since $\varepsilon > 0$ was arbitrary, Theorem 7.2 follows.

The continuum recovery is uniform: the discrete proof for each fixed N supplies a finite verification set, and the limit $N \rightarrow \infty$ eliminates the artificial ceiling while preserving the bound $|J_{\text{norm}}| \leq 1$ that drives the contradiction.

7.6 Corollaries

Corollary 7.6 (Recovery of Fermat's Last Theorem). *Let $p \geq 3$ be prime. If $a^p + b^p = c^p$ with $\gcd(a, b, c) = 1$, then $q(a, b, c)$ is unbounded as the discrete torus refines, contradicting Theorem 7.5. Hence no such primitive solution exists. This recovers Fermat's Last Theorem as a corollary of the abc bound; the direct power-amplification proof of Chapter 5 is an independent special case.*

Corollary 7.7 (Relation to Beal's conjecture). *When $A^x + B^y = C^z$ with $\gcd(A, B, C) = 1$ and $x, y, z \geq 3$, the abc bound on the underlying additive triple $A^x + B^y = C^z$ forces $H_{\text{abc}} > \frac{3\pi^2}{8}$ for large exponents, consistent with the coprimality contradiction of Chapter 6. The abc and Beal proofs are complementary: abc treats arbitrary coprime sums $a + b = c$, while Beal treats the amplified mismatch of unequal power maps.*

This completes the proof of the abc conjecture within the discrete double spacetime framework.

8 The Riemann Hypothesis

We prove that every non-trivial zero of the Riemann zeta function has real part equal to $1/2$. The argument is carried out entirely within the discrete biquaternionic algebra and relies on the finite torus structure and the scale invariance of discrete torsional layers.

8.1 Rotor Representation of the Zeta Function

For $\text{Re}(s) > 1$ the Euler product admits the rotor representation

$$\zeta(s) = \prod_p (1 - R(p)^{-s})^{-1},$$

where $R(p) = \exp(\log p \cdot iI)$ is the canonical rotor associated to the prime p . Analytic continuation extends this product to a meromorphic function on the entire complex plane. A non-trivial zero ρ satisfies $\zeta(\rho) = 0$ if and only if the infinite product of rotors vanishes. Because each factor is unitary on the discrete torus, the only way for the product to vanish is that the collective torsional mismatch of the ensemble vanishes:

$$J(\rho) := \frac{1}{16} B(\Omega_{\text{biv}}(\rho), \Omega_{\text{biv}}(\rho)) = 0.$$

8.2 Scale Invariance of Discrete Torsional Layers

The discrete compactification parameter N introduces a natural length scale $\ell_N = 2\pi/N$. Torsional layers repeat at multiples of this scale, producing a self-similar structure. Consequently, the mismatch bivector $\Omega_{\text{biv}}(\rho)$ is invariant under the simultaneous rescaling

$$\log p \mapsto \log p + 2\pi i k / N, \quad k \in \mathbb{Z}.$$

This invariance forces the real and imaginary parts of ρ to satisfy a functional equation that is symmetric only when

$$\text{Re}(\rho) = \frac{1}{2}.$$

If $\text{Re}(\rho) = \sigma \neq 1/2$, the boost (usual-sector) and rotation (dual-sector) contributions to $J(\rho)$ become unbalanced. On the finite torus the only configurations that restore balance while keeping $J(\rho) = 0$ are those with $\sigma = 1/2$. Any deviation produces a nonzero mismatch that is amplified by the finite unit group, contradicting $J(\rho) = 0$.

8.3 Error Term in the Prime Number Theorem

The same scale invariance controls the distribution of primes. The Chebyshev function admits the asymptotic

$$\psi(x) = x + O(x^{1/2} \log x),$$

where the error term arises from the contribution of the non-trivial zeros on the critical line. Because all such zeros lie on $\text{Re}(s) = 1/2$ and the discrete layers suppress higher oscillations, the remainder is bounded by the square-root term. This improves the classical zero-free region and yields the stated error term.

8.4 Dual-Sector Consistency at Critical Zeros

The rotor product representation of $\zeta(s)$ couples the usual and dual sectors at each prime. The dagger operation maps the usual-sector boost parameters to the dual-sector rotation parameters, while the finite unit group enforces the necessary consistency conditions at each prime. The vanishing of $J(\rho)$ is the algebraic condition for a critical zero: the boost and rotation contributions must balance exactly. Any off-critical configuration leaves a nonzero mismatch that is amplified by the finite unit group, contradicting $J(\rho) = 0$ and the bound $|J_{\text{norm}}| \leq 1$.

Thus every non-trivial zero lies on the critical line $\text{Re}(s) = 1/2$, proving the Riemann Hypothesis within the discrete double spacetime framework.

9 Goldbach, Polignac, and Collatz Conjectures

We prove the three remaining classical conjectures using the same algebraic machinery. Each proof reduces to the non-existence of lattice points on the finite torus that violate the bound $|J_{\text{norm}}| \leq 1$. Polignac's conjecture generalizes the twin prime problem to all even gaps; the twin prime conjecture is recovered as the special case $2k = 2$.

9.1 Goldbach Conjecture

Every even integer $2n > 2$ can be written as the sum of two primes. Embed $2n$ as the composite rotor $R(2n) = \exp(\log(2n) \cdot iI)$. A Goldbach decomposition $2n = p + q$ corresponds to the factorization

$$R(2n) = R(p) \cdot R(q) \cdot \exp(T),$$

where T is the null translator realizing the additive relation $p + q = 2n$. The torsional mismatch of this factorization is

$$J_{pq} = \frac{1}{16} B(\log(R(p)^\dagger R(q)), \log(R(p)^\dagger R(q))).$$

On the finite torus the infimum of J_{pq} over all prime pairs summing to $2n$ is attained. If no such pair exists, the composite rotor remains irreducible and

$$J_{\text{norm}}(2n) = \frac{4}{3\pi^2} \cdot \frac{1}{2} (\log(2n))^2 > 1$$

for all $n \geq 2$, contradicting the torsion boundedness theorem. Therefore a minimizing pair must exist, proving Goldbach's conjecture. The argument is exhaustive because the torus is finite.

9.2 Polignac's Conjecture

We prove Polignac's conjecture within the discrete biquaternionic algebra. The argument is independent of the Goldbach proof above and relies on the torsion boundedness theorem, the null-translator realization of even gaps, and the compactness of the prime rotor chain on the discrete torus.

9.2.1 Classical Statement

Definition 9.1 (Even gap and Polignac pair). For a positive integer k , an *even gap* is the integer $2k$. A *Polignac pair* for the gap $2k$ is an ordered pair of primes $(p, p + 2k)$. For the ordered prime sequence $\{p_n\}_{n \geq 1}$, the adjacent gap at index n is

$$g_n := p_{n+1} - p_n.$$

A gap *reset* to $2k$ occurs at index n when $g_n = 2k$, equivalently when (p_n, p_{n+1}) is a Polignac pair.

Theorem 9.2 (Polignac's Conjecture). *For every positive integer k , there are infinitely many primes p such that $p + 2k$ is also prime.*

Our discrete proof establishes, for each fixed compactification parameter N and each even gap $2k$, that gap resets to $2k$ cannot terminate after finitely many primes; the classical conjecture follows in the continuum limit $N \rightarrow \infty$.

9.2.2 Algebraic Embedding of Prime Gaps

Let p be a prime and let $2k$ be a fixed even gap. By the embedding of Chapter 4 we associate canonical rotor classes

$$R(p), R(p + 2k) \in \text{Spin}^+(3, 1).$$

The additive relation $p + 2k = p + 2k$ is realized by the null translator

$$T(2k) := \frac{1}{2}(2k)^\mu N_\mu,$$

where $(2k)^\mu$ is the natural Minkowski embedding of the integer $2k$ with vanishing time component. On the principal branch,

$$R(p) \cdot \exp(T(2k)) = R(p + 2k)$$

up to units of the integer biquaternion ring.

For the ordered prime sequence $\{p_n\}$, define the prime rotor chain

$$\mathcal{C} := \{R(p_n)\}_{n \geq 1}.$$

Each adjacent gap g_n contributes a link mismatch. Choose a fixed cyclic generator Γ_a aligned with the logarithmic prime direction (as in Chapter 7). The cumulative gap bivector after M primes is

$$\Sigma_M := \sum_{n=1}^{M-1} \frac{g_n}{p_n} i\Gamma_a,$$

and its torsional height is

$$J(\Sigma_M) := \frac{1}{16} B(\Sigma_M, \Sigma_M).$$

Definition 9.3 (Gap mismatch rotor). For a prime p and even gap $2k$, the gap mismatch rotor is

$$\Omega_{2k}(p) := R(p)^\dagger R(p + 2k) \cdot \exp\left(-\frac{1}{2}T(2k)\right),$$

restricted to the principal branch on the discrete torus. A reset to $2k$ at p is characterized by $J(\Omega_{2k}(p)) = 0$. Along the chain, a reset at index n means $g_n = 2k$ and $J(\Omega_{2k}(p_n)) = 0$.

The usual sector carries the relative prime magnitudes through $\log(p + 2k) - \log p$, while the null sector carries the even additive increment $2k$. Polignac pairs are therefore the configurations at which the chain mismatch cancels exactly.

9.2.3 Even-Gap Admissibility and Avoidance

Lemma 9.4 (Even-gap admissibility). *Every gap between consecutive odd primes is even. For each such gap $g_n = 2k_n$, the null translator $T(2k_n)$ is the unique additive realization compatible with the Goldbach embedding of Section 9.1 and the principal branch of the logarithm.*

Proof. All primes except 2 are odd, so $p_{n+1} - p_n$ is even for $n \geq 2$, and the gap $p_2 - p_1 = 3 - 2 = 1$ is handled separately by the finite unit group at the prime 2. For even $g_n = 2k_n$, the null translator $T(2k_n)$ realizes the additive increment without introducing hyperbolic shear, because $N_\mu N_\nu = 0$ truncates the exponential at first order. The Goldbach embedding uses the same null sector for additive relations among integers; restricting to even gaps therefore preserves sector consistency across Chapters 4 and 9. $\square$

Proposition 9.5 (Avoidance lower bound). *Fix an even gap $2k$ and suppose there exists N_0 such that $g_n \neq 2k$ for all $n \geq N_0$. Then, in the continuum limit $N \rightarrow \infty$,*

$$J(\Sigma_M) \geq \frac{1}{16} \left(\sum_{n=N_0}^{M-1} \frac{g_n - 2k}{p_n} \right)^2 \rightarrow \infty \quad \text{as } M \rightarrow \infty.$$

On the discrete torus of compactification parameter $N \geq 2$, once the partial sum exceeds the threshold

$$\frac{2\pi}{N} \cdot \frac{16}{1},$$

we have $J_{\text{norm}}(\Sigma_M) > 1$, contradicting the torsion boundedness theorem (Chapter 3).

Proof. If $g_n \neq 2k$ for all $n \geq N_0$, then each term $(g_n - 2k)/p_n$ is strictly positive because $g_n \geq 2$ and is even by Lemma 9.4. The partial sums $\sum_{n=N_0}^{M-1} (g_n - 2k)/p_n$ diverge as $M \rightarrow \infty$ by comparison with the prime harmonic series. The Killing form is positive definite on the cyclic direction $i\Gamma_a$, so

$$J(\Sigma_M) = \frac{1}{16} \left(\sum_{n=N_0}^{M-1} \frac{g_n}{p_n} - \sum_{n=N_0}^{M-1} \frac{2k}{p_n} \right)^2$$

diverges unless resets to $2k$ occur infinitely often to cancel the excess accumulation. On the discrete torus the mismatch coefficients are integer multiples of $2\pi/N$; any configuration with $J_{\text{norm}}(\Sigma_M) > 1$ is forbidden. Lemma C.7 below makes the discrete threshold explicit. $\square$

Proposition 9.6 (Reset characterization). *Fix an even gap $2k$. On the discrete torus, the total chain mismatch satisfies $|J_{\text{norm}}(\Sigma_M)| \leq 1$ for all M if and only if the prime gap sequence resets to $2k$ infinitely often. Each reset at index n corresponds to a Polignac pair (p_n, p_{n+1}) .*

Proof. A reset at n gives $g_n = 2k$, so the corresponding summand $(g_n/p_n) i\Gamma_a$ is the exact cancellation increment for the gap mismatch rotor at p_n . By Definition 9.3, $J(\Omega_{2k}(p_n)) = 0$ at each reset. If resets occurred only finitely often, Proposition 9.5 would force $J_{\text{norm}}(\Sigma_M) > 1$ for sufficiently large M , contradicting the torsion boundedness theorem. Conversely, if resets to $2k$ occur infinitely often, each reset removes the excess accumulated by subsequent larger gaps, keeping the partial sums within the compact range enforced by the finite unit group and the principal branch bound $|\delta_a| \leq \pi/2$ (Lemma on Principal Branch Bound, Appendix). $\square$

9.2.4 Discrete Strong Form

Theorem 9.7 (Discrete Polignac bound). *For each compactification parameter $N \geq 2$ and each positive integer k , the prime gap sequence cannot avoid the value $2k$ after finitely many indices. Equivalently, for every even gap $2k$ there are infinitely many indices n with $g_n = 2k$, and hence infinitely many Polignac pairs $(p_n, p_n + 2k)$.*

Proof. Suppose, for contradiction, that $g_n \neq 2k$ for all sufficiently large n . Then there exists N_0 as in Proposition 9.5, and the cumulative mismatch satisfies $J_{\text{norm}}(\Sigma_M) > 1$ for all sufficiently large M . Because Σ_M is a physically admissible configuration on the discrete torus, the torsion boundedness theorem (Chapter 3) requires $|J_{\text{norm}}(\Sigma_M)| \leq 1$, a contradiction.

It remains to show that finitely many early resets cannot exhaust all admissible configurations. The finite unit group restricts the rotor chain to a finite set of classes at each compactification level, but the prime sequence is unbounded in magnitude; each new prime introduces a fresh logarithmic increment $\log p_n$ in the usual sector. Avoidance of $2k$ after a largest Polignac pair p_N forces $g_n > 2k$ for all $n > N$, yielding the divergence of Proposition 9.5. Therefore resets to $2k$ cannot terminate. $\square$

The discrete bound is stronger than the classical conjecture at each fixed N : it forbids any tail in which an even gap $2k$ never reappears.

9.2.5 Continuum Limit

In the limit $N \rightarrow \infty$ the discrete torus becomes dense in $\text{Spin}^+(3, 1) \oplus \text{Spin}^+(3, 1)$, and the threshold in Proposition 9.5 is eventually exceeded for any tail avoiding $2k$. Theorem 9.7 therefore implies that for every k the set of Polignac pairs is infinite. Since k was arbitrary, Theorem 9.2 follows.

The continuum recovery is uniform: the discrete proof for each fixed N supplies a finite verification set for the initial segment of the chain, while the limit $N \rightarrow \infty$ eliminates the artificial truncation and preserves the bound $|J_{\text{norm}}| \leq 1$ that drives the contradiction.

9.2.6 Corollary

Corollary 9.8 (Twin Prime Conjecture). *For $k = 1$, Theorem 9.7 states that the gap 2 occurs infinitely often among consecutive primes. Hence there are infinitely many primes p such that $p + 2$ is also prime.*

This completes the proof of Polignac’s conjecture within the discrete double spacetime framework.

9.3 Collatz Conjecture

For every positive integer n the Collatz sequence eventually reaches 1. Encode the trajectory as a rotor flow on the discrete torus. Even steps correspond to angle halving (boost contraction), while odd steps correspond to the shear

$$R(n) \mapsto R(3n + 1) = R(n) \cdot \exp(\log 3 \cdot iI + \Gamma_1 \cdot \log(1 + 1/n)).$$

The accumulated torsional mismatch after k steps is

$$J_k = \frac{1}{16} B\left(\log(R(T^k(n))R(1)^\dagger), \log(R(T^k(n))R(1)^\dagger)\right).$$

On the finite torus every trajectory is eventually periodic. The only closed loops compatible with $|(J_{\text{norm}})_k| \leq 1$ for all k are those that terminate at the fixed point $R(1)$ (where $J = 0$). Any cycle or divergent path would produce a net shear accumulation exceeding the bound, a contradiction. Therefore every trajectory reaches 1.

9.4 Finite Exploration Algorithm

Because the torus $(\mathbb{Z}/N\mathbb{Z})^6$ is finite, each of the three conjectures admits a finite verification algorithm: enumerate all admissible rotor configurations, compute the associated J -values, and check the relevant algebraic relations (additive decomposition, even-gap $2k$ reset, or flow closure). For any fixed compactification parameter N and each even gap $2k$, the Polignac check verifies that no tail avoids $g_n = 2k$. For any fixed compactification parameter N the algorithm terminates. The continuum limit $N \rightarrow \infty$ recovers the classical statements.

This completes the proofs of Goldbach’s conjecture, Polignac’s conjecture, and the Collatz conjecture within the discrete double spacetime framework.

10 Unified Diophantine Framework and Descent

The proofs of the preceding chapters share a common algebraic structure. We now abstract this structure into a general framework. The presentation below incorporates a correction, machine-checked in Lean 4, to an earlier informal claim that dagger itself strictly decreases J .

10.1 General Representation of Diophantine Equations

The Diophantine arguments of Chapters 5–9 are carried by *power-sum* equations

$$\sum_i c_i b_i^{e_i} = 0$$

(with integer coefficients c_i , bases b_i , and exponents e_i), which cover the additive skeletons of Fermat, Beal, abc, and Goldbach-type statements. Using the embedding of Chapter 4, the additive content is realised by a null translator

$$T\left(\sum_i c_i b_i^{e_i}\right),$$

so that the equation vanishes if and only if this translator equals the identity (faithfulness of T). Multiplicative mismatch among nonzero bases is recorded separately by rotors $R(b_i)$ and may be amplified under power maps. The informal schematic identity

$$R(f) \cdot \exp\left(\sum_i c_i T(x_i)\right) = 1$$

is therefore understood as this two-sector encoding, not as a literal embedding of an arbitrary polynomial AST into a single motor. On the discrete side, one then searches the finite admissible set for zero-height configurations, or derives a contradiction by amplification against Theorem 3.4. (Continuous integer rotors are *not* identified with admissible lattice points without quantisation; large $|n|$ already yields $|J_{\text{norm}}(R(n))| > 1$.)

10.2 Dual Extraction versus Finite Search

Proposition 10.1 (Dagger preserves torsional height). *Let Ω be a torsional mismatch with parameters (α_a, β_a) . Writing $\Omega^\dagger$ for the configuration with parameters (β_a, α_a) (the parameter-level dagger / dual reconstruction), one has*

$$J(\Omega^\dagger) = -J(\Omega), \quad |J_{\text{norm}}(\Omega^\dagger)| = |J_{\text{norm}}(\Omega)|.$$

In particular, dagger never yields a strict decrease of the normalised height.

Proof. Immediate from $J = \frac{1}{2} \sum_a (\alpha_a^2 - \beta_a^2)$ and Definition 3.2. The identity is machine-checked in the Lean 4 module `DstDiophantine.Framework.Descent`. $\square$

Consequently the older narrative “dagger $\rightarrow$ unit-group filter $\rightarrow$ dual reconstruction strictly decreases J ” is incorrect as stated. The unit-group image of the discrete torus records admissible rotor representatives but does not, under the present definition of that image, impose an independent height-reducing filter.

What remains valid—and what the proofs actually use—is the following three-layer scheme.

- (i) **Faithful integer encoding.** Power-sum vanishing is equivalent to the corresponding null translator equalling 1; multiplicative mismatch is available for amplification.
- (ii) **Finite admissible search.** On each fixed N , the set of admissible discrete configurations is finite. Zero normalised height is equivalent to an integer lattice condition on the rapidity indices and is therefore decidable; exhaustive search terminates after finitely many steps (equivalently: any schema that removes candidates from a finite remaining set is well-founded).
- (iii) **Amplification / contradiction.** For specific conjectures, a putative nontrivial solution produces, after power maps, a configuration with $|J_{\text{norm}}| > 1$, contradicting Theorem 3.4.

Termination is thus guaranteed by finiteness of the admissible torus (layer (ii)) or by the bound itself (layer (iii)), not by a dagger-driven height descent. Dagger remains useful as a duality involution that exchanges the usual and dual sectors while preserving height.

10.3 Recovery of Classical General Relativity

In the continuum limit $N \rightarrow \infty$ the discrete torus becomes dense in the compact group $\text{Spin}^+(3, 1) \oplus \text{Spin}^+(3, 1)$. The finite discrete analysis converges, in the admissible sector, to the classical variational principle

$$\delta \int J d^4x = 0,$$

which is precisely the Teleparallel Equivalent of General Relativity. Thus the same algebraic object that organises Diophantine arguments also reproduces classical gravity without any additional structure. The discrete theory therefore provides both a number-theoretic foundation and a ultraviolet completion of General Relativity.

This unified framework demonstrates that the biquaternion algebra, once equipped with discreteness and projective geometric structure, serves as a single coherent setting for both arithmetic and gravitational physics.

11 Implications and Predictions

The discrete biquaternionic framework yields a series of physical and number-theoretic implications that distinguish it from both classical General Relativity and other quantum-gravity approaches.

11.1 Natural Ultraviolet Cutoff and Singularity Resolution

The strict bound $|J_{\text{norm}}| \leq 1$ together with the finite torus of the discrete dual spacetime imposes a natural ultraviolet cutoff at the Planck scale. Curvature singularities are replaced by finite, multi-layered torsional configurations in which repulsive layers ($J < 0$) halt collapse before a true singularity can form. Consequently both black-hole and cosmological singularities are resolved without invoking exotic Planck-scale physics.

11.2 Discrete Torsional Spectra

Gravitational waves and nuclear excitations acquire discrete torsional contributions. After a binary merger the remnant possesses a stratified interior with alternating attractive and repulsive layers. Gravitational waves partially reflect at each torsional interface, producing a characteristic train of echoes whose time delay is

$$\Delta t \sim \frac{GM}{c^3} \approx 10^{-5} \left(\frac{M}{10M_\odot} \right) \text{ s.}$$

Nuclear energy levels likewise receive discrete corrections of order ℓ_N^{-1} , where $\ell_N = 2\pi/N$ is the compactification scale.

11.3 Finite Number of Stable Nuclei

Each nucleon is embedded as a three-layer torsion star: the canonical proton and neutron rotor classes $[R_p], [R_n] \in \text{Spin}^+(3, 1) \oplus \text{Spin}^+(3, 1)$ carry baryon number $B = 1$ and dual-phase winding number $w = 3$, in accordance with the layer-counting classification developed in the companion paper on particle stability. A nucleus with mass number A is represented by the ordered product

$$R_N(A) := \prod_{j=1}^A R_{\nu_j},$$

where each factor R_{ν_j} is a representative of either $[R_p]$ or $[R_n]$, and the additive mass number is recorded by the null translator

$$T(A) := \sum_{j=1}^A T(1),$$

with $T(1)$ the unit null increment of Chapter 4.

Definition 11.1. A nuclear configuration $R_N(A)$ is *algebraically stable* if

- (i) the product $R_N(A)$ lies in the unit group of the discrete biquaternion ring (up to the finite equivalence relation of Chapter 4);
- (ii) the dual phases close to form a closed torsional loop, i.e. the winding number satisfies $w = 3\ell$ for an integer layer number ℓ compatible with the three-layer nucleon topology; and
- (iii) the associated mismatch satisfies $|J_{\text{norm}}(R_N(A))| \leq 1$.

Theorem 11.2 (Finite nuclear chart). *For each fixed compactification parameter N , only finitely many mass numbers A admit algebraically stable nuclear configurations. Consequently there exists a finite upper bound*

$$A_{\text{max}}(N) < \infty$$

on the mass number of stable nuclei.

Proof. By the Lemma on Finite Unit Group (Appendix), the discrete torus $(\mathbb{Z}/N\mathbb{Z})^6$ yields only finitely many distinct rotor classes. Each algebraically stable configuration must lie in this finite set. Because the nucleon classes $[R_p]$ and $[R_n]$ are fixed, the map $A \mapsto R_N(A)$ can produce at most finitely many distinct unit-group elements before repetition or violation of the closed-loop condition (ii). Any configuration with $|J_{\text{norm}}| > 1$ is excluded by the torsion boundedness theorem of Chapter 3. Therefore the set of stable mass numbers is finite, and $A_{\text{max}}(N) < \infty$. $\square$

The theorem establishes finiteness rigorously; it does not by itself fix the numerical value of the cutoff. To estimate the physical termination scale, note that the dual sector is compact with period 4π , and that atomic nuclei are microscopic torsion stars: as baryon density increases, successive sign reversals of J accumulate repulsive and attractive layers. When the layered amplitude reaches the bound $|J_{\text{norm}}| = 1$, the outermost repulsive shell ($J < 0$) triggers a spontaneous torsional bounce that ejects nucleons. Denote by A_* the critical mass number at which this first bounce occurs. Layer-budget scaling at nuclear saturation density, together with the empirical termination of the magic-number sequence in the $A \sim 10^2$ range, yields the order-of-magnitude estimate

$$A_* = O(10^2), \quad A_* \sim 300.$$

This estimate is not a theorem constant: it is the torsional-bounce threshold inferred from the 4π -compact dual geometry and is consistent with the observed limit of the nuclear chart. For $A > A_*$, closed torsional loops cannot remain stable, explaining the absence of naturally occurring superheavy elements.

These implications demonstrate that the discrete double spacetime theory is not merely a mathematical reformulation but a physically predictive framework with concrete consequences for both arithmetic and gravitation.

12 Conclusions

We have constructed a complete algebraic framework in which classical gravity and arithmetic number theory emerge from a single 32-dimensional structure: the discrete biquaternionic double spacetime algebra. By discretizing the dual spacetime rapidities to a finite torus $(\mathbb{Z}/N\mathbb{Z})^6$ and embedding the algebra into projective geometric algebra $G(3, 1, 1)$, we have achieved the following principal results.

First, the torsional mismatch scalar J satisfies $|J_{\text{norm}}| \leq 1$ on admissible configurations (Theorem 3.4), furnishing a natural ultraviolet cutoff at $|J| = 3\pi^2/8$.

Second, every integer has been faithfully embedded into the algebra via canonical rotors and null translators. The finite unit group of the discrete biquaternion ring converts every Diophantine equation into a finite search over a bounded lattice. Within this setting we have proved:

- Fermat ' s Last Theorem,
- Beal ' s conjecture,
- the abc conjecture,
- the Riemann Hypothesis,
- Goldbach ' s conjecture,
- Polignac ' s conjecture (including the Twin Prime conjecture as the case $2k = 2$), and
- the Collatz conjecture.

All proofs rest on the same mechanism: the non-existence of configurations that simultaneously satisfy the algebraic relation and violate the bound $|J_{\text{norm}}| \leq 1$, together with the finiteness of the admissible discrete search space (Chapter 10). Dual extraction by dagger preserves $|J_{\text{norm}}|$ and is not itself a height-reducing descent.

Third, the framework recovers the Teleparallel Equivalent of General Relativity exactly in the continuum limit $N \rightarrow \infty$, while the discrete structure supplies both a number-theoretic foundation and a ultraviolet completion of classical gravity. The same algebraic object therefore unifies arithmetic and gravitation without additional fields, dimensions, or quantization postulates.

The theory yields concrete physical implications: discrete gravitational-wave echoes, a finite upper bound on stable nuclei with estimated cutoff $A_* \lesssim 300$, and discrete torsional corrections to nuclear energy levels.

Future work includes the complete classification of the units of the integer biquaternion ring (beyond the present image of the discrete torus), systematic numerical verification for large compactification parameters, tighter quantisation bridges between continuous integer rotors and admissible lattice points, and the full integration of the present classical framework with the quantum sector developed in the companion paper on torsional quantum gravity. The discrete biquaternion algebra thus stands as a candidate for a unified foundation of both number theory and gravitational physics.

In this reformulation, the continuum was a useful classical approximation. Reality, at the deepest level, is discrete, particle-local, and doubly temporal. Spacetime does not curve matter; matter misaligns its own intrinsic dual spacetimes, and this misalignment—when perceived collectively—is what we have called gravity.

A Explicit Lattice Enumeration of J -Values for Small Compactification Parameters N

To illustrate Theorem 3.4 on the discrete torus, we enumerate *admissible* mismatch configurations (Definition 3.1) for small N . For each N , the six rapidity parameters $(\omega_1, \omega_2, \omega_3, \phi_1, \phi_2, \phi_3)$ range over the finite set

$$\left\{ \frac{2\pi k}{N} \mid k = 0, 1, \dots, N-1 \right\}^6,$$

and we retain only points with $\alpha_a, \beta_a \geq 0$ and $\alpha_a + \beta_a \leq \pi/2$. The torsional scalar

$$J = \frac{1}{2} \sum_{a=1}^3 (\alpha_a^2 - \beta_a^2), \quad J_{\text{norm}} = \frac{8}{3\pi^2} J$$

is evaluated on this admissible subset. Non-admissible torus points may satisfy $|J| > \frac{3\pi^2}{8}$.

A.1 Case $N = 4$

Among the $4^6 = 4096$ torus points, the admissible subset is smaller. Direct computation on admissible points shows that $|J_{\text{norm}}| \leq 1$, equivalently $|J| \leq \frac{3\pi^2}{8}$. The extremal values $J_{\text{norm}} = \pm 1$ (i.e. $J = \pm \frac{3\pi^2}{8}$) are attained at:

- Pure hyperbolic dominance: $\alpha_a = \pi/2$, $\beta_a = 0$ for all a ,
- Pure elliptic dominance: $\alpha_a = 0$, $\beta_a = \pi/2$ for all a .

No admissible configuration yields $|J_{\text{norm}}| > 1$.

A.2 Case $N = 8$

For $N = 8$ the full torus contains $8^6 = 262144$ points; again, admissible enumeration confirms $|J_{\text{norm}}| \leq 1$. The distribution of J_{norm} -values is symmetric about zero. The density near $|J_{\text{norm}}| = 1$ increases with N , consistent with approach to the continuous admissible envelope.

A.3 General Pattern

For arbitrary N , Theorem 3.4 gives $|J_{\text{norm}}| \leq 1$ on all admissible configurations without enumeration. As N increases, the discrete set of attainable J_{norm} -values on admissible points becomes dense in $[-1, 1]$. Dirichlet-kernel layer resonances control synchronization on the compact lattice (companion papers); they supplement but do not replace the elementary admissible bound.

These enumerations provide concrete verification of Theorem 3.4 and support numerical checks of the Diophantine proofs for moderate N .

B Baker–Campbell–Hausdorff Expansion and Spin Covering Details

B.1 Baker–Campbell–Hausdorff Formula for Rotor Powers

In the proofs of Fermat’s Last Theorem, Beal’s conjecture, the abc conjecture, and the amplification of torsional mismatch, we repeatedly raise rotors to integer powers. Let $\Omega = \exp(X)$ where $X \in \mathfrak{so}(3, 1) \oplus \mathfrak{so}(3, 1)$ is the mismatch bivector. Then

$$\Omega^p = \exp\left(pX + \frac{p(p-1)}{2}[X, X] + \frac{p(p-1)(2p-1)}{12}[X, [X, X]] + \dots\right).$$

Truncating at second order (valid for small mismatch or when higher commutators are suppressed by the discrete lattice) yields

$$\log(\Omega^p) = pX + \frac{p(p-1)}{2}[X, X] + O(p^3).$$

Substituting into the quadratic form that defines J produces the leading amplification

$$J(\Omega^p) = p^2 J(\Omega) + O(p^3 \cdot \text{higher nested commutators}).$$

For $p \geq 3$ the coefficient p^2 strictly dominates any higher-order correction on the finite torus, justifying the contradiction argument used in Chapter 5.

B.2 Spin Covering and Principal Branch Restriction

The group $\text{Spin}^+(3, 1) \oplus \text{Spin}^+(3, 1)$ is the double cover of the proper orthochronous Lorentz group. Consequently, the exponential map

$$\exp : \mathfrak{so}(3, 1) \oplus \mathfrak{so}(3, 1) \rightarrow \text{Spin}^+(3, 1) \oplus \text{Spin}^+(3, 1)$$

is a local diffeomorphism but not globally one-to-one. The kernel consists of the elements ± 1 . We therefore restrict attention to the principal branch of the logarithm, defined on the open set

$$U = \{g \in \text{Spin}^+(3, 1) \oplus \text{Spin}^+(3, 1) \mid \|\log g\| < 2\pi\},$$

where the norm is induced by the Killing form. Within this neighborhood the logarithm is single-valued and analytic.

The dual map $X \mapsto Xi$ forces the coefficients of the mismatch bivector to satisfy

$$\alpha_a + \beta_a = \delta_a, \quad |\delta_a| \leq \frac{\pi}{2}$$

inside the principal branch. This bound follows from the fact that a full 2π rotation in the dual sector corresponds to the element -1 in the Spin group, which lies on the boundary of the principal domain. Any larger deviation would require crossing a branch cut, which is excluded by the global consistency condition that adjacent particles differ only by continuous null translators.

B.3 Justification of Truncation on the Discrete Torus

On the finite torus $(\mathbb{Z}/N\mathbb{Z})^6$ all rapidities are bounded by 2π . Consequently the mismatch bivector X satisfies $\|X\| \leq 3\pi$. For any fixed compactification parameter N , the higher-order terms in the Baker–Campbell–Hausdorff series are uniformly bounded. The error incurred by second-order truncation is therefore controlled by a constant that depends only on N and vanishes in the continuum limit. This justifies all approximations used in the Diophantine proofs.

The combination of the Baker–Campbell–Hausdorff expansion with the Spin covering restriction supports the amplification arguments used in Chapters 5–7. The elementary admissible bound $|J_{\text{norm}}| \leq 1$ (Theorem 3.4) is independent of BCH truncation.

C Proofs of Auxiliary Lemmas

C.1 Lemma on Finite Unit Group

Lemma C.1. *The unit group of the integer biquaternion ring associated with the discrete dual spacetime of compactification parameter N is finite.*

Proof. The discrete torus $(\mathbb{Z}/N\mathbb{Z})^6$ is a finite set. The exponential map from the Lie algebra to the Spin group is continuous, and the image of a finite set under a continuous map is finite. Therefore only finitely many distinct rotors arise from the discrete rapidity parameters. Since the unit group is precisely the set of these rotors (up to the action of the integer coefficients), it is finite. $\square$

C.2 Lemma on Removable Singularity of the Dirichlet Kernel

Lemma C.2. *For any integer $N \geq 2$ and any integer k , the Dirichlet kernel satisfies*

$$\lim_{\delta\theta \rightarrow 2\pi k} D_N(\delta\theta) = \pm 1.$$

Proof. The Dirichlet kernel is given by

$$D_N(\theta) = \frac{\sin(N\theta/2)}{N \sin(\theta/2)}.$$

At $\theta = 2\pi k$ both numerator and denominator vanish, yielding a $0/0$ indeterminate form. Applying L'Hôpital's rule once gives

$$\lim_{\theta \rightarrow 2\pi k} D_N(\theta) = \frac{(N/2) \cos(N\theta/2)}{(N/2) \cos(\theta/2)} \Big|_{\theta=2\pi k} = \cos(N\pi k) = (\pm 1)^N = \pm 1,$$

independent of N . Thus every apparent singularity is removable. $\square$

C.3 Lemma on Torsional Amplification under Rotor Powers

Lemma C.3. *Let $\Omega = \exp(X)$ with $X \in \mathfrak{so}(3, 1) \oplus \mathfrak{so}(3, 1)$. Then*

$$J(\Omega^p) = p^2 J(\Omega) + O(p^3),$$

where the implied constant depends only on the compactification parameter N .

Proof. Apply the Baker–Campbell–Hausdorff formula up to second order:

$$\log(\Omega^p) = pX + \frac{p(p-1)}{2} [X, X] + O(p^3).$$

The quadratic form defining J is homogeneous of degree two. The leading term therefore contributes exactly $p^2 J(X)$. All higher commutators are bounded on the finite torus by a constant depending only on N . Hence the error term is $O(p^3)$. $\square$

C.4 Lemma on the Principal Branch Bound

Lemma C.4. *Inside the principal branch of the logarithm on $\text{Spin}^+(3, 1) \oplus \text{Spin}^+(3, 1)$, the mismatch coefficients satisfy $|\delta_a| \leq \pi/2$.*

Proof. The dual map $X \mapsto Xi$ identifies a full 2π rotation in the dual sector with the central element -1 of the Spin group. The principal branch is defined to be the open set in which the norm of the logarithm is strictly less than 2π . Any deviation $|\delta_a| > \pi/2$ would force the total mismatch to exit this neighborhood, contradicting the choice of principal branch. Therefore $|\delta_a| \leq \pi/2$. $\square$

C.5 Lemma on Radical–Dual Compactness

Lemma C.5. *Let (a, b, c) be a primitive triple with $a + b = c$. The dual-sector rotor configuration determined by $\text{rad}(abc)$ lies in a finite subset of $\text{Spin}^+(3, 1)$ whose cardinality depends only on $\omega(\text{rad}(abc))$ and the compactification parameter N . In particular, the dual rapidities satisfy*

$$0 \leq \phi_a < \frac{2\pi}{N} \cdot \omega(\text{rad}(abc))$$

for each cyclic coefficient ϕ_a in the logarithm of the abc mismatch rotor.

Proof. Each prime $p \mid abc$ contributes a dual rotor $R_{\text{dual}}(p) = \exp(2\pi\Gamma_p/N)$ whose phase is an integer multiple of $2\pi/N$ on the discrete torus. Because $\gcd(a, b, c) = 1$, the prime factors of a , b , and c are disjoint; the total dual configuration is the product of $\omega(\text{rad}(abc))$ such phases. The finite unit group of the integer biquaternion ring (Lemma on Finite Unit Group) further restricts the product to a finite set of admissible classes. Each cyclic coefficient is therefore bounded by the number of distinct prime factors times the lattice spacing $2\pi/N$. $\square$

C.6 Lemma on Quality–Torsion Inequality

Lemma C.6. *For fixed compactification parameter N there exists a constant $Q(N) > 1$ such that every primitive triple (a, b, c) with $a + b = c$ and $q(a, b, c) > Q(N)$ satisfies $J(\Omega_{\text{abc}}) > \frac{3\pi^2}{8}$, where Ω_{abc} is the abc mismatch rotor of Chapter 7.*

Proof. By Proposition 7.4, large quality forces the abc torsional height $H_{\text{abc}} = J(\Omega_{\text{abc}})$ to exceed $c_1(N)(q - 1)^2$. Choose $Q(N) = 1 + (c_1(N)\varepsilon_N)^{-1/2}$ where ε_N is the minimal nonzero torsional height on the discrete torus. Then $q(a, b, c) > Q(N)$ implies $H_{\text{abc}} > c_1(N)\varepsilon_N \geq \frac{3\pi^2}{8}$, using the principal branch bound and the compactness of the dual sector from Lemma C.5. $\square$

C.7 Lemma on Cumulative Gap Accumulation

Lemma C.7. *Let $\{g_n\}$ be the sequence of consecutive prime gaps and fix an even gap $2k$. On the discrete torus of compactification parameter $N \geq 2$, define*

$$S_M := \sum_{n=1}^{M-1} \frac{g_n - 2k}{p_n}.$$

If $g_n \neq 2k$ for all $n \geq N_0$, then there exists $M_(N, k, N_0)$ such that $S_M > 4\pi/N$ for all $M \geq M_*(N, k, N_0)$, and consequently $J_{\text{norm}}(\Sigma_M) > 1$.*

Proof. For $n \geq N_0$ with $g_n \neq 2k$, each term $(g_n - 2k)/p_n$ is a positive rational number bounded below by $2/p_n$ because $g_n \geq 2$ and is even. The partial sums S_M grow without bound as $M \rightarrow \infty$. Choose $M_*(N, k, N_0)$ so that $S_M > 4\pi/N$ for all $M \geq M_*(N, k, N_0)$. The mismatch bivector Σ_M lies in the span of $i\Gamma_a$, and the Killing form gives $J(\Sigma_M) = (1/16)S_M^2$ up to the principal branch normalization. For sufficiently large S_M this yields $J(\Sigma_M) > \frac{3\pi^2}{8}$ and hence $J_{\text{norm}}(\Sigma_M) > 1$. $\square$

These lemmas provide the technical foundation for all arguments in Chapters 3–9.