

CSC236 Worksheet 7 Solution

Hyungmo Gu

May 9, 2020

Question 1

- First, I need to find the value of k .

The definition tells us k is the non-recursive cost.

Since the non-recursive part of call occurs when $\text{len}(s) < 2$ and it returns the input as output, it has cost of 1.

Second, I need to find the value of b .

The definition tells us b is the number of almost-equal parts the input is divided into.

Since the input s is divided into three roughly equal parts, we can conclude $b = 3$.

Third, I need to find the value of a .

The definition tells us a is the number of recursive calls.

Since the recursive calls in this problem are $r(s_1)$, $r(s_2)$ and $r(s_3)$, there are three of them, so $a = 3$.

Fourth, I need to find the value of f .

The definition tells us f is the cost of splitting and recombining

Since the cost of splitting and recombining is $\text{len}(s_3) + \text{len}(s_2) + \text{len}(s_1) = n$, the value of f is n .

Fifth, I need to evaluate asymptotic time complexity of function r using master's theorem.

Since $f \in \Theta(n^d)$ where $d = 1$ and $a = 3 = 3^d = b^d$, the master's theorem tells us $r(s) \in \Theta(\text{len}(s) \log_3 \text{len}(s))$.

Finally, I need to compare its time complexity to copying the string elements in reverse order, using loop.

In comparison to $\Theta(\text{len}(s) \log_3 \text{len}(s))$ by divide and conquer method, copying the string elements has cost of $\Theta(n)$.

Notes:

- **Divide and Conquer:** Partitions problem into b roughly equal subproblems, solve, and recombine:

$$T(n) = \begin{cases} k & \text{if } n \leq B \\ a_1 T(\lceil n/b \rceil) + a_2 T(\lfloor n/b \rfloor) + f(n) & \text{if } n > B \end{cases} \quad (1)$$

$$T(n) = \begin{cases} k & \text{if } n \leq B \\ a T(n/b) + f(n) & \text{if } n > B \end{cases} \quad (2)$$

where $b, k > 0$, $a_1, a_2 \geq 0$, and $a = a_1 + a_2 > 0$. $f(n)$ is the cost of splitting and recombining.

Note:

k : non-recursive cost, when $n < b$

b : number of almost-equal parts we divide problem into

a_1 : number of recursive calls to ceiling

a_2 : number of recursive calls to floor

a : number of recursive calls

f : cost of splittig and later recombining (should be n^d for master theorem)

- **Divide and Conquer Master Theorem:**

If $f \in \Theta(n^d)$, then

$$T(n) \in \begin{cases} \Theta(n^d) & \text{if } a < b^d \\ \Theta(n^d \log_b n) & \text{if } a = b^d \\ \Theta(n^{\log_b a}) & \text{if } a > b^d \end{cases} \quad (3)$$

- The master theorem is for master method.
- The master method provides a cookbook method for solving recurrences of the form

$$T(n) = aT(n/b) + f(n) \tag{4}$$

where $a \geq 1$ and $b > 1$.