# CSC373 Worksheet 3 Solution

July 28, 2020

## 1. Notes:

- Dynamic Programming
  - Is applied to optimization problems
  - Applies when the subproblems overlap
  - Uses the following sequence of steps
    - 1. Characterize the structure of an optimal solution
    - 2. Recursively define the value of an optimal solution
    - 3. Construct an optimal solution from computed information
- Matrix-chain Multiplication
  - Is an optimization problem solved using dynamic programming
  - Goal is to find matrix parenthesis with fewest number of operations

#### Example:

Given chain of matrices  $\langle A, B, C \rangle$ , it's fully parenthesized product is:

- \* (AB)C needs  $(10 \times 30 \times 5) + (10 \times 5 \times 60) = 1500 + 3000 = 4500$  operations
- \* A(BC) needs  $(30 \times 5 \times 60) + (10 \times 30 \times 60) = 27000$  operations

Thus, (AB)C performs more efficiently than A(BC).

- Is stated as: given a chain  $\langle A_1, A_2, ..., A_n \rangle$  of n matrices, where for i = 1, 2, ..., n matrix  $A_i$  has dimension  $p_{i-1} \times p_i$ , fully parenthesize the product  $A_1A_2...A_n$  in a way that minimizes the number of scalar multiplications.
- Steps

### 1. Check is the problem has Optimal Substructure

## Example (From CS Breakdown on Youtube):

Let us adopt the notation  $A_{i...j}$  where  $i \leq j$ , for the matrix that results from evaluating the product  $A_i A_{i+1} ... A_j$ .

Assume the solution has the following parentheses:

$$(A_{i...k})(A_{k+1...j})$$

If there is a better way to multiply  $(A_{i...k})$ , then we would have a more optimal solution.

This would be a contradiction, as we already stated that we have the optimal solution for  $A_{i...j}$ .

Therefore, this problem has optimal substructure.

#### 2. Recursive Solution

## Example (From CS Breakdown on Youtube):

Let M[i,j] be the cost of multiplying matrices from  $A_i$  to  $A_j$ 

We want to find out at which k' returns the fewest number of multiplications, or the minimum number of M.

The recursive formula for the cost of multiplying from  $A_i$  to  $A_j$  is

$$M[i,j] = \begin{cases} 0 & \text{if } i = j \\ \min_{i \le k \le j} M[i,k] + M[k+1,j] + p_{i-1}p_k p_j & \text{if } i < j \end{cases}$$
 (1)

- 3. Computing the Estimated Cost
- 4. Constructing the Optimal Solution