

Worksheet 9 Review

March 28, 2020

Question 1

- a. For every set of size 0 has 0 subsets of size 2.
- b. Let $n = 0$. Let S be an arbitrary set. Assume S has size 0.

Since S has size 0, empty subsets are the **only** subsets that can be included in S .

Then, because we know empty subsets have size 0, we can conclude there are 0 subsets of size 2.

It follows from above that the base case holds.

Attempt #2:

We want to show every set S of size 0 has 0 subsets of size 2.

Since S has size 0, empty subsets are the **only** subsets that can be included in S .

Then, because we know an empty subset have size 0, we can conclude there are 0 subsets of size 2.

Notes:

- Professor specifically mentions **We want to show every set S of size 0 has 0 subsets of size 2**
- Professor doesn't include conclusion at the end of proof
- Under which cases conclusion to a proof are included.

c. Now we will prove inductive step.

Let $k \in \mathbb{N}$. Assume every set of size k has $\frac{k(k-1)}{2}$ subsets of size 2 .

We want to show a set of size $k+1$ has $\frac{(k+1)k}{2}$ subsets of size 2 .

Part 1: counting subsets of S of size 2 that contain s_{k+1}

It follows from the table below,

k	Sets	Subsets of Size 2 with s_{k+1}
0	$\{0, 1\}$	1
1	$\{0, 1, 2\}$	2
2	$\{0, 1, 2, 3\}$	3
2	$\{0, 1, 2, 3, 4\}$	4

that the number of subsets of size 2 that contain s_{k+1} is $k+1$.

Part 2: counting subsets of S of size 2 that doesn't contain s_{k+1}

Because we know the subset of S that doesn't contain s_{k+1} is a set S of size k , we can conclude using induction hypothesis that there are

$$\frac{k(k-1)}{2} \tag{1}$$

subsets of size 2 .

Part 3: Putting the counts together

Then,

$$\frac{k(k+1)}{2} + (k+1) = \frac{(k+1)(k+2)}{2} \quad (2)$$

Then, it follows from proof by induction that the statement ' $\forall n \in \mathbb{N}$, every set of size n has $\frac{n(n-1)}{2}$ subsets of size 2' is true for all natural numbers n .

Attempt #2:

Part 1: counting subsets of S of size 2 that contain s_{k+1}

It follows from the table below,

k	Sets	Subsets of Size 2 with s_{k+1}
2	$\{0, 1\}$	1
3	$\{0, 1, 2\}$	2
4	$\{0, 1, 2, 3\}$	3
5	$\{0, 1, 2, 3, 4\}$	4

that the number of subsets of size 2 that contain s_{k+1} is k .

Part 2: counting subsets of S of size 2 that doesn't contain s_{k+1}

Because we know the subset of S that doesn't contain s_{k+1} is a set S of size k , we can conclude using induction hypothesis that there are

$$\frac{k(k-1)}{2} \quad (3)$$

subsets of size 2.

Part 3: Putting the counts together

Then,

$$\frac{k(k-1)}{2} + k = \frac{(k+1)k}{2} \quad (4)$$

Then, it follows from proof by induction that the statement ' $\forall k \in \mathbb{N}$, every set of size k has $\frac{k(k-1)}{2}$ subsets of size 2' is true for all natural numbers k .

Notes:

- I forgot that k represents number of elements in a set.

Question 2

- **Statement:** For every $n \in \mathbb{N}$, every finite set S of size n , has

$$\frac{n(n-1)(n-2)}{6} \quad (1)$$

subsets of size 3.

We will prove this statement by using induction on n .

Base Case:

Let $n = 0$.

Then, only the empty subsets can be included in S .

Because an empty subset has size 0, there are 0 subsets of size 3 in S .

Then, since

$$\frac{0 \cdot (0 - 1)(0 - 2)}{6} = 0 \quad (2)$$

the base case holds.

Inductive Case:

Let $k \in \mathbb{N}$. Assume every finite set S of size k has exactly $\frac{k(k-1)(k-2)}{6}$ subsets of size 3.

We want to show finite set S of size $k + 1$ contains $\frac{(k+1)k(k-1)}{6}$ subsets of size 3.

It follows from the table below

k	Sets	# of Subsets of Size 3	# of Subsets of Size 2
0	$\{\}$	0	0
1	$\{s_0\}$	0	0
2	$\{s_0, s_1\}$	0	1
3	$\{s_0, s_1, s_2\}$	1	3
4	$\{s_0, s_1, s_2, s_3\}$	4	6
5	$\{s_0, s_1, s_2, s_3, s_4\}$	10	10

we can deduce that given a set S size $k + 1$, the number of subsets of size 3 containing s_{k+1} is the sum of # of subsets of size 3 that doesn't contain s_{k+1} and # of subsets of size 2 that doesn't contain s_{k+1} .

Then,

$$\frac{k(k-1)(k-2)}{6} + \frac{k(k-1)}{2} = \frac{k(k-1)(k-2)}{6} + \frac{3k(k-1)}{6} \quad (3)$$

$$= \frac{k(k-1)(k-2+3)}{6} \quad (4)$$

$$= \frac{k(k-1)(k+1)}{6} \quad (5)$$

Attempt #2:

Statement: For every $n \in \mathbb{N}$, every finite set S of size n , has

$$\frac{n(n-1)(n-2)}{6} \tag{1}$$

subsets of size 3.

We will prove this statement by using induction on n .

Base Case:

Let $n = 0$.

Then, only the empty subsets can be included in S .

Because an empty subset has size 0, there are 0 subsets of size 3 in S .

Then, since

$$\frac{0 \cdot (0-1)(0-2)}{6} = 0 \tag{2}$$

the base case holds.

Inductive Case:

Let $k \in \mathbb{N}$. Assume every finite set S of size k has exactly $\frac{k(k-1)(k-2)}{6}$ subsets of size 3.

We want to show finite set S of size $k+1$ contains $\frac{(k+1)k(k-1)}{6}$ subsets of size 3. We will do that by determining the number of subsets of size 3 including s_{k+1} , and the number of subsets of size 3 not including s_{k+1} , and then combining them together.

First, we will show the number of subsets of size 3 including s_{k+1} is $\frac{n(n-1)}{2}$.

Because we know the number of subsets of size 3 (i.e $\{a_1, a_2, s_{k+1}\}$) containing s_{k+1} depends on the unique combination of first two elements a_1 and a_2 , and because we know

1. $a_1 \neq a_2$
2. $a_1, a_2 \in \{s_0, s_1, \dots, s_k\}$

, we can conclude that the number of subsets of size 3 containing s_{k+1} is exactly the number of subsets of size 2 in a set S of size k or

$$\frac{n(n-1)}{2} \tag{3}$$

Second, we will show the number of subsets of size 3 not including s_{k+1} is $\frac{n(n-1)(n-2)}{6}$.

Since the number of subsets of size 3 not containing s_{k+1} must contain 3 elements from $\{s_0, \dots, s_k\}$, this is exactly number of subsets of size 3 in a set S of size k , or

$$\frac{n(n-1)(n-2)}{6} \tag{4}$$

by induction hypothesis.

Then,

$$\frac{n(n-1)}{2} + \frac{n(n-1)(n-2)}{6} = \frac{3n(n-1)}{6} + \frac{n(n-1)(n-2)}{6} \quad (5)$$

$$= \frac{n(n-1)}{6} \cdot (3+n-2) \quad (6)$$

$$= \frac{n(n-1)(n+1)}{6} \quad (7)$$

Notes:

- I wonder if table like above can be used in a proof. If not, why can't it be done? If yes, when is the use of table not valid? How should it be constructed that it's valid?
- Can I put a subheader like 'Part 1: counting subsets of S of size 2 that contain s_{k+1} ' in a proof? If so, are there anything that I should be aware/be careful of?
- Is it alright to play with examples when I don't know how to proceed? What's the danger to this approach?
- Noticed that professor lays out the big ideas of proof (proof by induction evaluating number of s_{k+1} and s_k) and then fills in the missing detail

Question 3

a. Subsets that contain 3:

$$\{\{3\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$$

Subsets that doesn't contain 3:

$$\{\{\}, \{1\}, \{2\}, \{1, 2\}\}$$