

CSC236 Worksheet 4 Solution

Hyungmo Gu

May 6, 2020

Question 1

-

Notes:

- **Repeated Substitution:**

- Is a technique used to find a closed form formula
- **closed form formula** is a simple formula that allows evaluation of $T(n)$ without the need to evaluate, say $T(n/2)$

i.e. from

$$T(n) = \begin{cases} c & \text{if } n = 1 \\ 2T(n/2) + dn & \text{if } n > 1 \end{cases} \quad (1)$$

to

$$T(n) = cn + dn \log_2 n$$

Example:

Consider the recurrence

$$T(n) = \begin{cases} c & \text{if } n = 1 \\ 2T(n/2) + dn & \text{if } n > 1 \end{cases} \quad (2)$$

Find closed form formula for $T(n)$, where n is an arbitrary power of 2. That is $\exists k \in \mathbb{N}, n = 2^k$.

Let $n \in \mathbb{N}$ and assume that $\exists k \in \mathbb{N}^+, n = 2^k$, so $k = \log_2 n$.

Then,

$$T(n) = 2T(n/2) + dn \quad [\text{By 1}] \quad (3)$$

$$= 2\left(2T(n/2^2) + dn/2\right) + dn \quad [\text{By substituting } n/2 \text{ for } n \text{ in 1}] \quad (4)$$

$$= 2^2T(n/2^2) + 2dn \quad (5)$$

$$= 2^2\left(2T(n/2^3) + dn/2^2\right) + 2dn \quad [\text{By substituting } n/2^2 \text{ for } n \text{ in 1}] \quad (6)$$

$$= 2^3T(n/2^3) + 3dn \quad [\text{By substituting } n/2^2 \text{ for } n \text{ in 1}] \quad (7)$$

$$\vdots \quad (8)$$

$$= 2^kT(n/2^k) + kdn \quad [\text{After } k \text{ applications}] \quad (9)$$

$$= 2^{\log_2 n}T(n/2^{\log_2 n}) + (\log_2 n)dn \quad [\text{By replacing } k = \log_2 n] \quad (10)$$

$$= nT(1) + (\log_2 n)dn \quad (11)$$

$$= cn + (\log_2 n)dn \quad (12)$$

Question 2