

# Problem Set 4 Solution

Hyungmo Gu

April 7, 2020

## Question 1

- a. **Statement:**  $\forall f, g : \mathbb{N} \rightarrow \mathbb{R}^+, b \in \mathbb{R}^+, (g(n) \in \Theta(f(n))) \wedge (n_0 \in \mathbb{N}, n \geq n_0 \Rightarrow f(n) \geq b \wedge g(n) \geq b) \wedge (b > 1) \Rightarrow \log_b(g(n)) \in \Theta(\log_b(f(n)))$

**Statement Expanded:**  $\forall f, g : \mathbb{N} \rightarrow \mathbb{R}^+, b \in \mathbb{R}^+, \left( \exists c_1, c_2, n_0 \in \mathbb{R}^+, \forall n \in \mathbb{N}, n \geq n_0 \Rightarrow c_1 \cdot g(n) \leq f(n) \leq c_2 \cdot g(n) \right) \wedge \left( \exists n_1 \in \mathbb{N}, n \geq n_1 \Rightarrow f(n) \geq b \wedge g(n) \geq b \right) \wedge (b > 1) \Rightarrow \left( \exists d_1, d_2, n_2 \in \mathbb{R}^+, \forall n \in \mathbb{N}, n \geq n_2 \Rightarrow d_1 \cdot \log_b(g(n)) \leq \log_b(f(n)) \leq d_2 \cdot \log_b(g(n)) \right)$

Let  $f, g : \mathbb{N} \rightarrow \mathbb{R}^+$ , and  $b \in \mathbb{R}^+$ . Assume  $c_1 = 1$ ,  $c_2 = b$ , and  $n_0 = 1$ , and  $n \in \mathbb{N}$  such that  $n \geq n_0$  and  $c_1 \cdot g(n) \leq f(n) \leq c_2 \cdot g(n)$ . Assume  $f(n)$  and  $g(n)$  are eventually  $\geq b$ . Assume  $b > 1$ . Let  $d_1 = 1$ ,  $d_2 = 2$ , and  $n_2 = n_0$ . Assume  $n \geq n_2$ .

We need to show  $d_1 \cdot \log_b g(n) \leq \log_b f(n) \leq d_2 \cdot \log_b g(n)$ .

We will do so in two parts. One for  $(d_1 \cdot \log_b g(n) \leq \log_b f(n))$  and the other for  $(\log_b f(n) \leq d_2 \cdot \log_b g(n))$ .

**Part 1** ( $d_1 \cdot \log_b g(n) \leq \log_b f(n)$ ):

The assumption tell us

$$c_1 \cdot g(n) \leq f(n) \tag{1}$$

Then,

$$\log(c_1 \cdot g(n)) \leq \log(f(n)) \tag{2}$$

by the fact  $\forall x, y \in \mathbb{R}^+, x \geq y \Leftrightarrow \log x \geq \log y$ .

Then, using the fact  $b > 1$ , we can calculate

$$\frac{\log(c_1 \cdot g(n))}{\log b} \leq \frac{\log(f(n))}{\log b} \quad (3)$$

$$\frac{\log(c_1) + \log(g(n))}{\log b} \leq \frac{\log(f(n))}{\log b} \quad (4)$$

Then,

$$\frac{\log(g(n))}{\log b} \leq \frac{\log(f(n))}{\log b} \quad (5)$$

by the fact  $c_1 = 1$  and  $\log c_1 = 0$ .

Then, because we know  $\frac{\log f(x)}{\log b} = \log_b f(x)$ , we can conclude

$$\log_b(g(n)) \leq \log_b(f(n)) \quad (6)$$

**Part 2** ( $\log_b f(n) \leq d_2 \cdot \log_b g(n)$ ):

**Notes:**

- $\forall x, y \in \mathbb{R}^+, x \geq y \Leftrightarrow \log x \geq \log y$
- $\exists c_1, c_2, n_0 \in \mathbb{R}^+, \forall n \in \mathbb{N}, n \geq n_0 \Rightarrow c_1 \cdot g(n) \leq f(n) \leq c_2 \cdot g(n)$
- **Definition of Eventually:**  $\exists n_0 \in \mathbb{N}, n \geq n_0 \Rightarrow P$ , where  $P : \mathbb{N} \rightarrow \{\text{True}, \text{False}\}$

**Question 2**

**Question 3**

**Question 4**