

CSC373 Worksheet 1 Solution

July 22, 2020

```
1. Strassen_Algorithm(A,B):
2     n = A.rows
3     let C be a new n x n matrix
4
5     if n == 1
6         C_11 = A_11 * B_11
7
8     else partition as in step 3 of strassen's algorithm
9
10        p1 = Strassen_Algorithm(A_11, B_12) -
11             Strassen_Algorithm(A_11, B_22)
12
13        p2 = Strassen_Algorithm(A_11, B_22) +
14             Strassen_Algorithm(A_12, B_22)
15
16        p3 = Strassen_Algorithm(A_21, B_11) +
17             Strassen_Algorithm(A_22, B_11)
18
19        p4 = Strassen_Algorithm(A_22, B_21) -
20             Strassen_Algorithm(A_22, B_11)
21
22        p5 = Strassen_Algorithm(A_11, B_11) +
23             Strassen_Algorithm(A_11, B_22) +
24             Strassen_Algorithm(A_22, B_11) +
25             Strassen_Algorithm(A_22, B_22)
26
27        p6 = Strassen_Algorithm(A_12, B_21) +
28             Strassen_Algorithm(A_12, B_22) -
29             Strassen_Algorithm(A_22, B_21) -
30             Strassen_Algorithm(A_22, B_22)
31
32        p7 = Strassen_Algorithm(A_11, B_11) +
33             Strassen_Algorithm(A_11, B_12) -
34             Strassen_Algorithm(A_21, B_11) -
35             Strassen_Algorithm(A_21, B_12)
36
37        C_11 = p5 + p4 - p2 + p6
38        C_12 = p1 + p2
39        C_21 = p3 + p4
```

```

40         C_22 = p5 + p1 - p3 - p7
41
42     return C
43

```

Notes:

- Strassen's method for matrix multiplication
 - Reduces the time complexity of matrix multiplication from $O(n^3)$ to $O(n^{\log_2 7}) = O(n^{2.81})$
 - Has four steps

- 1) Divide the input matrices A and B and output matrix C into $n/2 \times n/2$ submatrices

$$A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}, \quad B = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}, \quad C = \begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix},$$

- 2) Create 10 matrices, S_1, S_2, \dots, S_{10} each of which is $n/2 \times n/2$ and is the sum or difference of two matrices created in step 1

$$\begin{aligned}
 S_1 &= B_{12} - B_{22} \\
 S_2 &= A_{11} + A_{12} \\
 S_3 &= A_{21} + A_{22} \\
 S_4 &= B_{21} - B_{11} \\
 S_5 &= A_{11} + A_{22} \\
 S_6 &= B_{11} + B_{22} \\
 S_7 &= A_{12} - A_{22} \\
 S_8 &= B_{21} + B_{22} \\
 S_9 &= A_{11} - A_{21} \\
 S_{10} &= B_{11} + B_{12}
 \end{aligned}$$

- 3) Recursively multiply $n/2 \times n/2$ matrices seven times to compute the following $n/2 \times n/2$ matrices

$$\begin{aligned}
 P_1 &= A_{11} \cdot S_1 = A_{11} \cdot B_{12} - A_{11} \cdot B_{22} \\
 P_2 &= S_2 \cdot B_{22} = A_{11} \cdot B_{22} + A_{12} \cdot B_{22} \\
 P_3 &= S_3 \cdot B_{11} = A_{21} \cdot B_{11} + A_{22} \cdot B_{11} \\
 P_4 &= A_{22} \cdot S_4 = A_{21} \cdot B_{11} + A_{22} \cdot B_{11} \\
 P_5 &= S_5 \cdot S_6 = A_{11} \cdot B_{11} + A_{11} \cdot B_{22} + A_{22} \cdot B_{11} + A_{22} \cdot B_{22} \\
 P_6 &= S_7 \cdot S_8 = A_{12} \cdot B_{21} + A_{12} \cdot B_{22} - A_{22} \cdot B_{21} - A_{22} \cdot B_{22} \\
 P_7 &= S_9 \cdot S_{10} = A_{11} \cdot B_{11} + A_{11} \cdot B_{12} - A_{21} \cdot B_{11} - A_{21} \cdot B_{12}
 \end{aligned}$$

- 4) Construct the four $n/2 \times n/2$ submatrices of the product C

$$\begin{aligned}
 C_{11} &= P_5 + P_4 - P_2 + P_6 = A_{11} \cdot B_{11} + A_{12} \cdot B_{12} \\
 C_{12} &= P_1 + P_2 = A_{11} \cdot B_{12} + A_{12} \cdot B_{22} \\
 C_{21} &= P_3 + P_4 = A_{21} \cdot B_{11} + A_{22} \cdot B_{21} \\
 C_{22} &= P_5 + P_1 - P_3 - P_7 = A_{22} \cdot B_{22} + A_{21} \cdot B_{12}
 \end{aligned}$$

Example: Use Strassen's algorithm to compute the matrix product

$$\begin{pmatrix} 1 & 3 \\ 7 & 5 \end{pmatrix} \begin{pmatrix} 6 & 8 \\ 4 & 2 \end{pmatrix}$$

* **STEP 1**

$$A_{11} = 1, A_{12} = 3, A_{21} = 7, A_{22} = 5$$

$$B_{11} = 6, B_{12} = 8, B_{21} = 4, B_{22} = 2$$

* **STEP 2**

$$S_1 = B_{12} - B_{22} = 4 - 2 = 2$$

$$S_2 = A_{11} + A_{12} = 1 + 3 = 4$$

$$S_3 = A_{21} + A_{22} = 7 + 5 = 12$$

$$S_4 = B_{21} - B_{11} = 4 - 6 = -2$$

$$S_5 = A_{11} + A_{22} = 1 + 5 = 6$$

$$S_6 = B_{11} + B_{22} = 6 + 2 = 8$$

$$S_7 = A_{12} - A_{22} = 3 - 5 = -2$$

$$S_8 = B_{21} + B_{22} = 4 + 2 = 6$$

$$S_9 = A_{11} - A_{21} = 1 - 7 = -6$$

$$S_{10} = B_{11} + B_{12} = 6 + 4 = 10$$

* **STEP 3**

$$P_1 = A_{11} \cdot S_1 = A_{11} \cdot B_{12} - A_{11} \cdot B_{22} = 1 \cdot 4 - 1 \cdot 2 = 2$$

$$P_2 = S_2 \cdot B_{22} = A_{11} \cdot B_{22} + A_{12} \cdot B_{22} = 1 \cdot 2 + 3 \cdot 2 = 8$$

$$P_3 = S_3 \cdot B_{11} = A_{21} \cdot B_{11} + A_{22} \cdot B_{11} = 7 \cdot 6 + 5 \cdot 6 = 72$$

$$P_4 = A_{22} \cdot S_4 = A_{22} \cdot B_{21} - A_{22} \cdot B_{11} = 5 \cdot 4 - 5 \cdot 6 = -10$$

$$P_5 = S_5 \cdot S_6 = A_{11} \cdot B_{11} + A_{11} \cdot B_{22} + A_{22} \cdot B_{11} + A_{22} \cdot B_{22} = 48$$

$$P_6 = S_7 \cdot S_8 = A_{12} \cdot B_{21} + A_{12} \cdot B_{22} - A_{22} \cdot B_{21} - A_{22} \cdot B_{22} = -20$$

$$P_7 = S_9 \cdot S_{10} = A_{11} \cdot B_{11} + A_{11} \cdot B_{12} - A_{21} \cdot B_{11} - A_{21} \cdot B_{12} = -20$$

* **STEP 4**

$$C_{11} = P_5 + P_4 - P_2 + P_6 = 48 - 10 - 8 - 20 = 10$$

$$C_{12} = P_1 + P_2 = 10$$

$$C_{21} = P_3 + P_4 = 62$$

$$C_{22} = P_5 + P_1 - P_3 - P_7 = 48 + 2 - 72 + 20 = -2$$

– Is not preferred in practical purposes

- 1) The constants used in Strassen's method are high and for a typical application Naive method works better.
- 2) For Sparse matrices, there are better methods especially designed for them.
- 3) The submatrices in recursion take extra space.
- 4) Because of the limited precision of computer arithmetic on noninteger values, larger errors accumulate in Strassen's algorithm than in Naive Method

References:

- 1) GeeksForGeeks, Divide and Conquer — Set 5 (Strassen's Matrix Multiplication), [link](#)
- Regular matrix multiplication
 -
 - The master method for solving recurrences
 - provides 'cookbook' method for solving recurrences of the form

$$T(n) = aT(n/b) + f(n)$$

– depends on the following theorem

- * Let $a \leq 1$ and $b > 1$ be constants, let $f(n)$ be a function and let $T(n)$ be defined on the nonnegative integers by the recurrence

$$T(n) = aT(n/b) + f(n)$$

where we interpret n/b to mean either $\lfloor n/b \rfloor$ or $\lceil n/b \rceil$. Then $T(n)$ has the following asymptotic bounds:

1. If $f(n) = O(n^{\log_b a - \epsilon})$ for some constant $\epsilon > 0$, then $T(n) = \Theta(n^{\log_b a})$
2. If $f(n) = \Theta(n^{\log_b a})$, then $T(n) = \Theta(n^{\log_b a} \lg n)$
3. If $f(n) = \Omega(n^{\log_b a + \epsilon})$ for some constant $\epsilon > 0$, and if $af(n/b) \leq cf(n)$ for some constant $c < 1$, and all sufficiently large n , then $T(n) = \Theta(f(n))$.

Example:

$$T(n) = 9T(n/3) + n$$

Here, $a = 9$, $b = 3$, and $f(n) = n = O(n^{\log_3 9 - 1})$ where $\epsilon = 1$.

Thus, $T(n) = \Theta(n^{\log_3 9})$ or $T(n) = \Theta(n^2)$

Example 2:

$$T(n) = T(2n/3) + 1$$

Here, $a = 1$, $b = 3/2$, $f(n) = 1 = \Theta(n^{\log_{3/2} 1})$.

Thus, $T(n) = \theta(\lg n)$

Example 3:

$$T(n) = T(n/4) + n \lg n$$

Here $a = 1$, $b = 4$, and $f(n) = n \lg n$ has asymptotic lowerbound of $f(n) = \Omega(n^{\log_4 3 + \epsilon}) = \Omega(n)$ where $\epsilon \approx 0.2$

Furthermore,

$$\begin{aligned} af(n/b) &= (3n/4) \lg n/4 \\ &= (3/4)n \lg n/4 \\ &= (3/4)n \lg n/4 \\ &= 3/4n \lg n - \lg 4 \\ &< 3/4n \lg n \\ &= cf(n) \end{aligned}$$

where $c = 3/4$.

Thus, $T(n) = \Theta(n \lg n)$

Example 4:

$$T(n) = 2T(n/2) + n \lg n$$

Here, $a = 2$, $b = 2$, $f(n) = n \lg n$.

2.

3.

4.

5. a) Here, $a = 2, b = 2, f(n) = 4$.

Since $f(n) = n^{\log_2 2+3} = n^{\log_b a+\epsilon}$ where $\epsilon = 3$, and $af(\frac{n}{b}) = 2\left(\frac{n^4}{16}\right) = \frac{n^4}{8} \leq cn^4$ where $c = \frac{1}{8}$ for sufficiently large n , the case 3 of master's theorem applies.

Thus, $T(n)$ has upper bound of $\mathcal{O}(n^4)$ and lower bounds of $\Omega(n^4)$, or $\Theta(n^4)$.

b) Here $a = 1, b = \frac{10}{7}, f(n) = n$.

Since $f(n) = 1 = n^{0+1} = n^{\log_{10/7}(1)+1} = n^{\log_b(a)+\epsilon}$, where $\epsilon = 1$, and $af(\frac{n}{b}) = \frac{7n}{10} \leq cn^4$ where $c = \frac{7}{10}$ for sufficiently large n , the case 3 of master's theorem applies.

Thus, $T(n)$ has upper bound of $\mathcal{O}(n)$ and lower bounds of $\Omega(n^4)$, or $\Theta(n)$.

c) Here we have $a = 16, b = 4, f(n) = n^2$.

Since $f(n) = n^2 = n^{\log_4 16} = n^{\log_b a}$, case 2 of master's theorem applies.

Thus, $T(n)$ has upper bound of $\mathcal{O}(n^2 \lg n)$ and lower bounds of $\Omega(n^2 \lg n)$.

d) Here we have $a = 7, b = 3, f(n) = n^2$.

Since $n^2 = n^{\log_3 7+\epsilon} = n^{\log_b a+\epsilon}$ where $\epsilon \approx 0.23$, and $af(\frac{n}{b}) \leq cn^2$ where $c = \frac{7}{9}$, the case 3 of master's theorem applies.

Thus, $T(n)$ has upper bound of $\mathcal{O}(n^2)$ and lower bounds of $\Omega(n^2)$, or $\Theta(n^2)$.

e) Here we have $a = 7, b = 2, f(n) = n^2$.

Since $f(n) = n^2 = n^{\log_2(7)-\epsilon}$, where $\epsilon \approx 0.81$, case 1 of master theorem applies.

Thus, $T(n)$ has upper bound of $\mathcal{O}(n^{\log_2 7})$ and lower bound of $\Omega(n^{\log_2 7})$, or $\Theta(n^{\log_2 7})$.

f) Here we have $a = 2, b = 4, f(n) = \sqrt{n}$.

Since $f(n) = \sqrt{n} = n^{\log_4 2} = n^{\log_b a}$, case 2 of master's theorem applies.

Thus $T(n)$ has upper bound of $\mathcal{O}(\sqrt{n} \lg n)$, and lower bound of $\Omega(\sqrt{n} \lg n)$, or $\Theta(\sqrt{n} \lg n)$.

g) **Solution:**

$T(n)$	n^2	n^2
$T(n - 2)$	$(n-2)^2$	$(n-2)^2$
$T(n - 4)$	$(n-4)^2$	$(n-4)^2$
$T(1)$	$T(1)$	$(n - 2i)^2$