

# CSC373 Worksheet 3 Solution

July 29, 2020

1. Using the following formula

$$M[i, j] = \begin{cases} 0 & \text{if } i = j \\ \min_{i \leq k \leq j} M[i, k] + M[k + 1, j] + p_{i-1}p_kp_j & \text{if } i < j \end{cases} \quad (1)$$

we have

C	1	2	3	4	5	6
1	0	350	770	612	1212	1422
2	x	0	840	462	1662	1362
3	x	x	0	252	1092	1098
4	x	x	x	0	1440	936
5	x	x	x	x	0	720
6	x	x	x	x	x	0

And an optimal parenthesization is

**My Work:**

$(A_1A_2)(A_3A_4)(A_5A_6)$

**Notes:**

- Sequence of Dimensions

The sequence of dimensions  $\langle p_0 = 5, p_1 = 10, p_2 = 3, p_3 = 12, p_4 = 5, p_5 = 50, p_6 = 6 \rangle$  means there are 6 matrices with dimensions  $p_{i-1} \times p_i$

–  $A_1 \rightarrow 5 \times 10$

- $A_2 \rightarrow 10 \times 3$
- $A_3 \rightarrow 3 \times 12$
- $A_4 \rightarrow 12 \times 5$
- $A_5 \rightarrow 5 \times 50$
- $A_6 \rightarrow 50 \times 6$
- Dynamic Programming
  - Is applied to optimization problems
  - Applies when the subproblems overlap
  - Uses the following sequence of steps
    1. Characterize the structure of an optimal solution
    2. Recursively define the value of an optimal solution
    3. Construct an optimal solution from computed information
- Matrix-chain Multiplication
  - Is an optimization problem solved using dynamic programming
  - Goal is to find matrix parenthesis with fewest number of operations

**Example:**

Given chain of matrices  $\langle A, B, C \rangle$ , it's fully parenthesized product is:

- \*  $(AB)C$  needs  $(10 \times 30 \times 5) + (10 \times 5 \times 60) = 1500 + 3000 = 4500$  operations
- \*  $A(BC)$  needs  $(30 \times 5 \times 60) + (10 \times 30 \times 60) = 27000$  operations

Thus,  $(AB)C$  performs more efficiently than  $A(BC)$ .

- Is stated as: given a chain  $\langle A_1, A_2, \dots, A_n \rangle$  of  $n$  matrices, where for  $i = 1, 2, \dots, n$  matrix  $A_i$  has dimension  $p_{i-1} \times p_i$ , fully parenthesize the product  $A_1 A_2 \dots A_n$  in a way that minimizes the number of scalar multiplications.
- Steps
  1. **Check is the problem has Optimal Substructure**

Let us adopt the notation  $A_{i\dots j}$  where  $i \leq j$ , for the matrix that results from evaluating the product  $A_i A_{i+1} \dots A_j$ .

Assume the solution has the following parentheses:

$$(A_{i\dots k})(A_{k+1\dots j})$$

If there is a better way to multiply  $(A_{i\dots k})$ , then we would have a more optimal solution.

This would be a contradiction, as we already stated that we have the optimal solution for  $A_{i\dots j}$ .

Therefore, this problem has optimal substructure.

## 2. Find the Recursive Solution

Let  $M[i, j]$  be the cost of multiplying matrices from  $A_i$  to  $A_j$

We want to find out at which ' $k$ ' returns the fewest number of multiplications, or the minimum number of  $M$ .

The recursive formula for the cost of multiplying from  $A_i$  to  $A_j$  is

$$M[i, j] = \begin{cases} 0 & \text{if } i = j \\ \min_{i \leq k \leq j} M[i, k] + M[k + 1, j] + p_{i-1}p_kp_j & \text{if } i < j \end{cases} \quad (2)$$

## 3. Computing the Estimated Cost

\* Steps

- 1) Fill the table for  $i = j$
- 2) Fill the table for  $i < j$  with a spread of 1
- 3) Repeat 2 with the increased value of spread

### Example:

Given

$\langle A_1, A_2, A_3, A_4, A_5 \rangle$

where

- \*  $A_1 \rightarrow 4 \times 10$
- \*  $A_2 \rightarrow 10 \times 3$
- \*  $A_3 \rightarrow 3 \times 12$
- \*  $A_4 \rightarrow 12 \times 20$
- \*  $A_5 \rightarrow 20 \times 7$

we have:

- 1) Fill the table for  $i = j$

1)  $i = j$

$i \backslash j$	1	2	3	4	5
1	0				
2	x	0			
3	x	x	0		
4	x	x	x	0	
5	x	x	x	x	0

$$M[i, j] = \begin{cases} 0 & \text{if } i = j \\ \min_{i \leq k \leq j} M[i, k] + M[k+1, j] + p_{i-1}p_kp_j & \text{if } i < j \end{cases}$$

2) Fill the table for  $i < j$  with a spread of 1

2)  $(i = 1, j = 2), (i = 2, j = 3), (i = 3, j = 4), (i = 4, j = 5)$

$i \backslash j$	1	2	3	4	5
1	0	120			
2	x	0	360		
3	x	x	0	720	
4	x	x	x	0	1680
5	x	x	x	x	0

since

\*  $i = 1, j = 2$

$$M[1, 2] = \min_{1 \leq k \leq 2} (M[1, 1] + M[1, 2] + p_{i-1}p_kp_j) \quad (3)$$

$$= \min_{1 \leq k \leq 2} (0 + 0 + p_0p_1p_2) \quad (4)$$

$$= \min_{1 \leq k \leq 2} (0 + 0 + 4 \cdot 10 \cdot 3) \quad (5)$$

$$= 120 \quad (6)$$

where  $p_0 = 3$  is from the dimension  $3 \times 10$  of  $A_1$ ,  $p_k = 10$  is from the dimension of  $3 \times 10$  of  $A_1$ .

\*  $i = 2, j = 3$

$$M[2, 3] = \min_{2 \leq k \leq 3} (M[2, 2] + M[3, 3] + p_{i-1}p_kp_j) \quad (7)$$

$$= \min_{2 \leq k \leq 3} (0 + 0 + p_1p_2p_3) \quad (8)$$

$$= \min_{2 \leq k \leq 3} (0 + 0 + 10 \cdot 3 \cdot 12) \quad (9)$$

$$= 360 \quad (10)$$

\*  $i = 3, j = 4$

$$M[3, 4] = \min_{3 \leq k \leq 4} (M[3, 3] + M[4, 4] + p_{i-1}p_kp_j) \quad (11)$$

$$= \min_{3 \leq k \leq 4} (0 + 0 + p_2p_3p_4) \quad (12)$$

$$= \min_{3 \leq k \leq 4} (0 + 0 + 3 \cdot 12 \cdot 20) \quad (13)$$

$$= 720 \quad (14)$$

\*  $i = 4, j = 5$

$$M[4, 5] = \min_{4 \leq k \leq 5} (M[4, 4] + M[5, 5] + p_{i-1}p_kp_j) \quad (15)$$

$$= \min_{4 \leq k \leq 5} (0 + 0 + p_3p_4p_5) \quad (16)$$

$$= \min_{4 \leq k \leq 5} (0 + 0 + 12 \cdot 20 \cdot 7) \quad (17)$$

$$= 1680 \quad (18)$$

3) Repeat 2 with the increased value of spread

2) ( $i = 1, j = 2$ ), ( $i = 2, j = 3$ ), ( $i = 3, j = 4$ ), ( $i = 4, j = 5$ )

i \ j	1	2	3	4	5
1	0	120	264	1080	1344
2	x	0	360	1320	1350
3	x	x	0	720	1140
4	x	x	x	0	1680
5	x	x	x	x	0

\*  $i = 1, j = 3$

$k = 1$

$$M[1, 3] = M[1, 1] + M[2, 3] + p_{i-1}p_kp_j \quad (19)$$

$$= 0 + 360 + p_0p_1p_3 \quad (20)$$

$$= 0 + 360 + 4 \cdot 10 \cdot 12 \quad (21)$$

$$= 0 + 360 + 480 \quad (22)$$

$$= 840 \quad (23)$$

$k = 2$

$$M[1, 3] = M[1, 2] + M[3, 3] + p_{i-1}p_kp_j \quad (24)$$

$$= 120 + 0 + p_0p_2p_3 \quad (25)$$

$$= 120 + 0 + 4 \cdot 10 \cdot 12 \quad (26)$$

$$= 120 + 0 + 144 \quad (27)$$

$$= 264 \quad (28)$$

Thus,  $\min_{1 \leq k \leq 3} M[1, 3] = 264$ .

\*  $i = 2, j = 4$

$k = 2$

$$M[2, 4] = M[2, 2] + M[3, 4] + p_{i-1}p_kp_j \quad (29)$$

$$= 0 + 720 + p_1p_2p_4 \quad (30)$$

$$= 0 + 720 + 10 \cdot 3 \cdot 20 \quad (31)$$

$$= 0 + 720 + 600 \quad (32)$$

$$= 1320 \quad (33)$$

$k = 3$

$$M[2, 4] = M[2, 2] + M[3, 4] + p_{i-1}p_kp_j \quad (34)$$

$$= 360 + 0 + p_1p_3p_4 \quad (35)$$

$$= 360 + 0 + 10 \cdot 12 \cdot 20 \quad (36)$$

$$= 360 + 0 + 2400 \quad (37)$$

$$= 2760 \quad (38)$$

Thus,  $\min_{2 \leq k \leq 4} M[2, 4] = 1320$ .

$$* \ i = 3, j = 5$$

$$\underline{k = 3}$$

$$M[3, 5] = M[3, 3] + M[3, 5] + p_{i-1}p_kp_j \quad (39)$$

$$= 0 + 1680 + p_2p_3p_5 \quad (40)$$

$$= 0 + 1680 + 3 \cdot 12 \cdot 7 \quad (41)$$

$$= 0 + 1680 + 252 \quad (42)$$

$$= 1932 \quad (43)$$

$$\underline{k = 4}$$

$$M[3, 5] = M[3, 4] + M[5, 5] + p_{i-1}p_kp_j \quad (44)$$

$$= 720 + 0 + p_2p_4p_5 \quad (45)$$

$$= 720 + 0 + 3 \cdot 20 \cdot 7 \quad (46)$$

$$= 720 + 420 \quad (47)$$

$$= 1140 \quad (48)$$

$$\text{Thus, } \min_{3 \leq k \leq 5} M[3, 5] = 1140.$$

$$* \ i = 2, j = 5$$

$$\underline{k = 2}$$

$$M[2, 5] = M[2, 2] + M[3, 5] + p_{i-1}p_kp_j \quad (49)$$

$$= 0 + 1140 + p_1p_2p_5 \quad (50)$$

$$= 0 + 1140 + 10 \cdot 3 \cdot 7 \quad (51)$$

$$= 0 + 1140 + 210 \quad (52)$$

$$= 1350 \quad (53)$$

$$\underline{k = 3}$$

$$M[2, 5] = M[2, 3] + M[4, 5] + p_{i-1}p_kp_j \quad (54)$$

$$= 360 + 1680 + p_1p_3p_5 \quad (55)$$

$$= 2040 + 10 \cdot 12 \cdot 7 \quad (56)$$

$$= 2040 + 840 \quad (57)$$

$$= 2880 \quad (58)$$

$$\underline{k = 4}$$

$$M[2, 5] = M[2, 4] + M[5, 5] + p_{i-1}p_kp_j \quad (59)$$

$$= 1320 + p_1p_3p_5 \quad (60)$$

$$= 1320 + 10 \cdot 20 \cdot 7 \quad (61)$$

$$= 1320 + 1400 \quad (62)$$

$$= 2720 \quad (63)$$

$$\text{Thus, } \min_{2 \leq k \leq 5} M[2, 5] = 1350.$$

$$* \ i = 1, j = 5$$

$$\underline{k = 1}$$

$$M[1, 5] = M[1, 1] + M[3, 5] + p_{i-1}p_kp_j \quad (64)$$

$$= 0 + 1350 + p_0p_1p_5 \quad (65)$$

$$= 0 + 1350 + 4 \cdot 10 \cdot 7 \quad (66)$$

$$= 0 + 1350 + 280 \quad (67)$$

$$= 1630 \quad (68)$$

$$\underline{k = 2}$$

$$M[1, 5] = M[1, 2] + M[3, 5] + p_{i-1}p_kp_j \quad (69)$$

$$= 120 + 1140 + p_0p_2p_5 \quad (70)$$

$$= 120 + 1140 + 4 \cdot 3 \cdot 7 \quad (71)$$

$$= 1260 + 84 \quad (72)$$

$$= 1344 \quad (73)$$

$$\underline{k = 3}$$

$$M[1, 5] = M[1, 3] + M[4, 5] + p_{i-1}p_kp_j \quad (74)$$

$$= 264 + 1680 + p_0p_3p_5 \quad (75)$$

$$= 264 + 1680 + 4 \cdot 12 \cdot 7 \quad (76)$$

$$= 1944 + 336 \quad (77)$$

$$= 2280 \quad (78)$$

$$\underline{k = 4}$$



$$M[1, 5] = M[1, 4] + M[5, 5] + p_{i-1}p_kp_j \quad (79)$$

$$= 1080 + 0 + p_0p_4p_5 \quad (80)$$

$$= 1080 + 4 \cdot 20 \cdot 7 \quad (81)$$

$$= 1080 + 560 \quad (82)$$

$$= 1640 \quad (83)$$

Thus,  $\min_{1 \leq k \leq 5} M[1, 5] = 1344$ .

#### 4. Constructing the Optimal Solution

3)

i \ j	1	2	3	4	5
1	0	120	264	1080	1344
2	x	0	360	1320	1350
3	x	x	0	720	1140
4	x	x	x	0	1680
5	x	x	x	x	0

$(A_1A_2)((A_3A_4)A_5)$

So, the optimal solution is  $(A_1A_2)((A_3A_4)A_5)$

#### References:

1)

#### 2. Example:

- *MATRIX-CHAIN-ORDER* computes the table  $s$  containing optimal costs