

CSC236 Worksheet 2 Solution

Hyungmo Gu

April 29, 2020

Question 1

- **Statement:** Any full binary tree with at least 1 node has more leaves than internal nodes.

Rough Work:

Let n be the total number of nodes in a full binary tree.

We will prove the statement by complete induction on n .

1. Base Case ($n = 1$)

Let $n = 1$.

We need to prove the full binary tree with 1 total number of nodes has more leaves than internal nodes.

The definition of leaf tells us that a node that has no children is a leaf.

Since the full binary tree with 1 node has no children, we can conclude the node is a leaf.

Using this fact, we can write the full binary tree has 1 leaf.

Now, the definition of an internal node tells us that a node is an internal node if it is not a leaf.

Since there is 1 leaf and 1 total node in the full binary tree, we can write there is 0 internal node.

So, since there is 1 leaf node and 0 internal node, we can conclude the full binary tree has more leaves than internal nodes.

2. Base Case ($n = 2$)

Let $n = 2$.

We need to prove the full binary tree with 2 total number of nodes has more leaves than internal nodes.

The definition of full binary tree tells us that a binary tree is a full binary tree if every internal node has two children.

Since we know by observation that the tree with 2 total number of nodes has 1 leaf and 1 internal node, using above fact, we can write the tree is not a full binary tree.

Then, by vacuous truth, we can conclude the tree has more leaves than internal nodes.

3. Base Case ($n = 3$)

4. Inductive Step

Notes:

– Complete Induction

- * **Statement:** $\forall i \in \mathbb{N}, \forall n \in \mathbb{N}, n < i \Rightarrow A(n) \Rightarrow \forall i \in \mathbb{N}, A(i)$
- * **Statement Alt.:** $\left(\forall n \in \mathbb{N}, \left[\bigwedge_{k=0}^{n-1} P(k) \right] \Rightarrow P(n) \right) \Rightarrow \forall n \in \mathbb{N}, P(n)$

* **Simple Example 1:**

Statement: $\forall n \in \mathbb{N}, n \geq 0 \Rightarrow 10 \mid (n^5 - n)$

We will prove the statement by strong induction on n .

1. Base Case ($n = 0$)

Let $n = 0$.

We need to prove $10 \mid (n^5 - n)$ is true when $n = 0$. That is, there exists $k \in \mathbb{Z}$ such that $(n^5 - n) = 10k$.

Let $k = 0$.

Starting from the left hand side, using the fact $n = 0$, we can write

$$(n^5 - n) = 0 \tag{1}$$

Then, because we know $10k = 0$, we can conclude

$$(n^5 - n) = 10k \tag{2}$$

2. Base Case ($n = 1$)

Let $n = 1$.

We need to prove $10 \mid (n^5 - n)$ is true when $n = 1$. That is, there exists $k \in \mathbb{Z}$ such that $(n^5 - n) = 10k$.

Let $k = 0$.

Starting from the left hand side, using the fact $n = 1$, we can write

$$(n^5 - n) = 1 - 1 \tag{3}$$

$$= 0 \tag{4}$$

Then, because we know $10k = 0$, we can conclude

$$(n^5 - n) = 10k \tag{5}$$

3. Inductive Step

Assume $k \geq 1$. Assume that for all natural number i satisfying $0 \leq i \leq k$, $10 \mid (i^5 - i)$. That is, $\exists d \in \mathbb{Z}$, $(i^5 - i) = 10d$.

We need to prove $\exists \tilde{d} \in \mathbb{Z}$ such that $((k+1)^5 - (k+1)) = 10\tilde{d}$.

Let $\tilde{d} = c + (k-1)^4 + 4 \cdot (k-1)^3 + 8 \cdot (k-1)^2 + 8 \cdot (k-1) + 3$.

Starting from $((k+1)^5 - (k+1))$, using binominal theorem, we can write,

$$(k+1)^5 - (k+1) = \left[(k-1) + 2\right]^5 - \left[(k-1) + 2\right] \quad (6)$$

$$= \sum_{b=0}^5 \binom{5}{b} (k-1)^{5-b} \cdot 2^b \quad (7)$$

$$= (k-1)^5 + 10 \cdot (k-1)^4 + 40 \cdot (k-1)^3 + 80 \cdot (k-1)^2 + 80 \cdot (k-1) + 32 - \left[(k-1) + 2\right] \quad (8)$$

$$= \left[(k-1)^5 - (k-1)\right] + 10 \cdot (k-1)^4 + 40 \cdot (k-1)^3 + 80 \cdot (k-1)^2 + 80 \cdot (k-1) + 30 \quad (9)$$

(The reason why $k-1$ is chosen instead of $k-2$ and $k-3$ is because of the last term $2^5 = 32$, i.e $32 - 2 = 30$)

Then, because we know $0 \leq k-1 \leq k$ and $10 \mid (k-1)^5 - (k-1)$ from the header, we can write $\exists c \in \mathbb{Z}$ such that $(k-1)^5 - (k-1) = 10c$, and

$$(k+1)^5 - (k+1) = 10c + 10 \cdot (k-1)^4 + 40 \cdot (k-1)^3 + 80 \cdot (k-1)^2 + 80 \cdot (k-1) + 30 \quad (10)$$

$$(k+1)^5 - (k+1) = 10 \cdot \left[c + (k-1)^4 + 4 \cdot (k-1)^3 + 8 \cdot (k-1)^2 + 8 \cdot (k-1) + 3 \right] \quad (11)$$

$$(12)$$

Then, because we know $\tilde{d} = c + (k-1)^4 + 4 \cdot (k-1)^3 + 8 \cdot (k-1)^2 + 8 \cdot (k-1) + 3$ from the header, we can conclude

$$(k+1)^5 - (k+1) = 10\tilde{d} \quad (13)$$

Question 2

Question 3