

# Worksheet 16 Review

April 2, 2020

## Question 1

a. Let  $k \in \mathbb{N}$ .

Here, the minimum possible change occurs for the loop variable in a single iteration when  $i = i + 1$ .

The maximum possible change occurs for the loop variable in a single iteration when  $i = i + 6$ .

The exact upper bound of the variable after  $k$  iteration is

$$i_k \leq 6k \tag{1}$$

The exact lower bound of the variable after  $k$  iteration is

$$k \leq i_k \tag{2}$$

Using the fact that the termination occurs when  $i_k = n$ , we can calculate that for the upper bound, the loop terminates when

$$6k \geq n \quad (3)$$

$$k \geq \frac{n}{6} \quad (4)$$

Because we know  $\frac{n}{6}$  may be a decimal, we can conclude the closest value at which the loop terminates is when

$$k = \left\lceil \frac{n}{6} \right\rceil \quad (5)$$

Using the same fact, we can calculate that for the lower bound, the loop terminates when

$$k \geq n \quad (6)$$

It follows from above that for the lower bound, the smallest value of  $k$  at which the loop termination occurs is when

$$k = n \quad (7)$$

Then, we can conclude the function has asymptotic lower bound of  $\Omega(n)$ , and asymptotic upper bound of  $\mathcal{O}(n)$ .

Then, since both  $\Omega$  and  $\mathcal{O}$  have the same value,  $\Theta(n)$  is also true.

**Correct Solution:**

Here, the minimum possible change occurs for the loop variable in a single iteration when  $i = i + 1$ .

The maximum possible change occurs for the loop variable in a single iteration when  $i = i + 6$ .

The exact upper bound of the variable after k iteration is

$$i_k \leq 6k \quad (8)$$

The exact lower bound of the variable after k iteration is

$$k \leq i_k \quad (9)$$

Using the fact that the termination occurs when  $i_k = n$ , we can calculate that for the upper bound, the loop terminates when

$$6k \geq n \quad (10)$$

$$k \geq \frac{n}{6} \quad (11)$$

Because we know  $\frac{n}{6}$  may be a decimal, we can conclude the closest value at which the loop terminates is when

$$k = \left\lceil \frac{n}{6} \right\rceil + 1 \quad (12)$$

Using the same fact, we can calculate that for the lower bound, the loop terminates when

$$k \geq n \quad (13)$$

It follows from above that for the lower bound, the smallest value of  $k$  at which the loop termination occurs is when

$$k = n + 1 \tag{14}$$

Then, we can conclude the function has asymptotic lower bound of  $\Omega(n)$ , and asymptotic upper bound of  $\mathcal{O}(n)$ .

Since both  $\Omega$  and  $\mathcal{O}$  have the same value,  $\Theta(n)$  is also true.

#### Notes:

- where is  $+1$  coming from? Is it coming from the loop variable  $i = 0$ ?

b. Let  $k \in \mathbb{N}$ .

#### Part 1 (Determining maximum and minimum possible change in a single iteration):

It follows from observation that the minimum possible change occurs when  $i = i \cdot 2$ , and the maximum possible change when  $i = i \cdot 3$ .

#### Part 2 (Determining lower bound and upper bound of loop iteration):

Because we know the smallest possible change occurs when  $i = i \cdot 2$  occurs repeatedly, we can conclude that at  $k^{th}$  iteration  $i_k$  has the lower bound of  $2^k$ .

Similarly, because we know largest possible change occurs when  $i = i \cdot 3$  occurs repeatedly, we can conclude that at  $k^{th}$  iteration,  $i_k$  has the upper bound of  $3^k$ .

Then, by putting together, we can conclude that

$$2^k \leq i_k \leq 3^k \tag{1}$$

**Part 3 (Determining exact number of iterations for the lower bound and upper bound):**

Because we know the loop runs until  $i_k < n$ , we can conclude that at lower bound, termination occurs when

$$i_k \geq n \tag{2}$$

$$2^k \geq n \tag{3}$$

$$\log_2 2^k \geq \log_2 n \tag{4}$$

$$k \geq \log_2 n \tag{5}$$

Using the fact that we are looking for smallest value of  $k$ , we can calculate that for lower bound

$$k = \lceil \log_2 n \rceil + 1 \tag{6}$$

Similarly, for the upper bound, loop terminates when

$$i_k \geq n \tag{7}$$

$$3^k \geq n \tag{8}$$

$$\log_3 3^k \geq \log_3 n \tag{9}$$

$$k \geq \log_3 n \tag{10}$$

Using the fact, we can calculate that for upper bound,

$$k = \lceil \log_3 n \rceil + 1 \tag{11}$$

**Part 4 (Determining Big-Oh and Omega):**

Because we know  $\log_2 n$  dominates  $\log_3 n$ , we can conclude  $\log_2 n$  is the asymptotic upper bound, and  $\log_3 n$  is the asymptotic lower bound.

Then, we can conclude the algorithm has  $\mathcal{O}(\log_2 n)$  and  $\Omega(\log_3 n)$ .

**Notes:**

- How come in solution, **+1** doesn't exist? What rules of thumb i can follow to better determine whether **+1** should be included?

**Question 2**

**Question 3**