

Worksheet 14 Review

April 1, 2020

Question 1

- a. Since the inner loop starts at $j = 0$ and finishes at $j = n - 1$ with j increasing by 1 per iteration, we can conclude that the inner loop has

$$\lceil n - 1 - 0 + 1 \rceil = n \tag{1}$$

iterations.

Since the inner loop takes 1 step per iteration, we can conclude that the inner loop has the total cost of

$$n \cdot 1 = n \tag{2}$$

steps.

For the outer loop, because it starts at $i = 0$ and ends at $i = n - 1$ with i increasing by 5 per iteration, we can conclude that the outer loop has

$$\left\lceil \frac{n - 1 - 0 + 1}{5} \right\rceil = \left\lceil \frac{n}{5} \right\rceil \tag{3}$$

iterations.

Since each iteration in the outer loop takes n steps, we can conclude the outer loop has the total cost of

$$n \cdot n = n^2 \quad (4)$$

steps.

Since we are ignoring the cost of the loop variables, the total cost of the algorithm is $n^2 + n$ steps.

Then, because we know the algorithm takes total of $n^2 + n$ steps, we can conclude the algorithm has the runtime of $\Theta(n^2)$.

- b. We will determine the exact cost and theta of this algorithm by first calculating the exact cost of inner loop 1

```
1  j = 1
2  while j < n:
3      j = j * 3
4
```

and then, calculating the exact cost of inner loop 2

```
1  k = 0
2  while k < n:
3      k = k + 2
4
```

and then, calculating the exact cost of the outer loop using the information from the exact cost of inner loop 1 and inner loop 2

```
1  i = 4
2  while i < n:
3      j = 1
4      while j < n:
5          j = j * 3
6      k = 0
7      while k < n:
8          k = k + 2
9      i = i + 1
10
```

and then, we will finish off by calculating the theta of the outer loop.

Question 2