

Midterm 2 Version 1 Solution

April 3, 2020

Question 1

a.

$100 \div 2 = 50$, Remainders **0**

$50 \div 2 = 25$, Remainders **0**

$25 \div 2 = 12$, Remainders **1**

$12 \div 2 = 6$, Remainders **0**

$6 \div 2 = 3$, Remainders **0**

$3 \div 2 = 1$, Remainders **1**

$1 \div 2 = 0$, Remainders **1**

Then, it follows from above that the binary representation of 100 is $(1100100)_2$.

b. The smallest number that can be expressed by an n -digit balanced ternary representation is

$$\sum_{i=0}^{n-1} d_i \cdot 3^i, \text{ where } d_i \in \{0, 1, 2\} \quad (1)$$

Correct Solution:

The smallest number that can be expressed by an n-digit balanced ternary representation is

$$-\left[\sum_{i=0}^{n-1} 3^i\right] \quad (1)$$

Notes:

- Realized professor is asking for an example of the smallest number.
- Ternary representation of a number

$$\sum_{i=0}^{n-1} d_i \cdot 3^i, \text{ where } d_i \in \{0, 1, 2\}$$

- Learned a negative number could be expressed in in ternary or binary representation of numbers.

c.

$f(n) \in \Omega(n)$	True	$g(n) \in \Omega(n)$	False	$f(n) \in \mathcal{O}(g(n))$	False
$f(n) \in \Theta(g(n))$	False	$g(n) \in \Theta(\log_3 n)$	True	$f(n) + g(n) \in \Theta(f(n))$	True

Notes:

- $\forall g : \mathbb{N} \rightarrow \mathbb{R}^{\geq 0}$, and all numbers $a \in \mathbb{R}^{\geq 0}$, if $g \in \mathcal{O}(f)$, then $f + g \in \mathcal{O}(f)$
- $g \in \Theta(f) : g \in \mathcal{O}(f) \wedge g \in \Omega(f)$
or
 $g \in \Theta(f) : \exists c_1, c_2, n_1 \in \mathbb{R}^+, \forall n \in \mathbb{N}, n \geq n_1 \Rightarrow c_1 g(n) \leq f(n) \leq c_2 g(n)$, where $f, g : \mathbb{N} \rightarrow \mathbb{R}^{\geq 0}$
- $g \in \Omega(f) : \exists c, n_o \in \mathbb{R}^+, \forall n \in \mathbb{N}, n \geq n_o \Rightarrow g(n) \geq cf(n)$, where $f, g : \mathbb{N} \rightarrow \mathbb{R}^{\geq 0}$

- $g \in \mathcal{O}(f) : \exists c, n_o \in \mathbb{R}^+, \forall n \in \mathbb{N}, n \geq n_o \Rightarrow g(n) \leq cf(n)$, where $f, g : \mathbb{N} \rightarrow \mathbb{R}^{\geq 0}$

d.

k	0	1	2
i_k	$3 = 3^1$	$9 = 3^2$	$81 = 3^4$

The value of i_k is

$$3^{2^k} \quad (1)$$

Notes:

- Realized we are only concerned with the lines $\mathbf{i} = \mathbf{i} * \mathbf{i}$ and $\mathbf{i} = \mathbf{3}$
- e. The number of iterations the function's loop will run is

$$\lceil \log_2 \log_3 n \rceil - 1 \quad (1)$$

Notes:

- The loop terminates when $3^{2^{(k+1)}} = i_{k+1} = i_k \cdot i_k \geq n$.
- $\forall x \in \mathbb{Z}, \forall y \in \mathbb{R}, \lfloor x + y \rfloor = x + \lfloor y \rfloor$
- Feel more confident there is no need to add an extra $+1$. Done by playing with examples (i.e is $\lceil \log \log_3(82) \rceil - 1$ true? Would the loop run only once?)

Question 2

- **Predicate Logic:** $\forall n \in \mathbb{N}, n \geq 3 \Rightarrow 5^n + 50 < 6^n$

Proof. Let $n \in \mathbb{N}$.

We will prove the statement by induction on n .

Base Case ($n = 3$):

Let $n = 3$.

We want to show $5^3 + 50 < 6^3$.

Starting from $5^3 + 50$, we can calculate

$$5^3 + 50 = 125 + 50 \tag{1}$$

$$= 175 \tag{2}$$

$$< 216 \tag{3}$$

$$< 6^3 \tag{4}$$

Inductive Case:

Let $n \in \mathbb{N}$. Assume $n \geq 3$ and $5^n + 50 < 6^n$.

We want to show $5^{n+1} + 50 < 6^{n+1}$.

Starting from $5^{n+1} + 50$, we can calculate

$$50^{n+1} + 50 = 5^n \cdot 5 + 50 \tag{5}$$

$$< 5^n \cdot 5 + 50 \cdot 5 \tag{6}$$

$$< 5(5^n + 50) \tag{7}$$

Then,

$$50^{n+1} + 5 < 5 \cdot 6^n \tag{8}$$

$$< 6 \cdot 6^n \tag{9}$$

$$< 6^{n+1} \tag{10}$$

by using inductive hypothesis (i.e $5^n + 50 < 6^n$)

□

Question 3

Question 4