

FIELD THEORY

December 14, 2025

CONTENTS

| | |
|---------------------------------------|---|
| 1. Definitions | 1 |
| 1.1. Algebraic field extensions | 1 |
| 1.2. Transcendence degrees | 6 |
| Appendix A. Miscellaneous definitions | 7 |
| References | 9 |

1. DEFINITIONS

1.1. Algebraic field extensions.

Definition 1.1.1 (Field). A *field* is commutative division ring. In other words, a field is a commutative ring for which all nonzero elements have a multiplicative inverse.

Definition 1.1.2. Let K and L be fields (Definition 1.1.1).

1. A map $\phi : K \rightarrow L$ is a *field homomorphism* if it satisfies the following axioms for all $x, y \in K$:
 - (a) $\phi(x + y) = \phi(x) + \phi(y)$ (additivity);
 - (b) $\phi(xy) = \phi(x)\phi(y)$ (multiplicativity);
 - (c) $\phi(1_K) = 1_L$ (unitality).

Equivalently, a field homomorphism is a ring homomorphism (Definition A.0.7) between fields.

If such a map exists, we often say that *K embeds into L* ; this terminology is justified because field homomorphisms are injective (Definition A.0.8) as set maps (Proposition 1.1.3).

2. A field homomorphism $\phi : K \rightarrow L$ is an *isomorphism* if it is bijective (Definition A.0.8).
3. The set of all homomorphisms from a field K to a field L is denoted by $\text{Hom}(K, L)$.
4. Assuming that K and L have a common subfield F , an *F -embedding of K into L* is an embedding $K \rightarrow L$ that acts as the identity map (Definition A.0.9) on F . The set of F -embeddings of K into L is denoted by $\text{Hom}_F(K, L)$.

If K and L are fields such that K embeds into L , then that means that there is a subfield (Definition 1.1.5) of L that is isomorphic to K .

Proposition 1.1.3. Let K and L be fields (Definition 1.1.1) and let $\phi : K \rightarrow L$ be a field homomorphism (Definition 1.1.2). Then ϕ is injective (Definition A.0.8).

Definition 1.1.4 (Vector space over a field). Let $(k, +, \cdot)$ be a field (Definition 1.1.1). A *vector space over k* or a *k -vector space* is a triple $(V, +, \cdot)^1$ where

1. $(V, +)$ is an abelian group, and
2. \cdot is a map $k \times V \rightarrow V$, called *scalar multiplication*

such that the following axioms hold for all $a, b \in k$ and all $u, v \in V$:

1. (Compatibility with field multiplication)

$$(ab) \cdot v = a \cdot (b \cdot v).$$

2. (Identity scalar)

$$1 \cdot v = v.$$

3. (Distributivity over vector addition)

$$a \cdot (u + v) = a \cdot u + a \cdot v.$$

4. (Distributivity over scalar addition)

$$(a + b) \cdot v = a \cdot v + b \cdot v.$$

Definition 1.1.5 (Field Extension). Let K be a field and let L be a field such that $K \subseteq L$ and the operations of K are the restrictions of those of L . Then L is called a *extension field* (or just an *extension*) of K . The notation L/K is often used synonymously; we say that L/K is a *field extension*. Moreover, K is said to be a *subfield of L* .

Definition 1.1.6. Let R be a ring. There exists a unique ring homomorphism (Definition A.0.7) $\psi : \mathbb{Z} \rightarrow R$ defined by mapping $n \mapsto n \cdot 1_K$. The non-negative generator of the kernel of this map, $\ker(\psi) = \langle p \rangle \subseteq \mathbb{Z}$, is called the *characteristic of R* , denoted by $\text{char}(R)$.

Proposition 1.1.7. Let K be a field (Definition 1.1.1). Then $\text{char}(K)$ (Definition 1.1.6) is either 0 or a prime number p .

Definition 1.1.8. (♠ TODO: finite field, rational numbers) Let K be a field. The *prime subfield of K* is the intersection of all subfields (Definition 1.1.5) of K .

- If $\text{char}(K) = p > 0$, the prime subfield is isomorphic to the finite field $\mathbb{F}_p \cong \mathbb{Z}/p\mathbb{Z}$.
- If $\text{char}(K) = 0$, the prime subfield is isomorphic to the field of rational numbers \mathbb{Q} .

Lemma 1.1.9. Let L/K be a field extension (Definition 1.1.5). Then L is a vector space (Definition 1.1.4) over K .

Definition 1.1.10 (Degree of an Extension, Finite Extension). Let L/K be a field extension (Definition 1.1.5). The *degree* of the extension $[L : K] := \dim_K(L)$, i.e. the dimension of L as a K -vector space (Lemma 1.1.9). If $[L : K] < \infty$, the extension L/K is called a *finite field extension* and L is said to be a *finite extension of K* .

¹Note that $+$ and \cdot are abuse of notation here as these are already used for the addition and multiplication of \cdot .

Definition 1.1.11 (Algebraic Element, Algebraic Extension). Let L/K be a field extension (Definition 1.1.5) and let $x \in L$.

- If there exists a nonzero polynomial $f(t) \in K[t]$ such that $f(x) = 0$, then x is called an *algebraic element over K* . There exists a unique such monic irreducible polynomial $f(t)$, which is called the *minimal polynomial of x over K* .
- Otherwise, x is called a *transcendental element over K* .

If every $x \in L$ is algebraic over K , then L/K is called an *algebraic extension*.

Definition 1.1.12. A field (Definition 1.1.1) F is said to be *algebraically closed* if every nonconstant polynomial $f(x) \in F[x]$ has a root in F , i.e., for every such $f(x)$ there exists $a \in F$ with $f(a) = 0$.

Definition 1.1.13. Let K be a field. A field extension L/K is called an *algebraic closure of K* if the following two conditions hold:

1. Every element $a \in L$ is algebraic over K (Definition 1.1.11).
2. The field L is algebraically closed (Definition 1.1.12).

One often writes an algebraic closure of K by \overline{K} .

Definition 1.1.14 (Field Generated by Elements). Let L/K be a field extension (Definition 1.1.5) and let $S \subseteq L$ be a subset. The *field generated by S over K* is the smallest subfield of L that contains K and S . Equivalently, it is the intersection of all subfields of L containing $K \cup S$. This field is denoted by $K(S)$.

In the special case where $S = \{x_1, \dots, x_n\}$ is finite, one writes $K(x_1, \dots, x_n)$. If $n = 1$, then $K(x)$ is called a *simple field extension*.

Definition 1.1.15 (Finitely generated field extension). Let E/F be a field extension (Definition 1.1.5). We say that E/F is *finitely generated* if there exist finitely many elements $\alpha_1, \dots, \alpha_n \in E$ such that $E = F(\alpha_1, \dots, \alpha_n)$ (Definition 1.1.14), where $F(\alpha_1, \dots, \alpha_n)$ denotes the smallest subfield of E containing F and $\{\alpha_1, \dots, \alpha_n\}$.

Definition 1.1.16 (Field Automorphism). Let L be a field. A *field automorphism of L* is a bijective ring homomorphism $L \rightarrow L$.

Definition 1.1.17 (Automorphism Group of a Field Extension). Let L/K be a field extension (Definition 1.1.5).

- An automorphism (Definition 1.1.16) $\sigma : L \rightarrow L$ is called a *K -automorphism* if σ is a field automorphism of L that fixes every element of K , i.e. $\sigma(a) = a$ for all $a \in K$.
- The set of all such K -automorphisms of L forms a group (Definition A.0.2) under composition, called the *automorphism group of the extension L/K* . It is usually denoted by $\text{Aut}(L/K)$ or $\text{Aut}_K(L)$.

Definition 1.1.18 (Normal Extension). Let K be a field and let $S \subseteq K[t]$ be a set of polynomials.

- A field $L \supseteq K$ is a *splitting field of S over K* if

- every polynomial in S splits completely into linear factors over L , and
- L is generated over K (Definition 1.1.14) by the roots of the polynomials in S , namely

$$L = K(\{\alpha \mid f(\alpha) = 0, f \in S\}).$$

- An algebraic extension L/K is called a **normal extension** if L is the splitting field of a family of polynomials in $K[t]$.

Definition 1.1.19 (Separable Element, Separable Extension). Let L/K be a field extension (Definition 1.1.5) and let $x \in L$ be algebraic over K (Definition 1.1.11) with minimal polynomial (Definition 1.1.11) $m_{x,K}(t) \in K[t]$.

- The element x is **separable over K** if $m_{x,K}(t)$ has distinct roots in a splitting field (Definition 1.1.18).
- The element x is **inseparable over K** otherwise.

An algebraic extension (Definition 1.1.11) L/K is called **separable extension** if every element $x \in L$ is separable over K .

See also Definition 1.2.6, which defines separable field extensions in greater generality.

Definition 1.1.20. A field F is said to be **separably closed** if every nonconstant separable polynomial $f(x) \in F[x]$ has a root in F , i.e., for every such $f(x)$ there exists $a \in F$ with $f(a) = 0$ (equivalently, every separable polynomial factors completely into linear factors over F).

Definition 1.1.21. Let K be a field (Definition 1.1.1). A field extension (Definition 1.1.5) L/K is called a **separable closure of K** if the following hold:

1. The extension L/K is separable algebraic (Definition 1.1.19) (i.e., every $a \in L$ is separable over K).
2. The field L is separably closed (Definition 1.1.20).

One often writes a separable closure of K by K^{sep} . Some also write separable closures of K by \overline{K} , which may conflict with the common notation for algebraic closures of K .

Definition 1.1.22 (Galois Extension). An extension L/K is called a **Galois extension** if it is both a normal extension (Definition 1.1.18) and a separable algebraic extension (Definition 1.1.19). Its **Galois group**, usually denoted by $\text{Gal}(L/K)$, is defined to be the automorphism group $\text{Aut}(L/K)$ (Definition 1.1.17).

Theorem 1.1.23 (Fundamental Theorem of Galois Theory). Let L/K be a finite (Definition 1.1.10) Galois extension of fields (Definition 1.1.22) with Galois group $G = \text{Gal}(L/K)$. Then there is an inclusion-reversing bijection between the set of subgroups $H \leq G$ and the set of intermediate fields $K \subseteq F \subseteq L$, given by (♠ TODO: L^H)

$$H \longmapsto L^H := \{x \in L \mid \sigma(x) = x \text{ for all } \sigma \in H\},$$

$$F \longmapsto \text{Gal}(L/F).$$

This correspondence has the following properties:

- (i) L^H is a subfield of L containing K for each subgroup $H \leq G$.
- (ii) $\text{Gal}(L/F)$ is a subgroup of G for each intermediate field F .
- (iii) $H_1 \subseteq H_2 \iff L^{H_2} \subseteq L^{H_1}$.
- (iv) $[L^H : K] = |G : H|$, $[L : L^H] = |H|$.
- (v) F/K is Galois if and only if $\text{Gal}(L/F) \trianglelefteq G$,
in which case $\text{Gal}(F/K) \cong G / \text{Gal}(L/F)$.

Corollary 1.1.24 (Normality of the whole extension). If L/K is a finite (Definition 1.1.10) Galois extension of fields (Definition 1.1.22) with Galois group G , then $K = L^G$. (♠ TODO: L^G)

Theorem 1.1.25 (Primitive element theorem). Let L/K be a finite (Definition 1.1.10) separable extension (Definition 1.1.19). Then there exists an element $\alpha \in L$ such that

$$L = K(\alpha).$$

(Definition 1.1.14) In other words, every finite separable extension is simple (Definition 1.1.14).

Lemma 1.1.26 (Normal extension characterization). (♠ TODO: embedding) Let L/K be a finite extension. Then L/K is normal (Definition 1.1.18) if and only if every K -embedding $\sigma : L \rightarrow \bar{K}$ into an algebraic closure (Definition 1.1.11) \bar{K} satisfies $\sigma(L) = L$.

Theorem 1.1.27 (Infinite Galois Theory). Let L/K be a (possibly infinite) Galois extension with Galois group $G = \text{Gal}(L/K)$. Endow G with the Krull topology, i.e. the unique compact, Hausdorff, totally disconnected topology whose neighborhood basis at the identity is given by the open subgroups $\text{Gal}(L/F)$, where F runs through the finite Galois intermediate extensions $K \subseteq F \subseteq L$.

Then there is an inclusion-reversing bijection between: - the closed subgroups $H \leq G$, and - the intermediate fields $K \subseteq F \subseteq L$,

given by

$$\begin{aligned} H &\longmapsto L^H := \{x \in L \mid \sigma(x) = x \text{ for all } \sigma \in H\}, \\ F &\longmapsto \text{Gal}(L/F). \end{aligned}$$

This correspondence satisfies:

- (i) $H \leq G$ closed $\iff F = L^H$ for some intermediate field F .
- (ii) F/K is finite Galois $\iff \text{Gal}(L/F)$ is open in G .
- (iii) F/K is Galois $\iff \text{Gal}(L/F) \trianglelefteq G$,
in which case $\text{Gal}(F/K) \cong G / \text{Gal}(L/F)$.

Proposition 1.1.28 (Krull topology characterization). For a (possibly infinite) Galois extension L/K , the Galois group $G = \text{Gal}(L/K)$ is a profinite group, i.e. a compact, Hausdorff, totally disconnected topological group isomorphic to an inverse limit of finite groups.

1.2. Transcendence degrees.

Definition 1.2.1 (Algebraically independent elements). Let E/F be a field extension (Definition 1.1.5), and let $S = \{x_i\}_{i \in I}$ be a subset of E indexed by some set I . We say that S is **algebraically independent over F** if for every finite subset $\{x_{i_1}, \dots, x_{i_n}\} \subseteq S$ and every non-zero polynomial $P \in F[X_1, \dots, X_n]$, we have $P(x_{i_1}, \dots, x_{i_n}) \neq 0$. Equivalently, S is algebraically independent over F if the natural ring homomorphism (Definition A.0.7)

$$F[X_i : i \in I] \rightarrow E$$

sending $X_i \mapsto x_i$ is injective, where $F[X_i : i \in I]$ denotes the polynomial ring (Definition A.0.4) over F in the indeterminates $\{X_i\}_{i \in I}$.

Definition 1.2.2 (Transcendence basis). Let E/F be a field extension (Definition 1.1.5). A subset $T \subseteq E$ is called a **transcendence basis** of E over F if:

1. T is algebraically independent over F , i.e., there is no non-trivial polynomial relation with coefficients in F among finitely many elements of T , and
2. E is algebraic (Definition 1.1.11) over $F(T)$, where $F(T)$ denotes the field generated by (Definition 1.1.14) F and T .

Definition 1.2.3 (Transcendence degree). Let E/F be a field extension (Definition 1.1.5). The **transcendence degree of E over F** , denoted $\text{tr. deg}_F(E)$ or $\text{trdeg}_F(E)$, is defined as the cardinality (Definition A.0.5) of any transcendence basis (Definition 1.2.2) of E over F . By a classical result, all transcendence bases of E over F have the same cardinality, so the transcendence degree is well-defined.

If E/F is algebraic (Definition 1.1.11), then $\text{tr. deg}_F(E) = 0$. If no transcendence basis is finite, then $\text{tr. deg}_F(E)$ is an infinite cardinal.

Definition 1.2.4 (Function field). Let k be a field (Definition 1.1.1). A field K is called a **function field over k** (or a **function field in n variables over k**) if K is a finitely generated field extension (Definition 1.1.15) of k with transcendence degree (Definition 1.2.3) n over k , where $n \geq 1$. Equivalently, the function field in n variables over k is isomorphic to the fraction field (Definition A.0.6) of the polynomial ring (Definition A.0.4) $k[x_1, \dots, x_n]$ where x_1, \dots, x_n are indeterminate variables. Accordingly, the function field in n variables over k is often denoted as $k(x_1, \dots, x_n)$.

When $n = 1$, we say that K is a **function field of one variable over k** or an **algebraic function field over k** and denote it by $k(x)$.

Definition 1.2.5 (Separating transcendence basis). Let E/F be a field extension (Definition 1.1.5), and let $T \subseteq E$ be a transcendence basis of E over F . We say that T is a **separating transcendence basis** if the algebraic extension $E/F(T)$ (Definition 1.1.14) is separable (Definition 1.1.19), i.e., every element of E is separable over $F(T)$ (meaning that its minimal polynomial (Definition 1.1.11) over $F(T)$ has no repeated roots in an algebraic closure).

Definition 1.2.6 (Separable field extension). Let E/F be a field extension (Definition 1.1.5). In general (allowing transcendental extensions), E/F is called **separable** if there exists a separating transcendence basis (Definition 1.2.5) for E over F . Equivalently, E/F is separable if every finitely generated (Definition 1.1.15) intermediate field $F \subseteq K \subseteq E$ has a separating

transcendence basis over F . In particular, a field extension E/F is algebraic and separable in the above sense if and only if it is a separable algebraic extension in the sense of Definition 1.1.19.

APPENDIX A. MISCELLANEOUS DEFINITIONS

Definition A.0.1. Let X and Y be sets. A **map** (or **function**) from X to Y is a rule f assigning to each element $x \in X$ exactly one element $f(x) \in Y$. We write $f : X \rightarrow Y$.

We say that X is the **domain** and that Y is the **codomain of f** .

Definition A.0.2 (Groups). A **group** is a pair (G, \cdot) where G is a set and $\cdot : G \times G \rightarrow G$ is a binary operation, subject to the following conditions:

1. (Associativity) For all $g, h, k \in G$ one has

$$(g \cdot h) \cdot k = g \cdot (h \cdot k).$$

2. (Identity element) There exists an element $e \in G$ such that for all $g \in G$,

$$e \cdot g = g \cdot e = g.$$

3. (Inverse element) For all $g \in G$ there exists an element $g^{-1} \in G$ such that

$$g \cdot g^{-1} = g^{-1} \cdot g = e.$$

The element e is called the **identity element of G** , and g^{-1} is called the **inverse of g** .

Equivalently, a group is a monoid with inverse elements.

A group (G, \cdot) is often simply written as G , when the notation for the binary operation \cdot is clear.

An **abelian group** or synonymously, a **commutative group**, is a group (G, \cdot) whose binary operation \cdot is **abelian** or **commutative**, i.e. satisfies

$$g \cdot h = h \cdot g$$

for all $g, h \in G$.

Definition A.0.3. Let $(R, +, \cdot)$ and $(S, +, \cdot)$ be rings, not assumed to be commutative. A function $f : R \rightarrow S$ is called a **ring homomorphism** if for all $r_1, r_2 \in R$ the following properties hold:

1. $f(r_1 + r_2) = f(r_1) + f(r_2)$,
2. $f(r_1 r_2) = f(r_1) f(r_2)$,
3. $f(1_R) = 1_S$ where 1_R and 1_S denote the multiplicative identities in R and S , respectively.

A ring homomorphism is said to be a **ring isomorphism** if it is invertible as a map of sets.

An **R -ring** refers to a ring S equipped with a ring homomorphism $f : R \rightarrow S$.

We note that a ring homomorphism $f : R \rightarrow S$ yields a natural left R -module structure on S and a natural right R -module structure on S respectively as follows for $r \in R$ and $s \in S$:

$$r \cdot s = f(r) \cdot s$$

$$s \cdot r = s \cdot f(r).$$

However, these left and right module structures need not yield a two-sided R -module structure.

Definition A.0.4 (Polynomial ring over a commutative ring). Let R be a commutative ring. For a set of variables $\{x_i\}_{i \in I}$, the *polynomial ring in variables $\{x_i\}$ over R* , denoted by notations such as $R[x_i \mid i \in I]$ or $R[x_i]_{i \in I}$, is defined as the commutative R -algebra whose elements are finite R -linear combinations of monomials in the variables x_i , where the variables commute with each other and with elements of R .

That is, $R[x_i \mid i \in I]$ is the free commutative R -algebra generated by the set $\{x_i\}$.

In the case that I is a finite set, and writing y_1, \dots, y_n for the variables x_i , it is customary to let $R[y_1, \dots, y_n]$ denote the polynomial ring.

Definition A.0.5. An ordinal number κ is a *cardinal number* (or simply a *cardinal*) if for every ordinal $\alpha < \kappa$, there is no bijection (Definition A.0.8) between α and κ . Equivalently, a cardinal is an initial ordinal—an ordinal that is not equinumerous with any smaller ordinal.

The *cardinality of an arbitrary set X* , denoted by $|X|$, $\text{card}(X)$, or $\#X$, is the unique cardinal number κ such that there exists a bijection between X and κ . (The existence of such a κ for every set requires the Axiom of Choice).

Definition A.0.6. Let R be an integral domain, and consider the set $R \times (R \setminus \{0\})$ as above. Define a relation \sim on $R \times (R \setminus \{0\})$ by declaring that

$$(a, b) \sim (c, d) \quad \text{if and only if} \quad ad = bc,$$

for $a, c \in R$ and $b, d \in R \setminus \{0\}$. This relation is an equivalence relation. Its equivalence classes are denoted by $\frac{a}{b}$.

The set of equivalence classes

$$\left\{ \frac{a}{b} \mid a \in R, b \in R \setminus \{0\} \right\}$$

under the relation \sim defined above is called the *field of fractions of R* , and is denoted by $\text{Frac}(R)$.

The operations on $\text{Frac}(R)$ are defined by

$$\begin{aligned} \frac{a}{b} + \frac{c}{d} &= \frac{ad+bc}{bd}, \\ \frac{a}{b} \cdot \frac{c}{d} &= \frac{ac}{bd}, \end{aligned}$$

for $a, c \in R$ and $b, d \in R \setminus \{0\}$. With these operations, $\text{Frac}(R)$ is a field (Definition 1.1.1).

Definition A.0.7. Let $(R, +, \cdot)$ and $(S, +, \cdot)$ be rings, not assumed to be commutative. A function $f : R \rightarrow S$ is called a *ring homomorphism* if for all $r_1, r_2 \in R$ the following properties hold:

1. $f(r_1 + r_2) = f(r_1) + f(r_2)$,
2. $f(r_1 r_2) = f(r_1) f(r_2)$,
3. $f(1_R) = 1_S$ where 1_R and 1_S denote the multiplicative identities in R and S , respectively.

A ring homomorphism is said to be a **ring isomorphism** if it is invertible as a map of sets.

An **R -ring** refers to a ring S equipped with a ring homomorphism $f : R \rightarrow S$.

We note that a ring homomorphism $f : R \rightarrow S$ yields a natural left R -module structure on S and a natural right R -module structure on S respectively as follows for $r \in R$ and $s \in S$:

$$r \cdot s = f(r) \cdot s$$

$$s \cdot r = s \cdot f(r).$$

However, these left and right module structures need not yield a two-sided R -module structure.

Definition A.0.8. Let X and Y be sets and let $f : X \rightarrow Y$ be a function.

- The function f is said to be **injective** (or **one-to-one**) if for all $x_1, x_2 \in X$, $f(x_1) = f(x_2)$ implies $x_1 = x_2$.
- The function f is said to be **surjective** (or **onto**) if for every $y \in Y$ there exists $x \in X$ such that $f(x) = y$.
- The map f is **bijective** if it is both injective and surjective. In this case, there exists a unique **inverse map** $f^{-1} : Y \rightarrow X$ such that for all $x \in X$ and $y \in Y$,

$$f^{-1}(f(x)) = x \text{ and } f(f^{-1}(y)) = y.$$

Definition A.0.9. Let X be a set. The **identity function on X** , denoted by **id_X** , is the function (Definition A.0.1) $\text{id}_X : X \rightarrow X$ defined by

$$\text{id}_X(x) = x \quad \text{for all } x \in X.$$

It is the unique function on X satisfying $f \circ \text{id}_X = f = \text{id}_X \circ f$ for every function $f : X \rightarrow Y$ and every function $f : Y \rightarrow X$.

REFERENCES

- [Arm92] Brumer Armand. The average rank of elliptic curves i. *Inventiones mathematicae*, 109(1):445–472, 1992.
- [Ayo23] Joseph Ayoub. Counterexamples to F. Morel’s conjecture on $\pi_0^{\mathbb{D}^1}$. *Comptes Rendus. Mathématique*, 361:1087–1090, 2023.
- [BBD82] Alexander A. Beilinson, Joseph Bernstein, and Pierre Deligne. Analyse et topologie sur les espaces singuliers (i). *Astérisque*, 100, 1982.
- [BC19] Tilman Bauer and Magnus Carlson. Tensor products of affine and formal abelian groups. *Documenta Mathematica*, 24:2525–2582, 2019.
- [BFK⁺17] Valentin Blomer, Étienne Fouvry, Emmanuel Kowalski, Philippe Michel, and Djordje Milićević. Some applications of smooth bilinear forms with kloosterman sums. *Proceedings of the Steklov Institute of Mathematics*, 296:18–29, 2017.
- [BGI71] Pierre Berthelot, Alexander Grothendieck, and Luc Illusie. *Théorie des Intersections et Théorème de Riemann-Roch (SGA6)*, volume 225 of *Lecture Notes in Mathematics*. Springer-Verlag, 1971.

- [BH12] Salman Baig and Chris Hall. Experimental data for goldfeld’s conjecture over function fields. *Experimental Mathematics*, 21(4):362–374, 2012.
- [BLGHT11] Tom Barnet-Lamb, David Geraghty, Michael Harris, and Richard Taylor. A family of Calabi–Yau varieties and potential automorphy ii. *Publications of the Research Institute for Mathematical Sciences*, 47(1):29–98, 2011.
- [Bor12] Armand Borel. *Linear algebraic groups*, volume 126 of *Graduate Texts in Mathematics*. Springer New York, 2012.
- [BS15a] Manjul Bhargava and Arul Shankar. Binary quartic forms having bounded invariants, and the boundedness of the average rank of elliptic curves. *Annals of Mathematics*, pages 191–242, 2015.
- [BS15b] Manjul Bhargava and Arul Shankar. Ternary cubic forms having bounded invariants, and the existence of a positive proportion of elliptic curves having rank 0. *Annals of Mathematics*, pages 587–621, 2015.
- [BS15c] Bhargav Bhatt and Peter Scholze. The pro-étale topology for schemes. *Astérisque*, 360:99–201, 2015.
- [BSD65] Bryan John Birch and Peter Francis Swinnerton-Dyer. Notes on elliptic curves. ii. *Journal für die reine und angewandte Mathematik*, 218:79–108, 1965.
- [BSS18] Bhargav Bhatt, Christian Schnell, and Peter Scholze. Vanishing theorems for perverse sheaves on abelian varieties, revisited. *Selecta Mathematica*, 24:63–84, 2018.
- [Cho08] Utsav Choudhury. Homotopy theory of schemes and a^1 -fundamental groups. Master’s thesis, Università degli Studi di Padova, 2008.
- [CHT08] Laurent Clozel, Michael Harris, and Richard Taylor. Automorphy for some l -adic lifts of automorphic mod l galois representations. *Publications mathématiques*, 108:1–181, 2008.
- [DA73] Pierre Deligne and Michael Artin. *Théorie des Topos et Cohomologies Étale des Schémas. Séminaire de Géométrie Algébrique due Bois-Marie 1963-1964 (SGA 4)*. Lecture Notes in Mathematics. Springer Berlin, 1973.
- [DBG⁺77] Pierre Deligne, Jean-François Boutot, Alexander Grothendieck, Luc Illusie, and Jean-Louis Verdier. *Étale Cohomology. Séminaire de Géométrie Algébrique due Bois-Marie 1963-1964 (SGA 4 1/2)*. Lecture Notes in Mathematics. Springer-Verlag, 1977.
- [Del80] Pierre Deligne. La conjecture de Weil : II. *Publications Mathématiques de l’IHÉS*, 52:137–252, 1980.
- [Del89] Pierre Deligne. Le groupe fondamental de la droite projective moins trois points. In *Galois Groups over \mathbb{Q} : Proceedings of a Workshop Held March 23–27, 1987*, pages 79–297. Springer, 1989.
- [Dri89] Vladimir Gershonovich Drinfeld. Cohomology of compactified manifolds of modules of f -sheaves. *Journal of Soviet Mathematics*, 46(2):1789–1821, 1989.
- [DZ19] Alexander Dunn and Alexandru Zaharescu. Sums of Kloosterman sums over primes in an arithmetic progression. *The Quarterly Journal of Mathematics*, 70(1):319–342, 2019.
- [Eke07] Torsten Ekedahl. *On The Adic Formalism*, pages 197–218. Birkhäuser Boston, Boston, MA, 2007.
- [FFK24] Arthur Forey, Javier Fresán, and Emmanuel Kowalski. Arithmetic fourier transforms over finite fields: generic vanishing, convolution, and equidistribution, 2024.
- [FLR23] Tony Feng, Aaron Landesman, and Eric M. Rains. The geometric distribution of Selmer groups of elliptic curves over function fields. *Mathematische Annalen*, 387:615–687, 2023.
- [Fu15] Lei Fu. *Étale Cohomology Theory*, volume 14 of *Nankai Tracts in Mathematics*. World Scientific, 2015.
- [GL96] Ofer Gabber and François Loeser. Faisceaux pervers ℓ -adiques sur un tore. *Duke Math J.*, 83(3):501–606, 1996.
- [Gol06] Dorian Goldfeld. Conjectures on elliptic curves over quadratic fields. In *Number Theory Carbon-dale 1979: Proceedings of the Southern Illinois Number Theory Conference Carbondale, March 30 and 31, 1979*, pages 108–118. Springer, 2006.
- [GR04] Alexander Grothendieck and Michèle Raynaud. Revêtements étales et groupe fondamental (SGA 1). eprint arXiv math/0206203, 2004. Updated edition of the book of the same title published by Springer-Verlag in 1971 as volume 224 of the series Lecture Notes in Mathematics.

- [Gro77] Alexander Grothendieck. *Cohomologie l -adique et fonctions L* Séminaire de Géométrie Algébrique due Bois-Marie 1965-1966 (SGA 5), volume 589 of *Springer Lecture Notes*. Springer-Verlag, 1977. Avec la collaboration de I. Bucur, C. Houzel, L. Illusie, J.-P. Jouanolou, et J.-P. Serre.
- [GV72] Alexander Grothendieck and Jean-Louis Verdier. *Theorie des Topos et Cohomologie Etale des Schemas. Seminaire de Geometrie Algebrique du Bois-Marie 1963-1964 (SGA 4)*. Lecture Notes in mathematics. Springer-Verlag Berlin Heidelberg, 1 edition, 1972.
- [HB04] David Heath-Brown. The average analytic rank of elliptic curves. *Duke Math. J.*, 122:591–623, 2004.
- [HK25] Chris Hall and Hyun Jong Kim. Independence of ℓ (title to be determined). In progress, 2025.
- [HM73] Dale Husemoller and John Milnor. *Symmetric Bilinear Forms*, volume 73 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 2. Folge*. Springer Berlin Heidelberg, 1973.
- [HSBT05] Michael Harris, Nick Shepherd-Barron, and Richard Taylor. Ihara’s lemma and potential automorphy, 2005.
- [Hub97] Annette Huber. Mixed perverse sheaves for schemes over number fields. *Compositio Mathematica*, 108:107–121, 1997.
- [Jor05] Andrei Jorza. The birch and swinnerton-dyer conjecture for abelian varieties over number fields, 2005.
- [Kat90] Nicholas M. Katz. *Exponential sums and differential equations*, volume 124 of *Annals of Mathematics Studies*. Princeton University Press, 1990.
- [Kat96] Nicholas M. Katz. *Rigid Local Systems*, volume 139 of *annals of Mathematics Studies*. Princeton University Press, 1996.
- [Kat98] Nicholas M. Katz. *Gauss Sums, Kloosterman Sums, and Monodromy Groups*, volume 116 of *Annals of Mathematics Studies*. Princeton University Press, 1998.
- [Kat12] Nicholas M. Katz. *Convolution and Equidistribution Sato-Tate Theorems for Finite-Field Mellin Transforms*, volume 180 of *Annals of Mathematics Studies*. Princeton University Press, 2012.
- [Kim23] Hyun Jong Kim. `trouver`, 2023. GitHub repository: <https://github.com/hyunjongkimmath/trouver>.
- [KL85] Nicholas M. Katz and Gérard Laumon. Transformation de fourier et majoration de sommes exponentielles. *Publications Mathématiques de l’IHÉS*, 62:145–202, 1985.
- [KM23] Seoyoung Kim and M. Ram Murty. From the Birch and Swinnerton-Dyer conjecture to Nagao’s conjecture. *Mathematics of Computation*, 92(339):385–408, 2023.
- [KMS17] Emmanuel Kowalski, Philippe Michel, and Will Sawin. Bilinear forms with Kloosterman sums and applications. *Annals of Mathematics*, 186:413–500, 2017.
- [Krä14] Thomas Krämer. Perverse sheaves on semiabelian varieties. *Rendiconti del Seminario Matematico della Università di Padova*, 132:83–102, 2014.
- [KS99] Nicholas M. Katz and Peter Sarnak. *Random matrices, Frobenius eigenvalues, and monodromy*, volume 45. American Mathematical Society, 1999.
- [KW13] Reinhardt Kiehl and Rainer Weissauer. *Weil Conjectures, Perverse Sheaves and ℓ -adic Fourier Transform*, volume 42 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics*. Springer Berlin, Heidelberg, 2013.
- [KW15] Thomas Krämer and Rainer Weissauer. Vanishing theorems for constructible sheaves on abelian varieties. *J. Algebraic Geometry*, 24:531–568, 2015.
- [Laf02] Laurent Lafforgue. Chtoucas de Drinfeld et correspondance de Langlands. *Inventiones mathematicae*, 147:1–241, 2002.
- [May99] Jon Peter May. *A Concise Course in Algebraic Topology*. Chicago Lectures in Mathematics. University of Chicago Press, 1999.
- [Mil80] James S. Milne. *Etale cohomology*. Number 33 in Princeton Mathematical Series. Princeton university press, 1980.
- [Mil07] James S. Milne. Quotients of Tannakian categories. *Theory and Applications of Categories*, 18(21):654–664, 2007.
- [Mil13] James S. Milne. Lie algebras, algebraic groups, and lie groups, 2013. Available at www.jmilne.org/math/.

- [Mil17] James S. Milne. *Algebraic Groups*, volume 170 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 2017.
- [Mor06] Fabien Morel. A1-algebraic topology. In *International Congress of Mathematicians*, volume 2, pages 1035–1059, 2006.
- [Mor12] Fabien Morel. *A1-Algebraic Topology over a field*. Lecture Notes in Mathematics. Springer Berlin, Heidelberg, 2012.
- [MV99] Fabien Morel and Vladimir Voevodsky. A1-homotopy theory of schemes. *Publications Mathématiques de l’IHÉS*, 90:45–143, 1999.
- [Nag97] Koh-ichi Nagao. $Q(t)$ -rank of elliptic curves and certain limit coming from the local points. *Manuscripta mathematica*, 92(1):13–32, 1997.
- [nLa25a] nLab authors. geometric morphism. <https://ncatlab.org/nlab/show/geometric+morphism>, July 2025. Revision 61.
- [nLa25b] nLab authors. homotopy group of a spectrum. <https://ncatlab.org/nlab/show/homotopy+group+of+a+spectrum>, June 2025. Revision 7.
- [nLa25c] nLab authors. Introduction to Stable homotopy theory – 1-1. <https://ncatlab.org/nlab/show/Introduction+to+Stable+homotopy+theory+--+1-1>, June 2025. Revision 43.
- [nLa25d] nLab authors. model structure on topological sequential spectra. <https://ncatlab.org/nlab/show/model+structure+on+topological+sequential+spectra>, June 2025. Revision 61.
- [nLa25e] nLab authors. point of a topos. <https://ncatlab.org/nlab/show/point+of+a+topos>, July 2025. Revision 53.
- [nLa25f] nLab authors. sheafification. <https://ncatlab.org/nlab/show/sheafification>, September 2025. Revision 40.
- [nLa25g] nLab authors. stable homotopy category. <https://ncatlab.org/nlab/show/stable+homotopy+category>, June 2025. Revision 31.
- [Poo18] Bjorn Poonen. Heuristics for the arithmetic of elliptic curves. In *Proceedings of the International Congress of Mathematicians: Rio de Janeiro 2018*, pages 399–414. World Scientific, 2018.
- [PR12] BJORN POONEN and ERIC RAINS. Random maximal isotropic subspaces and selmer groups. *Journal of the American Mathematical Society*, 25(1):245–269, 2012.
- [Ros02] Michael Rosen. *Number Theory in Function Fields*, volume 210 of *Graduate Texts in Mathematics*. Springer-Verlag New York, 2002.
- [RS98] Michael Rosen and Joseph H. Silverman. On the rank of an elliptic surface. *Inventiones mathematicae*, 133:43–67, 1998.
- [Rud87] Walter Rudin. *Real and Complex Analysis*. Mathematics Series. McGraw-Hill Book Company, 3 edition, 1987.
- [Saw24] Will Sawin. General multiple dirichlet series from perverse sheaves. *Journal of Number Theory*, 262:408–453, 2024.
- [Ser72] Jean-Pierre Serre. Propriétés galoisiennes des points d’ordre fini des courbes elliptiques. *Invent. math*, 15:259–331, 1972.
- [SFFK23] Will Sawin, Arthur Forey, Javier Fresán, and Emmanuel Kowalski. Quantitative sheaf theory. *Journal of the American Mathematical Society*, 36(3):653–726, 2023.
- [Sil89] Joseph H. Silverman. Elliptic curves of bounded degree and height. *Proceedings of the American Mathematical Society*, 105(3):540–545, 1989.
- [Sil09] Joseph H. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Graduate Texts in Mathematics*. Springer-Verlag, 2 edition, 2009.
- [ST21] Will Sawin and Jacob Tsimerman. Bounds for the stalks of perverse sheaves in characteristic p and a conjecture of shende and tsimerman. *Inventiones mathematicae*, 224(1):1–32, 2021.
- [Sta25] The Stacks project authors. The stacks project. <https://stacks.math.columbia.edu>, 2025.
- [Tat65] John T. Tate. Algebraic cycles and poles of zeta functions. In *Arithmetical Algebraic Geometry (Proc. Conf. Purdue Univ., 1963)*, pages 93–110. Harper & Row, 1965. Also in *Collected works of John Tate* (2 vols.), Amer. Math. Soc. (2016), vol. 2.
- [Tat66] John Tate. On the conjectures of Birch and Swinnerton-Dyer and a geometric analog. In *Séminaire Bourbaki : années 1964/65 1965/66, exposés 277-312*, number 9 in *Astérisque*, pages 415–440. Société mathématique de France, 1966. talk:306.

- [Tay08] Richard Taylor. Automorphy for some l -adic lifts of automorphic mod l galois representations. ii. *Publications mathématiques*, 108:183–239, 2008.
- [Voe98] Vladimir Voevodsky. A1-homotopy theory. In *Proceedings of the international congress of mathematicians*, volume 1, pages 579–604. Berlin, 1998.
- [Wei94] Charles A. Weibel. *An Introduction to Homological Algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 1994. First paperback edition 1995 Reprinted 1997.
- [Wik25] Wikipedia contributors. Frobenius endomorphisms#frobenius for schemes — Wikipedia, the free encyclopedia, 2025. [Online; accessed 08-July-2025].
- [You06] Matthew Young. Low-lying zeros of families of elliptic curves. *Journal of the American Mathematical Society*, 19(1):205–250, 2006.