

ÉTALE SHEAVES ON SCHEMES

June 25, 2026

CONTENTS

1. Definitions	3
1.1. Étale morphisms of schemes	3
1.2. Étale covers	5
2. Galois categories	6
3. The étale fundamental group of a scheme at a geometric point	9
4. Sheaves on the small étale site of a scheme	12
4.1. The small and big Étale sites of a scheme	14
4.2. Other common sites of schemes	15
4.3. Sheafification of a presheaf on a site	20
4.4. Examples of sheaves on the étale site of a scheme	21
4.5. Stalk of a (pre)sheaf on the small étale site of a scheme	24
4.6. Constant and locally constant sheaves	25
5. Locally constant sheaves and local systems	26
6. Six functors	26
6.1. The functors on sheaves	26
6.1.1. Inverse and direct images of sheaves via continuous functors of general sites	27
6.1.2. Extension by zero of sheaves via open immersions of schemes	35
6.1.3. Hom and tensor product functors of sheaves	36
6.1.4. Stalks of sheaves under the functors	36
6.1.5. Exactness properties on sheaves of abelian groups	44
6.2. The functors on derived categories of sheaves	45
6.3. Verdier duality for bounded constructible complexes	47

6.4.	Torsion coefficients	47
6.5.	Integral adic coefficients	47
6.6.	Rational coefficients	48
6.7.	Algebraic coefficients	49
6.8.	Various coefficient systems	49
7.	Cohomology of literal sheaves for the small étale site on a scheme	50
7.1.	Sheaf cohomology of a general module over a sheaf of rings on a site	50
7.2.	Étale cohomology of literal sheaves	56
8.	Derived categories of λ -adic sheaves constructible through limits of categories	56
8.1.	ℓ -adic sheaves	57
8.2.	The categories of λ -adic sheaves and A-R λ -adic sheaves	57
8.3.	Constructible sheaves	57
8.4.	λ -adic sheaves	58
8.5.	Rational coefficients	62
8.5.1.	Geometric stalks of λ -adic sheaves	63
8.6.	The category $D_c^b(X, R)$ of integral coefficients	64
8.7.	Geometric stalks of objects of D_c^b categories	68
9.	Theorems	70
9.1.	ℓ -adic cohomology	70
9.2.	Base change theorem	71
9.3.	Poincaré duality	72
9.4.	Lefschetz trace formula	72
9.5.	Weil conjectures	73
10.	Ekedahl's formalism	74
11.	Pro-étale sheaves on schemes	80
11.1.	Pro-étale topology on a scheme	80
11.2.	Adically complete derived modules in replete topoi	81

11.3. Realizing the left-completion of the derived category on the étale site via the pro-étale site	81
11.3.1. Comparison to Ekedahl's theory	81
11.4. Comparison to classical ℓ -adic categories	82
.1. Categorical definitions	82
.1.1. Abelian categories	99
.2. Homological algebra	105
.3. Schemes	110
Appendix A. Presheaves and sheaves	112
Appendix B. Derived categories	115
B.1. Homotopy category of chain complexes of an additive category	115
B.2. Multiplicative systems in categories	119
B.3. Derived categories of abelian categories	120
B.4. Total derived functors	121
B.5. General derived functors for exact multifunctors between triangulated categories	123
B.5.1. Derived multifunctors of additive multifunctors between abelian categories	129
B.6. K -flat resolutions of chain complexes with respect to bi-additive functors	131
B.7. General conditions for defining derived functors	137
B.7.1. Adjunction of derived functors of adjoint functors	138
B.8. Having enough K -injectives and flats for sheaves of modules	138
B.9. Definitions of derived functors for modules of ringed sites	139
Appendix C. Miscellaneous definitions	146

1. DEFINITIONS

1.1. Étale morphisms of schemes.

Notation 1.1.1. Let S be a scheme. Let $\mathbf{Sch}/S$ denote the category of schemes over S (Definition .3.3).

Definition 1.1.2 (Locally of finite presentation morphism of schemes). Let $f : X \rightarrow Y$ be a morphism of schemes.

1. We say that f is *locally of finite presentation* if for every affine open subset $\text{Spec}(B) \subseteq Y$ (Definition .3.4), and every affine open subset $\text{Spec}(A) \subseteq f^{-1}(\text{Spec}(B))$, the induced ring homomorphism $B \rightarrow A$ presents A as a B -algebra of finite presentation (Definition .3.6); that is, A is isomorphic to a quotient of a polynomial ring in finitely many variables over B by a finitely generated ideal:

$$A \cong B[x_1, \dots, x_n]/(f_1, \dots, f_m),$$

for some finite n, m .

2. A morphism of schemes $f : X \rightarrow Y$ is of *finite presentation* if it is locally of finite presentation and quasi-compact and quasi-separated; in particular, f can be covered by finitely many affine opens satisfying the finite presentation condition above.

Definition 1.1.3 (Flat module over a ring). Let R be a (not necessarily commutative) ring (Definition C.0.8).

1. Let M be a left R -module. The module M is said to be *flat (with respect to the left R -module structure)* if the functor

$$- \otimes_R M : \text{Mod}_R \rightarrow \mathbf{Ab}$$

(♠ TODO: comment on how I might as well be talking about other more specific kinds of modules) (Definition C.0.12) from the category of right R -modules to abelian groups is exact; that is, for every exact sequence of right R -modules

$$0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0,$$

the induced sequence

$$0 \rightarrow N' \otimes_R M \rightarrow N \otimes_R M \rightarrow N'' \otimes_R M \rightarrow 0$$

is exact.

Equivalently, M is flat if $\text{Tor}_1^R(-, M) = 0(??)$.

2. Let M be a right R -module. The module M is said to be *flat (with respect to the right R -module structure)* if the functor

$$M \otimes_R - : {}_R\text{Mod} \rightarrow \mathbf{Ab}$$

from the category of left R -modules to abelian groups is exact; that is, for every exact sequence of left R -modules

$$0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0,$$

the induced sequence

$$0 \rightarrow M \otimes_R N' \rightarrow M \otimes_R N \rightarrow M \otimes_R N'' \rightarrow 0$$

is exact.

Equivalently, M is flat if $\text{Tor}_1^R(M, -) = 0(??)$.

Definition 1.1.4 (Flat morphism of schemes). Let $f : X \rightarrow Y$ be a morphism of schemes (Definition .3.2).

1. Let $x \in X$ be a point and let $y = f(x)$. We say that f is **flat at x** if the induced ring homomorphism on local rings (Definition C.0.21)

$$\mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{X,x}$$

makes $\mathcal{O}_{X,x}$ (Definition C.0.23) into a flat (Definition 1.1.3) $\mathcal{O}_{Y,y}$ -module.

2. We say f is **flat** if it is flat at every point $x \in X$.
3. f is **faithfully flat** if it is flat and surjective.

Definition 1.1.5 (Unramified morphism of schemes). A morphism of schemes (Definition .3.2) $f : X \rightarrow Y$ is **unramified** if it is locally of finite type (Definition C.0.19) and the sheaf of relative differentials $\Omega_{X/Y}$ is zero. Equivalently:

- For every $x \in X$, the induced ring map on stalks $\mathcal{O}_{Y,f(x)} \rightarrow \mathcal{O}_{X,x}$ is of finite type,
- and the module of Kähler differentials $\Omega_{\mathcal{O}_{X,x}/\mathcal{O}_{Y,f(x)}}$ is 0.

Definition 1.1.6. A morphism of schemes (Definition .3.2) $f : X \rightarrow Y$ is called **étale** if it satisfies the following conditions:

- f is locally of finite presentation (Definition 1.1.2),
- f is flat (Definition 1.1.4),
- f is unramified (Definition 1.1.5), i.e., the sheaf of relative differentials $\Omega_{X/Y}$ equals 0.

(♠ TODO: relative dimension) Equivalently, a morphism of schemes is étale if and only if it is smooth of relative dimension 0.

Definition 1.1.7. Let S be a scheme and let $f : X \rightarrow S$ be a morphism of schemes over S . The **automorphism group of X over S** , denoted $\text{Aut}(X/S) = \text{Aut}_S(X)$, is the group of all S -scheme (Definition .3.3) automorphisms of X , i.e., all isomorphisms of schemes $\varphi : X \rightarrow X$ such that the diagram

$$\begin{array}{ccc} X & \xrightarrow[\sim]{\varphi} & X \\ & \searrow f & \swarrow f \\ & S & \end{array}$$

commutes. The group operation is composition of morphisms.

1.2. Étale covers.

Definition 1.2.1. Let $f : X \rightarrow Y$ be an affine morphism of schemes (Definition .3.4). We say that f is a **finite morphism** if for every affine open $V = \text{Spec } B \subseteq Y$ with $U = f^{-1}(V) = \text{Spec } A$, the ring A is a finite B -algebra.

Definition 1.2.2. Let X be a scheme (Definition .3.1). A **finite étale cover of X** is a scheme Y equipped with a finite (Definition 1.2.1), étale morphism (Definition 1.1.6) $Y \rightarrow X$.

Such a cover is surjective and hence is genuinely an étale cover. (♠ TODO: prove)

Definition 1.2.3. Let X be a scheme. The **category of finite étale covers of X** is the category whose

- objects are finite étale covers (Definition 1.2.2) $Y \rightarrow X$

- morphisms $(Y_1 \rightarrow X) \rightarrow (Y_2 \rightarrow X)$ are scheme morphisms $Y_1 \rightarrow Y_2$ such that

$$\begin{array}{ccc} Y_1 & \xrightarrow{f} & Y_2 \\ & \searrow p_1 & \swarrow p_2 \\ & X & \end{array}$$

commutes.

In fact, the scheme morphisms $Y_1 \rightarrow Y_2$ are necessarily themselves finite étale (Definition 1.2.2) (♠ TODO: prove).

The category of finite étale covers of X is often denoted by notations such as $\mathbf{F\acute{E}t}(X)$, $\mathbf{F\acute{E}t}_X$, or $\mathbf{F\acute{E}t}/X$.

Definition 1.2.4. Let $f : X \rightarrow Y$ be a finite (Definition 1.2.1) morphism (Definition .3.2). (♠ TODO: define categorical quotient)

1. f is called **Galois** if there exists a finite group G acting on X over Y such that Y is the categorical quotient X/G and f is the natural quotient morphism.
In this case, $G \cong \mathrm{Aut}_Y(X)$ (Definition 1.1.7) if X is connected. A **Galois G -morphism** is a Galois morphism equipped with a fixed isomorphism $\phi : G \xrightarrow{\sim} \mathrm{Aut}_Y(X)$.
2. A **Galois G -morphism** is a Galois morphism equipped with a fixed isomorphism $\phi : G \xrightarrow{\sim} \mathrm{Aut}_Y(X)$.
3. f is called a **Galois cover** if it is a Galois morphism and is surjective.

(♠ TODO: describe Galois covers in terms of G -torsors)

In case f is an étale morphism, then f is equivalently a Galois morphism if there exists a finite group G acting on X over Y such that

- f is identified with the quotient map $X \rightarrow X/G$.
- G acts simply transitively on geometric fibres.

2. GALOIS CATEGORIES

Definition 2.0.1. (♠ TODO: recheck this definition) Let $\mathcal{C}$ be a locally small category (Definition .1.4). We say $\mathcal{C}$ is a **pre-Galois category** if it satisfies the following conditions:

1. $\mathcal{C}$ has a terminal object (Definition .1.10) and all finite (Definition 1.2.1) pullbacks (Definition C.0.15) exist. (♠ TODO: quotient of a category object by a group action or group of automorphisms)
2. $\mathcal{C}$ has finite coproducts (Definition .1.16), including an initial object (Definition .1.10), and the quotient of any object by a finite group of automorphisms exists. (♠ TODO: strict epimorphism) (♠ TODO: connected component)
3. Every morphism $f : X \rightarrow Y$ in $\mathcal{C}$ can be written as a composition $X \xrightarrow{p} Z \xrightarrow{i} Y$ where p is a strict epimorphism and i is a monomorphism (Definition .1.27) which is an isomorphism onto a connected component of Y .

Definition 2.0.2. (♠ TODO: category of finite sets) Let $\mathcal{C}$ be a pre-Galois category (Definition 2.0.1). A functor $F : \mathcal{C} \rightarrow \mathbf{FinSet}$ from $\mathcal{C}$ to the category of finite sets is called a *fiber functor* if:

1. F is exact (preserves finite limits (Definition .1.13) and finite colimits (Definition .1.13)).
2. F is faithful and reflects isomorphisms.

A *Galois category* is a pair $(\mathcal{C}, F)$ consisting of a pre-Galois category and a fiber functor.

Definition 2.0.3 (Connected object of a Galois category). Let $(\mathcal{C}, F)$ be a Galois category (Definition 2.0.2). An object $X \in \text{Ob}(\mathcal{C})$ is called *connected* if it satisfies the following equivalent conditions:

(♠ TODO: define non-emptiness and indecomposability in general)

1. X is non-empty (i.e., $F(X) \neq \emptyset$) and indecomposable: if $X \cong Y \amalg Z$ for objects $Y, Z \in \mathcal{C}$, then either Y or Z is an initial object (Definition .1.10) of $\mathcal{C}$.
2. The group $\text{Aut}(X)$ of automorphisms of X acts transitively on the finite set $F(X)$.

Every object X in a Galois category can be uniquely written as a finite coproduct (Definition .1.16) $X \cong \coprod_{i=1}^n X_i$ of connected objects X_i , unique up to isomorphism and reordering.

Definition 2.0.4 (Galois object of a Galois category). Let $(\mathcal{C}, F)$ be a Galois category (Definition 2.0.2). A *Galois object* of $(\mathcal{C}, F)$ is an object $X \in \text{Ob}(\mathcal{C})$ satisfying the following equivalent conditions:

1. X is a connected object (Definition 2.0.3) of $(\mathcal{C}, F)$ such that the natural group homomorphism $\text{Aut}_{\mathcal{C}}(X) \rightarrow \text{Perm}(F(X))$ is an isomorphism, i.e., $|\text{Aut}_{\mathcal{C}}(X)| = |F(X)|$.
2. The group $\text{Aut}_{\mathcal{C}}(X)$ of automorphisms of X acts simply transitively on the finite set $F(X)$, i.e., for any two elements $s, t \in F(X)$ there exists a unique element $\sigma \in \text{Aut}_{\mathcal{C}}(X)$ such that $\sigma(s) = t$.

A Galois object is also called a *Galois object with simply transitive monodromy*. Every Galois object is connected, but not every connected object is Galois.

Definition 2.0.5. Let $\mathcal{C}, \mathcal{D}$ be categories (Definition .1.1) and let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. (♠ TODO: identity natural transformation) A *natural automorphism of F* is a natural transformation $\eta : F \rightarrow F$ such that there exists a natural transformation $\xi : F \rightarrow F$ such that $\xi \circ \eta, \eta \circ \xi = \text{id}_F$.

Equivalently, a natural automorphism of F is an automorphism of F as an object in the functor category (Definition .1.33) $\text{Fun}(\mathcal{C}, \mathcal{D})$.

Definition 2.0.6. Let $(\mathcal{C}, F)$ be a Galois category (Definition 2.0.2). The *fundamental group of $(\mathcal{C}, F)$* is the group of natural automorphisms (Definition 2.0.5) of $F : \mathcal{C} \rightarrow \mathbf{FinSets}$. It may be denoted by notations such as $\pi(\mathcal{C}, F)$, $\Pi(\mathcal{C}, F)$, or $\text{Aut}(\mathcal{C}, F)$. If the fiber functor F is clear from context, then the F may be suppressed from the notation.

It is profinite (Proposition 2.0.7).

Proposition 2.0.7. Let $(\mathcal{C}, F)$ be a Galois category (Definition 2.0.2). The fundamental group (Definition 2.0.6) of $(\mathcal{C}, F)$ is profinite.

Theorem 2.0.8. Let $(\mathcal{C}, F)$ be a Galois category (Definition 2.0.2). Let $G = \pi(\mathcal{C}, F)$ be the fundamental group (Definition 2.0.6) of $(\mathcal{C}, F)$.

Let **F- G -Set** denote the category of finite sets equipped with a continuous action of G . The functor F induces an equivalence of categories (Definition .1.6)

$$\Phi : \mathcal{C} \xrightarrow{\sim} \mathbf{F-}G\text{-}\mathbf{Set}$$

defined by mapping an object $X \in \mathcal{C}$ to the set $F(X)$ with the natural G -action.

Proposition 2.0.9. Let $(\mathcal{C}, F)$ be a Galois category (Definition 2.0.2).

1. The fiber functor (Definition 2.0.2) F induces a bijection
 - (a) Isomorphism classes of connected (Definition 2.0.3) Galois objects (Definition 2.0.4) $X \in \mathcal{C}$ whose automorphism group is isomorphic to Γ .
 - (b) Open (??) normal subgroups $H \subseteq G$ such that $G/H \cong \Gamma$.
2. An object X of $\mathcal{C}$ is connected if and only if the image of $G \rightarrow \mathrm{Sym}(F(X))$ acts transitively on the set $F(X)$.

(♠ TODO: category of finite etales over a scheme is galois category)

Definition 2.0.10 ([?, Exposé V, Proposition 6.1.]). Let $\mathcal{C}$ and $\mathcal{C}'$ be pre-Galois categories (Definition 2.0.1), and let F' be a fiber functor (Definition 2.0.2) of $\mathcal{C}'$. Let $H : \mathcal{C} \rightarrow \mathcal{C}'$ be a covariant functor. Let $F = F' \circ H$. We say that H is an *exact functor* $(\mathcal{C}, F) \rightarrow (\mathcal{C}', F')$ if the following equivalent conditions hold:

1. H is exact as a functor.
2. H is left exact, transforms finite coproducts (Definition .1.16) to finite coproducts, and epimorphisms (Definition .1.27) into epimorphisms.
3. F is a fiber functor of $\mathcal{C}$.

(♠ TODO: stacks 0BN5)

Lemma 2.0.11 (see [?, 0BN5]). Let $(\mathcal{C}, F)$ and $(\mathcal{C}', F')$ be Galois categories (Definition 2.0.2).

1. Let $H : \mathcal{C} \rightarrow \mathcal{C}'$ be an exact functor (Definition 2.0.10). There exists an isomorphism $t : F' \circ H \rightarrow F$. The choice of t determines a continuous homomorphism $h : G' = \mathrm{Aut}(F') \rightarrow \mathrm{Aut}(F) = G$ (Definition 2.0.6) and a 2-commutative diagram (♠ TODO: 2-diagram)

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{H} & \mathcal{C}' \\ \downarrow & & \downarrow \\ \mathbf{Finite-}G\text{-}\mathbf{Sets} & \xrightarrow{h} & \mathbf{Finite-}G'\text{-}\mathbf{Sets} \end{array}$$

The map h is independent of t up to an inner automorphism of G .

2. Conversely, given a continuous homomorphism $h : G' \rightarrow G$ there is an exact functor $H : \mathcal{C} \rightarrow \mathcal{C}'$ and an isomorphism t recovering h as above.

Lemma 2.0.12 (see [?, 0BN6]). Let $(\mathcal{C}, F)$ and $(\mathcal{C}', F')$ be Galois categories (Definition 2.0.2). Let $H : \mathcal{C} \rightarrow \mathcal{C}'$ be an exact functor (Definition 2.0.10). Let $h : \pi(\mathcal{C}', F') \rightarrow \pi(\mathcal{C}, F)$ be the corresponding morphism of fundamental groups (Definition 2.0.6). Write $*$ and $*$ ' for the final objects (Definition 1.1.10) of $\mathcal{C}$ and $\mathcal{C}'$ respectively. (♠ TODO: the corresponding morphism; need 0BN5) The following are equivalent:

1. h is surjective
2. H is fully faithful
3. whenever $X \in \mathcal{C}$ is connected (Definition 2.0.3), $H(X) \in \mathcal{C}'$ is connected.
4. whenever $X \in \mathcal{C}$ is connected and there is a morphism $*' \rightarrow H(X)$ in $\mathcal{C}'$, there is a morphism $* \rightarrow X$.
5. for any object X of $\mathcal{C}$, the set map $\text{Hom}_{\mathcal{C}}(*, X) \rightarrow \text{Hom}_{\mathcal{C}'}(*', H(X))$ is bijective.

3. THE ÉTALE FUNDAMENTAL GROUP OF A SCHEME AT A GEOMETRIC POINT

Definition 3.0.1. Let S be a scheme (Definition 3.1). There are two inequivalent conventions for the definition of geometric points of S :

1. A *(algebraic) geometric point of S* is a morphism $\bar{s} : \text{Spec}(\Omega) \rightarrow S$ where Ω is a separably closed field.
2. A *(separable) geometric point of S* is a morphism $\bar{s} : \text{Spec}(\Omega) \rightarrow S$ where Ω is an algebraically closed field.

Care should be taken to specify which convention is used in a given context. In general, one should use the algebraic notion of geometric points for algebraic discussions, say those involving the Nullstellensatz, classical intersection theory/degrees, “geometric properties” of schemes over fields, etc., and the separable notion of geometric points for arithmetic discussions, say those involving Galois theory, Galois modules and cohomology, étale sites, étale fundamental groups, stalks of étale sheaves, etc.

The two theory of the two conventions are usually equivalent when “purely inseparable phenomena” are not present and usually inequivalent when purely inseparable phenomena are present.

(♠ TODO: define fiber functor for finite étale covers at a geometric point) (♠ TODO: state how Fet is a galois category) (♠ TODO: describe that connected schemes correspond to connected objects, geometric points correspond to fiber functors, geometric fibers correspond to $F(X)$.)

Definition 3.0.2. Let S be a connected scheme (Definition 3.1). The *étale fundamental group of S at a geometric point $\bar{s} : \text{Spec}(\Omega) \rightarrow S$* is the profinite group

$$\pi_1^{\text{ét}}(S, \bar{s}) := \varprojlim_{\bar{X} \rightarrow S} \text{Aut}_S(X),$$

(Definition 1.1.7) where the inverse limit is taken over all finite étale (Definition 1.1.6) covers $X \rightarrow S$ pointed above $\bar{s}$.

Equivalently, one may take the limit over all finite étale (Definition 1.2.2) Galois covers (Definition 1.2.4) $X \rightarrow S$ pointed above $\bar{s}$. The étale fundamental group is equipped with the profinite topology.

Also equivalently, the étale fundamental group of X at $\bar{x}$ is defined as the fundamental group (Definition 2.0.6) of the Galois category (Definition 2.0.2) $(\mathbf{F}\acute{\text{E}}\text{t}_X, F_{\bar{x}})$ where

$$F_{\bar{x}} : \mathbf{F}\acute{\text{E}}\text{t}_X \rightarrow \mathbf{FSet}$$

(Definition 1.2.3) is the fiber functor defined by $F_{\bar{x}}(Y \rightarrow X) = Y \times_X \text{Spec}(\Omega)$ (Definition C.0.15), where $\text{Spec}(\Omega) \rightarrow X$ is the geometric point (Definition 3.0.1) $\bar{x}$.

Definition 3.0.3. Let X be a scheme (Definition .3.1) and let $\bar{x}$ be a geometric point (Definition 3.0.1) of X . The functor

$$F_{\bar{x}} : \mathbf{F}\acute{\text{E}}\text{t}_X \rightarrow \mathbf{Sets}$$

sending $Y \in \mathbf{F}\acute{\text{E}}\text{t}_X$ to the underlying set of the geometric fiber $Y \times_X \bar{x}$ is called the *fiber functor or fundamental functor of $\mathbf{F}\acute{\text{E}}\text{t}_X$ at $\bar{x}$* .

Lemma 3.0.4 (see [?, 0BNB]). Let X be a connected scheme (Definition .3.1) and let $\bar{x}$ be a geometric point (Definition 3.0.1) of X . The fiber functor (Definition 3.0.3)

$$F_{\bar{x}} : \mathbf{F}\acute{\text{E}}\text{t}_X \rightarrow \mathbf{Sets}, \quad Y \mapsto |Y_{\bar{x}}|$$

sending Y to the underlying set of the geometric fiber $Y_{\bar{x}} = Y \times_X \bar{x}$ makes $\mathbf{F}\acute{\text{E}}\text{t}_X$ a Galois category (Definition 2.0.2).

Theorem 3.0.5. Let S be a connected scheme (Definition .3.1) with a chosen geometric point (Definition 3.0.1) $\bar{s}$.

1. The functor

$$\mathbf{F}\acute{\text{E}}\text{t}_S \rightarrow \mathbf{FinSets}, \quad X \mapsto \text{Hom}_S(\bar{s}, X).$$

is an equivalence of categories (Definition .1.6) between the category of finite étale covers (Definition 1.2.2) of S and the category of finite sets (equipped with the discrete topology) with a continuous (left) action of the étale fundamental group $\pi_1^{\text{et}}(S, \bar{s})$ (Definition 3.0.2) (Definition 3.0.2). We also note that the discrete set $\text{Hom}_S(\bar{s}, X)$ is identifiable with the geometric fiber $X \times_S \bar{s}$.

2. There is a natural bijection between isomorphism classes of finite étale (Definition 1.2.2) Galois covers (Definition 1.2.4) $f : X \rightarrow S$ with Galois group isomorphic to G and continuous group homomorphisms

$$(A) \quad \varphi : \pi_1^{\text{et}}(S, \bar{s}) \rightarrow G$$

up to conjugation in G . More explicitly, two homomorphism φ, φ' correspond to equivalent finite étale Galois G -covers of S if and only if there exists some $g \in G$ such that

$$\varphi'(\gamma) = g\varphi(\gamma)g^{-1}$$

for all $\gamma \in \pi_1^{\text{et}}(S, \bar{s})$.

Concretely, given a finite étale Galois cover $f : X \rightarrow S$, the fiber $f^{-1}(\bar{s}) \cong X \times_S \bar{s}$ has an action of $\pi_1^{\text{et}}(S, \bar{s})$. Since G acts simply transitively on this fiber, there is an

induced group homomorphism (A); the conjugacy class of this group homomorphism corresponds to f .

Furthmore, the covering scheme X is connected if and only if the homomorphism φ is surjective.

Definition 3.0.6. Let $f : X \rightarrow Y$ be a morphism of schemes (Definition .3.2). The *base change functor on $\mathbf{F}\acute{\text{E}}\text{t}_Y$* is the functor

$$\mathbf{F}\acute{\text{E}}\text{t}_Y \rightarrow \mathbf{F}\acute{\text{E}}\text{t}_X, \quad Z \mapsto Z \times_Y X$$

(Definition 1.2.3) (Definition C.0.15).

Lemma 3.0.7. Let $f : X \rightarrow Y$ be a morphism of connected schemes (Definition .3.1). Let $\bar{x}$ be a geometric point (Definition 3.0.1) of X and let $\bar{y} = f(\bar{x})$.

The *base change functor on $\mathbf{F}\acute{\text{E}}\text{t}_Y$*

$$\mathbf{F}\acute{\text{E}}\text{t}_Y \rightarrow \mathbf{F}\acute{\text{E}}\text{t}_X, \quad Z \mapsto Z \times_Y X$$

(Definition 1.2.3) (Definition C.0.15) is an exact functor of Galois categories (Definition 2.0.10) when $\mathbf{F}\acute{\text{E}}\text{t}_X$ and $\mathbf{F}\acute{\text{E}}\text{t}_Y$ are equipped with the fiber functors (Definition 3.0.3) $F_{\bar{x}}$ and $F_{\bar{y}}$ for any geometric point (Definition 3.0.1) $\bar{x}$ of X and $\bar{y} = f(\bar{x})$.

Lemma 3.0.8 (see [?, 0BND]). Let $f : X \rightarrow Y$ be a morphism of connected schemes (Definition .3.1). Let $\bar{x}$ be a geometric point (Definition 3.0.1) of X and let $\bar{y} = f(\bar{x})$. There is a canonical continuous homomorphism

$$f_* : \pi_1(X, \bar{x}) \rightarrow \pi_1(Y, \bar{y})$$

such that the diagram

$$\begin{array}{ccc} \mathbf{F}\acute{\text{E}}\text{t}_Y & \xrightarrow{\text{base change}} & \mathbf{F}\acute{\text{E}}\text{t}_X \\ F_{\bar{y}} \downarrow & & \downarrow F_{\bar{x}} \\ \text{Finite-}\pi_1(Y, \bar{y})\text{-Sets} & \xrightarrow{f_*} & \text{Finite-}\pi_1(X, \bar{x})\text{-Sets} \end{array}$$

(Definition 1.2.3) (Definition 3.0.3) (♠ TODO: pushforward) is commutative.

Proof. Apply Lemma 3.0.7 and obtain the continuous group homomorphism via Lemma 2.0.11 □

Lemma 3.0.9 (see [?, Tag 0BQE]). (♠ TODO: scheme theoretic image) Let $f : X \rightarrow Y$ be a morphism of schemes (Definition .3.2). If $f(X)$ is dense in Y , then the base change functor (Definition 3.0.6)

$$\mathbf{F}\acute{\text{E}}\text{t}_Y \rightarrow \mathbf{F}\acute{\text{E}}\text{t}_X$$

is faithful.

Lemma 3.0.10 (see [?, 0BQG], see also the various statements in [?, 0BQD]). Let X be a connected (Definition 2.0.3) scheme (Definition 3.1). Let $j : U \hookrightarrow X$ be the embedding of a dense open subscheme. Assume that

1. every quasi-compact open (Definition C.0.6) of X has finitely many irreducible components (♠ TODO: punctured spectrum)
2. for every $x \in X \setminus U$ the punctured spectrum of the strict henselization (Definition 4.5.7) of $\mathcal{O}_{X,x}$ (Definition C.0.23) is connected (Definition 2.0.3).

then the base change functor (Definition 3.0.6) $\text{F}\acute{\text{E}}\text{t}_X \rightarrow \text{F}\acute{\text{E}}\text{t}_U$ is fully faithful and hence (Lemma 2.0.12) the induced homomorphism Lemma 3.0.8 $\pi_1(U, \bar{x}) \rightarrow \pi_1(X, \bar{x})$ is surjective.

Proposition 3.0.11 ([?, Lemma 0BQI]). Let X be an irreducible, geometrically unibranch scheme. For any nonempty open $U \subseteq X$ and geometric point (Definition 3.0.1) $\bar{u}$ of U , the canonical map

$$\pi_1^{\acute{\text{e}}\text{t}}(U, \bar{u}) \rightarrow \pi_1^{\acute{\text{e}}\text{t}}(X, \bar{u})$$

is surjective.

Letting η be the generic point of X and letting $\bar{\eta}$ be a geometric point (Definition 3.0.1) of X with image η , the canonical map

$$\pi_1^{\acute{\text{e}}\text{t}}(U, \bar{\eta}) \rightarrow \pi_1^{\acute{\text{e}}\text{t}}(X, \bar{\eta})$$

is surjective as well.

Proposition 3.0.12 (see [?, Tag 0BQM]). Let X be a normal integral scheme function field K . Let η be the generic point of X and let $\bar{\eta}$ be a geometric point (Definition 3.0.1) of X whose image is η . The canonical continuous group homomorphism

$$\text{Gal}(K^{\text{sep}}/K) \cong \pi_1^{\acute{\text{e}}\text{t}}(\eta, \bar{\eta}) \rightarrow \pi_1(X, \bar{\eta})$$

(Definition 3.0.2) (♠ TODO: better describe the canonical map)

is identified with the quotient map

$$\text{Gal}(K^{\text{sep}}/K) \rightarrow \text{Gal}(M/K)$$

(♠ TODO: defined unramifiedness of X in L) where $M \subseteq K^{\text{sep}}$ is the union of the finite subextensions L such that X is unramified in L .

4. SHEAVES ON THE SMALL ÉTALE SITE OF A SCHEME

We define presheaves generally.

Definition 4.0.1 (Presheaf on a category). Let $\mathcal{C}$ and $\mathcal{A}$ be (large) categories (Definition 1.1.1).

1. A *presheaf* $\mathcal{F}$ on $\mathcal{C}$ with values in $\mathcal{A}$ is a functor

$$\mathcal{F} : \mathcal{C}^{\text{op}} \rightarrow \mathcal{A}.$$

In other words, a presheaf $\mathcal{F}$ on $\mathcal{C}$ with values in $\mathcal{A}$ is simply a contravariant functor from $\mathcal{C}$ to $\mathcal{A}$. Explicitly, for every object U in $\mathcal{C}$, one has an object $\mathcal{F}(U)$ in $\mathcal{A}$ (called

the U -valued sections/sections evaluated at U of $\mathcal{F}$, cf. Definition 7.1.1), and for every morphism $f : V \rightarrow U$ in C , one has a morphism (called the *restriction map*)

$$\mathcal{F}(f) : \mathcal{F}(U) \rightarrow \mathcal{F}(V)$$

in $\mathcal{A}$, such that for all composable morphisms $W \xrightarrow{g} V \xrightarrow{f} U$ in C , the following diagram in $\mathcal{A}$ commutes:

$$\begin{array}{ccccc} & & \mathcal{F}(f \circ g) & & \\ & \nearrow & & \searrow & \\ \mathcal{F}(U) & \xrightarrow{\mathcal{F}(f)} & \mathcal{F}(V) & \xrightarrow{\mathcal{F}(g)} & \mathcal{F}(W) \end{array}$$

That is,

$$\mathcal{F}(g) \circ \mathcal{F}(f) = \mathcal{F}(f \circ g),$$

and for every object U in C , $\mathcal{F}(\text{id}_U) = \text{id}_{\mathcal{F}(U)}$.

2. Let $\mathcal{F}, \mathcal{G} : C^{\text{op}} \rightarrow \mathcal{A}$ be two presheaves on C with values in $\mathcal{A}$. A *morphism of presheaves*

$$\varphi : \mathcal{F} \rightarrow \mathcal{G}$$

is a natural transformation of functors: for each object U of C , one has a morphism

$$\varphi_U : \mathcal{F}(U) \rightarrow \mathcal{G}(U)$$

in $\mathcal{A}$, such that for every morphism $f : V \rightarrow U$ in C , the diagram

$$\begin{array}{ccc} \mathcal{F}(U) & \xrightarrow{\mathcal{F}(f)} & \mathcal{F}(V) \\ \varphi_U \downarrow & & \downarrow \varphi_V \\ \mathcal{G}(U) & \xrightarrow{\mathcal{G}(f)} & \mathcal{G}(V) \end{array}$$

commutes, i.e.,

$$\varphi_V \circ \mathcal{F}(f) = \mathcal{G}(f) \circ \varphi_U$$

for all objects and morphisms in C .

3. Given a universe (Definition .1.2) U , a *U -presheaf on $\mathcal{C}$* typically refers to a presheaf of U -sets on C .
4. The *presheaf category/category of $\mathcal{A}$ -valued presheaves on $\mathcal{C}$* is the (large) category whose objects are the presheaves on C with values in $\mathcal{A}$ and whose morphisms are the presheaf morphisms. Common notations for the presheaf category include, but are not limited to: $\mathcal{A}^{C^{\text{op}}}$, $\text{PreShv}(\mathcal{C}, \mathcal{A})$, $[C^{\text{op}}, \mathcal{A}]$. If the value category $\mathcal{A}$ is clear from context, then notations such as $\text{PreShv}(\mathcal{C})$ are also common. Note that the presheaf category $\text{PreShv}(\mathcal{C}, \mathcal{A})$ is equivalent to the category of functors (Definition .1.33) $C^{\text{op}} \rightarrow \mathcal{A}$ and hence notations for the functor categories are applicable as notations for presheaf categories.

We can speak of sheaves on a general site.

We will be interested in sheaves of sets/abelian groups/ R -modules on the small étale site $X_{\text{ét}}$ (Definition 4.1.3) of any scheme as $X_{\text{ét}}$ is essentially small (Definition .1.26).

Definition 4.0.2 (Sheaf on a site). Let $(\mathcal{C}, J)$ be a site (Definition A.0.4). Let $\mathcal{A}$ be a (large) category (Definition .1.1).

1. A presheaf (Definition 4.0.1) $\mathcal{F} : \mathcal{C}^{\text{op}} \rightarrow \mathcal{A}$ is called a *sheaf on the site $(\mathcal{C}, J)$ valued in $\mathcal{A}$* if, for every object U of $\mathcal{C}$ and every covering sieve (Definition A.0.4) $S \in J(U)$, the limit (Definition .1.13)

$$\varprojlim_{(V \rightarrow U) \in (\mathcal{D}_S)^{\text{op}}} \mathcal{F}|_{\mathcal{D}_S}(V),$$

exists and the canonical natural morphism

$$\mathcal{F}(U) \rightarrow \varprojlim_{(V \rightarrow U) \in (\mathcal{D}_S)^{\text{op}}} \mathcal{F}|_{\mathcal{D}_S}(V)$$

is an isomorphism. Here, $\mathcal{D}_S \hookrightarrow \mathcal{C}/U$ (Definition A.0.6) is the full downward-closed subcategory such that $\text{Ob}(\mathcal{D}_S) = \{(f : V \rightarrow U) : f \in S(V)\}$,

In particular, when we are working with a Grothendieck pretopology (Definition 4.2.2) K on a category $\mathcal{C}$, we may speak of sheaves on the site whose Grothendieck topology is the one generated by (Definition 4.2.1) K .

2. Given sheaves $\mathcal{F}, \mathcal{G} : \mathcal{C}^{\text{op}} \rightarrow \mathcal{A}$ on the site $(\mathcal{C}, J)$, a *morphism between the sheaves* is a morphism (Definition 4.0.1) between $\mathcal{F}$ and $\mathcal{G}$ as presheaves.
3. Let U be a universe (Definition .1.2). A *U -sheaf* typically refers to a U -presheaf that is a sheaf for a U -site. In other words, a U -sheaf is a sheaf on a site whose underlying category is U -locally small (Definition .1.4) and which has a U -small topologically generating family such that the sheaf is valued in U -sets.
4. The *sheaf category/category of $\mathcal{A}$ -valued sheaves on $\mathcal{C}$* is the (large) category defined as the full subcategory of $\text{PreShv}(\mathcal{C}, \mathcal{A})$ whose objects are the sheaves on $\mathcal{C}$ with values in $\mathcal{A}$. Common notations for the sheaf category include $\text{Shv}(\mathcal{C}, \mathcal{A})$, $\text{Shv}(\mathcal{C}, J, \mathcal{A})$, $\text{Sh}(\mathcal{C}, \mathcal{A})$, $\text{Sh}(\mathcal{C}, J, \mathcal{A})$. If the value category $\mathcal{A}$ is clear from context, then notations such as $\text{Shv}(\mathcal{C})$, $\text{Shv}(\mathcal{C}, J)$, $\text{Sh}(\mathcal{C})$, $\text{Sh}(\mathcal{C}, J)$ are also common.

4.1. The small and big Étale sites of a scheme.

Definition 4.1.1 (e.g. see [?, II §1]). Let S be a scheme (Definition .3.1). Let E be some class of morphisms in Sch/S (Definition .3.3) satisfying the following:

- All isomorphisms are in E .
 - E is closed under compositions.
 - Any base change of a morphism in E is in E .
1. Let C/S be some full subcategory of Sch/S that is closed under fiber products such that for any $X \rightarrow S$ in C/S and any E -morphism $U \rightarrow X$, the composite $U \rightarrow S$ is in C/S . An *E -covering of an object X of C/S* is a family $(U_i \xrightarrow{g_i} X)_{i \in I}$ of E -morphisms such that $Y = \bigcup_i g_i(X_i)$. The class of all such coverings of all such objects is the *E -topology on C/S* ; it is a Grothendieck pretopology (Definition 4.2.2). The category C/S equipped with the E -topology (more precisely, the Grothendieck topology (Definition A.0.4) generated by (Definition 4.2.1) the E -topology) is the *E -site over S* .

2. Assuming that all morphisms in E are locally of finite-type, the **big E -site on S** is the site (Definition A.0.4) whose underlying category is the category of locally of finite type schemes (Definition C.0.19) over S and whose Grothendieck topology (Definition A.0.4) is the one generated by (Definition 4.2.1) the pretopology (Definition 4.2.2) whose coverings are the E -coverings.
3. The **small E -site on S** is the site whose underlying category is the full subcategory of Sch/S of S -schemes whose structure morphisms are E -morphisms and whose coverings are E -coverings.

Definition 4.1.2 (Big étale site of a scheme). Let S be a fixed scheme. The **big étale site on S** , denoted by $(\text{Sch}/S)_{\text{ét}}$, is defined as the following site (Definition A.0.4):

- The underlying category is the category Sch/S (Notation 1.1.1) of all schemes over S (Notation 1.1.1). That is, objects are morphisms of schemes $X \rightarrow S$, and morphisms are S -morphisms between such X .
- The Grothendieck topology (Definition A.0.4) is the one generated by (Definition 4.2.1) the pretopology (Definition 4.2.2) whose covering families are families $\{f_i : X_i \rightarrow X\}_{i \in I}$ in Sch/S such that each f_i is an étale morphism (Definition 1.1.6) and the family is jointly surjective on the underlying topological spaces. Such a cover is called an **étale cover of X** .

Equivalently, the big étale site on S is the big site for étale morphisms on S (Definition 4.1.1).

Definition 4.1.3 (Small étale site of a scheme). Let X be a fixed scheme (Definition 3.1). The **small étale site on X** , commonly denoted by notations including $X_{\text{ét}}$, $X_{\text{étale}}$, or Et/X , is defined as the following site (Definition A.0.4):

- The underlying category is the full subcategory of the big étale site $(\text{Sch}/X)_{\text{ét}}$ (Definition 4.1.2) (Definition 4.1.2) whose objects are schemes U equipped with an étale morphism $U \rightarrow X$.
- The Grothendieck topology is the one generated by (Definition 4.2.1) the pretopology (Definition 4.2.2) whose covering families are families $\{g_j : U_j \rightarrow U\}_{j \in J}$ of morphisms such that each g_j is an étale (Definition 1.1.6) and the family is jointly surjective on the underlying topological spaces.

Equivalently, the small étale site on X is the small site for étale morphisms on X (Definition 4.1.1). (♠ TODO: state this as a fact) $X_{\text{ét}}$ is an essentially small category (Definition 1.26).

4.2. Other common sites of schemes.

Definition 4.2.1. Let $\mathcal{C}$ be a category equipped with a Grothendieck pretopology (Definition 4.2.2) K . The **Grothendieck topology generated by K** , denoted J_K , is the smallest Grothendieck topology (Definition A.0.4) on $\mathcal{C}$ such that every family in $K(U)$ is a covering family for J_K .

Explicitly, a sieve (Definition A.0.1) S on an object U belongs to $J_K(U)$ if and only if there exists a covering family (Definition 4.2.2) $\{U_i \rightarrow U\}_{i \in I} \in K(U)$ such that for every $i \in I$, the

morphism $U_i \rightarrow U$ belongs to S , i.e. the sieve generated by (Definition A.0.3) $\{U_i \rightarrow U\}_{i \in I}$ is a sub-sieve of S .

Definition 4.2.2. Let $\mathcal{C}$ be a category (Definition .1.1). A *basis for a Grothendieck topology* (also called a *Grothendieck pretopology* or simply a *pretopology*) on $\mathcal{C}$ is a collection of families $K(U)$ of morphisms for each object $U \in \mathcal{C}$, called *coverings* or *covering families*, satisfying the following axioms:

1. **(Identity)** For every isomorphism $U' \rightarrow U$, the singleton family $\{U' \rightarrow U\}$ is in $K(U)$.
2. **(Base Change)** If $\{U_i \rightarrow U\}_{i \in I}$ is a covering family in $K(U)$ and $V \rightarrow U$ is any morphism in $\mathcal{C}$, then the fiber products (Definition C.0.15) $U_i \times_U V$ exist, and the family of projections $\{U_i \times_U V \rightarrow V\}_{i \in I}$ is in $K(V)$.
3. **(Composition)** If $\{U_i \rightarrow U\}_{i \in I}$ is in $K(U)$ and for each $i \in I$, $\{V_{ij} \rightarrow U_i\}_{j \in J_i}$ is in $K(U_i)$, then the composite family $\{V_{ij} \rightarrow U_i \rightarrow U\}_{i \in I, j \in J_i}$ is in $K(U)$.

Given a family $\{U_i \rightarrow U\}_{i \in I}$ in $K(U)$, the sieve (Definition A.0.1) that it generates (Definition A.0.3) is called a *covering sieve of U (generated by $\{U_i \rightarrow U\}_{i \in I}$)*.

Definition 4.2.3 (Big Zariski site). Let S be a scheme. The *big Zariski site of S* , denoted by $(\text{Sch}/S)_{\text{Zar}}$, is the site (Definition A.0.4) whose underlying category is the category of S -schemes and whose Grothendieck topology is the one generated by (Definition 4.2.1) the pretopology (Definition 4.2.2) whose coverings are families of morphisms

$$\{f_i : U_i \rightarrow U\}_{i \in I}$$

such that each f_i is an open immersion (Definition C.0.14) and the images $\{f_i(U_i)\}_{i \in I}$ form an open cover of U . Such a covering is called a *Zariski covering of U* .

Equivalently, the big Zariski site of S is the big site on the category of schemes over S (Definition 4.1.1) for the class of open immersions.

Definition 4.2.4 (Small Zariski site). Let X be a scheme. The *small Zariski site of X* , denoted by X_{Zar} , is the site (Definition A.0.4)

whose underlying category consists of open subschemes $U \subseteq X$, with inclusions as morphisms, and whose Grothendieck topology is the one generated by (Definition 4.2.1) the pretopology (Definition 4.2.2) whose coverings are families of open immersions (Definition C.0.14)

$$\{U_i \rightarrow U\}_{i \in I}$$

such that the U_i form an open cover of U in the usual topological sense.

Equivalently, the small Zariski site of X is the small site on the category of schemes over X (Definition 4.1.1) for the class of open immersions.

Also equivalently, the small Zariski site of X is the Site of opens of X (Definition .1.9) as a topological space.

Definition 4.2.5 (Faithfully flat morphism of schemes). Let $f : X \rightarrow Y$ be a morphism of schemes (Definition .3.2).

The morphism f is **faithfully flat** if it is flat (Definition 1.1.4) and surjective on the underlying topological spaces.

More precisely:

- f is **flat**, meaning for every $x \in X$ with $y = f(x)$, the local ring homomorphism (Definition C.0.5)

$$\mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{X,x}$$

makes $\mathcal{O}_{X,x}$ a flat (Definition 1.1.3) $\mathcal{O}_{Y,y}$ -module.

- f is **surjective** at the level of topological spaces, i.e., the continuous map $f : |X| \rightarrow |Y|$ is surjective.

Equivalently, f is faithfully flat if the functor $f^* : \mathrm{QCoh}(Y) \rightarrow \mathrm{QCoh}(X)$ on quasi-coherent sheaves is both exact (Definition 1.1.36) and faithful.

Definition 4.2.6 (big fppf site). Let S be a scheme. The **big fppf site of S** , denoted by $(\mathrm{Sch}/S)_{\mathrm{fppf}}$, is the following site (Definition A.0.4):

- The underlying category is the category Sch/S (Notation 1.1.1) of all schemes over S .
- The Grothendieck topology is the one generated by (Definition 4.2.1) the pretopology (Definition 4.2.2) whose coverings are families of morphisms $\{f_i : X_i \rightarrow X\}_{i \in I}$ in Sch/S is a covering family in $(\mathrm{Sch}/S)_{\mathrm{fppf}}$ if each f_i is a flat (Definition 1.1.4) morphism of locally of finite presentation (Definition 3.6) such that $\bigcup_i f_i(X_i) = f(X)$. Such a cover is called an **fppf cover of X** .

Equivalently, the big fppf site of S is the big site on S (Definition 4.1.1) for the class of flat (Definition 1.1.4) and locally of finite presentation (Definition 1.1.2) morphisms.

fppf stands for *fidèlement plate de présentation finie*, which translates to faithfully flat and of finite presentation; note however, that the morphisms in an fppf cover are flat and locally of finite presentation, not faithfully flat (Definition 4.2.5) and of finite presentation (Definition 3.6).

Definition 4.2.7 (small fppf site). Let X be a fixed scheme. The **small fppf site on X** , commonly denoted by notations including X_{fppf} , or fppf/X , is defined as the following site (Definition A.0.4):

- The underlying category is the full subcategory of the big étale site $(\mathrm{Sch}/X)_{\mathrm{fppf}}$ (Definition 4.2.6) whose objects are schemes U equipped with a (♠ TODO:)
- The Grothendieck topology is the one generated by (Definition 4.2.1) the pretopology (Definition 4.2.2) whose coverings are families of morphisms $\{g_j : U_j \rightarrow U\}_{j \in J}$ in this category is a covering family in $X_{\mathrm{ét}}$ if each g_j is étale (Definition 1.1.6) and the family is jointly surjective on the underlying topological spaces.

Equivalently, the small fppf site on X is the small site for morphisms that are flat and locally of finite presentation on X (Definition 4.1.1). (♠ TODO: state this as a fact) X_{fppf} is an essentially small category (Definition 1.1.26).

Definition 4.2.8 (big fpqc site). (♠ TODO:) Let S be a scheme. The *big fpqc site of S* , denoted by $(\text{Sch}/S)_{\text{fpqc}}$, is following site (Definition A.0.4):

1. The underlying category is the category Sch/S (Notation 1.1.1) of all schemes over S .
2. A family of morphisms $\{f_i : X_i \rightarrow X\}_{i \in I}$ in Sch/S is a covering family in $(\text{Sch}/S)_{\text{fpqc}}$ if
 - each f_i is a flat (Definition 1.1.4) morphism
 - for each affine open $U \subseteq X$, there exists a finite subset $J \subseteq I$ and affine open subsets $U_j \subseteq X_j$ for each $j \in J$ such that $U = \bigcup_{j \in J} f_j(U_j)$.

Such a cover is called an *fpqc cover of X* .

Definition 4.2.9 (Small fpqc site). (♠ TODO:) Let X be a scheme. The *small fpqc site of X* , denoted by X_{fpqc} , is the site whose underlying category consists of schemes U equipped with a morphism

$$U \longrightarrow X,$$

such that the morphism is *flat* and *quasi-compact* and whose Grothendieck topology is the one generated by (Definition 4.2.1) the pretopology (Definition 4.2.2) whose coverings are families of morphisms

$$\{f_i : U_i \rightarrow U\}_{i \in I}$$

over X such that

1. Each f_i is *faithfully flat*.
2. Each f_i is *quasi-compact*.
3. The images of the family jointly cover U ; that is,

$$\bigcup_{i \in I} f_i(U_i) = U.$$

4. For every affine open $V \subseteq U$, there exists a finite subset $J \subseteq I$ and affine opens $V_j \subseteq U_j$, for $j \in J$, such that

$$V = \bigcup_{j \in J} f_j(V_j).$$

This topology is called the *fpqc topology* (faithfully flat and quasi-compact).

Definition 4.2.10. [See [?, Definition 2.1], [?, Section 3 Definition 1.3]] Let S be a scheme (Definition .3.1). An *elementary distinguished square in the category Sm/S* of smooth schemes over S is a square of the form

$$(B) \quad \begin{array}{ccc} p^{-1}(U) & \longrightarrow & V \\ \downarrow p & & \downarrow p \\ U & \xrightarrow{j} & X \end{array}$$

(♠ TODO: open embedding, reduced subscheme, support) such that p is an étale morphism (Definition 1.1.6), j is an open embedding, and $p^{-1}(X - U) \rightarrow X - U$ is an isomorphism (where $X - U$ is the maximal reduced subscheme with support in the closed subset $X - U$).

Proposition 4.2.11 ([?, Section 3 Proposition 1.1]). Let S be a Noetherian (Definition C.0.16) scheme of finite dimension (Definition C.0.20). Let X be a scheme of finite type (Definition C.0.19) over S and let $\{U_i \rightarrow X\}$ be a finite family of étale morphisms (Definition 1.1.6) in Sch/S (Definition .3.3). The following conditions are equivalent: (♠ TODO: residue field)

1. For any point x of X there is an i and a point u of U_i over x such that the corresponding morphism of residue fields is an isomorphism which maps to x with the same residue field.
2. For any point $x \in X$, the morphism

$$\coprod_i (U_i \times_X \text{Spec } \mathcal{O}_{X,x}^h) \rightarrow \text{Spec } \mathcal{O}_{X,x}^h$$

of S -schemes admits a section.

Moreover, the collection of families of étale morphisms $\{U_i \rightarrow X\}$ in Sm/S satisfying the equivalent conditions above forms a pretopology on Sm/S .

Definition 4.2.12. [See [?, Section 3 Definition 1.2]] Let S be a Noetherian (Definition C.0.16) scheme of finite dimension (Definition C.0.20). The Grothendieck topology (Definition A.0.4) generated by the pretopology (Definition 4.2.2) of Proposition 4.2.11 is called the *Nisnevich topology on Sm/S* . The site whose underlying category is Sm/S and whose Grothendieck topology is the Nisnevich topology is called the *(big) Nisnevich site of S* and is denoted by notations such as $(\text{Sm}/S)_{\text{Nis}}$, $(\mathbf{Sm}/S)_{\text{Nis}}$, etc. A covering family in the Nisnevich site of S is called a *Nisnevich covering*.

(♠ TODO: establish that a nisnevich covering is a family of etale morphisms such that there is an isomorphism of residue fields at every point)

Definition 4.2.13 (Small Nisnevich site). Let X be a scheme. The *small Nisnevich site of X* , denoted by

$$X_{\text{Nis}},$$

is the site whose underlying category consists of schemes U equipped with an étale morphism (Definition 1.1.6)

$$U \rightarrow X,$$

and whose Grothendieck topology is the one generated by (Definition 4.2.1) the Grothendieck pretopology (Definition 4.2.2) whose coverings are given by Nisnevich coverings (Definition 4.2.12).

Definition 4.2.14 (crystalline site). (♠ TODO:) Let S be a scheme and p a prime number. Let $\mathcal{C} = \text{CRIS}(S/\mathbf{Z}_p)$ denote the category whose objects are triples (U, T, δ) where

- U is an open subscheme of S ,
- T is a scheme equipped with a divided power structure δ on an ideal $\mathcal{I} \subseteq \mathcal{O}_T$,
- together with a closed immersion $U \hookrightarrow T$ compatible with δ (called a PD-thickening).

The *crystalline site* $(S/\mathbf{Z}_p)_{\text{cris}}$ is this category equipped with the Grothendieck topology (Definition A.0.4) generated by (Definition 4.2.1) families of morphisms $\{(U_i, T_i, \delta_i) \rightarrow (U, T, \delta)\}$ such that the underlying morphisms of the $T_i \rightarrow T$ form a Zariski open cover of T .

Theorem 4.2.15 (Hierarchy of Grothendieck topologies on Sch/S). Let S be a scheme and consider the following Grothendieck topologies (Definition A.0.4) on the category Sch/S of schemes over S (Definition .3.3):

$$\text{Zar} \quad (\text{Zariski}), \quad \text{Nis} \quad (\text{Nisnevich}), \quad \text{Ét} \quad (\text{étale}), \quad \text{fppf}, \quad \text{fpqc}.$$

(Definition 4.2.3, Definition 4.2.12, Definition 4.1.2, Definition 4.2.6, Definition 4.2.8)

Then these topologies satisfy the chain of refinements (fineness) relations

$$\text{Zar} \prec \text{Nis} \prec \text{Ét} \prec \text{fppf} \prec \text{fpqc},$$

meaning that each topology is strictly finer (Definition .1.3) than the previous one, i.e., every covering in a coarser topology is a covering in any finer topology.

More explicitly, for every object $X \in \text{Sch}/S$,

$$\text{Cov}_{\text{Zar}}(X) \subsetneq \text{Cov}_{\text{Nis}}(X) \subsetneq \text{Cov}_{\text{Ét}}(X) \subsetneq \text{Cov}_{\text{fppf}}(X) \subsetneq \text{Cov}_{\text{fpqc}}(X).$$

Theorem 4.2.16 (Essential smallness of small sites). Let X be a scheme. Consider the following sites associated to X :

- small Zariski site X_{Zar} (Definition 4.2.4),
- small Nisnevich site X_{Nis} (Definition 4.2.13),
- small étale site $X_{\text{Ét}}$ (Definition 4.1.3),
- small fppf site X_{fppf} (Definition 4.2.7).
- small fpqc site X_{fpqc} (Definition 4.2.9).

These sites are essentially small (Definition .1.26).

4.3. Sheafification of a presheaf on a site.

Definition 4.3.1. Let $\mathcal{C}$ be a site (Definition A.0.4) and let $\mathcal{A}$ be a (large) category (Definition .1.1).

Assuming that the presheaf (Definition 4.0.1) category $\text{PreShv}(\mathcal{C}, \mathcal{A})$ (and hence the sheaf (Definition 4.0.2) category $\text{Shv}(\mathcal{C}, \mathcal{A})$) is locally small (Definition .1.4) (or U -locally small if a Grothendieck universe (Definition .1.2) U is available), a *sheafification functor* refers to a functor

$$a : \text{PreShv}(\mathcal{C}, \mathcal{A}) \rightarrow \text{Shv}(\mathcal{C}, \mathcal{A})$$

that is left adjoint (Definition .1.8) to the inclusion functor

$$i : \text{Shv}(\mathcal{C}, \mathcal{A}) \hookrightarrow \text{PreShv}(\mathcal{C}, \mathcal{A}).$$

If such a sheafification functor exists, then it is unique up to unique natural isomorphism. Given a presheaf P , the sheafification $a(P)$ is also sometimes called the *sheaf associated to P* . See Theorem 4.3.2 for common conditions under which sheafification exists.

Theorem 4.3.2. cf. [?, Exposé II, Théorème 3.4]

1. Let U be a universe. Let $\mathcal{C}$ be a U -site (Definition A.0.4). A sheafification functor

$$a : \text{PreShv}(\mathcal{C}, U\text{-}\mathbf{Sets}) \rightarrow \text{Shv}(\mathcal{C}, U\text{-}\mathbf{Sets}).$$

exists.

2. Let $\mathcal{C}$ be a site whose underlying category is locally small (Definition .1.4) and which has a topologically generating family (Definition A.0.4) that is a set (rather than a proper class). A sheafification functor

$$a : \text{PreShv}(\mathcal{C}, \mathbf{Sets}) \rightarrow \text{Shv}(\mathcal{C}, \mathbf{Sets})$$

exists.

(♠ TODO: It may be the case that I need to assume $\mathcal{A}$ to be locally small; if so, recheck all the statements that use this.)

3. (see e.g. [?, 3], [?, Theorem 17.4.7]) Let $(\mathcal{C}, J)$ be a site (Definition A.0.4) on an essentially small category (Definition .1.26) $\mathcal{C}$. Suppose that the category $\mathcal{A}$ is complete, cocomplete, that small filtered colimits (Definition .1.14) in $\mathcal{A}$ are exact, and that $\mathcal{A}$ satisfies the IPC-property. A sheafification functor (Definition 4.3.1)

$$a : \text{PreShv}(\mathcal{C}, \mathcal{A}) \rightarrow \text{Shv}(\mathcal{C}, \mathcal{A})$$

exists.

(♠ TODO: state as a fact that the category of groups and k -algebras are complete, cocomplete, with small filtered colimits that are exact) This is true for instance of $\mathcal{A} = \mathbf{Set}, \mathbf{Grp}, k\text{-}\mathbf{Alg}$ for a field k , or $\mathbf{Mod}_R$ for a (not necessarily commutative unital) ring R (Definition C.0.8).

Remark 4.3.3. If the presheaf is valued in nice “algebraic category”, e.g. groups, abelian groups, rings, modules over a ring, etc., then the sheafification is again valued in that category. (♠ TODO: Make this more precise.)

Corollary 4.3.4. Let X be a scheme. For the categories $\mathcal{A} = \mathbf{Set}, \mathbf{Grp}, k\text{-}\mathbf{Alg}$ for a field k , or $\mathbf{Mod}_R$ for a (not necessarily commutative unital) ring R (Definition C.0.8), and for the sites $\mathcal{C} = X_{\text{Zar}}, X_{\text{Nis}}, X_{\text{Ét}}, X_{\text{fppf}}, X_{\text{fpqc}}$, there exists a sheafification functor (Theorem 4.3.2)

$$a : \text{PreShv}(\mathcal{C}, \mathcal{A}) \rightarrow \text{Shv}(\mathcal{C}, \mathcal{A}).$$

Proof. This follows from Theorem 4.3.2 and Theorem 4.2.16. □

4.4. Examples of sheaves on the étale site of a scheme. Constant and locally constant sheaves are discussed not just for $X_{\text{ét}}$, but also for more general sites.

Definition 4.4.1 (Constant sheaf on a site). Let $\mathcal{C}$ be a category (Definition .1.1), let $\mathcal{A}$ be a category, and let A be an object of $\mathcal{A}$.

1. The *constant presheaf on $\mathcal{C}$ with value A* is the presheaf (Definition 4.0.1) P defined by

$$P(U) = A$$

for every object U of $\mathcal{C}$ such that every morphism $f : V \rightarrow U$ in $\mathcal{C}$ induces the identity map $A = P(U) \rightarrow P(V) = A$.

2. Let $\mathcal{C}$ be a site (Definition A.0.4) and assume that a sheafification functor (Definition 4.3.1)

$$a : \text{Shv}(\mathcal{C}, \mathcal{A}) \rightarrow \text{PreShv}(\mathcal{C}, \mathcal{A})$$

exists (e.g. see Theorem 4.3.2) . The *constant sheaf on $\mathcal{C}$ with value A* , or the *constant sheaf on $\mathcal{C}$ associated to A* commonly denoted $\underline{A}$ or sometimes just A by abuse of notation, is the sheaf associated to (Theorem 4.3.2) the constant presheaf P with value A above.

3. Let $\mathcal{C}$ be a site. Let $\mathcal{O}$ be a sheaf of (not-necessarily commutative) rings on $\mathcal{C}$. Assume that the global sections ring $\Gamma(\mathcal{O})$ (Definition 7.1.1) exists. A *constant $\mathcal{O}$ -module* is an $\mathcal{O}$ -module (Definition 7.1.3) $\mathcal{F}$ which is isomorphic as a sheaf to the constant sheaf on $\mathcal{C}$ with value M where M is a module of the ring $\Gamma(\mathcal{O})$. Note that sheafification functors exist for presheaves/sheaves valued in $\mathbf{Ab}$ (Theorem 4.3.2).

In case that $\mathcal{O}$ is the constant sheaf associated to A for some (not-necessarily commutative) ring A , a constant $\mathcal{O}$ -module is simply called a *constant A -module*.

Definition 4.4.2 (Locally constant sheaf on a site). Let $(\mathcal{C}, J)$ be a site (Definition A.0.4).

1. Let $\mathcal{A}$ be a (large) category, and let A be an object of $\mathcal{A}$. (♠ TODO: If such a sheafification functor exist, does a sheafification functor exist when restricted to an object U ?) Assume that a sheafification functor (Definition 4.3.1)

$$a : \mathbf{PreShv}(\mathcal{C}, J, \mathcal{A}) \rightarrow \mathbf{Shv}(\mathcal{C}, J, \mathcal{A})$$

(Definition 4.0.2) (Definition 4.0.1) exists (e.g. see Theorem 4.3.2) . Let U be an object of $\mathcal{C}$.

A sheaf $\mathcal{F}$ on $\mathcal{C}$ with values in $\mathcal{A}$ is said to be a *locally constant sheaf on U with value A* if there exists a covering sieve (Definition A.0.4) $\{U_i \rightarrow U\}$ of every object U in $\mathcal{C}$ such that for each i , the restriction $\mathcal{F}|_{U_i}$ is isomorphic to the constant sheaf (Definition C.0.10) $\underline{A}$ on the slice site $\mathcal{C}_{/U_i}$ (Definition A.0.8).

If $\mathcal{C}$ has a final object, then we may say that $\mathcal{F}$ is a *locally constant sheaf with value A* if it is a locally constants sheaf on the final object with value A .

2. Let $\mathcal{O}$ be a sheaf of (not-necessarily commutative) rings on $\mathcal{C}$. A *locally constant $\mathcal{O}$ -module* is a $\mathcal{O}$ -module (Definition 7.1.3) $\mathcal{F}$ such that there exists a covering sieve (Definition A.0.4) $\{U_i \rightarrow U\}$ for every object U in $\mathcal{C}$ such that for each i , the restriction $\mathcal{F}|_{U_i}$ is isomorphic, as an $\mathcal{O}|_{U_i}$ -module, to a constant $\mathcal{O}|_{U_i}$ -module (Definition C.0.10)¹.

We additionally say that $\mathcal{F}$ is

- (a) *locally free of rank r over $\mathcal{O}$* if there exists a covering $\{U_i \rightarrow U\}$ such that $\mathcal{F}|_{U_i}$ is isomorphic as a $\mathcal{O}|_{U_i}$ -module to $(\mathcal{O}|_{U_i})^{\oplus r}$ for each i .
- (b) *free of rank r over $\mathcal{O}$* if $\mathcal{F} \cong \mathcal{O}^{\oplus r}$ as $\mathcal{O}$ -modules.
- (c) *of finite type* if there exists a covering $\{U_i \rightarrow U\}$ such that $\mathcal{F}|_{U_i}$ generated by finitely many sections over U_i as an $\mathcal{O}|_{U_i}$ -module. In other words, there is an epimorphism

$$(\mathcal{O}|_{U_i})^{\oplus n_i} \rightarrow \mathcal{F}|_{U_i}$$

of $\mathcal{O}|_{U_i}$ -modules for each i .

In case that $\mathcal{O}$ is the constant sheaf on $\mathcal{C}$ associated to A (Definition C.0.10) for some (not-necessarily commutative) ring A , a locally constant $\mathcal{O}$ -module is simply called a *locally constant A -module*.

¹The global sections rings $\Gamma(\mathcal{O}|_{U_i})$ (Definition 7.1.1) of the sheaves $\mathcal{O}|_{U_i}$ on the slice sites $\mathcal{C}_{/U_i}$ (Definition A.0.8) exist because each $\mathcal{C}_{/U_i}$ has a final object (Definition .1.10)(Lemma A.0.7), so we may speak of constant $\mathcal{O}|_{U_i}$ -modules.

Definition 4.4.3. Let C be a category enriched in a monoidal category $\mathcal{V}$. Given an object X of C , the *functor of points* h_X is the functor/presheaf (Definition 4.0.1) $C^{\text{op}} \rightarrow \mathcal{V}$ given by $T \mapsto \text{Hom}_C(T, X)$. A functor $C^{\text{op}} \rightarrow \mathcal{V}$ (or equivalently, a presheaf on C valued in $\mathcal{V}$) is said to be *representable* if it is naturally isomorphic to some functor h_X of points for an object X of C .

Dually, a functor $C \rightarrow \mathcal{V}$ is called *co-representable* if it is naturally isomorphic to a functor $T \mapsto \text{Hom}_C(X, T)$ for an object X in C .

For instance, we may speak of these notions when $\mathcal{V}$ is the monoidal category **Sets**, i.e. C is a locally small category (Definition 1.4).

Proposition 4.4.4 (Representable functors are étale sheaves). (♠ TODO: This is probably more generally true for fppf, fpqc sites, etc.) Let X be a scheme (Definition 3.1) and consider the étale site $X_{\text{ét}}$ (Definition 4.1.3).

If a functor

$$F : (X_{\text{ét}})^{\text{op}} \rightarrow \mathbf{Sets}$$

is representable (Definition 4.4.3) by an X -scheme Y , i.e.,

$$F(U) \cong \text{Hom}_X(U, Y)$$

for all étale $U \rightarrow X$, then the presheaf F is an étale (Definition 4.1.3) sheaf (Definition 4.0.2).

Definition 4.4.5. (♠ TODO: regular function on a scheme) For a scheme (Definition 3.1) X , the *additive sheaf* $\mathbb{G}_a$ on $X_{\text{ét}}$ is defined by

$$\mathbb{G}_a(U) := \mathcal{O}_U(U),$$

the ring of regular functions on U for each étale $U \rightarrow X$. It may also be called the *additive group scheme*, on account of the fact that the sheaf $\mathbb{G}_a$ is representable by (Definition 4.4.3) the scheme $\mathbb{G}_a = \mathbb{A}_X^1$ (Definition 3.7). (♠ TODO: affine line over a scheme) In particular, it is indeed a sheaf (Proposition 4.4.4).

Definition 4.4.6. For a scheme X , the *multiplicative sheaf* $\mathbb{G}_m$ on $X_{\text{ét}}$ is defined by

$$\mathbb{G}_m(U) := \mathcal{O}_U(U)^\times,$$

the group of invertible regular functions on U for each étale $U \rightarrow X$. It may also be called the *multiplicative group scheme*, on account of the fact that the sheaf $\mathbb{G}_m$ is representable by (Definition 4.4.3) the scheme $\mathbb{G}_m = \mathbb{A}_X^1 \setminus \{0\}$. (♠ TODO: affine line over a scheme) In particular, it is indeed a sheaf (Proposition 4.4.4).

Definition 4.4.7 (The sheaf of n -th roots of unity μ_n). (♠ TODO: invertible integer on a scheme) (♠ TODO: kernel for sheaves) Let n be a positive integer invertible on the scheme (Definition 3.1) X . The sheaf (Definition 4.0.2) μ_n on $X_{\text{ét}}$ (Definition 4.1.3) is defined as the kernel of the multiplication-by- n map on $\mathbb{G}_m$:

$$\mu_n := \ker \left(\mathbb{G}_m \xrightarrow{(\cdot)^n} \mathbb{G}_m \right).$$

In other words,

$$\mu_n(U) = \{f \in \mathcal{O}_U(U)^\times \mid f^n = 1\}$$

for each étale (Definition 1.1.6) $U \rightarrow X$. The sheaf μ_n is often called the *n -Kummer sheaf* on $X_{\text{ét}}$,

We can use the sheaves μ_{ℓ^n} for a prime ℓ to construct ℓ -adic sheaves $\mathbb{Z}_{\ell}(d)$ (Definition 8.4.11) called *Tate twists*; we note that ℓ -adic sheaves (Definition 8.4.9) are not actual sheaves on $X_{\text{ét}}$, but rather a certain type of inverse system of sheaves on $X_{\text{ét}}$.

4.5. Stalk of a (pre)sheaf on the small étale site of a scheme. As with Zariski (pre)sheaves (or more generally locally ringed spaces), one can talk about stalks at points — for the étale site the stalks are not just as closed points, but rather at *geometric* points.

Definition 4.5.1 (Stalk of an étale sheaf at a geometric point). Let X be a scheme (Definition .3.1), $\bar{x} : \text{Spec}(\Omega) \rightarrow X$ a (separable) geometric point (Definition 3.0.1), and let $\mathcal{F}$ be a presheaf (Definition 4.0.1) of sets (resp. abelian groups, etc.) on the étale site $X_{\text{ét}}$ (Definition 4.1.3).

The *(geometric) stalk of $\mathcal{F}$ at $\bar{x}$* is defined as the colimit (Definition .1.13)

$$\mathcal{F}_{\bar{x}} := \varinjlim_{(U,u)} \mathcal{F}(U),$$

where the limit runs over the filtering category (Definition .1.12) of pairs (U, u) consisting of an étale morphism (Definition 1.1.6) $U \rightarrow X$ together with a lift $u : \text{Spec}(\Omega) \rightarrow U$ of $\bar{x}$, i.e., such that the composed morphism $u : \text{Spec}(\Omega) \rightarrow U \rightarrow X$ equals $\bar{x}$.

This construction makes $\mathcal{F}_{\bar{x}}$ a set (resp. an abelian group, etc.), intuitively representing the "germ" of sections of $\mathcal{F}$ near the geometric point $\bar{x}$.

In fact, letting x be a point of X with residue field $\kappa(x)$ and letting $\bar{x} : \text{Spec}(\kappa(x)^s) \rightarrow X$ be an associated geometric point, automorphisms on $\text{Spec}(\kappa(x)^s)$ over $\text{Spec}(\kappa(x))$ correspond to field automorphisms $\kappa(x)^s/\kappa(x)$, and in turn the absolute Galois group $\text{Gal}(\kappa(x)^s/\kappa(x))$ naturally acts on $\mathcal{F}_{\bar{x}}$. In fact, the action is continuous for the discrete topology on $\mathcal{F}_{\bar{x}}$. Moreover, the stalk construction is functorial — for every choice of such x and $\bar{x}$, we have a functor

$$\mathcal{F} \mapsto \mathcal{F}_{\bar{x}} : \text{Sh}(X_{\text{ét}}, \mathbf{Sets}) \rightarrow \text{Gal}(\kappa(x)^s/\kappa(x)) - \mathbf{Sets}$$

(♠ TODO: category of sets with continuous group action)

If $\mathcal{F}$ is a sheaf of abelian groups, then the stalk functor is in fact a functor to the category of discrete abelian groups with continuous $\text{Gal}(\kappa(x)^s/\kappa(x))$ -action.

Theorem 4.5.9 describes stalks of étale sheaves in terms of strict henselizations (Definition 4.5.7). We define the notions of Henselian and strictly Henselian local rings.

Definition 4.5.2 (Henselian ring). Let $(R, \mathfrak{m})$ be a local ring (Definition C.0.21) with maximal ideal $\mathfrak{m}$ (Definition C.0.4). The ring R is called *henselian* if for every monic polynomial $f(x) \in R[x]$ and factorization

$$\bar{f}(x) = \bar{g}(x) \cdot \bar{h}(x)$$

in $(R/\mathfrak{m})[x]$ into coprime monic polynomials $\bar{g}, \bar{h}$, there exist monic polynomials $g(x), h(x) \in R[x]$ lifting $\bar{g}, \bar{h}$ respectively such that

$$f(x) = g(x) \cdot h(x).$$

Theorem 4.5.3. All complete local rings are henselian (Definition 4.5.2).

Definition 4.5.4 (Strictly henselian ring). A *strictly henselian (local) ring* or a *strictly local ring* is a henselian local ring (Definition 4.5.2) $(R, \mathfrak{m})$ whose residue field $k = R/\mathfrak{m}$ is separably closed.

Theorem 4.5.5. All complete local rings with separably closed residue fields are strictly henselian (Definition 4.5.4).

Definition 4.5.6 (Henselization). Let $(R, \mathfrak{m})$ be a local ring (Definition C.0.21). The *henselization of R* is the initial object among henselian local (Definition 4.5.2) R -algebras $(R^h, \mathfrak{m}^h)$ equipped with a local homomorphism (Definition C.0.5) $R \rightarrow R^h$ inducing an isomorphism on residue fields. Concretely, it is a henselian local ring R^h together with a morphism $R \rightarrow R^h$ such that any local ring homomorphism from R to a henselian local ring factors uniquely through R^h .

Definition 4.5.7 (Strict henselization). Let $(R, \mathfrak{m})$ be a local ring (Definition C.0.21) with residue field k . Fix a separable closure k^{sep} of k . The *strict henselization R^{sh}* of R is the henselization of the local ring of the étale site of $\text{Spec}(R)$ at the geometric point corresponding to k^{sep} . It is a strictly henselian local ring equipped with a local homomorphism $R \rightarrow R^{sh}$ inducing the embedding $k \subseteq k^{\text{sep}}$ on residue fields.

Theorem 4.5.8 (Examples of Henselizations and Strict Henselizations of Specific Local Rings). Let p be a prime number

1. The ring of integers localized at p , $\mathbb{Z}_{(p)}$, has Henselization (Definition 4.5.6) given by the p -adic integers $\mathbb{Z}_p$. More precisely,

$$\mathbb{Z}_p \cong \text{Henselization of } \mathbb{Z}_{(p)}$$

and is a complete discrete valuation ring that is Henselian.

2. The strict Henselization (Definition 4.5.7) of $\mathbb{Z}_{(p)}$ is the maximal unramified extension of $\mathbb{Z}_p$, obtained by adjoining all roots of unity of order prime to p . It is strictly Henselian and can be viewed as the ring of integers of the maximal unramified extension of the p -adic field $\mathbb{Q}_p$.

Theorem 4.5.9 (Stalks of étale sheaves and strict henselizations). Let X be a scheme (Definition 3.1), and let $\bar{x} : \text{Spec}(\Omega) \rightarrow X$ be a geometric point (Definition 3.0.1), where Ω is an algebraically closed field. Let $\mathcal{F}$ be a sheaf on the étale site $X_{\text{ét}}$ (Definition 4.1.3).

Writing $(\mathcal{O}_{X,x})^{sh}$ for the strict henselization (Definition 4.5.7) of the local ring at the point $x \in X$ which is the image of $\bar{x}$, there is a natural isomorphism

$$\mathcal{F}_{\bar{x}} \cong \Gamma(\text{Spec}(\mathcal{O}_{X,x}^{sh}), g^* \mathcal{F}),$$

(Definition 6.1.20) (Definition 7.1.1) where g is the canonical scheme map $\text{Spec}(\mathcal{O}_{X,x}^{sh}) \rightarrow X$. One often denotes $\Gamma(\text{Spec}(\mathcal{O}_{X,x}^{sh}), g^* \mathcal{F})$ as $\mathcal{F}(\mathcal{O}_{X,x}^{sh})$. (♠ TODO: describe this canonical scheme map.)

4.6. Constant and locally constant sheaves.

5. LOCALLY CONSTANT SHEAVES AND LOCAL SYSTEMS

Definition 5.0.1. (♠ TODO: distinguish between the case where Λ is finite and where Λ is a limit of finite rings) Let X be a scheme and let Λ be a commutative ring. A *local system of Λ -modules on X* is a locally constant sheaf of finite free Λ -modules (Definition 4.4.2) on the (small) étale site $X_{\text{ét}}$ (Definition 4.1.3) (Definition 4.1.3).

Theorem 5.0.2 (See [?, Tags 0DV5, 0GIY]). Let X be a connected (Definition 2.0.3) scheme and let $\bar{x} \in X$ be a geometric point (Definition 3.0.1).

1. There is an equivalence of categories (♠ TODO: define finite in this context)

$$\left\{ \begin{array}{l} \text{finite locally constant} \\ \text{sheaves of sets on } X_{\text{étale}} \end{array} \right\} \longleftrightarrow \left\{ \text{finite } \pi_1^{\text{ét}}(X, \bar{x})\text{-sets} \right\}.$$

(Definition 4.4.2) (Definition 4.1.3) (Definition 3.0.2)

2. There is an equivalence of categories

$$\left\{ \begin{array}{l} \text{finite cardinality locally constant} \\ \text{sheaves of abelian groups on } X_{\text{étale}} \end{array} \right\} \longleftrightarrow \left\{ \text{finite cardinality } \pi_1^{\text{ét}}(X, \bar{x})\text{-modules} \right\}.$$

3. For a finite ring Λ , there is an equivalence of categories

$$\left\{ \begin{array}{l} \text{finite type, locally constant} \\ \text{sheaves of } \Lambda\text{-modules on } X_{\text{étale}} \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \text{finite } \pi_1^{\text{ét}}(X, \bar{x})\text{-modules endowed} \\ \text{with commuting } \Lambda\text{-module structure} \end{array} \right\}.$$

(Definition 4.4.2)

♠ TODO: in the below, the geometrically unibranch hypothesis might not be needed, but it was in the stacks project

4. Assume that X is irreducible and geometrically unibranch. For a ring Λ , there is an equivalence of categories (Definition .1.6)

$$\left\{ \begin{array}{l} \text{finite type, locally constant} \\ \text{sheaves of } \Lambda\text{-modules on } X_{\text{étale}} \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{l} \text{finite type } \Lambda\text{-modules } M \text{ endowed} \\ \text{with a continuous } \pi_1^{\text{ét}}(X, \bar{x})\text{-action} \end{array} \right\}.$$

See also Theorem 8.5.3.

Proposition 5.0.3. (♠ TODO: galois reps, modules, galois actions) Let X be an integral normal scheme. Let $U \subseteq X$ be a non-empty open subscheme. Let $x \in X \setminus U$ be a point. Let $\mathcal{F}$ be a locally constant sheaf (Definition 4.4.2) on U corresponding to (Theorem 5.0.2) a (discrete) set with a continuous $\pi_1^{\text{ét}}(U, \bar{x})$ -action. The sheaf $\mathcal{F}$ is unramified at x if and only if the action is unramified at any (equivalently all) valuations of the function field of X centered at x . (♠ TODO: check the correspondence)

6. SIX FUNCTORS

6.1. The functors on sheaves.

6.1.1. Inverse and direct images of sheaves via continuous functors of general sites.

Definition 6.1.1 ([?, Exposé III, Définition 1.1]). Let $(\mathcal{C}, J)$ and $(\mathcal{D}, K)$ be sites (Definition A.0.4) (or $\mathcal{U}$ -sites (Definition A.0.5) if a Grothendieck universe (Definition .1.2) $\mathcal{U}$ is available).

(♠ TODO: the pullback sieve definition here is probably not the correct definition that is supposed to be here) A functor $u : \mathcal{C} \rightarrow \mathcal{D}$ is said to be a *continuous functor of sites* if, for every object $U \in \text{Ob}(\mathcal{D})$, every covering sieve (Definition A.0.4) $S \in K(U)$, every object $V \in \text{Ob}(\mathcal{C})$, and every morphism $f : u(V) \rightarrow U$ in $\mathcal{D}$, the pullback sieve f^*S (Definition A.0.2) belongs to $J(V)$.

$$f^*S : \{g : W \rightarrow V \text{ in } \mathcal{C} : f \circ u(g) \in S\}.$$

Equivalently, u is continuous if for every sheaf (Definition 4.0.2) of sets F on $\mathcal{D}$, the presheaf (Definition 4.0.1) $\mathcal{C}^{\text{op}} \rightarrow \mathbf{Sets}, X \mapsto F(u(X))$ is a sheaf on $\mathcal{C}$. See Proposition 6.1.2. (♠ TODO: define morphism of sites and recheck ref's to this definition)

Proposition 6.1.2. Let $(\mathcal{C}, J)$ and $(\mathcal{D}, K)$ be sites (Definition A.0.4) (or $\mathcal{U}$ -sites (Definition A.0.5) if a Grothendieck universe (Definition .1.2) $\mathcal{U}$ is available). Let $u : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. The following conditions are equivalent:

1. (Pullback sieve condition) For every object $U \in \text{Ob}(\mathcal{D})$, every covering sieve (Definition A.0.4) S on U , every object $V \in \text{Ob}(\mathcal{C})$, and every morphism $f : u(V) \rightarrow U$ in $\mathcal{D}$, the pullback sieve (Definition A.0.2) f^*S belongs to $J(V)$. This is the condition of Definition 6.1.1.
2. (Covering-sieve pushforward condition) For every object $V \in \text{Ob}(\mathcal{C})$ and every covering sieve (Definition A.0.4) S on V , the sieve generated by (Definition A.0.3) the image family $\{u(g) \mid g \in S\}$ is a covering sieve on $u(V)$ in K .
3. (Sheaf preservation condition) For every sheaf of sets (Definition 4.0.2) F on $\mathcal{D}$, the composite presheaf $F \circ u : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Sets}$ is a sheaf on $\mathcal{C}$. This is the equivalent condition mentioned in Definition 6.1.1.

A functor satisfying any (hence all) of these conditions is called a continuous functor of sites (Definition 6.1.1).

Definition 6.1.3. Let (X, τ_X) and (Y, τ_Y) be topological spaces (Definition C.0.6), and let $f : X \rightarrow Y$ be a continuous map. Let $\text{Open}(X)$ and $\text{Open}(Y)$ be their respective categories of open sets with inclusion morphisms, equipped with the canonical (Definition .1.9) Grothendieck topologies (Definition A.0.4) given by open coverings.

Define the functor

$$f^{-1} : \text{Open}(Y) \rightarrow \text{Open}(X), \quad U \mapsto f^{-1}(U).$$

It is a continuous functor of sites (Definition 6.1.1) from $\text{Open}(Y)$ to $\text{Open}(X)$ which induces a site morphism

$$f : (\text{Open}(X), \text{can}) \rightarrow (\text{Open}(Y), \text{can})$$

.

Definition 6.1.4. Let $f : X \rightarrow Y$ be a morphism of schemes, and consider one of the common Grothendieck topologies on schemes such as the Zariski, étale, Nisnevich, fppf, fpqc, or crystalline topology. Denote by $\mathbf{C}(X)$ and $\mathbf{C}(Y)$ the corresponding small sites (Definition A.0.4) of X and Y (i.e., categories of morphisms to X and Y respectively equipped with one of these topologies).

Then the base change functor

$$f^{-1} : \mathbf{C}(Y) \rightarrow \mathbf{C}(X), \quad (V \rightarrow Y) \mapsto (V \times_Y X \rightarrow X)$$

(Definition C.0.15) is a continuous functor. It in fact induces a morphism of sites

$$f : (\mathbf{C}(X), \tau_{\mathbf{C}}) \rightarrow (\mathbf{C}(Y), \tau_{\mathbf{C}})$$

where $\tau_{\mathbf{C}}$ denotes the chosen topology (Zariski, étale, Nisnevich, fppf, fpqc, crystalline).

Definition 6.1.5. (♠ TODO: there are places where sites and sheaves of rings on them are used, but it would be better to just have them be ringed sites.)

A *ringed site* is a site (Definition A.0.4) $(\mathcal{C}, J)$ with a small topological generating family (Definition A.0.4) equipped with a sheaf (Definition 4.0.2) of (not necessarily commutative) rings $\mathcal{O}$. If the Grothendieck topology J is clear in context, one may even write that $(\mathcal{C}, \mathcal{O})$ is a ringed site.

Definition 6.1.6. Let $((\mathcal{C}, J), \mathcal{O}_{\mathcal{C}})$ and $((\mathcal{D}, K), \mathcal{O}_{\mathcal{D}})$ be ringed sites (Definition 6.1.5).

A *morphism of ringed sites* $f : ((\mathcal{C}, J), \mathcal{O}_{\mathcal{C}}) \rightarrow ((\mathcal{D}, K), \mathcal{O}_{\mathcal{D}})$ is a pair $(f, f^{\#})$ where:

1. $f : (\mathcal{C}, J) \rightarrow (\mathcal{D}, K)$ is a morphism of sites.
2. $f^{\#} : \mathcal{O}_{\mathcal{D}} \rightarrow f_* \mathcal{O}_{\mathcal{C}}$ (Definition 6.1.7) is a morphism of sheaves of rings (Definition 4.0.2) on the site $(\mathcal{D}, K)$.

Equivalently, the morphism $f^{\#}$ a morphism can be described by the adjoint morphism $(??)$ $f^{\#} : f^{-1} \mathcal{O}_{\mathcal{D}} \rightarrow \mathcal{O}_{\mathcal{C}}$ of sheaves of rings on $(\mathcal{C}, J)$, where f^{-1} is the inverse image functor (Definition 6.1.10) of sheaves.

Definition 6.1.7. Let $(\mathcal{C}, J)$ and $(\mathcal{D}, K)$ be sites (Definition A.0.4) with small topological generating families (Definition A.0.4) (or U -small topologically generating families if a universe (Definition .1.2) U is available), and let $u : \mathcal{C} \rightarrow \mathcal{D}$ be a continuous functor of sites (Definition 6.1.1).

Let $\mathcal{A}$ be a (large) category which has all small (or U -small) products (Definition .1.16).

For any sheaf (Definition 4.0.2)

$$\mathcal{F} \in \mathrm{Sh}(\mathcal{D}, K; \mathcal{A}),$$

Define the *pushforward/direct image sheaf* $u^s \mathcal{F}$ by

$$u^s \mathcal{F} := \mathcal{F} \circ u : \mathcal{C}^{\mathrm{op}} \rightarrow \mathcal{A}.$$

The continuity of u implies that $u^s \mathcal{F}$ is indeed a sheaf on $(\mathcal{C}, J)$ (Lemma 6.1.8). The assignment $\mathcal{F} \mapsto u^s \mathcal{F}$ defines the *direct image/pushforward functor*

$$u^s : \mathrm{Sh}(\mathcal{D}, K; \mathcal{A}) \rightarrow \mathrm{Sh}(\mathcal{C}, J; \mathcal{A}).$$

If u is the functor underlying a site morphism $f : (\mathcal{D}, K) \rightarrow (\mathcal{C}, J)$, we may alternatively denote $u^s \mathcal{F}$ by $f_* \mathcal{F}$ and call it the *direct image/pushforward of $\mathcal{F}$ under f* ; the assignment $\mathcal{F} \mapsto f_* \mathcal{F}$ is then the *direct image/pushforward functor*.

$$f_* : \mathrm{Sh}(\mathcal{D}, K; \mathcal{A}) \rightarrow \mathrm{Sh}(\mathcal{C}, J; \mathcal{A}).$$

Note that while the continuous functor u and the site morphism f point in opposite directions, the definition $f_* := u^s$ ensures that f_* corresponds to the standard geometric pushforward used in topology and algebraic geometry.

Note that the “inverse image of a cocontinuous functor of sites” and the “direct image of a continuous functor of sites” share the same constructions, i.e. by precomposition. More precisely, if $u : \mathcal{C} \rightarrow \mathcal{D}$ is a continuous (Definition 6.1.1) and cocontinuous functor of sites, then the direct image u_* of u as a continuous functor and the inverse image u^* of u as a cocontinuous functor are exactly the same.

Lemma 6.1.8. Let $(\mathcal{C}, J)$ and $(\mathcal{D}, K)$ be sites (Definition A.0.4) with small topological generating families (Definition A.0.4) (or U -small topologically generating families if a universe (Definition .1.2) U is available), and let $u : \mathcal{C} \rightarrow \mathcal{D}$ be a continuous functor of sites (Definition 6.1.1). Let $\mathcal{A}$ be a (large) category.

For any sheaf $\mathcal{F} \in \mathrm{Sh}(\mathcal{D}, K; \mathcal{A})$ (Definition 4.0.2), the pushforward/direct image sheaf (Definition 6.1.7) $u^s \mathcal{F}$ is indeed a sheaf; more precisely, it is an object of $\mathrm{Sh}(\mathcal{C}, J; \mathcal{A})$.

Proof. Suppose that U is an object of $\mathcal{C}$ and that $S \in J(U)$ is a covering sieve (Definition A.0.4). Since u is continuous, the sieve $u(S)^+$ on $u(U)$ generated by (Definition A.0.3) the image $u(S)$ of S under u is a covering sieve (Definition A.0.4) of $u(U)$ in K (♠ TODO: why). Write $\mathcal{D}_S \hookrightarrow \mathcal{C}/U$ (Definition A.0.6) for the full downward-closed subcategory such that $\mathrm{Ob}(\mathcal{D}_S) = \{(f : V \rightarrow U) : f \in S(V)\}$. Write $\mathcal{D}_{u(S)^+} \hookrightarrow \mathcal{D}/u(U)$ for the analogous subcategory. The functor u induces a functor $u_U : \mathcal{D}_S \rightarrow \mathcal{D}_{u(S)^+}$, $(V \rightarrow U) \mapsto (u(V) \rightarrow u(U))$. Due to the continuity of u , this functor is cofinal (♠ TODO: cofinality) (♠ TODO: reference).

We want to show that the limit

$$(C) \quad \varprojlim_{(V \rightarrow U) \in (\mathcal{D}_S)^{\mathrm{op}}} (u^s \mathcal{F})|_{\mathcal{D}_S}(V)$$

exists and the canonical natural morphism

$$u^s \mathcal{F}(U) \rightarrow \varprojlim_{(V \rightarrow U) \in (\mathcal{D}_S)^{\mathrm{op}}} (u^s \mathcal{F})|_{\mathcal{D}_S}(V)$$

is an isomorphism. (C) equals

$$\varprojlim_{(V \rightarrow U) \in (\mathcal{D}_S)^{\mathrm{op}}} \mathcal{F}|_{\mathcal{D}_S}(u(V)).$$

Moreover, $u^s \mathcal{F}(U) = \mathcal{F}(u(U))$. Since $\mathcal{F}$ is a sheaf on $(\mathcal{D}, K)$, there is a canonical isomorphism

$$\mathcal{F}(u(U)) \cong \varprojlim_{W \rightarrow u(U) \in (\mathcal{D}_{u(S)^+})^{\mathrm{op}}} \mathcal{F}(W).$$

Because u_U is cofinal, the limit above, which is assumed to exist is isomorphic to

$$\varprojlim_{(V \rightarrow U) \in \mathcal{D}_S^{\text{op}}} \mathcal{F}(u(V))$$

(♠ TODO: I need a statement that says that a (co)final functor 1. guarantees the existence of one direct/inverse limit assuming the existence of the other direct/inverse limit and 2. induces isomorphisms of direct/inverse limits as long as at least one (and hence both) limit exists. The statement really should be a generalization of Theorem .1.21, which is written only for cofinal functors that are basically subcategories) as needed. $\square$

Definition 6.1.9. Let $(\mathcal{C}, J)$ and $(\mathcal{D}, K)$ be sites (Definition A.0.4) with small topological generating families (Definition A.0.4) (or U -small topologically generating families if a universe (Definition .1.2) U is available), and let $u : \mathcal{C} \rightarrow \mathcal{D}$ be a continuous functor of sites (Definition 6.1.1).

As per Definition 6.1.7 and by Theorem 4.3.2, there is a pushforward/direct image functor

$$u^s : \text{Sh}(\mathcal{D}, K; \text{Ab}) \rightarrow \text{Sh}(\mathcal{C}, J; \text{Ab})$$

As discussed in Definition 6.1.7, if u is the functor underlying a site morphism $f : (\mathcal{D}, K) \rightarrow (\mathcal{C}, J)$, then we may alternatively denote u^s by

$$f_* : \text{Sh}(\mathcal{D}, K; \text{Ab}) \rightarrow \text{Sh}(\mathcal{C}, J; \text{Ab})$$

and call this a *direct image/pushforward functor via f* .

Definition 6.1.10. Let $(\mathcal{C}, J)$ and $(\mathcal{D}, K)$ be sites (Definition A.0.4) with small topological generating families (Definition A.0.4), and let $u : \mathcal{C} \rightarrow \mathcal{D}$ be a continuous functor of sites (Definition 6.1.1). Let $\mathcal{A}$ be a (large) category such that the presheaf category (Definition 4.0.1) $\text{PreSh}(\mathcal{D}, K; \mathcal{A})$ has sheafification (Definition 4.3.1).

For any sheaf (Definition 4.0.2)

$$\mathcal{G} \in \text{Sh}(\mathcal{C}, J; \mathcal{A}),$$

the *inverse image/pullback sheaf of $\mathcal{G}$ under u* is defined, assuming that all colimits below exist, as:

$$u_s \mathcal{G} : \mathcal{D}^{\text{op}} \rightarrow \mathcal{A}, \quad V \mapsto a \left(\varinjlim_{(V \downarrow u)} \mathcal{G}(U) \right),$$

where a is the sheafification (Definition 4.3.1) functor of presheaves and the colimit (Definition .1.13) is taken over the comma category $(V \downarrow u)$ of pairs $(U, V \rightarrow u(U))$ with $U \in \mathcal{C}$.

The assignment $\mathcal{G} \mapsto u_s \mathcal{G}$ defines the *inverse image/pullback functor*

$$u_s : \text{Sh}(\mathcal{C}, J; \mathcal{A}) \rightarrow \text{Sh}(\mathcal{D}, K; \mathcal{A}).$$

If u is the functor underlying a site morphism $f : (\mathcal{D}, K) \rightarrow (\mathcal{C}, J)$, we may alternatively denote $u_s \mathcal{G}$ by $f^* \mathcal{G}$ (or sometimes by $f^{-1} \mathcal{G}$) and call it the *inverse image/pullback of $\mathcal{G}$ under f* .

Note that while the continuous functor u and the site morphism f point in opposite directions, the identification $f^* := u_s$ ensures that f^* corresponds to the standard geometric pullback. In the case of topological spaces, this recovers the usual construction involving colimits over open neighborhoods to obtain stalks followed by sheafification.

The inverse image f^* of a continuous functor of sites should not be confused with the notion of the inverse image of a cocontinuous functor of sites, as these two notions are different constructions, even when f is both continuous and cocontinuous.

Definition 6.1.11. Let $(\mathcal{C}, J)$ and $(\mathcal{D}, K)$ be sites (Definition A.0.4) with small topological generating families (Definition A.0.4) (or U -small topologically generating families if a universe (Definition .1.2) U is available), and let $u : \mathcal{C} \rightarrow \mathcal{D}$ be a continuous functor of sites (Definition 6.1.1).

As per Definition 6.1.10 and by Theorem 4.3.2, there is a pullback/inverse image functor

$$u_s : \mathrm{Sh}(\mathcal{C}, J; \mathrm{Ab}) \rightarrow \mathrm{Sh}(\mathcal{D}, K; \mathrm{Ab})$$

As discussed in Definition 6.1.10, if u is the functor underlying a site morphism $f : (\mathcal{D}, K) \rightarrow (\mathcal{C}, J)$, then we may alternatively denote u_s by

$$f^* : \mathrm{Sh}(\mathcal{C}, J; \mathrm{Ab}) \rightarrow \mathrm{Sh}(\mathcal{D}, K; \mathrm{Ab})$$

and call this a *inverse image/pullback functor via f* .

Theorem 6.1.12. Let (C, J) and (D, K) be sites (Definition A.0.4) with small (Definition .1.4) topologically generating families, and let $u : C \rightarrow D$ be a continuous functor of sites (Definition 6.1.1).

Let $\mathcal{A}$ be a locally small (Definition .1.4) (or U -locally small) category such that the presheaf category (Definition 4.0.1) $\mathrm{PreSh}(\mathcal{D}, K; \mathcal{A})$ has sheafification (Definition 4.3.1). In particular, $\mathrm{Sh}(D, K; \mathcal{A})$ and $\mathrm{Sh}(C, J; \mathcal{A})$ are locally small².

The inverse image (Definition 6.1.7) and direct image (Definition 6.1.10) functors

$$u_s : \mathrm{Sh}(D, K; \mathcal{A}) \rightarrow \mathrm{Sh}(C, J; \mathcal{A})$$

$$u^s : \mathrm{Sh}(C, J; \mathcal{A}) \rightarrow \mathrm{Sh}(D, K; \mathcal{A})$$

are adjoint (Definition .1.8), i.e. for any sheaves (Definition 4.0.2),

$$\mathcal{F} \in \mathrm{Sh}(C, J; \mathcal{A}), \quad \mathcal{G} \in \mathrm{Sh}(D, K; \mathcal{A}),$$

there is a natural isomorphism

$$\mathrm{Hom}_{\mathrm{Sh}(C, J; \mathcal{A})}(u_s \mathcal{G}, \mathcal{F}) \cong \mathrm{Hom}_{\mathrm{Sh}(D, K; \mathcal{A})}(\mathcal{G}, u^s \mathcal{F}).$$

In particular, if u is the functor underlying a site morphism $f : (\mathcal{D}, K) \rightarrow (\mathcal{C}, J)$, then there is a natural isomorphism

$$\mathrm{Hom}_{\mathrm{Sh}(C, J; \mathcal{A})}(f^* \mathcal{G}, \mathcal{F}) \cong \mathrm{Hom}_{\mathrm{Sh}(D, K; \mathcal{A})}(\mathcal{G}, f_* \mathcal{F}).$$

²as a consequence of Lemma .1.7

Theorem 6.1.13. Let (C, J) and (D, K) be sites (Definition A.0.4) with small (Definition .1.4) topologically generating families, and let $u : C \rightarrow D$ be a continuous functor of sites (Definition 6.1.1).

The inverse image (Definition 6.1.11) and direct image (Definition 6.1.9) functors

$$\begin{aligned} u_s : \mathrm{Sh}(D, K; \mathrm{Ab}) &\rightarrow \mathrm{Sh}(C, J; \mathrm{Ab}) \\ u^s : \mathrm{Sh}(C, J; \mathrm{Ab}) &\rightarrow \mathrm{Sh}(D, K; \mathrm{Ab}) \end{aligned}$$

are adjoint (Definition .1.8), i.e. for any sheaves (Definition 4.0.2),

$$\mathcal{F} \in \mathrm{Sh}(C, J; \mathrm{Ab}), \quad \mathcal{G} \in \mathrm{Sh}(D, K; \mathrm{Ab}),$$

there is a natural isomorphism

$$\mathrm{Hom}_{\mathrm{Sh}(C, J; \mathrm{Ab})}(u_s \mathcal{G}, \mathcal{F}) \cong \mathrm{Hom}_{\mathrm{Sh}(D, K; \mathrm{Ab})}(\mathcal{G}, u^s \mathcal{F}).$$

In particular, if u is the functor underlying a site morphism $f : (\mathcal{D}, K) \rightarrow (\mathcal{C}, J)$, then there is a natural isomorphism

$$\mathrm{Hom}_{\mathrm{Sh}(C, J; \mathrm{Ab})}(f^* \mathcal{G}, \mathcal{F}) \cong \mathrm{Hom}_{\mathrm{Sh}(D, K; \mathrm{Ab})}(\mathcal{G}, f_* \mathcal{F}).$$

See also Theorem 6.1.16

Definition 6.1.14. Let $f : (\mathcal{C}, J, \mathcal{O}_C) \rightarrow (\mathcal{D}, K), \mathcal{O}_D$ be a morphism of ringed sites (Definition B.9.6) with small (Definition .1.4) topological generating families (Definition A.0.4).

Given a (right) $\mathcal{O}_C$ -module (Definition 7.1.3) $\mathcal{F}$, its direct image (Definition 6.1.9) $f_* \mathcal{G}$ has the natural structure of a $f_* \mathcal{O}_C$ -module. Since there is a natural map

$$f^\# : \mathcal{O}_D \rightarrow f_* \mathcal{O}_C$$

of sheaves of rings on $(\mathcal{D}, K)$ coming from the morphism of ringed sites, $f^* \mathcal{G}$ can be regarded as a $\mathcal{O}_D$ -module. This $\mathcal{O}_D$ -module is also called the *direct image/pushforward of $\mathcal{G}$ under f* and is still denoted by $f_* \mathcal{G}$.

Definition 6.1.15. Let $f : (\mathcal{C}, J, \mathcal{O}_C) \rightarrow (\mathcal{D}, K), \mathcal{O}_D$ be a morphism of ringed sites (Definition B.9.6) with small (Definition .1.4) topological generating families (Definition A.0.4). Write f_{Ab}^* for the inverse image functor (Definition 6.1.11)

$$f_{\mathrm{Ab}}^* : \mathrm{Sh}(\mathcal{D}, K, \mathrm{Ab}) \rightarrow \mathrm{Sh}(\mathcal{C}, J, \mathrm{Ab}).$$

Given a (right) $\mathcal{O}_D$ -module (Definition 7.1.3) $\mathcal{G}$, its *inverse image*, notated $f^* \mathcal{G}$ by abuse of notation and which should not be mistaken for f_{Ab}^* , is the $\mathcal{O}_C$ -module constructed as

$$f_{\mathrm{Ab}}^* \mathcal{G} \otimes_{f^* \mathcal{O}_D} \mathcal{O}_C.$$

Note that there is a morphism $f^* \mathcal{O}_D \rightarrow \mathcal{O}_C$ corresponding under the adjunction of f^* and f_* (??) (Theorem 6.1.13)

Theorem 6.1.16. Let $f : (\mathcal{C}, J, \mathcal{O}_C) \rightarrow (\mathcal{D}, K), \mathcal{O}_D$ be a morphism of ringed sites (Definition B.9.6) with small (Definition .1.4) topological generating families (Definition A.0.4).

The inverse image (Definition 6.1.15) and direct image (Definition 6.1.14) functors

$$f^* : \mathrm{Mod}(\mathcal{O}_D) \rightarrow \mathrm{Mod}(\mathcal{O}_C)$$

$$f_* : \text{Mod}(\mathcal{O}_C) \rightarrow \text{Mod}(\mathcal{O}_D)$$

are adjoint (Definition .1.8), i.e. for any

$$\mathcal{F} \in \text{Mod}(\mathcal{O}_C), \quad \mathcal{G} \in \text{Mod}(\mathcal{O}_D),$$

there is a natural isomorphism

$$\text{Hom}_{\mathcal{O}_C}(f^*\mathcal{G}, \mathcal{F}) \cong \text{Hom}_{\mathcal{O}_D}(\mathcal{G}, f_*\mathcal{F}).$$

Definition 6.1.17 (Pullback (inverse image) of a sheaf). Let $f : X \rightarrow Y$ be a continuous map between topological spaces (Definition C.0.6). Let $\mathcal{D}$ be a category with a terminal object (Definition .1.10).

1. Let $\mathcal{G}$ be a presheaf on Y valued in a $\mathcal{D}$ (Definition C.0.22). The *pullback* or *inverse image presheaf* $f^{-1}\mathcal{G}$ on X is defined as the presheaf

$$U \mapsto \varinjlim_{V \supseteq f(U)} \mathcal{G}(V)$$

provided these colimits (Definition .1.13) exist for all U , where U ranges over open subsets of X and the colimit is taken over all open subsets $V \subseteq Y$ containing $f(U)$. This construction admits a natural isomorphism

$$(f^{-1}\mathcal{G})_x \rightarrow \mathcal{G}_{f(x)}$$

of stalks (Definition C.0.23) for every $x \in X$.

2. If $\mathcal{G}$ is a sheaf valued in $\mathcal{D}$, then we can define the *pullback* or *inverse image sheaf* $f^*\mathcal{G}$ on X as the sheaf associated to the presheaf (Definition 4.3.1) $f^{-1}\mathcal{G}$, assuming it exists.

Assuming that a sheafification functor (Definition 4.3.1) exists, one may equivalently define $f^*\mathcal{G}$ via Definition 6.1.10 — More concretely, $f^*\mathcal{G}$ is the following equivalent constructions:

- The inverse image (Definition 6.1.10) $(f^{-1})_s\mathcal{G}$ of $\mathcal{G}$ under the continuous functor $f^{-1} : \text{Open } Y \rightarrow \text{Open } X$, $W \mapsto f^{-1}(W)$ (Definition .1.9) (Definition 6.1.3).
- The inverse image (Definition 6.1.10) of $\mathcal{G}$ under the site morphism $\text{Open } X \rightarrow \text{Open } Y$ whose underlying continuous functor (Definition 6.1.1) is f^{-1}

If $\mathcal{D}$ is the category of sets, of groups, of abelian groups, of modules over rings, for example, then the relevant colimits (Definition .1.13) needed to define the inverse image presheaves and the sheafification needed to define the inverse image sheaves exist.

Definition 6.1.18 (Pushforward (direct image) of a sheaf). Let $f : X \rightarrow Y$ be a continuous map between topological spaces (Definition C.0.6), and let $\mathcal{F}$ be a presheaf on X valued in a category $\mathcal{D}$ (Definition C.0.22) with a terminal object. The *pushforward* or *direct image presheaf* $f_*\mathcal{F}$ on Y is the presheaf valued in $\mathcal{D}$ on Y (Definition 4.0.1) defined as follows: For every open set $V \subseteq Y$, the value of the pushforward is given by

$$f_*\mathcal{F}(V) := \mathcal{F}(f^{-1}(V)).$$

For an inclusion of open sets $V' \subseteq V$ in Y , the restriction morphism

$$\text{res}_{V, V'}^{f_*\mathcal{F}} : f_*\mathcal{F}(V) \rightarrow f_*\mathcal{F}(V')$$

is defined as the restriction morphism of $\mathcal{F}$ associated with the inclusion of preimages $f^{-1}(V') \subseteq f^{-1}(V)$ in X :

$$\text{res}_{f^{-1}(V), f^{-1}(V')}^{\mathcal{F}} : \mathcal{F}(f^{-1}(V)) \rightarrow \mathcal{F}(f^{-1}(V')).$$

If $\mathcal{F}$ is a sheaf, then so is $f_*\mathcal{F}$. In this case, it is equivalent to define $f_*\mathcal{F}$ as the direct image (Definition 6.1.7) $(f^{-1})^s\mathcal{F}$ of $\mathcal{F}$ under the continuous functor $f^{-1} : \text{Open } Y \rightarrow \text{Open } X$ (Definition 6.1.3) (Definition 6.1.3) (Definition .1.9), which is also equivalent to the direct image (Definition 6.1.7) of $\mathcal{F}$ under the site morphism $\text{Open } X \rightarrow \text{Open } Y$ whose underlying continuous functor is f^{-1} .

(♠ TODO: comment on what is happening on the stalks)

Theorem 6.1.19. Let $f : X \rightarrow Y$ be a continuous map between topological spaces (Definition C.0.6). Let $\mathcal{A}$ be a locally small category (Definition .1.4) such that $\mathcal{A}$ is cocomplete, i.e. admits small colimits, and such that $\text{PreShv}(Y, \mathcal{A})$ admits sheafification; for example, the categories of sets, of groups, of abelian groups, of rings, of modules over a ring, etc. satisfy these hypotheses.

The inverse image functor (Definition 6.1.17)

$$f^* : \text{Sh}(Y, \mathcal{A}) \rightarrow \text{Sh}(X, \mathcal{A})$$

and the direct image functor (Definition 6.1.18)

$$f_* : \text{Sh}(X, \mathcal{A}) \rightarrow \text{Sh}(Y, \mathcal{A})$$

form an adjoint pair (Definition .1.8) $f^* \dashv f_*$, i.e., for any sheaves $\mathcal{F} \in \text{Sh}(Y, \mathcal{A})$ and $\mathcal{G} \in \text{Sh}(X, \mathcal{A})$, there is a natural isomorphism

$$\text{Hom}_{\text{Sh}(X, \mathcal{A})}(f^*\mathcal{F}, \mathcal{G}) \cong \text{Hom}_{\text{Sh}(Y, \mathcal{A})}(\mathcal{F}, f_*\mathcal{G}).$$

Definition 6.1.20. (♠ TODO: think about if assumptions on the small site) Let $f : X \rightarrow Y$ be a morphism of schemes, and let $\mathbf{C}(X)$ and $\mathbf{C}(Y)$ be small sites (Definition 4.1.1) associated to X and Y respectively, equipped with a Grothendieck topologies for which the continuous functor (Definition 6.1.1) $f^{-1} : \mathbf{C}(Y) \rightarrow \mathbf{C}(X)$ (Definition 6.1.4) on sites underlies a site morphism $f : \mathbf{C}(X) \rightarrow \mathbf{C}(Y)$. For example, the common Grothendieck topologies such as Zariski (Definition 4.2.4), étale (Definition 4.1.3), Nisnevich (Definition 4.2.13), fppf (Definition 4.2.7), fpqc (Definition 4.2.9), or crystalline satisfy this hypothesis(♠ TODO: cite these).

1. Let $\mathcal{A}$ be a (large) category. Given a sheaf $\mathcal{F} \in \text{Sh}(\mathbf{C}(X); \mathcal{A})$, the *direct image along f* is the sheaf $f_*\mathcal{F}$ defined as the direct image (Definition 6.1.7) of $\mathcal{F}$ under site morphism $f : \mathbf{C}(X) \rightarrow \mathbf{C}(Y)$. In particular, it is an object of $\text{Sh}(\mathbf{C}(Y); \mathcal{A})$ if it exists. If $f_*\mathcal{F}$ exists for all sheaves $\mathcal{F} \in \text{Sh}(\mathbf{C}(X); \mathcal{A})$, then f_* yields a functor

$$f_* : \text{Sh}(\mathbf{C}(X); \mathcal{A}) \rightarrow \text{Sh}(\mathbf{C}(Y); \mathcal{A}).$$

2. We have (Definition 6.1.9) a direct image functor

$$f_* : \text{Sh}(\mathbf{C}(X); \text{Ab}) \rightarrow \text{Sh}(\mathbf{C}(Y); \text{Ab}).$$

In fact, if f yields a ringed site morphism $f : (\mathbf{C}(X), \mathcal{O}_X) \rightarrow (\mathbf{C}(Y), \mathcal{O}_Y)$, then this induces (Definition 6.1.14) a functor

$$f_* : \text{Mod}(\mathcal{O}_X) \rightarrow \text{Mod}(\mathcal{O}_Y).$$

3. Let $\mathcal{A}$ be a (large) category such that the presheaf category (Definition 4.0.1) $\text{PreSh}(\mathcal{D}, K; \mathcal{A})$ has sheafification (Definition 4.3.1). Given a sheaf $\mathcal{G} \in \text{Sh}(\mathbf{C}(Y); \mathcal{A})$, the *inverse image of $\mathcal{G}$ along f* is the sheaf $f^*\mathcal{G}$ defined as the inverse image (Definition 6.1.10) of $\mathcal{G}$ under the continuous functor (Definition 6.1.1) $f^{-1} : \mathbf{C}(Y) \rightarrow \mathbf{C}(X)$ (Definition 6.1.4) on sites. In particular, it is an object of $\text{Sh}(\mathbf{C}(X); \mathcal{A})$ and f^* yields a functor

$$f^* : \text{Sh}(\mathbf{C}(Y); \mathcal{A}) \rightarrow \text{Sh}(\mathbf{C}(X); \mathcal{A}),$$

4. We have (Definition 6.1.11) an inverse image functor

$$f^* : \text{Sh}(\mathbf{C}(Y); \text{Ab}) \rightarrow \text{Sh}(\mathbf{C}(X); \text{Ab}).$$

In fact, if f yields a ringed site morphism $f : (\mathbf{C}(X), \mathcal{O}_X) \rightarrow (\mathbf{C}(Y), \mathcal{O}_Y)$, then this induces (Definition 6.1.15) a functor

$$f^* : \text{Mod}(\mathcal{O}_Y) \rightarrow \text{Mod}(\mathcal{O}_X).$$

Proposition 6.1.21. Let $j : U \hookrightarrow X$ be an open embedding (Definition C.0.14) of schemes. Let Λ be a ring. Recall that (Theorem 6.1.16) there is an adjunction (Definition .1.8) $j^* \dashv j_*$ between the functors

$$j^* : \text{Sh}(X_{\text{ét}}, \Lambda) \Rightarrow \text{Sh}(U_{\text{ét}}, \Lambda)$$

(Definition 6.1.20) (Definition 4.1.3) (Definition 7.1.3) and

$$j_* : \text{Sh}(U_{\text{ét}}, \Lambda) \Rightarrow \text{Sh}(X_{\text{ét}}, \Lambda)$$

The counit (Definition .1.8) natural morphism

$$j^*j_* \Rightarrow \text{id}$$

is in fact a natural isomorphism.

6.1.2. Extension by zero of sheaves via open immersions of schemes.

Definition 6.1.22 (Extension by zero functor for open immersions of topological spaces). Let X be a topological space and let $j : U \hookrightarrow X$ be an open immersion, where U is an open subset of X . Consider the category $\text{Sh}(X)$ of sheaves of abelian groups on X and $\text{Sh}(U)$ on U . The *extension by zero functor*

$$j_! : \text{Sh}(U) \rightarrow \text{Sh}(X)$$

is defined as follows: for any sheaf $\mathcal{F}$ on U , the sheaf $j_!\mathcal{F}$ on X is given by

$$(j_!\mathcal{F})(V) = \{s \in \mathcal{F}(V \cap U) \mid s \text{ extends by zero outside } U\}$$

for each open subset $V \subseteq X$. Concretely, sections over V are sections over $V \cap U$, and sections supported outside U are identified with zero.

Definition 6.1.23 (Extension by zero functor for open immersions and the étale site). Let X be a scheme (Definition .3.1), $j : U \hookrightarrow X$ an open immersion (Definition C.0.14) of schemes, and let Et/X (resp. Et/U) denote the small étale site (Definition 4.1.3) of X (resp. U).

The *extension by zero functor via the open immersion j* is the functor

$$j_! : \mathrm{Sh}(Et/U) \rightarrow \mathrm{Sh}(Et/X),$$

also written $j_{U!}$, is defined as follows: for any sheaf $\mathcal{F}$ of abelian groups on Et/U and any object $V \rightarrow X$ of Et/X ,

- if $V \rightarrow X$ factors through U (i.e., $V \times_X U \cong V$), then $(j_!\mathcal{F})(V) = \mathcal{F}(V \times_X U)$,
- otherwise, $(j_!\mathcal{F})(V) = 0$.

The functor $j_!$ is exact (Definition .1.36). (♠ TODO: Prove this)

Proposition 6.1.24 (Compactly supported direct image under closed immersion and finite morphism for étale sheaves). (♠ TODO: AI generated, verify) Let $i : Z \hookrightarrow X$ be a closed immersion (Definition C.0.14) of schemes, and let $\mathcal{F}$ be a sheaf on the étale site $Z_{\text{ét}}$ (Definition 4.1.3). Since closed immersions are proper, the direct image with compact support coincides with the ordinary direct image:

$$i_! = i_*$$

In particular, $(i_!\mathcal{F})_{\bar{x}} = \mathcal{F}_{\bar{x}}$ if $\bar{x}$ lands in Z , and $(i_!\mathcal{F})_{\bar{x}} = 0$ otherwise.

Similarly, let $f : X \rightarrow Y$ be a finite morphism (Definition 1.2.1) (hence proper). For any sheaf $\mathcal{F}$ on $X_{\text{ét}}$ and geometric point $\bar{y}$ of Y :

$$(f_!\mathcal{F})_{\bar{y}} = \prod_{\bar{x} \in f^{-1}(\bar{y})} \mathcal{F}_{\bar{x}}$$

Proof. (♠ TODO: AI generated, verify) Closed immersions and finite morphisms are proper. For proper morphisms, the forget-support map $f_! \rightarrow f_*$ is an isomorphism, as compactness of support is automatic for sections over proper preimages. The stalk formulas then follow from the corresponding propositions for f_* . $\square$

6.1.3. Hom and tensor product functors of sheaves. (♠ TODO:)

6.1.4. Stalks of sheaves under the functors.

Lemma 6.1.25 (Decomposition of a finite morphism over a Henselian local ring). (♠ TODO: AI generated, verify) Let $(R, \mathfrak{m})$ be a Henselian local ring (Definition 4.5.2) with residue field $k = R/\mathfrak{m}$. Let A be a finite R -algebra. Then the canonical map (♠ TODO: what is this canonical map? Are you saying that there is a map $A \rightarrow A_{\mathfrak{n}}$ for every $\mathfrak{n}$ and then we have a map $A \rightarrow \prod$ by the universal property of products?) induces an isomorphism of rings

$$A \cong \prod_{\mathfrak{n} \in \mathrm{Spec}(A), \mathfrak{n} \cap R = \mathfrak{m}} A_{\mathfrak{n}},$$

(♠ TODO: write the definition of lying over for primes in rings and schemes) where the product runs over all prime ideals (Definition C.0.4) of A lying over $\mathfrak{m}$. Since A is finite over R , there are finitely many such primes.

Equivalently, for any scheme (Definition 3.1) Y and any finite morphism (Definition 1.2.1) $f : X \rightarrow Y$, let $\bar{y} : \text{Spec}(\Omega) \rightarrow Y$ be a geometric point (Definition 3.0.1) of Y . There exists an étale neighborhood $(V, \bar{v})$ of $\bar{y}$ such that the fiber product (Definition C.0.15) decomposes as a finite disjoint union of open-and-closed subschemes

$$X \times_Y V = \coprod_{\bar{x} \in f^{-1}(\bar{y})} Z_{\bar{x}},$$

where $f^{-1}(\bar{y})$ denotes the set of geometric points $\text{Spec}(\Omega) \rightarrow X$ lying over $\bar{y}$ (which is finite since f is finite), and each $Z_{\bar{x}}$ is an affine scheme containing exactly one geometric point $\bar{x}$ mapping to $\bar{v}$. Each $Z_{\bar{x}} \rightarrow V$ is finite (Definition 1.2.1) and flat (Definition 1.1.4) over a possibly smaller étale neighborhood of $\bar{y}$. (♠ TODO: define what it means for an étale neighborhood to be "smaller" than another and define what it means to for a condition to hold for all sufficiently small étale neighborhoods)

Proof. (♠ TODO: AI generated, verify) **Algebraic form.** Since A is a finite R -algebra, the fiber $A \otimes_R k = A/\mathfrak{m}A$ is a finite-dimensional k -algebra and hence Artinian. An Artinian ring decomposes uniquely as a finite product of local Artinian rings:

$$A/\mathfrak{m}A \cong \prod_{i=1}^n (A/\mathfrak{m}A)_{\mathfrak{n}_i},$$

where $\mathfrak{n}_1, \dots, \mathfrak{n}_n$ are the prime ideals of A lying over $\mathfrak{m}$ (these are precisely the maximal ideals of $A/\mathfrak{m}A$). The idempotents of this product lift uniquely to orthogonal idempotents of A because R is Henselian: a factorization of 1 into orthogonal idempotents modulo $\mathfrak{m}$ lifts to A (Definition 4.5.2). These lifted idempotents yield the decomposition $A \cong \prod_{i=1}^n Ae_i$, where each Ae_i is a local finite R -algebra with maximal ideal corresponding to $\mathfrak{n}_i$.

Geometric form. Since f is finite, it is affine. After replacing Y and X by affine open neighborhoods of $\bar{y}$ and $f^{-1}(\bar{y})$, we may assume $Y = \text{Spec}(R)$ and $X = \text{Spec}(A)$ for some finite R -algebra A . Let $(R^{sh}, \mathfrak{m}^{sh})$ be the strict henselization of R at $\bar{y}$; this is a strictly henselian local ring (Definition 4.5.4) (Definition 4.5.7), hence Henselian. By the algebraic form, we have $A \otimes_R R^{sh} \cong \prod_{\bar{x} \in f^{-1}(\bar{y})} (A \otimes_R R^{sh})_{\mathfrak{n}_{\bar{x}}}$ where each factor is local.

Since the decomposition is given by a finite set of orthogonal idempotents in $A \otimes_R R^{sh}$, and R^{sh} is a filtered colimit of étale R -algebras, these idempotents descend to some étale R -algebra S . That is, there exists an étale morphism $V = \text{Spec}(S) \rightarrow Y$ and a point $\bar{v} : \text{Spec}(\Omega) \rightarrow V$ lifting $\bar{y}$ such that the idempotents defining the decomposition exist in $A \otimes_R S$. Shrinking V to an affine étale neighborhood if necessary, we obtain

$$X \times_Y V = \text{Spec}(A \otimes_R S) \cong \coprod_{\bar{x} \in f^{-1}(\bar{y})} Z_{\bar{x}}$$

as a disjoint union of open-and-closed subschemes. Each $Z_{\bar{x}}$ is affine, finite over V , and contains exactly one geometric point $\bar{x}$ lying over $\bar{v}$. $\square$

Theorem 6.1.26 (Zariski's Main Theorem, Grothendieck form). Let $f : X \rightarrow Y$ be a quasi-finite morphism of schemes that is also separated. Then f factors as

$$X \xrightarrow{j} \tilde{X} \xrightarrow{\pi} Y$$

where j is an open immersion (Definition C.0.14) and π is a finite morphism (Definition 1.2.1).

Moreover, for every point $y \in Y$, the open immersion j induces a bijection between $X_y = X \times_Y \operatorname{Spec}(\kappa(y))$ and the connected component of $\tilde{X}_y = \tilde{X} \times_Y \operatorname{Spec}(\kappa(y))$ containing the image of X_y . In particular, if f is also surjective with geometrically connected fibers, then j is an isomorphism and f itself is finite.

Proof. (♠ TODO: AI generated, verify) Since f is separated, the diagonal morphism (Definition 11.1.1) $\Delta_f : X \rightarrow X \times_Y X$ is a closed immersion. Since f is quasi-finite, the diagonal is also an open immersion (the locus where two maps into a separated scheme agree is closed, and quasi-finiteness implies it is locally an isomorphism on a neighborhood of the diagonal). Hence Δ_f identifies X with an open subscheme of $X \times_Y X$.

One constructs $\tilde{X}$ by taking the normalization of Y in the function field of X (when Y is integral and normal) or, more generally, by taking the closure of the graph of f inside $X \times_Y \mathbb{P}^n$ for some embedding and then normalizing. The resulting morphism $\pi : \tilde{X} \rightarrow Y$ is finite by construction, and $j : X \hookrightarrow \tilde{X}$ is an open immersion by the quasi-finiteness of f .

The fiberwise statement follows from the fact that for a finite morphism, the fibers are finite schemes, and the connected components of a finite scheme over a field correspond to the local factors in its decomposition. Since j is an open immersion and induces a bijection on geometric points (by quasi-finiteness), it identifies X_y with the connected component of $\tilde{X}_y$ containing it. $\square$

Lemma 6.1.27. Let G be a group. Let $f : X \rightarrow Y$ be a a Galois G -morphism of schemes (Definition 1.2.4) where Y is a connected scheme (Definition .3.1). Let $\mathcal{F}$ be a sheaf on (Definition 4.0.2) $X_{\text{ét}}$ (Definition 4.1.3) valued in a (large) category $\mathcal{A}$ that has all small (Definition .1.4) products (Definition .1.16) (or U -small (Definition .1.4) products (Definition .1.16) if a universe (Definition .1.2) U is available). The direct image (Definition 6.1.20) $f_*\mathcal{F}$ is a sheaf on $Y_{\text{ét}}$ equipped with an action by (Definition .1.23) G . In fact, the direct image functor is a functor

$$f_* : \operatorname{Sh}(X_{\text{ét}}, \mathcal{A}) \rightarrow G - \operatorname{Sh}(Y_{\text{ét}}, \mathcal{A})$$

where $G - \operatorname{Sh}(Y_{\text{ét}}, \mathcal{A})$ (Definition .1.25) is the category of objects of $\operatorname{Sh}(Y_{\text{ét}}, \mathcal{A})$ equipped with a G -action.

Proof. For any étale morphism (Definition 1.1.6) $U \rightarrow Y$, we have

$$\Gamma(U, f_*\mathcal{F}) = \Gamma(U \times_Y X, \mathcal{F})$$

(Definition 7.1.1). Due to identification of $\operatorname{Aut}_Y(X)$ (Definition 1.1.7) with G the action of G on X induces an action of G on $U \times_Y X$ — for all $g \in G$, there is an automorphism $g : X \rightarrow X$ over Y , which in turn yields an automorphism $U \times_Y X \xrightarrow{(\operatorname{id}, g)} U \times_Y X$. In turn, each such automorphism (contravariantly) induces an automorphism $g : \Gamma(U \times_Y X, \mathcal{F}) \rightarrow \Gamma(U \times_Y X, \mathcal{F})$

$X, \mathcal{F}$). That G acts on X implies that these automorphisms on $\Gamma(U \times_Y X, \mathcal{F})$ together form a group action by G — for all $g, h \in G$, the automorphism $(g \cdot h)$ on $\Gamma(U \times_Y X, \mathcal{F})$ coincides with the composition of the automorphisms g and h , and $1 \in G$ induces the identity automorphism.

Moreover, for any morphism $p : (U \rightarrow Y) \rightarrow (V \rightarrow Y)$ in $Y_{\text{ét}}$ and for any $g \in G$, we have a commuting diagram

$$\begin{array}{ccc} U \times_Y X & \xrightarrow{(\text{id}, g)} & U \times_Y X \\ \downarrow (p, \text{id}) & & \downarrow (p, \text{id}) \\ V \times_Y X & \xrightarrow{(\text{id}, g)} & V \times_Y X \end{array}$$

which contravariantly induces a commuting diagram

$$\begin{array}{ccc} \Gamma(V \times_Y X, \mathcal{F}) & \xrightarrow{g} & \Gamma(V \times_Y X, \mathcal{F}) \\ \downarrow & & \downarrow \\ \Gamma(U \times_Y X, \mathcal{F}) & \xrightarrow{g} & \Gamma(U \times_Y X, \mathcal{F}). \end{array}$$

Therefore, for each $g \in G$, there is an automorphism $g : f_* \mathcal{F} \rightarrow f_* \mathcal{F}$, and in fact that the automorphisms on $\Gamma(U \times_Y X, \mathcal{F})$ together form an action by G naturally in U implies that the automorphisms on $f_* \mathcal{F}$ together form an action by G as well.

To see that f_* is a functor

$$f_* : \text{Sh}(X_{\text{ét}}, \mathcal{A}) \rightarrow G\text{-}\text{Sh}(Y_{\text{ét}}, \mathcal{A}),$$

let $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ be any morphism in $\text{Sh}(X_{\text{ét}}, \mathcal{A})$. We want to show that the induced morphism $f_* \varphi : f_* \mathcal{F} \rightarrow f_* \mathcal{G}$ is G -equivariant, i.e. the following diagram commutes for all $g \in G$:

$$\begin{array}{ccc} f_* \mathcal{F} & \xrightarrow{g} & f_* \mathcal{F} \\ \downarrow f_* \varphi & & \downarrow f_* \varphi \\ f_* \mathcal{G} & \xrightarrow{g} & f_* \mathcal{G}. \end{array}$$

Equivalently, we are to show that

$$\begin{array}{ccc} \Gamma(U, f_* \mathcal{F}) & \xrightarrow{g} & \Gamma(U, f_* \mathcal{F}) \\ \downarrow f_* \varphi & & \downarrow f_* \varphi \\ \Gamma(U, f_* \mathcal{G}) & \xrightarrow{g} & \Gamma(U, f_* \mathcal{G}). \end{array}$$

commutes for all $U \rightarrow Y$ in $Y_{\text{ét}}$. This diagram is identifiable with

$$\begin{array}{ccc}
\Gamma(U \times_Y X, \mathcal{F}) & \xrightarrow{g} & \Gamma(U \times_Y X, \mathcal{F}) \\
\downarrow \varphi & & \downarrow \varphi \\
\Gamma(U \times_Y X, \mathcal{G}) & \xrightarrow{g} & \Gamma(U \times_Y X, \mathcal{G}).
\end{array}$$

That φ is a morphism of (pre)sheaves implies that the above commutes, because $g : U \times_Y X \rightarrow U \times_Y X$ is a (auto)morphism in $X_{\text{ét}}$. Thus, f_* is indeed a functor into $G\text{-Sh}(Y_{\text{ét}}, \mathcal{A})$. $\square$

Proposition 6.1.28 (Stalk of direct image under finite morphism for étale sheaves). Let $f : X \rightarrow Y$ be a finite morphism of schemes (Definition 1.2.1), and let $\mathcal{F}$ be a sheaf on étale site $X_{\text{ét}}$ (Definition 4.1.3) of (♠ TODO: try to change this to a more general category with direct limits)

- sets,
- abelian groups, or
- $\mathcal{O}$ -modules (Definition 7.1.3) for a sheaf (Definition 4.0.2) $\mathcal{O}$ of rings (Definition C.0.8) on $X_{\text{ét}}$.

Let $\bar{y} : \text{Spec}(\Omega) \rightarrow Y$ be a geometric point (Definition 3.0.1) of Y .

1. The stalk (Definition 4.5.1) of the direct image (Definition 6.1.20) at $\bar{y}$ decomposes as a product over the geometric fiber:

$$(f_*\mathcal{F})_{\bar{y}} = \prod_{\bar{x} \in f^{-1}(\bar{y})} \mathcal{F}_{\bar{x}},$$

where $f^{-1}(\bar{y})$ is the set of geometric points $\text{Spec}(\Omega) \rightarrow X$ lying over $\bar{y}$, which is finite since f is finite. The product runs over the (finite) set of such points.

2. Suppose that $\mathcal{F} = \underline{\Lambda}$ is a constant sheaf on X where Λ is a set, an abelian group, or a module over a ring. For any sufficiently small étale neighborhood $V \rightarrow Y$ of $\bar{y}$ such that $X \times_Y V$ decomposes into a disjoint union of connected components $X \times_Y V = \coprod_{i=1}^n Z_i$ (Lemma 6.1.25), the fiber $f^{-1}(\bar{y})$ consists of one geometric point $\bar{x}_i$ for each component Z_i . Note that we have isomorphisms

$$\Gamma(X \times_Y V, \underline{\Lambda}) \cong \prod_{i=1}^n \Gamma(Z_i, \underline{\Lambda}|_{Z_i}) \cong \prod_{i=1}^n \Lambda.$$

We then have natural isomorphism

$$\Gamma(X \times_Y V, \underline{\Lambda}) \cong \prod_{i=1}^n \Lambda \xrightarrow{\sim} \prod_{\bar{x} \in f^{-1}(\bar{y})} \Lambda \cong \prod_{\bar{x} \in f^{-1}(\bar{y})} \mathcal{F}_{\bar{x}} \cong (f_*\mathcal{F})_{\bar{y}},$$

where the i -th factor of the first product maps as the identity to the factor corresponding to the point $\bar{x}_i$ contained in the component Z_i , and maps as zero to the other factors.

Proof. Let $f : X \rightarrow Y$ be a finite morphism (Definition 1.2.1). Since f is finite, it is affine (Definition 3.4), and thus for any étale neighborhood $(V, \bar{v}) \rightarrow (Y, \bar{y})$, the fiber product

(Definition C.0.15) $X_V := X \times_Y V$ is an affine V -scheme (Definition .3.4), and the projection $f_V : X_V \rightarrow V$ is finite.

By Lemma 6.1.25, there exists a cofinal system (Definition .1.20) of étale neighborhoods $(V, \bar{v})$ such that the fiber product decomposes into a finite disjoint union of connected components:

$$X \times_Y V = \coprod_{i=1}^n Z_i$$

where each Z_i is an affine scheme containing exactly one geometric point $\bar{x}_i$ mapping to $\bar{v}$ (which sits over $\bar{y}$).

The stalk $(f_*\mathcal{F})_{\bar{y}}$ is defined as the filtered colimit over the category of étale neighborhoods of $\bar{y}$:

$$(f_*\mathcal{F})_{\bar{y}} = \varinjlim_{(V, \bar{v})} \Gamma(X \times_Y V, \mathcal{F}).$$

Restricting to the cofinal subcategory of neighborhoods where $X \times_Y V$ decomposes as above, and using the fact that the section functor Γ converts disjoint unions into products, we have:

(♠ TODO: this fact)

$$(f_*\mathcal{F})_{\bar{y}} \cong \varinjlim_{(V, \bar{v})} \prod_{i=1}^n \Gamma(Z_i, \mathcal{F}|_{Z_i}).$$

(♠ TODO: filtered colimit commutes with finite products) Since the filtered colimit commutes with finite products, we obtain:

$$(f_*\mathcal{F})_{\bar{y}} \cong \prod_{i=1}^n \left(\varinjlim_{(V, \bar{v})} \Gamma(Z_i, \mathcal{F}|_{Z_i}) \right).$$

For each i , as $(V, \bar{v})$ varies over the cofinal system, Z_i runs through a cofinal system of étale neighborhoods of $\bar{x}_i$. By the definition of the stalk at a geometric point, the term in the parentheses is precisely $\mathcal{F}_{\bar{x}_i}$. Thus:

$$(f_*\mathcal{F})_{\bar{y}} \cong \prod_{\bar{x} \in f^{-1}(\bar{y})} \mathcal{F}_{\bar{x}}.$$

Finally, in the case of a constant sheaf $\mathcal{F} = \underline{\Lambda}$, we have $\mathcal{F}_{\bar{x}} \cong \Lambda$ for each point. The isomorphism $\Gamma(X \times_Y V, \underline{\Lambda}) \cong \prod_{i=1}^n \Lambda$ arises because $\underline{\Lambda}(Z_i) \cong \Lambda$ for each connected component Z_i , and the natural map to the product is given by restricting the constant section to each connected component. $\square$

Proposition 6.1.29 (Pushforward of constant sheaf along finite étale G -cover has group ring structure). (♠ TODO: AI generated, verify) Let Λ be a ring. Let $f : X \rightarrow Y$ be a finite étale Galois cover (Definition 1.2.4) with group G . Let $\underline{\Lambda}$ be the constant sheaf (Definition C.0.10) on $X_{\text{ét}}$ (Definition 4.1.3) with value Λ .

(♠ TODO: write about how pushing forward a locally constant sheaf under a finite étale morphism yields a locally constant sheaf)

1. The action of G on X over Y induces an action of G on the direct image sheaf $f_*\underline{\Lambda}$, making it a locally constant sheaf (Definition 4.4.2) of Λ -algebras on the étale site (Definition 4.1.3) $Y_{\text{ét}}$, equipped with a G -action by Λ -algebra automorphisms.
2. Let $\bar{y} : \text{Spec}(\Omega) \rightarrow Y$ be a geometric point. The geometric stalk of the direct image is canonically isomorphic to regular representation of G over Λ :

$$(f_*\underline{\Lambda})_{\bar{y}} \cong \Lambda[G]$$

where G acts on $\Lambda[G]$ by left translation. This isomorphism is induced by taking the isomorphism

$$(f_*\underline{\Lambda})_{\bar{y}} \cong \prod_{\bar{x} \in f^{-1}(\bar{y})} \underline{\Lambda}_{\bar{x}} \cong \prod_{\bar{x} \in f^{-1}(\bar{y})} \Lambda$$

(Proposition 6.1.28), picking any $\bar{x}_0 \in f^{-1}(\bar{y})$, and defining the map $\Phi_{x_0} : \prod_{\bar{x} \in f^{-1}(\bar{y})} \Lambda \rightarrow \Lambda[G]$ via the identification of the copy of Λ corresponding to the point $g \cdot \bar{x}_0$ with basis element $g \in \Lambda[G]$.³

3. The invariant subsheaf recovers the constant sheaf on the base:

$$(f_*\underline{\Lambda})^G = \underline{\Lambda}_Y$$

where $\underline{\Lambda}_Y$ is the constant sheaf on $Y_{\text{ét}}$ with value Λ .

Proof. 1.

2. By Proposition 6.1.28, for any sufficiently small étale neighborhood $V \rightarrow Y$ of $\bar{y}$ such that $X \times_Y V$ decomposes into a disjoint union of connected components $X \times_Y V = \coprod_{i=1}^n Z_i$ (Lemma 6.1.25), the fiber $f^{-1}(\bar{y})$ consists of one geometric point $\bar{x}_i$ for each component Z_i and we have the natural isomorphism

$$\Gamma(X \times_Y V, \underline{\Lambda}) \cong \prod_{i=1}^n \Lambda \xrightarrow{\sim} \prod_{\bar{x} \in f^{-1}(\bar{y})} \Lambda \cong (f_*\mathcal{F})_{\bar{y}},$$

where the i -th factor of the first product maps as the identity to the factor corresponding to the point $\bar{x}_i$ contained in the component Z_i , and maps as zero to the other factors. Letting $V = X$, we have that $X \times_Y X \cong \coprod_{g \in G} X$, and the components of this decomposition correspond to the geometric points in $f^{-1}(\bar{y})$ (♠ **TODO: reference**). The action of G on the components of $\coprod_{g \in G} X$ then corresponds to the action of G on $f^{-1}(\bar{y})$. Since the factors of $\prod_{i=1}^n \Lambda$ correspond to the components of $\coprod_{g \in G} X$, it has a natural action by G tantamount to the regular representation. In particular, we have isomorphisms the composition

$$(f_*\mathcal{F})_{\bar{y}} \rightarrow \prod_{\bar{x} \in f^{-1}(\bar{y})} \Lambda \rightarrow \prod_{i=1}^n \Lambda \rightarrow \Lambda[G]$$

and the composition of the latter two is tantamount to the described map Φ_{x_0} .

³While this identification depends on the choice of $\bar{x}_0$, any two choices differ by an element of G and the resulting map remains the same up to the G -action, rendering the isomorphism of G -modules canonical; $f^{-1}(y)$ is a G -torsor and the structure of G as a G -torsor determines the structure of $\Lambda[G]$ as a G -module.

3.

(♠ TODO: AI generated, verify) (a) Since G acts on X over Y , for each $g \in G$ the automorphism $g : X \rightarrow X$ induces a pullback $(g^{-1})^*$ on sheaves on $X_{\text{ét}}$. Because $\underline{\Lambda}$ is constant, $(g^{-1})^*\underline{\Lambda} = \underline{\Lambda}$, so there are natural maps

$$f_*\underline{\Lambda} \xrightarrow{(f \circ g)_*\underline{\Lambda}} f_*\underline{\Lambda}$$

for each $g \in G$, giving a well-defined action of G on $f_*\underline{\Lambda}$ by sheaf automorphisms. Since $\underline{\Lambda}$ is a sheaf of Λ -algebras and the maps are induced by pullbacks along automorphisms, each g acts by Λ -algebra automorphisms. The sheaf $f_*\underline{\Lambda}$ is locally constant because after pulling back to Y via f , it becomes trivial: $f^*f_*\underline{\Lambda} \cong \underline{\Lambda}[G]$ as a trivial local system.

(b) By the proper base change theorem for finite morphisms, the geometric fiber $(f_*\underline{\Lambda})_{\bar{y}}$ is canonically isomorphic to the global sections of $\underline{\Lambda}$ on the fiber $X_{\bar{y}} = X \times_Y \text{Spec}(\Omega)$. Since f is a finite étale Galois cover with group G , the fiber $X_{\bar{y}}$ consists of $|G|$ points permuted simply transitively by G . Fixing a geometric point $\bar{x}$ in the fiber identifies $X_{\bar{y}}$ with G as a right G -set. The sections of the constant sheaf $\underline{\Lambda}$ on this finite discrete scheme are the functions $G \rightarrow \Lambda$, which is canonically $\Lambda[G]$. The left G -action on the direct image corresponds to left translation on G , hence on $\Lambda[G]$.

(c) For any étale $V \rightarrow Y$, the sections $(f_*\underline{\Lambda})(V) = \underline{\Lambda}(X \times_Y V)$ are the locally constant functions $X \times_Y V \rightarrow \Lambda$. The G -invariant sections are precisely those functions constant on G -orbits, which descend to functions on $(X \times_Y V)/G = V$ (since f is the quotient map $X \rightarrow X/G = Y$). Hence $(f_*\underline{\Lambda})^G(V) = \underline{\Lambda}_Y(V)$ naturally in V . $\square$

Proposition 6.1.30 (Stalk of direct image under Galois morphism for étale sheaves). (♠ TODO: AI generated, verify) Let $f : X \rightarrow Y$ be a Galois morphism of schemes (Definition 1.2.4) with Galois group G , and let $\mathcal{F}$ be a sheaf on the étale site $X_{\text{ét}}$ (Definition 4.1.3). Let $\bar{y} : \text{Spec}(\Omega) \rightarrow Y$ be a geometric point of Y , and fix a geometric point $\bar{x}$ in the fiber $f^{-1}(\bar{y})$.

The group G acts on the stalk $\mathcal{F}_{\bar{x}}$ (up to conjugacy, depending on the choice of $\bar{x}$). The stalk of the direct image is the invariant part:

$$(f_*\mathcal{F})_{\bar{y}} = \mathcal{F}_{\bar{x}}^G$$

More precisely, if f is also finite (so that $X \times_Y \text{Spec}(\Omega) \cong \coprod_{g \in G} \text{Spec}(\Omega)$), then the decomposition of Proposition 6.1.28 identifies $(f_*\mathcal{F})_{\bar{y}}$ with $\prod_{g \in G} \mathcal{F}_{g\bar{x}}$, and the G -action permutes the factors transitively, yielding the identification with $\mathcal{F}_{\bar{x}}^G$.

Proof. (♠ TODO: AI generated, verify) Since f is Galois with group G , the fiber over $\bar{y}$ consists of $|G|$ geometric points permuted simply transitively by G . By Proposition 6.1.28, the stalk decomposes as a product over these points. The G -action on X/Y induces an action on each factor, and taking invariants under this permutation recovers the fixed-point subobject of a single stalk. $\square$

Proposition 6.1.31 (Stalk of direct image under closed immersion for étale sheaves). (♠ TODO: AI generated, verify) Let $i : Z \hookrightarrow X$ be a closed immersion (Definition C.0.14) of

schemes, and let $\mathcal{F}$ be a sheaf on the étale site $Z_{\text{ét}}$ (Definition 4.1.3). Let $\bar{x}$ be a geometric point. (♠ TODO: the statement needs to specify the value of the sheaf; in particular, the notion of 0 does not make sense unless the sheaf is valued in abelian groups.)

1. If the image of $\bar{x}$ lies in Z , then

$$(i_*\mathcal{F})_{\bar{x}} = \mathcal{F}_{\bar{x}}$$

2. If the image of $\bar{x}$ lies in $X \setminus Z$, then

$$(i_*\mathcal{F})_{\bar{x}} = 0$$

Proof. This is a consequence of Proposition 6.1.28, as closed immersions are finite morphisms. □

Proposition 6.1.32 (Stalk of direct image under open immersion for étale sheaves). (♠ TODO: AI generated, verify) Let $j : U \hookrightarrow X$ be an open immersion (Definition C.0.14) of schemes, and let $\mathcal{F}$ be a sheaf on the étale site $U_{\text{ét}}$ (Definition 4.1.3). Let $\bar{x}$ be a geometric point.

1. If the image of $\bar{x}$ lies in U , then

$$(j_*\mathcal{F})_{\bar{x}} = \mathcal{F}_{\bar{x}}$$

2. If the image of $\bar{x}$ lies in $X \setminus U$, then

$$(j_*\mathcal{F})_{\bar{x}} = \varinjlim_{(V, \bar{v})} \mathcal{F}(V \times_X U),$$

where the colimit runs over étale neighborhoods $(V, \bar{v})$ of $\bar{x}$ in X . The fiber product $V \times_X U$ is an open subscheme of V , and may be empty for small enough V if $\bar{x}$ maps to the boundary.

Proof. (♠ TODO: AI generated, verify) If $\bar{x}$ lands in U , then every étale neighborhood $(V, \bar{v})$ of $\bar{x}$ has $V \times_X U$ as an open-neighborhood of the geometric point induced by $\bar{x}$ in U . The two directed systems are cofinal.

If $\bar{x}$ lands outside U , the formula follows directly from $(j_*\mathcal{F})(V) = \mathcal{F}(V \times_X U)$ and taking the colimit over étale neighborhoods. □

6.1.5. Exactness properties on sheaves of abelian groups.

Theorem 6.1.33 (Exactness of stalk functors on étale sheaves of abelian groups). Let X be a scheme (Definition 3.1). For any geometric point (Definition 3.0.1) $\bar{x} : \text{Spec}(\Omega) \rightarrow X$, the stalk (Definition 4.5.1) functor

$$\mathcal{F} \mapsto \mathcal{F}_{\bar{x}}$$

is an exact functor (Definition 4.1.36) from the category of sheaves (Definition 4.0.2) of abelian groups on $X_{\text{ét}}$ (Definition 4.1.3) (Proposition B.9.10) to the category of abelian groups.

That is, for every short exact sequence of sheaves of abelian groups

$$0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0,$$

the induced sequence of stalks at $\bar{x}$

$$0 \rightarrow \mathcal{F}'_{\bar{x}} \rightarrow \mathcal{F}_{\bar{x}} \rightarrow \mathcal{F}''_{\bar{x}} \rightarrow 0$$

is exact.

Proposition 6.1.34 (Exactness of inverse image functors). Let $f : Y \rightarrow X$ be a morphism of schemes. Then the inverse image functor (Definition 6.1.20)

$$f^{-1} : \mathrm{Sh}(X_{\text{ét}}, \mathrm{Ab}) \rightarrow \mathrm{Sh}(Y_{\text{ét}}, \mathrm{Ab})$$

on sheaves of abelian groups on the small étale sites is exact (Definition .1.36).

Proposition 6.1.35 (Left exactness and exactness criteria for direct image functors). Let $f : Y \rightarrow X$ be a morphism of schemes. The direct image functor (Definition 6.1.20)

$$f_* : \mathrm{Sh}(Y_{\text{ét}}, \mathrm{Ab}) \rightarrow \mathrm{Sh}(X_{\text{ét}}, \mathrm{Ab})$$

is left exact (Definition .1.36) in general.

(♠ TODO: pin down precise conditions under which f_* is exact.) Moreover, if f is an étale morphism, or more generally if f is a morphism satisfying certain finiteness and flatness conditions (e.g., finite étale), then f_* is exact. In particular, if f is an open immersion (Definition C.0.14) or étale morphism (Definition 1.1.6), then f_* is exact.

6.2. The functors on derived categories of sheaves. (♠ TODO: Take the derived functors here and incorporate them in the definition of things for $D_c^b(X, R)$.)

Definition 6.2.1 (Torsion sheaf of abelian groups). (♠ TODO: must the site have a small topological generating family?) Let $(\mathcal{C}, J)$ be a site (Definition A.0.4) with a small topological generating family (Definition A.0.4) (or a U -small topologically generating family if a universe (Definition .1.2) U is available). A sheaf (Definition 7.1.10) of abelian groups $\mathcal{F}$ on $(\mathcal{C}, J)$ is called a *torsion sheaf of abelian groups* if for every object U in $\mathcal{C}$ and every section $s \in \mathcal{F}(U)$, there exists a nonzero integer n such that $ns = 0$ in $\mathcal{F}(U)$.

Notation 6.2.2. Let X/S be a scheme over a base scheme S , and choose some small site (Definition 4.1.1) X_{site} on X associated to a big site on (Sch/S) . Let $\mathcal{O}$ be a sheaf of rings on X_{site} . It is standard to write the following:

- $D(X_{\text{site}}, \mathcal{O})$, or simply $D(X, \mathcal{O})$ if the small site X_{site} is understood, for the derived category (Definition B.3.1) of the category of sheaves of $\mathcal{O}$ -modules (Definition 7.1.3).
- $D_{\text{tor}}(X_{\text{site}}, \mathcal{O})$ or $D_{\text{tors}}(X_{\text{site}}, \mathcal{O})$ for the full subcategory of $D(X)$ of complexes whose cohomology objects (Definition .2.3) are torsion sheaves (Definition 6.2.1) (♠ TODO: continue)
- In the case that $\mathcal{O} = \mathbb{Z}$ (Definition C.0.10), i.e. for derived categories of categories of sheaves of abelian groups, one may simply write $D(X_{\text{site}})$, $D(X)$, etc.

Superscripts of $+$, $-$, b may also be used accordingly — for example, for $? \in \{+, -, b\}$, it makes sense to write $D^?(X_{\text{site}}, \mathcal{O})$, see Definition B.3.1. Compare with Notation B.9.1.

(♠ TODO: morphism of schemes induce morphisms of sites)

Definition 6.2.3. e.g. see [?, End of 6.3] Let $f : X \rightarrow Y$ be a morphism of schemes (Definition .3.2). Let (♠ TODO: need to choose sites)

The (right total) derived pushforward/direct image functor of f is the right derived functor

$$Rf_* : D(X) \rightarrow D(Y)$$

(Notation 6.2.2)

The functor Rf_*

(♠ TODO: define $Rf_* : D^+(X) \rightarrow D^+(Y)$, f^*)

(♠ TODO: state adjunction)

Definition 6.2.4. Let S be a scheme (Definition .3.1) and let $f : X \rightarrow S$ be a morphism of schemes (Definition .3.2). A compactification of f is a commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & S \\ & \searrow j & \nearrow \bar{f} \\ & \bar{X} & \end{array}$$

in which $j : X \hookrightarrow \bar{X}$ is an open immersion (Definition C.0.14) and $\bar{f} : \bar{X} \rightarrow S$ is a proper morphism.

A morphism $f : X \rightarrow S$ is said to be compactifiable over S or S -compactifiable if it admits a compactification. For example, if S is quasi-compact and quasi-separated and f is separated and of finite type (Definition C.0.19), then f is compactifiable by Nagata's compactification theorem (Theorem 6.2.5).

Theorem 6.2.5 (Nagata's compactification theorem, cf. [?, Tag 0F41]). Let S be a quasi-compact and quasi-separated scheme and let $f : X \rightarrow S$ be a separated and of finite type (Definition C.0.19) morphism of schemes. Then, f admits a compactification (Definition 6.2.4)

$$\begin{array}{ccc} X & \xrightarrow{f} & S \\ & \searrow j & \nearrow \bar{f} \\ & \bar{X} & \end{array} .$$

Definition 6.2.6. Let S be a scheme (Definition .3.1) and let $f : X \rightarrow S$ be a morphism of schemes (Definition .3.2). Let two compactifications (Definition 6.2.4)

$$\begin{array}{ccc} X & \xrightarrow{f} & S \\ & \searrow j & \nearrow \bar{f} \\ & \bar{X} & \end{array} \quad \text{and} \quad \begin{array}{ccc} X & \xrightarrow{f} & S \\ & \searrow k & \nearrow \bar{g} \\ & \bar{Y} & \end{array}$$

of f be given. We say that the second compactification dominates the first if there exists a morphism $\phi : \bar{Y} \rightarrow \bar{X}$ of schemes over S such that $\phi \circ k = j$.

For any two compactification of a mapping f , there is always a third that dominates them both (♠ TODO: justify)

Definition 6.2.7. e.g. see (♠ TODO:) Let $f : X \rightarrow Y$ be an S -compactifiable morphism of schemes. (♠ TODO: define $Rf_! : D^+(X) \rightarrow D^+(Y)$, $f^!$)

(♠ TODO: state adjunction)

Definition 6.2.8. e.g. see (♠ TODO:) Let X/S be a scheme over a base scheme. (♠ TODO: define Ext , sheaf Ext , Tor)

(♠ TODO: state adjunction)

6.3. Verdier duality for bounded constructible complexes. The Verdier dual functor provides the duality needed to define self-dual (middle) perverse t -structures. We record the dualizing complex and Verdier dual for each coefficient system.

6.4. Torsion coefficients.

Definition 6.4.1 (Dualizing complex for Verdier duality on a noetherian scheme with torsion coefficients). Let S be a noetherian scheme (Definition C.0.16). Let A be a noetherian torsion ring. Consider sheaves (Definition 4.0.2) of A -modules (Definition C.0.11) on the (small) étale site (Definition 4.1.3) of S .

1. The *dualizing complex for Verdier duality on S with coefficients in A* , denoted by $\mathbb{D}_S^\bullet$, is the constant sheaf A_S concentrated in degree 0.
2. Let X be a finite type scheme over S . The *(relative) dualizing complex for Verdier duality on X/S with coefficients in A* , denoted by $\mathbb{D}_{X/S}^\bullet$, is defined as the object in $D^+(X_{\text{ét}}, A)$ (Notation 10.0.7) defined as

$$\mathbb{D}_{X/S}^\bullet = Rf^! \mathbb{D}_S^\bullet$$

Definition 6.4.2 (Verdier dual of a constructible complex with integral coefficients). Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let S be a noetherian scheme (Definition C.0.16). Let X be an S -scheme of finite type (Definition C.0.19). Consider sheaves on the (small) étale sites (Definition 4.1.3) of X and S .

For any complex $M \in D_c^b(X, R)$ (Definition 8.6.2), the *Verdier dual of M on X relative to S with integral coefficients* is the object of $D_c^b(X, R)$ defined by

$$\mathbb{D}_X(M) := R\mathcal{H}om_R(M, \mathbb{D}_{X/S}^\bullet(R))$$

6.5. Integral adic coefficients.

Definition 6.5.1 (Dualizing complex for Verdier duality on a noetherian scheme with integral coefficients). Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let S be a noetherian scheme (Definition C.0.16). Consider sheaves on the (small) étale site of S .

1. The *dualizing complex for Verdier duality on S with integral coefficients*, denoted by $\mathbb{D}_S^\bullet(R)$, is the object of $D_c^b(X, R)$ (Definition 8.6.2) given as the constant inverse system $(R/\lambda^n)_n$ where each R/λ^n is regarded as an object of $D_c^b(X, R/\lambda^n)$ concentrated in degree 0.
2. The *(relative) dualizing complex for Verdier duality on X/S with integral coefficients*, denoted by $\mathbb{D}_{X/S}^\bullet(R)$, is defined by

$$\mathbb{D}_{X/S}^\bullet(R) = Rf^! \mathbb{D}_S^\bullet(R)$$

Definition 6.5.2 (Verdier dual of a constructible complex with integral coefficients). Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let S be a noetherian scheme (Definition C.0.16). Let X be an S -scheme of finite type (Definition C.0.19). Consider sheaves on the (small) étale sites (Definition 4.1.3) of X and S .

For any complex $M \in D_c^b(X, R)$ (Definition 8.6.2), the *Verdier dual of M on X relative to S with integral coefficients* is the object of $D_c^b(X, R)$ defined by

$$\mathbb{D}_X(M) := R\mathcal{H}om_R(M, \mathbb{D}_{X/S}^\bullet(R))$$

6.6. Rational coefficients.

Definition 6.6.1 (Dualizing complex for Verdier duality on a noetherian scheme with rational coefficients). Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let E be the fraction field of R . Let S be a noetherian scheme (Definition C.0.16). Consider sheaves on the (small) étale site of S .

1. The *dualizing complex for Verdier duality on S with rational coefficients*, denoted by $\mathbb{D}_S^\bullet(E)$, is the object of $D_c^b(S, E)$ (Definition 8.6.10) given as the dualizing complex (Definition 6.5.1) $\mathbb{D}_S^\bullet(R)$ in $D_c^b(S, R)$ (Definition 8.6.2).
2. The *(relative) dualizing complex for Verdier duality on X/S with rational coefficients*, denoted by $\mathbb{D}_{X/S}^\bullet(E)$, is the object of $D_c^b(X, E)$ defined by

$$\mathbb{D}_{X/S}^\bullet(E) = Rf^! \mathbb{D}_S^\bullet(E).$$

Equivalently, as an object of $D_c^b(X, R)$, it is given as the dualizing complex (Definition 6.5.1) $\mathbb{D}_{X/S}^\bullet(R)$

Definition 6.6.2 (Verdier dual of a constructible complex with rational coefficients). Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let E be the fraction field of R . Let S be a noetherian scheme (Definition C.0.16). Let X be an S -scheme of finite type (Definition C.0.19). Consider sheaves on the (small) étale sites of X and S .

For any object $M \in D_c^b(X, E)$, the *Verdier dual of M on X relative to S with rational coefficients* is the object in $D_c^b(X, E)$ given by

$$\mathbb{D}_X(M) := R\mathcal{H}om_E(M, \mathbb{D}_{X/S}^\bullet(E)).$$

Equivalently, its underlying object in $D_c^b(X, R)$ (Definition 8.6.2) is the Verdier dual (Definition 6.5.2) $\mathbb{D}_X(M_0)$ where $M_0 \in D_c^b(X, R)$ is the underlying object of M .

6.7. Algebraic coefficients.

Definition 6.7.1 (Dualizing complex for Verdier duality on a noetherian scheme with algebraic coefficients). Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let E be the fraction field of R , and let K/E be a algebraic extension (Definition 8.6.9). Let S be a noetherian scheme (Definition C.0.16). Consider sheaves on the (small) étale site of S .

1. The *dualizing complex for Verdier duality on S with algebraic coefficients*, denoted by $\mathbb{D}_S^\bullet(K)$, is the image of $\mathbb{D}_S^\bullet(E)$ in $D_c^b(S, K)$ under the transition functor $D_c^b(S, E) \rightarrow D_c^b(S, K)$.
2. Let X be a finite type scheme over S . The *(relative) dualizing complex for Verdier duality on X/S with algebraic coefficients*, denoted by $\mathbb{D}_{X/S}^\bullet(K)$, is defined by

$$\mathbb{D}_{X/S}^\bullet(K) = Rf^! \mathbb{D}_S^\bullet(K).$$

Equivalently, it is the image of the dualizing complex (Definition 6.6.1) $\mathbb{D}_{X/S}^\bullet(E)$ under the transition functor $D_c^b(S, E) \rightarrow D_c^b(S, K)$.

Definition 6.7.2 (Verdier dual of a constructible complex with algebraic coefficients). Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let E be the fraction field of R , and let K/E be a algebraic extension (Definition 8.6.9). Let S be a noetherian scheme (Definition C.0.16). Let X be an S -scheme of finite type (Definition C.0.19). Consider sheaves on the (small) étale sites of X and S .

For any object $M \in D_c^b(X, K)$, the *Verdier dual of M on X relative to S with algebraic coefficients* is

$$\mathbb{D}_X(M) := R\mathcal{H}om_K(M, \mathbb{D}_{X/S}^\bullet(K)).$$

(Definition 6.7.1)

6.8. Various coefficient systems.

Definition 6.8.1 (Dualizing complex for Verdier duality on a noetherian scheme for various coefficient systems). (♠ TODO: AI generated, verify) Let S be a noetherian scheme (Definition C.0.16). Consider sheaves on the (small) étale site (Definition 4.1.3) of S .

Let the coefficient system be one of the following:

1. A noetherian torsion ring (torsion coefficients)
2. Assuming that S is noetherian (Definition C.0.16) and with R, E, K as in ??

In each case, the *dualizing complex for Verdier duality on S* , denoted by $\mathbb{D}_S^\bullet$ (notation suppresses coefficient dependence), is the object in the corresponding category characterized by the following property: for every S -scheme X of finite type (Definition C.0.19) with structure morphism $f : X \rightarrow S$, the relative dualizing complex satisfies

$$\mathbb{D}_{X/S}^\bullet = Rf^! \mathbb{D}_S^\bullet$$

where $Rf^!$ is the exceptional inverse image functor appropriate to the coefficient system.

See Definition 6.4.1 for torsion coefficients, Definition 6.5.1 for integral adic coefficients, Definition 6.6.1 for rational coefficients, and Definition 6.7.1 for algebraic coefficients.

Definition 6.8.2 (Verdier dual of a constructible complex for various coefficient systems). (♠ TODO: AI generated, verify) Let S be a noetherian scheme (Definition C.0.16). Let X be an S -scheme of finite type (Definition C.0.19). Consider sheaves on the (small) étale sites (Definition 4.1.3) of X and S .

Let the coefficient system be one of the following:

1. A a noetherian torsion ring (torsion coefficients)
2. Assuming that S is noetherian (Definition C.0.16) and with R, E, K as in ??
 - (a) (R, λ) (integral adic coefficients)
 - (b) $E = \text{Frac}(R)$ (rational coefficients)
 - (c) K (algebraic coefficients)

For any complex $\mathcal{K}$ in the corresponding category on X , the *Verdier dual of $\mathcal{K}$ on X relative to S* is

$$\mathbb{D}_X(\mathcal{K}) := R\mathcal{H}om(\mathcal{K}, \mathbb{D}_{X/S}^\bullet)$$

where $\mathbb{D}_{X/S}^\bullet$ is the dualizing complex on X relative to S for the given coefficient system, and $R\mathcal{H}om$ is the derived internal Hom functor appropriate to the coefficient system (Definition B.9.3).

See Definition 6.5.2 for torsion coefficients, Definition 6.5.2 for integral adic coefficients, Definition 6.6.2 for rational coefficients, and Definition 6.7.2 for algebraic coefficients.

7. COHOMOLOGY OF LITERAL SHEAVES FOR THE SMALL ÉTALE SITE ON A SCHEME

In this section, we define the étale cohomology of a *literal* sheaf (Definition 4.0.2) of abelian groups; we emphasize that the sheaves here as *literal* because λ -adic sheaves (Definition 8.4.9) as defined later are not actual sheaves but rather projective systems of them.

7.1. Sheaf cohomology of a general module over a sheaf of rings on a site.

Definition 7.1.1. Let $\mathcal{C}$ be a (large) category (Definition .1.1), and let $\mathcal{D}$ be a (large) category (Definition .1.1). Let $\mathcal{F} : \mathcal{C}^{op} \rightarrow \mathcal{D}$ be a presheaf valued in $\mathcal{D}$ (Definition 4.0.1).

1. For an object $U \in \mathcal{C}$, the *sections functor evaluated at U* is the functor

$$\Gamma(U, -) : \text{PSh}(\mathcal{C}, \mathcal{D}) \rightarrow \mathcal{D}$$

defined by

$$\Gamma(U, \mathcal{F}) := \mathcal{F}(U),$$

i.e., the value of the presheaf $\mathcal{F}$ at the object U .

2. The *global sections of $\mathcal{F}$* is the object $\Gamma(\mathcal{F})$ of $\mathcal{D}$ defined as the limit (Definition .1.13)

$$\Gamma(\mathcal{F}) = \varprojlim_{U \in \mathcal{C}^{\text{op}}} \mathcal{F}(U)$$

assuming that such a limit exists, where the limit is taken over objects $U \in \mathcal{C}$ and the restriction morphisms $\mathcal{F}(V) \rightarrow \mathcal{F}(U)$ in $\mathcal{D}$ for morphisms $U \rightarrow V$ in $\mathcal{C}$.

If a final object (Definition .1.10) $*$ $\in \mathcal{C}$ exists, then $\Gamma(\mathcal{F})$ exists and coincides with $\Gamma(*, \mathcal{F}) = \mathcal{F}(*)$. The construction $\Gamma(\mathcal{F})$ is functorial; in particular, if $\Gamma(\mathcal{F})$ exists for all $\mathcal{F}$ in $\text{PSh}(\mathcal{C}, \mathcal{D})$, e.g. if limits of (Definition .1.13) diagrams in $\mathcal{D}$ indexed by $\mathcal{C}$ exist, then Γ is a functor

$$\Gamma : \text{PSh}(\mathcal{C}, \mathcal{D}) \rightarrow \mathcal{D}$$

called the *global sections functor on $\text{PSh}(\mathcal{C}, \mathcal{D})$* .

Proposition 7.1.2. Let $\mathcal{C}$ be an essentially small category (Definition .1.26) and let $\mathcal{A}$ be an abelian category (Definition .1.34).

1. Let $U \in \text{Ob}(\mathcal{C})$ be some fixed object. The sections functor (Definition 7.1.1)

$$\Gamma(U, -) : \text{PSh}(\mathcal{C}, \mathcal{A}) \rightarrow \mathcal{A}$$

is left exact (Definition .1.36). (♠ TODO: state that presheaves and sheaves valued in an abelian category form abelian categories)

2. Assume that $\Gamma(\mathcal{F})$ exists for all $\mathcal{F}$ in $\text{PSh}(\mathcal{C}, \mathcal{A})$ so that Γ is a functor

$$\Gamma : \text{PSh}(\mathcal{C}, \mathcal{A}) \rightarrow \mathcal{A}.$$

The functor Γ is left exact.

Proof. Recall that $\text{PSh}(\mathcal{C}, \mathcal{D})$ is an abelian category (Proposition B.9.10) since $\mathcal{C}$ is essentially small. (♠ TODO: Talk about how limits are left exact and) $\square$

Definition 7.1.3. 1. Let $\mathcal{C}$ be a site, and let $\mathcal{A}$ and $\mathcal{B}$ be sheaves (Definition 4.0.2) of (not necessarily commutative) rings (Definition C.0.8) on $\mathcal{C}$.

- (a) An $(\mathcal{A}, \mathcal{B})$ -bimodule (or a bimodule over $(\mathcal{A}, \mathcal{B})$) is a sheaf (Definition 4.0.2) $\mathcal{M}$ of abelian groups on $\mathcal{C}$ equipped with a left $\mathcal{A}$ -module structure given by a morphism of sheaves (Definition 4.0.2) of sets

$$\lambda : \mathcal{A} \times \mathcal{M} \longrightarrow \mathcal{M},$$

and a right $\mathcal{B}$ -module structure given by a morphism of sheaves of sets

$$\rho : \mathcal{M} \times \mathcal{B} \longrightarrow \mathcal{M},$$

such that the actions are compatible. Specifically, for every object U in $\mathcal{C}$, every section $m \in \mathcal{M}(U)$, every $a \in \mathcal{A}(U)$, and every $b \in \mathcal{B}(U)$, the equality

$$\lambda_U(a, \rho_U(m, b)) = \rho_U(\lambda_U(a, m), b)$$

holds in $\mathcal{M}(U)$. In standard multiplicative notation where $\lambda(a, m)$ is denoted $a \cdot m$ and $\rho(m, b)$ is denoted $m \cdot b$, this condition is the associativity axiom

$$(a \cdot m) \cdot b = a \cdot (m \cdot b).$$

In particular, for every object $U \in \mathcal{C}$, the abelian group $\mathcal{M}(U)$ has the structure of an $\mathcal{A}(U) - \mathcal{B}(U)$ -bimodule (Definition C.0.11).

- (b) Assume that a sheafification functor (Definition 4.3.1)

$$\mathrm{PreShv}(\mathcal{C}, \mathbf{Rings}) \rightarrow \mathrm{Shv}(\mathcal{C}, \mathbf{Rings})$$

exists. For rings A and B , a A - B -bimodule or bimodule over (A, B) on $\mathcal{C}$ is a bimodule over $(\underline{A}, \underline{B})$ where $\underline{A}$ and $\underline{B}$ are the constant sheaves (Definition C.0.10) on $\mathcal{C}$ with values A and B respectively.

- (c) Let $\mathcal{M}$ and $\mathcal{N}$ be $(\mathcal{A}, \mathcal{B})$ -bimodules. A $homomorphism of (\mathcal{A}, \mathcal{B})$ -bimodules (or an $(\mathcal{A}, \mathcal{B})$ -linear morphism) is a morphism of sheaves of abelian groups $f : \mathcal{M} \rightarrow \mathcal{N}$ such that for every object U of $\mathcal{C}$, every section $m \in \mathcal{M}(U)$, every $a \in \mathcal{A}(U)$, and every $b \in \mathcal{B}(U)$, the following compatibility conditions hold:

$$f_U(a \cdot m) = a \cdot f_U(m) \quad \text{and} \quad f_U(m \cdot b) = f_U(m) \cdot b.$$

We denote the category of $(\mathcal{A}, \mathcal{B})$ -bimodules, with morphisms being morphisms of sheaves of abelian groups compatible with both the left $\mathcal{A}$ -action and the right $\mathcal{B}$ -action, by $\mathcal{A}\text{-}\mathcal{B}\text{-Mod}$ or sometimes by ${}_A\mathrm{Mod}_B$ (♠ TODO: talk about how bimodules can be identifies with left/right modules)

2. Let $(\mathcal{C}, J)$ be a site (Definition A.0.4). Let $\mathcal{O}$ be a sheaf of (not necessarily commutative) rings on $(\mathcal{C}, J)$ (Definition 4.0.2), i.e. $((\mathcal{C}, J), \mathcal{O})$ is a ringed site (Definition 6.1.5).

- (a) An $(left/right/two-sided) \mathcal{O}$ -module consists of the following data:

- A sheaf $\mathcal{F}$ of abelian groups on $(\mathcal{C}, J)$,
- for every object $U \in \mathcal{C}$, the structure of an $(left/right/two-sided) \mathcal{O}(U)$ -module on $\mathcal{F}(U)$,

such that for every morphism $f : V \rightarrow U$ in $\mathcal{C}$, the restriction map

$$\rho_{U,V} : \mathcal{F}(U) \rightarrow \mathcal{F}(V)$$

is $\mathcal{O}(U)$ -linear when the $\mathcal{O}(U)$ -action on $\mathcal{F}(V)$ is defined via the natural ring homomorphism

$$\mathcal{O}(U) \rightarrow \mathcal{O}(V)$$

induced by f .

- (b) Let $\mathcal{F}$ and $\mathcal{G}$ be $\mathcal{O}$ -modules (Definition 7.1.3).

A $morphism of \mathcal{O}$ -modules $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ is a morphism of sheaves (Definition 4.0.2) of abelian groups such that, for every object $U \in \mathcal{C}$, the component map

$$\varphi_U : \mathcal{F}(U) \rightarrow \mathcal{G}(U)$$

is $\mathcal{O}(U)$ -linear, i.e. it satisfies

$$\varphi_U(r \cdot s) = r \cdot \varphi_U(s) \quad \text{for all } r \in \mathcal{O}(U), s \in \mathcal{F}(U).$$

The collection of all $\mathcal{O}$ -modules together with their morphisms of $\mathcal{O}$ -modules forms the $category of \mathcal{O}$ -modules, denoted $\mathbf{Mod}(\mathcal{O})$. The Hom functors $\mathrm{Hom}_{\mathbf{Mod}(\mathcal{O})}$ of this category may be denoted by $\mathrm{Hom}_{\mathcal{O}}$.

In case that a sheafification functor (Definition 4.3.1)

$$\mathrm{PreShv}(\mathcal{C}, \mathbf{Rings}) \rightarrow \mathrm{Shv}(\mathcal{C}, \mathbf{Rings})$$

exists, a left, right, two-sided $\mathcal{O}$ -module (and morphisms thereof) is equivalent to a $(\mathcal{O}, \mathbb{Z})$ -bimodule, $(\mathbb{Z}, \mathcal{O})$ -bimodule, and $(\mathcal{O}, \mathcal{O})$ -bimodule (and morphisms thereof) respectively, where $\mathbb{Z}$ is the constant sheaf (Definition C.0.10) of the integer ring $\mathbb{Z}$.

In particular, the category of $(\mathbb{Z}, \mathbb{Z})$ -bimodules is equivalent to the category of sheaves of abelian groups on $\mathcal{C}$.

Definition 7.1.4 (Tensor product of sheaves of bimodules). Let $\mathcal{C}$ be a site (Definition A.0.4). Let $\mathcal{R}, \mathcal{S}, \mathcal{T}$ be (not necessarily commutative) sheaves (Definition 4.0.2) of rings (Definition C.0.8) on $\mathcal{C}$. Let $\mathcal{M}$ be an $(\mathcal{R}, \mathcal{S})$ -bimodule (Definition 7.1.3), and let $\mathcal{N}$ be an $(\mathcal{S}, \mathcal{T})$ -bimodule.

(♠ TODO: Consolidate with)

1. Define the **presheaf tensor product** $\mathcal{M} \otimes_{p, \mathcal{S}} \mathcal{N}$ to be the presheaf (Definition 4.0.1) of abelian groups given by

$$U \mapsto \mathcal{M}(U) \otimes_{\mathcal{S}(U)} \mathcal{N}(U).$$

(Definition C.0.12) As a presheaf, it has the structure of a $\mathcal{R} - \mathcal{T}$ -bimodule (♠ TODO: bimodule for a presheaf).

2. Assume that a sheafification functor (Definition 4.3.1)

$$\mathrm{Shv}(\mathcal{C}, \mathbf{Sets}) \rightarrow \mathrm{PreShv}(\mathcal{C}, \mathbf{Sets})$$

exists. The **tensor product of sheaves** $\mathcal{M}$ and $\mathcal{N}$, denoted $\mathcal{M} \otimes_{\mathcal{S}} \mathcal{N}$, is defined to be the sheafification of the presheaf $\mathcal{M} \otimes_{p, \mathcal{S}} \mathcal{N}$. This tensor product becomes naturally an $(\mathcal{R}, \mathcal{T})$ -bimodule.

The left $\mathcal{R}$ -action and right $\mathcal{T}$ -action are induced by the presheaf actions defined section-wise by

$$\begin{aligned} r \cdot (m \otimes n) &= (r \cdot m) \otimes n, \\ (m \otimes n) \cdot t &= m \otimes (n \cdot t), \end{aligned}$$

for all $r \in \mathcal{R}(U)$, $t \in \mathcal{T}(U)$, $m \in \mathcal{M}(U)$, and $n \in \mathcal{N}(U)$. The sheafification functor preserves these actions, yielding the required bimodule structure on $\mathcal{M} \otimes_{\mathcal{S}} \mathcal{N}$.

Inductively, given sheaves of rings $\mathcal{R}_0, \dots, \mathcal{R}_k$ and $(\mathcal{R}_{i-1}, \mathcal{R}_i)$ -bimodules $\mathcal{M}_i$ for $i = 1, \dots, k$, we may speak of the tensor product

$$\mathcal{M}_0 \otimes_{\mathcal{R}_1} \mathcal{M}_1 \otimes_{\mathcal{R}_2} \cdots \otimes_{\mathcal{R}_{k-1}} \mathcal{M}_k.$$

Tensor products of sheaves are associative up to canonical isomorphism(♠ TODO:), allowing us to omit parentheses. This iterated tensor product carries a natural $(\mathcal{R}_0, \mathcal{R}_k)$ -bimodule structure.

In general, the assignment $(\mathcal{M}_0, \dots, \mathcal{M}_k) \mapsto \mathcal{M}_0 \otimes_{\mathcal{R}_1} \cdots \otimes_{\mathcal{R}_{k-1}} \mathcal{M}_k$ defines a $(k+1)$ -ary additive functor (Definition C.0.13)

$$\mathcal{R}_0 \mathbf{Mod}_{\mathcal{R}_1} \times \cdots \times \mathcal{R}_{k-1} \mathbf{Mod}_{\mathcal{R}_k} \rightarrow \mathcal{R}_0 \mathbf{Mod}_{\mathcal{R}_k}$$

()Proposition B.9.10).

Given a sheaf of rings $\mathcal{R}$ and a two-sided $\mathcal{R}$ -module $\mathcal{M}$, we may also speak of the **n -fold tensor product** $\mathcal{M}^{\otimes n} = \mathcal{M}^{\otimes_{\mathcal{R}} n}$.

Definition 7.1.5. Let $(\mathcal{C}, J)$ be a site (Definition 4.0.2), and let $\mathcal{A}$ be a category in which sheaves (Definition 4.0.2) on $(\mathcal{C}, J)$ can be defined.

(♠ TODO: subsheaf on a topological space) For a sheaf $\mathcal{F} \in \text{Sh}(\mathcal{C}, J; \mathcal{A})$, a *subsheaf* $\mathcal{G}$ of $\mathcal{F}$ is a sheaf equipped with a monomorphism (Definition .1.27) of sheaves

$$\iota : \mathcal{G} \hookrightarrow \mathcal{F}$$

in $\text{Sh}(\mathcal{C}, J; \mathcal{A})$. Equivalently, a subsheaf of $\mathcal{F}$ is simply a subobject of $\mathcal{F}$ in the category $\text{Sh}(\mathcal{C}, J; \mathcal{A})$.

Concretely, for every object $U \in \mathcal{C}$, the morphism

$$\iota(U) : \mathcal{G}(U) \rightarrow \mathcal{F}(U)$$

is a monomorphism in $\mathcal{A}$, and the collection $\{\iota(U)\}_U$ is compatible with restriction morphisms so that $\mathcal{G}$ is a subsheaf in the usual sense.

Definition 7.1.6. Let $(\mathcal{C}, J)$ be a site (Definition A.0.4), and let $\mathcal{O}$ be a sheaf of (not necessarily comutative) rings (Definition 4.0.2) on $(\mathcal{C}, J)$ with values in a category $\mathcal{A}$ for which sheaves are defined.

A *sheaf of (left/right/two-sided) ideals* $\mathcal{I}$ of the sheaf of rings $\mathcal{O}$ is a subsheaf of $\mathcal{O}$ (Definition 7.1.5) in the category of sheaves of (left/right/two-sided) $\mathcal{O}$ -modules, i.e., a sheaf of $\mathcal{O}$ -modules $\mathcal{I}$ together with a monomorphism (Definition .1.27) of sheaves of $\mathcal{O}$ -modules $\iota : \mathcal{I} \hookrightarrow \mathcal{O}$ such that for every object U in $\mathcal{C}$, the morphism $\iota(U) : \mathcal{I}(U) \rightarrow \mathcal{O}(U)$ identifies $\mathcal{I}(U)$ as an ideal of the ring $\mathcal{O}(U)$.

Equivalently, a sheaf of ideals $\mathcal{I}$ is a subsheaf (Definition 7.1.5) of modules of $\mathcal{O}$ as a sheaf of modules (Definition 7.1.3) of $\mathcal{O}$

Theorem 7.1.7 (e.g. see [?, Tag 01DU]). For any site (Definition A.0.4) $(\mathcal{C}, J)$ on an essentially small category (Definition .1.26) $\mathcal{C}$ and a sheaf of rings (Definition 4.0.2) $\mathcal{O}$ on $\mathcal{C}$, the category $\mathbf{Mod}(\mathcal{O})$ of $\mathcal{O}$ -modules (Definition 7.1.3) is an abelian category (Definition .1.34) that has enough injectives (Definition .1.37). In fact, there is a functorial injective embedding (♠ TODO: what does this mean?)

We can describe the sections of a sheaf over an object by the direct image from the object to the final object, if it exists.

Theorem 7.1.8. Let $(\mathcal{C}, J)$ be a site (Definition 4.0.2) with final object x (Definition .1.10), and let F be a sheaf on $(\mathcal{C}, J)$ (Definition 4.0.2). Let $f : X \rightarrow x$ be any object X over x (regarded as a morphism in $\mathcal{C}$), and let f_*F denote the direct image (pushforward) of F along f (Definition 6.1.10). Then

$$(f_*F)(x) = F(X) = \Gamma(X, F).$$

(Definition 7.1.1)

Corollary 7.1.9. (♠ TODO: the sites) Let $f : X \rightarrow S$ be a scheme over a scheme S and let $\mathbf{C}(X)$ and $\mathbf{C}(S)$ be small sites associated to X and S respectively, equipped with any common Grothendieck topologies such as Zariski, étale, Nisnevich, fppf, fpqc, or crystalline.

Let $\mathcal{A}$ be a (large) category and let $\mathcal{F} \in \mathrm{Sh}(\mathbf{C}(X), \mathcal{A})$ (Definition 4.0.2). We have a natural isomorphism

$$\Gamma(\mathcal{F}) \cong (f_*\mathcal{F})(S)$$

(Definition 6.1.20) (♠ TODO: natural in both $\mathcal{F}$ and S perhaps?)

Definition 7.1.10. (♠ TODO: generalize to more general abelian value categories)

Let $(\mathcal{C}, J)$ be a site (Definition A.0.4) on a locally small category (Definition .1.4) or a U -site for some universe (Definition .1.2) U . Let $\mathcal{O}$ be a sheaf of rings (Definition 4.0.2) on $\mathcal{C}$, so that $(\mathcal{C}, J, \mathcal{O})$ is a ringed site (Definition 6.1.5). Recall that the category $\mathbf{Mod}(\mathcal{O})$ (Definition 7.1.3) of $\mathcal{O}$ -modules is abelian and has enough injectives (Definition .1.37) (Theorem 7.1.7).

Assume that global sections objects $\Gamma(\mathcal{G})$ (Definition 7.1.1) exist for all objects $\mathcal{G}$ of $\mathrm{Sh}(\mathcal{C}, \mathbf{Ab})$ ⁴ so that Γ is a functor

$$\mathrm{Sh}(\mathcal{C}, \mathbf{Ab}) \rightarrow \mathbf{Ab},$$

which is a left exact functor (Definition .1.36) (Proposition 7.1.2). Note that Γ restricts to a left exact functor

$$\mathbf{Mod}(\mathcal{O}) \rightarrow \mathbf{Ab}.$$

If $\mathcal{C}$ has a final object (Definition .1.10) $*$ as well, then recall that $\Gamma(\mathcal{F}) = \mathcal{F}(*)$.

Let $\mathcal{F}$ be an object of $\mathbf{Mod}(\mathcal{O})$.

1. For each integer $n \geq 0$, the *n -th (abelian) (global) sheaf cohomology group of $\mathcal{F}$* is

$$H^n(\mathcal{C}, J; \mathcal{F}) := R^n\Gamma(\mathcal{F}),$$

where $R^n\Gamma$ is the n -th right derived functor (Definition .1.39) of the global sections functor Γ (Definition 7.1.1).

In particular, each H^n is a functor

$$H^n : \mathbf{Mod}(\mathcal{O}) \rightarrow \mathbf{Ab}.$$

2. Given an object $U \in \mathcal{C}$ and for each integer $n \geq 0$, the *n -th (abelian) sheaf cohomology group of $\mathcal{F}$ of sections at U* is

$$H^n(U, \mathcal{F}) := (R^n\Gamma(U, -))(\mathcal{F}),$$

where $R^n\Gamma(U, -)$ is the n -th right derived functor (Definition .1.39) of the sections functor $\Gamma(U, -)$ evaluated at U (Definition 7.1.1).

In particular, $H^n(U, \mathcal{F})$ can be regarded as the n th global sheaf cohomology group of the restriction $\mathcal{F}|_U$ of $\mathcal{F}$ to U .

⁴for example, this occurs when $\mathcal{C}$ is essentially small (Definition .1.26)

7.2. Étale cohomology of literal sheaves.

Definition 7.2.1. Let X be a scheme, let $\mathcal{O}$ be a sheaf of rings (Definition 4.0.2) on $X_{\text{ét}}$ (Definition 4.1.3) and let $\mathcal{F}$ be an $\mathcal{O}$ -module (Definition 7.1.3), which is sheaf of abelian groups on the small étale site $X_{\text{ét}}$ (Definition 4.1.3). Note that the category of $\mathcal{O}$ -modules has enough injectives (Definition 7.1.37) (Theorem 7.1.7)

The *étale cohomology groups of X with coefficients in $\mathcal{F}$* are defined as the right derived functors (Definition 7.1.39) of the global sections functor (Definition 7.1.1), which is left exact (Definition 7.1.36) (Proposition 7.1.2). Explicitly, they are the abelian groups

$$H_{\text{ét}}^i(X, \mathcal{F}) := H^i(X_{\text{ét}}, \mathcal{F}) = R^i\Gamma(X, \mathcal{F}), \quad i \geq 0.$$

Equivalently, $H_{\text{ét}}^i(X, \mathcal{F})$ is defined as the sheaf cohomology groups of the sheaf $\mathcal{F}$ on the essentially small site $X_{\text{ét}}$ (Definition 7.1.10)

Definition 7.2.2. Let X be a scheme. Let A be an abelian group and write $\underline{A}$ for the constant sheaf (Definition C.0.10) on $X_{\text{ét}}$ (Definition 4.1.3) associated to A . In this case, the *étale cohomology groups of X with coefficients in A* are defined as

$$H_{\text{ét}}^i(X, A) := H_{\text{ét}}^i(X, \underline{A}).$$

(Definition 7.2.1); here $\underline{A}$ is regarded as a module of the constant sheaf $\mathbb{Z}$ on $X_{\text{ét}}$.

8. DERIVED CATEGORIES OF λ -ADIC SHEAVES CONSTRUCTIBLE THROUGH LIMITS OF CATEGORIES

(♠ TODO: Give an outline of how $D_c^b(X, \overline{\mathbb{Q}}_\ell)$ is defined) Classically, given a noetherian scheme X and a prime ℓ invertible on X , the category $D_c^b(X, \overline{\mathbb{Q}}_\ell)$, which is intuitively treated as a cohomologically bounded derived category of constructible complexes of sheaves with $\overline{\mathbb{Q}}_\ell$ coefficients, is constructed via the following steps:

(♠ TODO: Cite the statements of the constructions)

1. Write E for a finite extension of $\mathbb{Q}_\ell$, and write R for the integral closure of $\mathbb{Z}_\ell$ in E . Note that R is a complete discrete valuation ring. Write λ for a uniformizer of R .
2. Construct $D_c^b(X, R)$ (♠ TODO: Explain how $D_c^b(X, R)$ is constructed.).
3. Define $D_c^b(X, E) = D_c^b(X, R)[\lambda^{-1}]$, i.e. construct $D_c^b(X, E)$ as the localization of $D_c^b(X, R)$ by the maps $\lambda^m : \mathcal{F} \rightarrow \mathcal{F}$ in $D_c^b(X, R)$.
4. Observe that for finite extensions E'/E , there are functors

$$D_c^b(X, E) \rightarrow D_c^b(X, E')$$

5. Construct $D_c^b(X, \overline{\mathbb{Q}}_\ell)$ as the 2-direct limit of the $D_c^b(X, E)$ as E runs over the finite extensions of $\mathbb{Q}_\ell$ in $\overline{\mathbb{Q}}_\ell$:

$$D_c^b(X, \overline{\mathbb{Q}}_\ell) = \varinjlim_{\substack{\overline{\mathbb{Q}}_\ell \supset E \supset \mathbb{Q}_\ell \\ E/\mathbb{Q}_\ell \text{ finite}}} D_c^b(X, E)$$

8.1. **ℓ -adic sheaves.** ℓ -adic sheaves do not refer to literal sheaves on $X_{\text{ét}}$ with coefficients in ℓ -adic rings such as $\mathbb{Z}_\ell$ or $\mathbb{Q}_\ell$. In a purely formal sense, one could talk about sheaves on $X_{\text{ét}}$ with such coefficients, but the categories of such sheaves are too ill-behaved to be of arithmetic use.

8.2. **The categories of λ -adic sheaves and A-R λ -adic sheaves.** Historically, the theory of λ -adic sheaves (as opposed to “derived categories”/categories of complexes of λ -adic sheaves) was first studied in SGA5 [?, Exposé V]. Note that SGA5 [?, Exposé III] did study derived categories of sheaves with *finite* coefficients.

This subsection considers four abelian categories whose objects are inverse systems of sheaves:

1. The category of inverse systems of λ -torsion sheaves (Definition 8.4.1)
2. The A-R category of inverse systems of λ -torsion sheaves (Definition 8.4.8)
3. The category of λ -adic sheaves (Definition 8.4.9)
4. The category of A-R λ -adic sheaves (Definition 8.4.9)

These categories are related via the following commutative diagram of functors:

$$\begin{array}{ccc}
 (\text{inverse system of } \lambda\text{-torsion sheaves}) & \xrightarrow{\text{localization}} & (\text{A-R category}) \\
 \uparrow \text{full subcategory} & & \uparrow \text{full subcategory} \\
 (\lambda\text{-adic sheaves}) & \xrightarrow{\cong} & (\text{A-R } \lambda\text{-adic sheaves})
 \end{array}$$

8.3. Constructible sheaves.

Definition 8.3.1 (Constructible sheaf). Let X be either

1. A topological space (Definition C.0.6) (made into a site (Definition A.0.4) with the natural Grothendieck topology (Definition .1.9)) or (♠ TODO: This second case needs to be formulated more precisely; in particular, a notion of “restriction” is necessary, and there needs to be a more precise way to formulate that the site is a site associated to X .)
2. A scheme (Definition .3.1) along with a small site $(\mathcal{C}, J)$ on X (Definition 4.1.1). Assume that there is a sheafification functor (Definition 4.3.1) for the presheaves/sheaves valued in abelian groups so that locally constant sheaves (Definition 4.4.2) of abelian groups are defined. For example, it suffices for the underlying category of the site to be essentially small (Definition .1.26)(Theorem 4.3.2).

Let $\mathcal{O}$ be a sheaf of rings (Definition 4.0.2) on the chosen site $(\mathcal{C}, J)$ associated to X . Let $\mathcal{F}$ be a sheaf of $\mathcal{O}$ -modules (Definition 7.1.3) on X .

The sheaf $\mathcal{F}$ is called *constructible* if there exists a finite partition of X into locally closed subsets (Definition C.0.7)

$$X = \bigsqcup_{\alpha \in A} X_\alpha$$

such that the restriction $\mathcal{F}|_{X_\alpha}$ is a locally constant sheaf of $\mathcal{O}|_{X_\alpha}$ -modules of finite type (Definition 4.4.2) (Definition 4.4.2) on the site on X_α induced by $(\mathcal{C}, J)$ and the embedding $X_\alpha \hookrightarrow X$ (♠ TODO: The site on X_α needs to be elaborated on.) for each $\alpha \in A$.

Definition 8.3.2 (Full subcategory). Let $\mathcal{C}$ be a (large) category (Definition .1.1). A *full subcategory* $\mathcal{D}$ of $\mathcal{C}$ is a subcategory such that for every pair of objects $X, Y \in \text{Ob}(\mathcal{D})$, the morphism classes coincide:

$$\text{Hom}_{\mathcal{D}}(X, Y) = \text{Hom}_{\mathcal{C}}(X, Y).$$

In other words, a full subcategory includes all morphisms between its objects that exist in the ambient category $\mathcal{C}$.

Definition 8.3.3. Let X be either

1. A topological space (Definition C.0.6) (made into a site (Definition A.0.4) with the natural Grothendieck topology (Definition .1.9)) or
2. A scheme (Definition .3.1) along with a small site $(\mathcal{C}, J)$ on X (Definition 4.1.1). Assume that there is a sheafification functor (Definition 4.3.1) for the presheaves/sheaves valued in abelian groups so that locally constant sheaves (Definition 4.4.2) of abelian groups are defined. For example, it suffices for the underlying category of the site to be essentially small (Definition .1.26)(Theorem 4.3.2).

Note that the sites are essentially small (Definition .1.26) (see Theorem 4.2.16).

The category $D_c(X, \mathcal{O})$ is the full subcategory (Definition 8.3.2) of the derived category $D(X, \mathcal{O})$ (Notation 10.0.7) of objects $K^\bullet$ whose (co)homology objects (Definition .2.3) are constructible sheaves (Definition 8.3.1). It is customary to speak of the subcategories $D^?(X, \mathcal{O})$ for $? \in \{+, -, b\}$ as usual.

8.4. λ -adic sheaves.

Definition 8.4.1 (λ -torsion sheaf on a scheme). Let $(\mathcal{C}, J)$ be a site (Definition A.0.4). Let $\mathcal{O}$ be a sheaf of commutative rings (Definition 4.0.2) on $(\mathcal{C}, J)$. Let $\lambda \subseteq \mathcal{O}$ be a sheaf of ideals (Definition 7.1.6).

A sheaf $\mathcal{F}$ of $\mathcal{O}$ -modules (Definition 7.1.3) on $(\mathcal{C}, J)$ is called a *λ -torsion sheaf* if

$$\mathcal{F} = \bigcup_{n \geq 1} \ker(\lambda^n : \mathcal{F} \rightarrow \mathcal{F})$$

or equivalently if for every section (Definition 7.1.1) $s \in \mathcal{F}(U)$ for some object $U \in \text{Ob}(\mathcal{C})$, there exists some $n \geq 1$ such that $\lambda^n \cdot s = 0$.

For instance, we may speak of λ -torsion sheaves by letting $\mathcal{O} = \underline{R}$ (Definition C.0.10) where R is a commutative ring, $\lambda = \underline{I}$ where $I \subseteq R$ is an ideal, and $(\mathcal{C}, J)$ be the small étale site of a scheme (Definition 4.1.3).

λ -adic sheaves (Definition 8.4.9) are specific inverse systems of torsion sheaves; imposing the Mittag-Leffler condition (Definition 8.4.4) ensures the existence of the limit as a sheaf and the stronger Artin-Rees-Mittag-Leffler condition (Definition 8.4.7) yields an abelian category.

Definition 8.4.2. Let $\mathcal{C}$ be a category (Definition .1.1) and let I be the poset (Definition .1.28) of non-negative integers $\mathbb{Z}_{\geq 0}$ with the standard ordering $\geq$ (i.e. I has arrows $\cdots \rightarrow 2 \rightarrow 1 \rightarrow 0$). A **tower in $\mathcal{C}$** is a functor $X : I \rightarrow \mathcal{C}$. Explicitly, a tower consists of a sequence of objects $\{A_i\}_{i \in \mathbb{Z}_{\geq 0}}$ and morphisms $f_i : A_i \rightarrow A_{i-1}$ for each $i \geq 1$:

$$\cdots \xrightarrow{f_3} A_2 \xrightarrow{f_2} A_1 \xrightarrow{f_1} A_0.$$

Moreover a **morphism of towers** is a morphism of towers as objects of the functor category (Definition .1.33) $\mathcal{C}^I$.

The notions of towers and their morphisms are equivalent to those of inverse sequences (Definition 10.0.1) up to indexing.

Definition 8.4.3. Let $\mathcal{C}$ be a category (Definition .1.1) and let I be the poset (Definition .1.28) of non-negative integers $\mathbb{N}$ with the standard ordering $\geq$ (i.e. I has arrows $\cdots \rightarrow 2 \rightarrow 1 \rightarrow 0$) (or more generally let I be the poset of integers $\mathbb{Z}$ with similar ordering).

Given a tower (Definition 8.4.2) $A = (A_i, u_i)_{i \in I}$ where $u_i : A_i \rightarrow A_{i-1}$ (for $i \geq 1$ if $\mathbb{N}$ is the collection of objects of I) and $r \geq 0$, the **shifted tower $A[r]$** is the tower $A[r] = (A[r]_i, u[r]_i)$ given by $A[r]_i = A_{r+i}$ and whose transition maps are given by $u[r]_i : A[r]_i = A_{r+i} \xrightarrow{u_{r+i}} A_{r+i-1} = A[r]_{i-1}$.

Note that there is a morphism of towers (Definition 8.4.2) $A[r] \rightarrow A$ where $A[r]_i \rightarrow A_i$ is given by the transition morphism $A_{r+i} \rightarrow A_i$.

Definition 8.4.4 (cf. [?, Definition 3.5.6]). Let I be a directed set (Definition .1.12) and let $\{A_i, \phi_{ji}\}_{i \in I}$ be an inverse system (Definition .1.18) of objects in a category $\mathcal{C}$ where images are well-defined (such as the category of sets, abelian groups, or modules).

The system is said to satisfy the **Mittag-Leffler condition** if for every index $i \in I$, there exists an index $j \geq i$ such that for all $k \geq j$, the image of the transition map $\phi_{ki} : A_k \rightarrow A_i$ is equal to the image of $\phi_{ji} : A_j \rightarrow A_i$.

For a fixed i , let $I_{k,i} = \text{im}(\phi_{ki}) \subseteq A_i$ for all $k \geq i$. The condition states that the decreasing family of subobjects

$$A_i \supseteq I_{i,i} \supseteq I_{i+1,i} \supseteq I_{i+2,i} \supseteq \cdots$$

becomes stationary.

Definition 8.4.5 (cf. [?, Definition 3.5.6]). Let I be a directed set (Definition .1.12) and let $\{A_i, \phi_{ji}\}_{i \in I}$ be an inverse system (Definition .1.18) of objects in a pointed category $\mathcal{C}$.

The system is said to satisfy the **trivial Mittag-Leffler condition** if for every index $i \in I$, there exists an index $j > i$ such that the map $A_j \rightarrow A_i$ is a zero morphism.

Definition 8.4.6. Let I be the poset (Definition .1.28) of non-negative integers $\mathbb{N}$ with the standard ordering $\geq$ (i.e. I has arrows $\cdots \rightarrow 2 \rightarrow 1 \rightarrow 0$) and let $\{A_i, \phi_{ji}\}_{i \in I}$ be an inverse system (Definition .1.18) of objects (so a tower (Definition 8.4.2)) in a pointed category $\mathcal{C}$ where images are well-defined (such as the category of sets, abelian groups, or modules).

The system is said to be a **null system** if there exists $r \geq 0$ such that for every $i \in I$, the transition map $\phi_{(i+r)i} : A_{i+r} \rightarrow A_i$ is the zero morphism.

Equivalently, the tower is a null system if there exists an integer $r \geq 0$ such that the morphism of towers (Definition 8.4.2) $A[r] \rightarrow A$ (Definition 8.4.3) is the zero morphism. (♠ TODO: The category of towers $\mathcal{C}^I$ is pointed as $\mathcal{C}$ is pointed; the zero morphism is defined component-wise.)

Definition 8.4.7 (See e.g. [?, Exposé V, VI], [?, 1.1] [?, 10.1]). Let $A = \{A_i, \varphi_{ji}\}$ be a tower (Definition 8.4.2) in a category $\mathcal{C}$ where images are well-defined.

The tower is said to satisfy the *Artin-Rees-Mittag-Lejfler (ARML) condition* if there exists an integer $r \geq 0$ such that for every index $i \in \mathbb{N}$ and every $k \geq i + r$, the image of the transition map $\phi_{ki} : A_k \rightarrow A_i$ is equal to the image of $\phi_{(i+r)i} : A_{i+r} \rightarrow A_i$.

Equivalently, the tower satisfies the ARML condition if and only if for any integer n , there exists an integer $r \geq 0$ such that

$$\mathrm{im}(A[r] \rightarrow A) = \mathrm{im}(A[t] \rightarrow A)$$

Definition 8.4.3 for all $t \geq r$.

Definition 8.4.8 (A-R category of inverse systems of λ -torsion sheaves). Let X be a scheme. Let

$$\mathcal{F} = (\mathcal{F}_n, u_n)_{n \in \mathbb{Z}}, \quad u_n : \mathcal{F}_n \rightarrow \mathcal{F}_{n-1}$$

be an inverse system (Definition .1.18) of sheaves of abelian groups on the (small) étale site $X_{\text{ét}}$ (Definition 4.1.3). Let R be a commutative ring and let $\lambda \subseteq R$ be an ideal.

The *A-R category of inverse systems of λ -torsion sheaves on $X_{\text{ét}}$* , denoted $\mathrm{AR}(\lambda\text{-tors}(X))$, is the localization of the multiplicative system consisting of the family of morphisms of the form $\mathcal{F}[r] \rightarrow \mathcal{F}$. In other words, the objects of the category are inverse systems of λ -torsion sheaves (Definition 8.4.1) and for any objects $\mathcal{F}$ and $\mathcal{G}$, we have

$$\mathrm{Hom}_{\mathrm{AR}(\lambda\text{-tors}(X))} = \varinjlim_{r \geq 0} \mathrm{Hom}(\mathcal{F}[r], \mathcal{G}).$$

(♠ TODO: morphism of inverse systems; more generally diagrams) In particular, a morphism $\mathcal{F} \rightarrow \mathcal{G}$ in $\mathrm{AR}(\lambda\text{-tors}(X))$ is represented by a morphism $\mathcal{F}[r] \rightarrow \mathcal{G}$ of inverse systems for some $r \geq 0$, and the kernel and cokernel of $\mathcal{F}[r] \rightarrow \mathcal{G}$ are the kernel and cokernel of $\mathcal{F} \rightarrow \mathcal{G}$ in $\mathrm{AR}(\lambda\text{-tors}(X))$

It is worth noting that $\mathrm{AR}(\lambda\text{-tors}(X))$ is an abelian category (Definition .1.34) and that $\mathcal{F} \in \mathrm{AR}(\lambda\text{-tors}(X))$ is zero if and only if it is a null system (Definition 8.4.6).

Definition 8.4.9 (λ -adic sheaf on a scheme). Let R be a commutative ring (Definition C.0.1) and let $\lambda \subseteq R$ be an ideal (Definition C.0.2). Let X be a scheme (Definition .3.1). Let

$$\mathcal{F} = (\mathcal{F}_n, u_n)_{n \in \mathbb{Z}_{\geq 1}}, \quad u_n : \mathcal{F}_n \rightarrow \mathcal{F}_{n-1}$$

be an inverse system (Definition .1.18) of λ -torsion sheaves (Definition 8.4.1) on the (small) étale site $X_{\text{ét}}$ (Definition 4.1.3).

1. The inverse system is called a *λ -adic sheaf on X* or a *sheaf on X with coefficients in R* if

- $\mathcal{F}_n$ is constructible (Definition 8.3.1) for all n ,
- $\mathcal{F}_n$ is annihilated by λ^n for all n ,

- there are isomorphisms $\mathcal{F}_n \cong \mathcal{F}_{n+1}/\lambda^n \mathcal{F}_{n+1}$ compatible with transition maps.
2. A λ -adic sheaf is called *lisse* if each $\mathcal{F}_n$ is locally constant (Definition 4.4.2).
 3. A λ -adic sheaf is called an *A - R λ -adic sheaf* if it satisfies the Artin–Rees–Mittag–Leffler condition (Definition 8.4.7).

The category of λ -adic sheaves may be denoted by notations such as $\mathbf{Sh}(X, R)$. In particular, the *category of ℓ -adic sheaves* is the category $\mathbf{Sh}(X, \mathbb{Z}_\ell)$. Note however, that a λ -adic sheaves is not a sheaf (Definition 4.0.2) in the literal sense but only an inverse system of sheaves.

One typically works with this theory only when X is noetherian (Definition C.0.16) so that constructibility is stable under basic operations. One also typically works with this theory when R is a complete DVR and λ its maximal ideal. Under nice enough circumstances, the categories of λ -adic sheaves and $A - R$ λ -adic sheaves are equivalent (Theorem 8.4.13).

Here is a basic example of a lambda adic sheaf:

Definition 8.4.10 (ℓ -adic sheaf $\mathbb{Z}_\ell$ on the small étale site). (♠ TODO: prime invertible on a scheme) Let X be a scheme (Definition .3.1) and let ℓ be a prime invertible on X .

The *ℓ -adic sheaf* $\mathbb{Z}_\ell$ on the small étale site $X_{\text{ét}}$ (Definition 4.1.3) is defined as the inverse system (Definition .1.18)

$$\mathbb{Z}_\ell := \varprojlim_k \mathbb{Z}/\ell^k \mathbb{Z},$$

where each $\mathbb{Z}/\ell^k \mathbb{Z}$ is the constant sheaf (Definition C.0.10) of abelian groups on $X_{\text{ét}}$, and the transition maps are the canonical projections.

These inverse limits define objects in the category of ℓ -adic sheaves (Definition 8.4.9), rather than single sheaves in the classical étale topology.

Definition 8.4.11 (The ℓ -adic Tate twists $\mathbb{Z}_\ell(1)$ and $\mathbb{Z}_\ell(d)$). Let X be a scheme and let ℓ be a prime invertible on X , and set

$$\mathbb{Z}_\ell(1) := \varprojlim_k \mu_{\ell^k},$$

the inverse limit (projective system) (Definition .1.13) of sheaves of ℓ^k -th roots of unity (Definition 4.4.7) on $X_{\text{ét}}$ (Definition 4.1.3), equipped with its natural $\mathbb{Z}_\ell$ -module structure.

For an integer $d \geq 1$, define the *d -th Tate twist*

$$\mathbb{Z}_\ell(d) := \underbrace{\mathbb{Z}_\ell(1) \otimes_{\mathbb{Z}_\ell} \cdots \otimes_{\mathbb{Z}_\ell} \mathbb{Z}_\ell(1)}_{d \text{ times}}.$$

For $d = 0$ set $\mathbb{Z}_\ell(0) := \mathbb{Z}_\ell$ (Definition 8.4.10).

Lemma 8.4.12. Let $(\mathcal{C}, J)$ be a site (Definition A.0.4). Let $\mathcal{O}$ be a sheaf of commutative rings (Definition 4.0.2) on $(\mathcal{C}, J)$. Let $\lambda \subseteq \mathcal{O}$ be a sheaf of ideals (Definition 7.1.6). Let $\{\mathcal{G}_n\}_{n \geq 1}$ be an inverse system of constructible (♠ TODO: generalize the notion of constructible sheaves to a sheaf of $\mathcal{O}$ -modules)

Let R be a commutative ring and let $\lambda \subseteq R$ be an ideal. (♠ TODO: Try to generalize stacks 03UN as an exercise)

Theorem 8.4.13. See e.g. [?, 10.1] Let X be a noetherian scheme (Definition C.0.16). Let ℓ be a prime number that is invertible on X . Let R be the integral closure of $\mathbb{Z}_\ell$ in a finite extension E of $\mathbb{Q}_\ell$. Note that R is a complete discrete valuation ring and let λ be the maximal ideal of R . The category of λ -adic sheaves (Definition 8.4.9) is equivalent to the category of A-R λ -adic sheaves (Definition 8.4.9).

8.5. Rational coefficients.

Definition 8.5.1. (♠ TODO: what are the minimal hypotheses needed for the following) Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let E be the fraction field of R . Let X be a noetherian scheme (Definition C.0.16), and consider sheaves (Definition 4.0.2) on the (small) étale site on X (Definition 4.1.3).

The *category of sheaves with coefficients in E* is the category $\text{Sh}(X, E)$ that is the localization (Definition B.2.2) of $\text{Sh}(X, R)$ (Definition 8.4.9) by the class

$$\{\lambda^n \cdot \text{id}_{\mathcal{F}} : \mathcal{F} \in \text{Sh}(X, R), n \in \mathbb{Z}_{\geq 0}\}$$

of morphisms.

Equivalently, $\text{Sh}(X, E)$ is the category with the same objects as $\text{Sh}(X, R)$ and whose morphism sets are given by

$$\text{Hom}_{\text{Sh}(X, E)}(\mathcal{F}, \mathcal{G}) = \text{Hom}_{\text{Sh}(X, R)}(\mathcal{F}, \mathcal{G}) \otimes_R E.$$

One case of particular interest is the case that $R = \mathbb{Z}_\ell$ and $E = \mathbb{Q}_\ell$.

As is the case with $\text{Sh}(X, R)$, note that the objects of $\text{Sh}(X, E)$ are not literal sheaves but rather inverse systems of sheaves.

$\text{Sh}(X, E)$ is a full subcategory of $D_c^b(X, E)$ (Definition 8.6.10).

Definition 8.5.2. Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let E be the fraction field of R . Let K be an algebraic extension of E . Let X be a noetherian scheme (Definition C.0.16), and consider sheaves (Definition 4.0.2) on the (small) étale site on X (Definition 4.1.3).

For each finite extension field E' of E in K , note that there is a category $\text{Sh}(X, E')$ (Definition 8.5.1) and that these categories form a direct system — given $E'_1 \subseteq E'_2$, we have a functor $\text{Sh}(X, E'_1) \rightarrow \text{Sh}(X, E'_2)$.

The *category of sheaves with coefficients in K* is the category $\text{Sh}(X, K)$ defined as the 2-limit of this direct system:

$$\text{Sh}(X, K) := \varinjlim_{E \subseteq E' \subseteq K, [E':E] < \infty} \text{Sh}(X, E').$$

More concretely, an object $\mathcal{F}$ of $\text{Sh}(X, K)$ is represented by an object $\mathcal{F}_{E'}$ of $\text{Sh}(X, E')$ for some finite extension E' of E in K . Two objects $\mathcal{F}, \mathcal{G} \in \text{Sh}(X, K)$ represented by $\mathcal{F}_{E'_1}$ and

$\mathcal{G}_{E'_2}$ are considered the same if there exists some finite extension E'' of E containing both E'_1 and E'_2 such that the pushforwards of $\mathcal{F}_{E'_1}$ and $\mathcal{G}_{E'_2}$ into $\mathrm{Sh}(X, E'')$ are isomorphic objects.

Furthermore, for objects $\mathcal{F}, \mathcal{G}$ of $\mathrm{Sh}(X, K)$ represented by $\mathcal{F}_{E'}, \mathcal{G}_{E'} \in \mathrm{Sh}(X, E')$ for some finite extension E' of E in K , their Hom-set is given by

$$\mathrm{Hom}_{\mathrm{Sh}(X, K)}(\mathcal{F}, \mathcal{G}) = \varinjlim_{E' \subseteq E'' \subseteq K, [E'': E] < \infty} \mathrm{Hom}_{\mathrm{Sh}(X, E'')}(\mathcal{F}_{E'}, \mathcal{G}_{E'})$$

where $\mathcal{F}_{E'}$, $\mathcal{G}_{E'}$ are respectively the pushforwards of $\mathcal{F}, \mathcal{G}$ in $\mathrm{Sh}(X, E')$. Equivalently,

$$\mathrm{Hom}_{\mathrm{Sh}(X, K)}(\mathcal{F}, \mathcal{G}) = \mathrm{Hom}_{\mathrm{Sh}(X, E')}(\mathcal{F}, \mathcal{G}) \otimes_{E'} K$$

One case of particular interest is the case that $R = \mathbb{Z}_\ell$, $E = \mathbb{Q}_\ell$, and $K = \overline{\mathbb{Q}_\ell}$.

As is the case with $\mathrm{Sh}(X, R)$ or $\mathrm{Sh}(X, E)$, note that the objects of $\mathrm{Sh}(X, K)$ are not literal sheaves but rather inverse systems of sheaves.

Note that the definition of $\mathrm{Sh}(X, K)$ simulatenously uses and generalizes the definition of $\mathrm{Sh}(X, E)$.

$\mathrm{Sh}(X, K)$ is a full subcategory of $D_c^b(X, K)$ (Definition 8.6.10).

Theorem 8.5.3. Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let E be the fraction field of R . Let K be an algebraic extension of E . Let X be a connected noetherian scheme (Definition C.0.16). Let $\bar{x} \in X$ be a geometric point (Definition 3.0.1).

1. There is an equivalence of categories between:

$$\{\text{lisse } \lambda\text{-adic sheaves on } X_{\text{étale}}\} \longleftrightarrow \left\{ \begin{array}{l} \text{finitely generated free } R\text{-modules } M \\ \text{with a continuous } \pi_1^{\text{ét}}(X, \bar{x})\text{-action} \end{array} \right\}.$$

(Definition 8.4.9)

2. There is an equivalence of categories between:

$$\{\text{lisse objects of } \mathrm{Sh}(X, E)\} \longleftrightarrow \left\{ \begin{array}{l} \text{finite-dimensional } E\text{-vector spaces } V \\ \text{with a continuous } \pi_1^{\text{ét}}(X, \bar{x})\text{-action} \end{array} \right\}.$$

(Definition 8.5.1)

3. Let K be an algebraic extension of E . There is an equivalence of categories between:

$$\{\text{lisse objects of } \mathrm{Sh}(X, K)\} \longleftrightarrow \left\{ \begin{array}{l} \text{finite-dimensional } K\text{-vector spaces } V \\ \text{with a continuous } \pi_1^{\text{ét}}(X, \bar{x})\text{-action} \end{array} \right\}.$$

(Definition 8.5.2)

8.5.1. Geometric stalks of λ -adic sheaves.

Definition 8.5.4. (♠ TODO: emergency) Let R be a commutative ring and let $\lambda \subseteq R$ be an ideal. Let X be a scheme. Let $\bar{x} \in X$ be a geometric point. Let

$$\mathcal{F} = (\mathcal{F}_n, u_n)_{n \in \mathbb{Z}_{\geq 1}}, \quad u_n : \mathcal{F}_n \rightarrow \mathcal{F}_{n-1}$$

be a λ -adic sheaf (Definition 8.4.9) (on the small étale site of X).

The *(geometric) stalk of $\mathcal{F}$ at $\bar{x}$* is the inverse limit

$$\mathcal{F}_{\bar{x}} := \varprojlim_n \mathcal{F}_{n, \bar{x}}.$$

(Definition 4.5.1) It is naturally a $\widehat{R}$ module where $\widehat{R}$ is the λ -adic completion of R . In particular, every geometric point $\bar{x} \in X$ yields a functor

$$\mathcal{F} \mapsto \mathcal{F}_{\bar{x}} : \mathrm{Sh}(X, R) \rightarrow \mathrm{Mod}(\widehat{R})$$

that is in fact exact.

Definition 8.5.5. Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$, let E be the fraction field of R , let K be an algebraic extension of E , let X be a noetherian scheme (Definition C.0.16) equipped with its (small) étale site (Definition 4.1.3), let $\mathcal{F}$ be an object of $\mathrm{Sh}(X, K)$ (Definition 8.5.2) (Definition 8.5.1), and let $\bar{x} \in X$ be a geometric point.

By definition, $\mathcal{F}$ is representable by an object $\mathcal{F}_{E'}$ of $\mathrm{Sh}(X, E')$ for some finite extension E' of E in K . In turn, $\mathcal{F}_{E'}$ is an object of $\mathrm{Sh}(X, R')$ where R' is the integral closure of R in E' or equivalently the ring of integers of E' . The *(geometric) stalk of $\mathcal{F}$* is defined as

$$\mathcal{F}_{\bar{x}} := \mathcal{F}_{E', \bar{x}} \otimes_{E'} K.$$

(Definition 8.5.4) (Definition C.0.12) It is not dependent on the choice of E' or the representative $\mathcal{F}_{E'}$.

For each geometric point $\bar{x}$ of X , we have an exact functor

$$\mathcal{F} \mapsto \mathcal{F}_{\bar{x}}, \quad \mathrm{Sh}(X, K) \rightarrow \mathrm{Mod}(K).$$

8.6. The category $D_c^b(X, R)$ of integral coefficients. For a scheme X , a prime ℓ invertible on X , and the ring of integers R in a finite extension E of $\mathbb{Q}_\ell$, Deligne [?, I.I.2] defined “derived” categories $D_c^b(X, R)$ (Definition 8.6.2), $D_c^b(X, E)$, and $D_c^b(X, \overline{\mathbb{Q}}_\ell)$, intuitively of “integral”, “rational”, and $\overline{\mathbb{Q}}_\ell$ coefficients respectively. These categories are not derived categories themselves but constructed λ -adically through the genuine derived categories $D_c^b(X, R/\lambda^n)$ of finite coefficients. loc. cit. also established that the six functor formalisms on the categories $D_c^b(X, R/\lambda^n)$ induce a six functor formalism on $D_c^b(X, R)$ under suitable finiteness conditions, e.g. X is of finite type over a regular base S of dimension ≤ 1 , and that $D_c^b(X, R)$ is a triangulated category under other finiteness conditions, e.g. X is of finite type over a finite field or an algebraically closed field.

Definition 8.6.1. Let X be a scheme (Definition 4.1.3) equipped with its (small) étale site (Definition 4.1.3). Let R be a (commutative) ring (Definition C.0.1) equipped with an ideal (Definition C.0.2) λ .

Define a category $D^-(X, R)$ as follows:

- Objects are families $K = (K_n, u_n)_{n \geq 0}$ where $K_n \in \text{Ob } D^-(X, R/\lambda^{n+1})$ (Notation B.9.11) (Definition B.3.1) ⁵ and u_n are isomorphisms

$$u_n : K_{n+1} \overset{\text{L}}{\otimes}_{R/\lambda^{n+2}} R/\lambda^{n+1} \cong K_n$$

(Definition B.9.4) in $D^-(X, R/\lambda^{n+1})$.

- Morphisms $f : K = (K_n, u_n)_{n \geq 0} \rightarrow K' = (K'_n, u'_n)_{n \geq 0}$ are families (f_n) of morphisms $f_n : K_n \rightarrow K'_n$ in $D(X, R/\lambda^{n+1})$ such that

$$f_n u_n = u'_n (f_{n+1} \overset{\text{L}}{\otimes}_{R/\lambda^{n+2}} \text{id}_{R/\lambda^{n+1}}).$$

This category $D^-(X, R)$ should not in general not be mistaken for the category also denoted $D^-(X, R)$, which is a derived category of sheaves of R -modules on $X_{\text{ét}}$.

K_n is also often denoted by $K \overset{\text{L}}{\otimes} R/\lambda^{n+1}$.

Definition 8.6.2 (see [?, I.I.2], cf. [?, Between Propositions 10.1.16 and 10.1.17], for a discussion in the case that R is the ring of integers in a finite extension of $\mathbb{Q}_\ell$ and λ is the maximal ideal of R). (♠ TODO: it may be reasonable to divide this definition into two parts, one just for $D^-(X, R)$ where X is not assumed to be noetherian and maybe R is assumed to not be a DVR)

Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$.

Let X be a noetherian scheme (Definition C.0.16), and consider sheaves (Definition 4.0.2) on the (small) étale site on X (Definition 4.1.3).

Let $D_c^b(X, R)$ be the full subcategory of $D^-(X, R)$ (Definition 8.6.1) (Notation B.9.11) whose objects $K = (K_n, u_n)_{n \geq 0}$ satisfy $K_0 \in \text{Ob } D_c^b(X, R/(\lambda))$ Definition 8.3.3.

Proposition 8.6.3 (See, e.g. [?, Proposition 10.1.16]). Let X be a noetherian scheme. Let ℓ be a prime number that is invertible on X . Let R be the integral closure of $\mathbb{Z}_\ell$ in a finite extension E of $\mathbb{Q}_\ell$. Note that R is a complete discrete valuation ring and let λ be the maximal ideal of R .

For $K = (K_n, u_n)_{n \geq 0}$ in $D_c^b(X, R)$ (Definition 8.6.2), we have that $K_n \in D_{\text{ctf}}^b(X, R/(\lambda^{n+1}))$ (♠ TODO: ctf) for all n , and the inverse systems $(H^i(K_n)_{n \in \mathbb{Z}})$ are A-R λ -adic (Definition 8.4.9).

Remark 8.6.4. Let X be a noetherian scheme. Let ℓ be a prime number that is invertible on X . Let R be the integral closure of $\mathbb{Z}_\ell$ in a finite extension E of $\mathbb{Q}_\ell$. If $\mathcal{F} = (\mathcal{F}_n)$ is a torsion free λ -adic sheaf (♠ TODO: define torsion free), then $\mathcal{F}$ defines an objects in $D_c^b(X, R)$.

Definition 8.6.5 (cf. [?, Proposition 10.1.17.]). (♠ TODO: AI generated, verify) (♠ TODO: cite these things) Let X be a noetherian (Definition C.0.16) scheme. Let ℓ be a prime number that is invertible on X . Let R be the integral closure (Definition C.0.17) of $\mathbb{Z}_\ell$ in a finite extension E of $\mathbb{Q}_\ell$.

(♠ TODO: Define the six functors on $D_c^b(X, R/\lambda^n)$) Let $f : X \rightarrow Y$ be a morphism between noetherian-schemes over a base scheme S . Consider sheaves (Definition 4.0.2) on

⁵Note that [?, I.I.2] uses slightly different numbering conventions, letting K_n be an object of $D^-(X, R/\lambda^n)$ instead.

the small étale sites (Definition 4.1.3) on X and Y . Take objects $K = (K_n)$ and $L = (L_n)$ of $D_c^b(X, R)$ (Definition 8.6.2) and $M = (M_n)$ of $D_c^b(Y, R)$. Define the following:

1.

$$f^*K = (f^*K_n)_n$$

(♠ TODO: definition explaining why f^* doesn't need Lf^*) (Definition 6.1.20) (Definition 6.1.10) (Definition B.9.8)

2. If f is an S -compactifiable morphism (Definition 6.2.4) over some base scheme S , define

$$Rf_!K = (Rf_!K_n)_n.$$

(Definition 6.2.4) This construction is independent of the choice of compactification due to ??.

3. For any morphism f , define

$$Rf_*K = (Rf_*K_n)_n.$$

(Definition B.9.7)

4. On a single scheme X , for $K, L \in D_c^b(X, R)$, define

$$K \overset{L}{\otimes}_R L = (K_n \overset{L}{\otimes}_{R/\lambda^{n+1}} L_n), \quad R\mathcal{H}om(K, L) = (R\mathcal{H}om)(K_n, L_n).$$

(Definition B.9.15) (Definition B.9.3)

5. If f is an S -compactifiable morphism where X and Y are S -schemes of finite type (Definition C.0.19), define

$$Rf^!K = (Rf^!K_n)_n.$$

(♠ TODO: exceptional inverse image)

When they are defined, Rf_*K and $Rf_!K$ are objects in $D_c^b(Y, R)$ and $f^*M, Rf^!M, K \overset{L}{\otimes}_R L$ and $R\mathcal{H}om(K, L)$ are objects in $D_c^b(X, R)$. $Rf^!K, K \overset{L}{\otimes}_R L$, and $R\mathcal{H}om(K, L)$ may alternatively be denoted by notations such as $f^!K, K \otimes_R L$, and $\mathbf{RHom}(K, L)$ respectively.

Proposition 8.6.6. Let $f : X \rightarrow Y$ be a morphism between noetherian schemes. Let ℓ be a prime number that is invertible on X and Y . Let R be the integral closure of $\mathbb{Z}_\ell$ in a finite extension E of $\mathbb{Q}_\ell$. Note that R is a complete discrete valuation ring and let λ be the maximal ideal of R .

1. For any $K = (K_n) \in D_c^b(Y, R)$, we have $f^*k = (f^*K_n) \in D_c^b(X, R)$.

Definition 8.6.7. Let $(R, +, \cdot)$ be a not-necessarily commutative ring (Definition C.0.8).

1. An element $a \in R$ is a **left zero-divisor** if there exists a nonzero $x \in R$ such that $ax = 0$. Otherwise, a is called **left regular** or **left cancellable**.
2. An element $a \in R$ is a **right zero-divisor** if there exists a nonzero $x \in R$ such that $xa = 0$. Otherwise, a is called **right regular** or **right cancellable**.
3. An element $a \in R$ is a **zero-divisor** if it is a left zero-divisor or a right zero-divisor.
4. An element $a \in R$ is a **two-sided zero-divisor** if it is both a left zero-divisor and a right zero-divisor.

5. An element $a \in R$ is *regular*, *cancellable*, or a *non-zero-divisor* if it is both left and right regular.

A zero-divisor of any kind that is not itself 0 is said to be a *nonzero zero divisor* or a *nontrivial zero divisor* of its kind.

Definition 8.6.8 (cf. [?, Between Corollary 10.1.21 and Lemma 10.1.22]). Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let E be the fraction field of R . Let X be a noetherian scheme (Definition C.0.16), and consider sheaves (Definition 4.0.2) on the (small) étale site on X (Definition 4.1.3).

1. On the category $D_c^b(X, R)$ (Definition 8.6.2), the morphisms defined by multiplications by λ^m for $m \geq 0$ form a multiplicative system (Definition B.2.1) (♠ TODO: why). Define the category $D_c^b(X, E)$ as the localization (Definition B.2.2) of $D_c^b(X, R)$ by this system. In particular, the objects of $D_c^b(X, E)$ are those of $D_c^b(X, R)$ and

$$\mathrm{Hom}_{D_c^b(X, R)}(K, L) \otimes_R E \cong \mathrm{Hom}_{D_c^b(X, E)}(K, L)$$

for all objects K, L of $D_c^b(X, R)$.

For example, the above notion of $D_c^b(X, E)$ is often discussed for $E = \mathbb{Q}_\ell$ where ℓ is a prime invertible on X .

2. Let K/E be some algebraic extension (Definition 8.6.9). The category $D_c^b(X, K)$ is defined as the 2-direct limit of the categories $D_c^b(X, L)$ where L run over the finite extensions of K :

$$D_c^b(X, K) = \varinjlim_{L/K \text{ finite extension}} D_c^b(X, L).$$

In other words, each object of $D_c^b(X, K)$ is represented by some object of $D_c^b(X, L)$ for some L/K and given two objects $M_1 \in \mathrm{Ob}(D_c^b(X, L_1))$ and $M_2 \in \mathrm{Ob}(D_c^b(X, L_2))$, we have

$$\mathrm{Hom}_{D_c^b(X, K)}(M_1, M_2) \cong \varinjlim_{L/K \text{ finite extension, } L_1, L_2 \subseteq L} \mathrm{Hom}(M_1 \otimes_{L_1} L, M_2 \otimes_{L_2} L).$$

(♠ TODO: define base change)

For example, the above notion of $D_c^b(X, K)$ is often discussed for $K = \overline{\mathbb{Q}}_\ell$ where ℓ is a prime invertible on X .

Definition 8.6.9 (Algebraic Element, Algebraic Extension). Let L/K be a field extension and let $x \in L$.

- If there exists a nonzero polynomial $f(t) \in K[t]$ such that $f(x) = 0$, then x is called an *algebraic element over K* . There exists a unique such monic irreducible polynomial $f(t)$, which is called the *minimal polynomial of x over K* .
- Otherwise, x is called a *transcendental element over K* .

If every $x \in L$ is algebraic over K , then L/K is called an *algebraic extension*.

Definition 8.6.10 (cf. [?, Between Corollary 10.1.21 and Lemma 10.1.22]). Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let E be the fraction field of R . Let X

be a noetherian scheme (Definition C.0.16), and consider sheaves (Definition 4.0.2) on the (small) étale site on X (Definition 4.1.3).

1. On the category $D_c^b(X, R)$ (Definition 8.6.2), the morphisms defined by multiplications by λ^m for $m \geq 0$ form a multiplicative system (Definition B.2.1) (♠ TODO: why). Define the category $D_c^b(X, E)$ as the localization (Definition B.2.2) of $D_c^b(X, R)$ by this system. In particular, the objects of $D_c^b(X, E)$ are those of $D_c^b(X, R)$ and

$$\mathrm{Hom}_{D_c^b(X, R)}(K, L) \otimes_R E \cong \mathrm{Hom}_{D_c^b(X, E)}(K, L)$$

for all objects K, L of $D_c^b(X, R)$.

For example, the above notion of $D_c^b(X, E)$ is often discussed for $E = \mathbb{Q}_\ell$ where ℓ is a prime invertible on X .

2. Let K/E be some algebraic extension (Definition 8.6.9). The category $D_c^b(X, K)$ is defined as the 2-direct limit of the categories $D_c^b(X, L)$ where L run over the finite extensions of K :

$$D_c^b(X, K) = \varinjlim_{L/K \text{ finite extension}} D_c^b(X, L).$$

In other words, each object of $D_c^b(X, K)$ is represented by some object of $D_c^b(X, L)$ for some L/K and given two objects $M_1 \in \mathrm{Ob}(D_c^b(X, L_1))$ and $M_2 \in \mathrm{Ob}(D_c^b(X, L_2))$, we have

$$\mathrm{Hom}_{D_c^b(X, K)}(M_1, M_2) \cong \varinjlim_{L/K \text{ finite extension, } L_1, L_2 \subseteq L} \mathrm{Hom}(M_1 \otimes_{L_1} L, M_2 \otimes_{L_2} L).$$

♠ TODO: define base change

For example, the above notion of $D_c^b(X, K)$ is often discussed for $K = \overline{\mathbb{Q}}_\ell$ where ℓ is a prime invertible on X .

8.7. Geometric stalks of objects of D_c^b categories.

Definition 8.7.1 (Geometric stalk of literal derived category of étale sheaves of abelian groups). Let X be a scheme (Definition .3.1). Let $x \in X$ be a point. Let $\bar{x} : \mathrm{Spec} \kappa(x)^s \rightarrow X$ be a geometric point (Definition 3.0.1) over X . Let $K \in D(X_{\mathrm{\acute{e}t}}, \mathrm{Ab})$ be an object in the derived category (Definition B.3.1) of sheaves (Definition 4.0.2) of abelian groups on $X_{\mathrm{\acute{e}t}}$ (Definition 4.1.3).

Recall that there is a stalk functor

$$\mathcal{F} \mapsto \mathcal{F}_{\bar{x}} : \mathrm{Sh}(X_{\mathrm{\acute{e}t}}, \mathrm{Ab}) \rightarrow \mathrm{Gal}(\kappa(x)^s / \kappa(x)) - \mathrm{Ab}$$

♠ TODO: category of sets with continuous group action and that it is exact. It thus induces (Proposition B.4.2) a *stalk functor*

$$K \mapsto K_{\bar{x}} : D(X_{\mathrm{\acute{e}t}}, \mathrm{Ab}) \rightarrow D(\mathrm{Gal}(\kappa(x)^s / \kappa(x)) - \mathrm{Ab})$$

♠ TODO: the exactness to the derived category of the abelian caetgory of discrete abelian groups with a continuous $\mathrm{Gal}(\kappa(x)^s / \kappa(x))$ -action. In fact, for a ring Λ , this stalk functor restricts to a functor

$$K \mapsto K_{\bar{x}} : D(X_{\mathrm{\acute{e}t}}, \Lambda) \rightarrow D(\mathrm{Gal}(\kappa(x)^s / \kappa(x)) - \Lambda - \mathrm{Mod})$$

from the derived category $D(X_{\text{ét}}, \Lambda)$ of the abelian category of Λ -modules (Definition 7.1.3) on $X_{\text{ét}}$ to the derived category of the abelian category of $\text{Gal}(\kappa(x)^s/\kappa(x)) - \Lambda - \text{Mod}$.

Definition 8.7.2 (Geometric stalk of object of $D^-(X, R)$ for adic R). Let X be a scheme (Definition .3.1) equipped with its (small) étale site (Definition 4.1.3). Let R be a (commutative) ring (Definition C.0.1) equipped with an ideal (Definition C.0.2) λ . Let $x \in X$ be a point. Let $\bar{x} : \text{Spec } \kappa(x)^s \rightarrow X$ be a geometric point (Definition 3.0.1) over X .

Given an object $K = (K_n, u_n)_{n \geq 0}$ of $D^-(X, R)$ (Definition 8.6.1), define its **stalk at $\bar{x}$** as the family $K_{\bar{x}} := (K_{n, \bar{x}}, u_{n, \bar{x}})$ where $K_{n, \bar{x}} \in D^-(\text{Gal}(\kappa(x)^s/\kappa(x)) - (R/\lambda^{n+1}) - \text{Mod})$ is the stalk (Definition 8.7.1) of K_n at $\bar{x}$ and $u_{n, \bar{x}} : K_{n+1, \bar{x}} \otimes_{R/\lambda^{n+2}}^L R/\lambda^{n+1} \cong K_{n, \bar{x}}$ (Definition B.9.4) is the morphism induced from u_n .

For every such x and $\bar{x}$, we have a functor

$$D^-(X, R) \rightarrow \text{Pro}(\text{Gal}(\kappa(x)^s/\kappa(x)) - \text{Ab})$$

In the case that X is noetherian, (R, λ) is a complete DVR such that R/λ^n is finite for all $n \geq 1$, and $K \in D_c^b(X, R)$ (Definition 8.6.2), this notion of stalk should not be mistaken for its projective limit (Definition .1.14), which is more usually referred to as the stalk of K at $\bar{x}$, see Definition 8.7.3.

Definition 8.7.3 (Geometric stalk of object of $D_c^b(X, R)$ for integral R). Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let X be a noetherian scheme (Definition C.0.16), and consider sheaves (Definition 4.0.2) on the (small) étale site on X (Definition 4.1.3). Let $x \in X$ be a point. Let $\bar{x} : \text{Spec } \kappa(x)^s \rightarrow X$ be a geometric point (Definition 3.0.1) over X .

Given an object $K = (K_n, u_n)_{n \geq 0}$ of $D_c^b(X, R)$ (Definition 8.6.2), one may speak of the stalk (Definition 8.7.2) $(K_{n, \bar{x}}, u_{n, \bar{x}})$ of K at $\bar{x}$. However, by the **stalk of K at $\bar{x}$** , one more usually means the inverse limit (Definition .1.14)

$$K_{\bar{x}} := \varprojlim_n K_{n, \bar{x}}$$

which is a R -module with a continuous $\text{Gal}(\kappa(x)^s/\kappa(x))$ -action when equipped when $K_{\bar{x}}$ is equipped with the profinite topology.

For every such x and $\bar{x}$, we have a functor

$$D_c^b(X, R) \rightarrow \text{Gal}(\kappa(x)^s/\kappa(x)) - R - \text{Mod}$$

Definition 8.7.4 (Geometric stalk of object of $D_c^b(X, E)$ for rational E). Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let E be the fraction field of R . Let X be a noetherian scheme (Definition C.0.16), and consider sheaves (Definition 4.0.2) on the (small) étale site on X (Definition 4.1.3). Let $x \in X$ be a point. Let $\bar{x} : \text{Spec } \kappa(x)^s \rightarrow X$ be a geometric point (Definition 3.0.1) over X .

Given an object $L \in D_c^b(X, E)$ (Definition 8.6.10) which is given by an object $K \in D_c^b(X, R)$ (Definition 8.6.2), define the **stalk of L at $\bar{x}$** as

$$L_{\bar{x}} := K_{\bar{x}} \otimes_R E = (\varprojlim_n K_{n, \bar{x}}) \otimes_R E$$

For every such x and $\bar{x}$, we have a functor (♠ TODO: category of vector spaces with continuous galois action)

$$D_c^b(X, E) \rightarrow \text{Gal}(\kappa(x)^s/\kappa(x)) - \text{Vec}_E$$

Definition 8.7.5 (Geometric stalk of object of $D_c^b(X, K)$ for algebraic K). Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let E be the fraction field of R . Let K be an algebraic extension of E . Let X be a noetherian scheme (Definition C.0.16), and consider sheaves (Definition 4.0.2) on the (small) étale site on X (Definition 4.1.3). Let $x \in X$ be a point. Let $\bar{x} : \text{Spec } \kappa(x)^s \rightarrow X$ be a geometric point (Definition 3.0.1) over X .

Given an object $M \in D_c^b(X, K)$ (Definition 8.6.10) which is represented by an object $L \in D_c^b(X, E')$ (Definition 8.6.10), define the *stalk of M at $\bar{x}$* as

$$M_{\bar{x}} := L_{\bar{x}} \otimes_{E'} K$$

(Definition 8.7.4) This is independent of the choice of E' and the choice of L .

For every such x and $\bar{x}$, we have a functor (♠ TODO: category of vector spaces with continuous galois action)

$$D_c^b(X, K) \rightarrow \text{Gal}(\kappa(x)^s/\kappa(x)) - \text{Vec}_K$$

9. THEOREMS

Corollary 9.0.1 (Finiteness of cohomology). (♠ TODO: read) Let X be a variety of finite type over a separably closed field k , and let $\ell \neq \text{char}(k)$. Then each $H_{\text{ét}}^i(X, \mathbb{Q}_\ell)$ is a finite-dimensional $\mathbb{Q}_\ell$ -vector space, and vanishes for $i > 2 \dim(X)$.

9.1. ℓ -adic cohomology. Deligne's ℓ -adic cohomology does not refer to étale cohomology with coefficients (Definition 7.2.2) in the rings $\mathbb{Z}_\ell$ or $\mathbb{Q}_\ell$. Rather, it takes limits of étale cohomology with coefficients in finite rings.

Definition 9.1.1. Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let X be a noetherian scheme (Definition C.0.16), and consider sheaves (Definition 4.0.2) on the (small) étale site on X (Definition 4.1.3).

Given an object M of $D^-(X, R)$ (Definition 8.6.1), the *λ -adic cohomology groups with coefficients in M* are the groups $H^q(X, M)$ defined as

$$H^q(X, M) := \varprojlim_n H^q(X, M_n)$$

(Definition 7.2.2) (Definition 4.1.13)

In particular, this notion is defined on the subcategory $D_c^b(X, R)$ (Definition 7.2.2) and on the subcategory of λ -adic sheaves (Definition 8.4.9) on X .

Definition 9.1.2. Let (R, λ) be a complete DVR such that R/λ^n is finite for all $n \geq 1$. Let X be a noetherian scheme (Definition C.0.16), and consider sheaves (Definition 4.0.2) on the (small) étale site on X (Definition 4.1.3).

1. Let E be the fraction field of R .

(♠ TODO: try to see if $D^-(X, E)$ is definable and if this notion of cohomology is definable there.) Given an object N of $D_c^b(X, E)$ (Definition 8.6.10) represented by an object M of $D_c^b(X, R)$ (Definition 7.2.2), the *(rational) λ -adic cohomology groups* $H^q(X, N)$ are defined as

$$H^q(X, N) := H^q(X, M) \otimes_R E$$

(Definition 9.1.1) (Definition .1.13)

2. Let K/E be some algebraic extension (Definition 8.6.9).

Given an object P of $D_c^b(X, K)$ (Definition 8.6.10) represented by an object N of $D_c^b(X, L)$ for a finite extension L/K , define the *(rational) λ -adic cohomology group* $H^q(X, P)$ as

$$H^q(X, P) := H^q(X, N) \otimes_L K.$$

9.2. Base change theorem.

Theorem 9.2.1 (Smooth base change theorem for étale sheaves with finite coefficients on schemes).

Proposition 9.2.2 (Smooth base change). (♠ TODO: read) Let $f : X \rightarrow S$ be a smooth, proper morphism of schemes, and let ℓ be a prime invertible on S . Then the formation of étale cohomology commutes with base change: for any cartesian square

$$\begin{array}{ccc} Y & \longrightarrow & X \\ \downarrow & & \downarrow f \\ T & \longrightarrow & S \end{array}$$

one has natural isomorphisms

$$H_{\text{ét}}^i(Y_{\bar{t}}, \mathbb{Q}_\ell) \cong H_{\text{ét}}^i(X_{\bar{s}}, \mathbb{Q}_\ell),$$

where $\bar{t}$ lies over $t \in T$, and $\bar{s}$ its image in S .

Theorem 9.2.3 (Proper base change theorem). Let $f : X \rightarrow Y$ be a proper morphism of schemes, and let ℓ be a prime invertible on Y . Then for any sheaf of $\mathbb{Z}_\ell$ -modules $\mathcal{F}$ on $X_{\text{ét}}$, the higher direct image sheaves $R^i f_* \mathcal{F}$ on $Y_{\text{ét}}$ satisfy

$$(R^i f_* \mathcal{F})_y \cong H_{\text{ét}}^i(X_y, \mathcal{F}|_{X_y})$$

for each geometric point y of Y , where X_y is the fiber over y .

Theorem 9.2.4 (Smooth and proper base change compatibility). Under suitable smooth and proper morphisms, the formation of étale cohomology commutes with base change on the target scheme. More precisely, given a cartesian square

$$\begin{array}{ccc} X' & \rightarrow & X \\ \downarrow & & \downarrow \\ Y' & \rightarrow & Y \end{array}$$

with $f : X \rightarrow Y$ smooth and proper, and a prime ℓ invertible on Y , the natural base change morphism induces isomorphisms in étale cohomology

$$H_{\text{ét}}^i(X_y, \mathbb{Q}_\ell) \cong H_{\text{ét}}^i(X'_{y'}, \mathbb{Q}_\ell)$$

for geometric points y of Y and y' over y in Y' .

9.3. Poincaré duality.

Theorem 9.3.1 (Generalized Poincaré Duality in Étale Cohomology). (♠ TODO: read this) Let k be a separably closed field, and let X be a smooth, separated scheme of finite type over k , of pure dimension d . Let ℓ be a prime different from $\text{char}(k)$.

Let Λ be one of the following coefficient rings:

- a finite extension of $\mathbb{Z}_\ell$,
- the ring of integers $\mathcal{O}_E$ of a finite extension E of $\mathbb{Q}_\ell$,
- E itself, or
- an algebraic closure $\overline{\mathbb{Q}_\ell}$.

Let $\mathcal{F}^\bullet$ be a bounded constructible complex of Λ -modules on the étale site $X_{\text{ét}}$.

Then there is a canonical perfect pairing in the derived category of Λ -modules:

$$R\Gamma_{\text{ét}}(X, \mathcal{F}^\bullet) \otimes_{\Lambda}^{\mathbf{L}} R\Gamma_c^{\text{ét}}(X, R\mathcal{H}om_{\Lambda}(\mathcal{F}^\bullet, \Lambda(d)[2d])) \longrightarrow \Lambda[-2d],$$

where $R\Gamma_c^{\text{ét}}$ denotes étale cohomology with compact support, and

$$\Lambda(d) := \underbrace{\Lambda(1) \otimes_{\Lambda} \cdots \otimes_{\Lambda} \Lambda(1)}_{d \text{ times}}$$

is the d -th Tate twist.

In particular, this induces a perfect duality between finite generated Λ -modules

$$H_{\text{ét}}^i(X, \mathcal{F}^\bullet) \cong \text{Hom}_{\Lambda}(H_c^{2d-i}(X, R\mathcal{H}om_{\Lambda}(\mathcal{F}^\bullet, \Lambda(d)[2d])), \Lambda).$$

Remarks:

- For projective X , $R\Gamma_c^{\text{ét}}$ can be replaced by the usual cohomology $R\Gamma_{\text{ét}}$.
- The duality is a manifestation of Verdier duality and requires the formalism of the six operations and dualizing complexes in étale cohomology.
- This generalizes classical Poincaré duality by allowing more general coefficient complexes and non-projective varieties with compact support.

9.4. Lefschetz trace formula.

Theorem 9.4.1 (Lefschetz trace formula in étale cohomology). (♠ TODO: read) Let X be a separated scheme of finite type over a finite field $\mathbb{F}_q$, and let ℓ be a prime number

not dividing q . Then the number of $\mathbb{F}_q$ -rational points of X is given by the Lefschetz trace formula:

$$|X(\mathbb{F}_q)| = \sum_{i=0}^{2 \dim(X)} (-1)^i \operatorname{Tr}(\operatorname{Frob}_q^* | H_{\text{ét}}^i(X_{\overline{\mathbb{F}}_q}, \mathbb{Q}_\ell)),$$

where Frob_q is the geometric Frobenius acting on the ℓ -adic étale cohomology groups of X .

Theorem 9.4.2 (Grothendieck, see [?, Rapport 3 Théorème 3.1]; see also [?, Theorem 10.5.1] for a statement and cf. [?, Theorem I.1.1]). Let X_0 be a separated scheme (Definition 3.1) of finite type (Definition C.0.19) over the finite field $\mathbb{F}_q$, and let $K_0 \in D_c^b(X_0, \overline{\mathbb{Q}}_\ell)$ (Definition 9.1.2). Write $X = X_0 \times_{\mathbb{F}_q} \overline{\mathbb{F}}_q$ for the base change, and K for the pullback of K_0 to X .

We have

$$L(X_0, K_0, t) = \prod_{i \in \mathbb{Z}} \det(1 - t \cdot \operatorname{Frob}_q | H_c^i(X, K))^{(-1)^{i+1}}.$$

(??) where Frob_q is the geometric Frobenius endomorphism

Theorem 9.4.3 (Grothendieck–Lefschetz trace formula in terms of traces). Let X_0 be a separated scheme of finite type over a finite field $\mathbb{F}_q$. Denote by $\overline{X} = X_0 \times_{\mathbb{F}_q} \overline{\mathbb{F}}_q$ the base change to an algebraic closure, and let $K_0 \in D_c^b(X_0, \overline{\mathbb{Q}}_\ell)$ be a bounded constructible complex of étale sheaves.

Then the number of fixed points of the n -th iterate of the geometric Frobenius on X_0 , weighted by the trace of the induced endomorphism on stalks of K_0 , satisfies the formula

$$\sum_{x \in X_0(\mathbb{F}_{q^n})} \operatorname{tr}((\operatorname{Frob}_{q^n})_x | K_{0,\overline{x}}) = \sum_{i \in \mathbb{Z}} (-1)^i \operatorname{tr}(\operatorname{Frob}_{q^n} | H_c^i(\overline{X}, K)),$$

where K denotes the pullback of K_0 to $\overline{X}$, and $(\operatorname{Frob}_{q^n})_x$ is the induced action on the stalk of K_0 at a geometric point over x .

In particular, this expresses the weighted count of $\mathbb{F}_{q^n}$ -rational points on X_0 in terms of traces of Frobenius on étale cohomology with compact support, encapsulating the Grothendieck–Lefschetz trace formula in terms of traces rather than directly using determinant or Frobenius polynomials.

9.5. Weil conjectures.

Theorem 9.5.1 (Weil Conjectures via Étale Cohomology). Let X be a smooth projective variety of pure dimension d defined over a finite field $\mathbb{F}_q$. Let ℓ be a prime number different from the characteristic of $\mathbb{F}_q$. Let F denote the geometric Frobenius endomorphism acting on the étale cohomology groups $H_{\text{ét}}^i(\overline{X}, \mathbb{Q}_\ell)$, where $\overline{X} = X \times_{\mathbb{F}_q} \overline{\mathbb{F}}_q$.

Then the Weil conjectures state the following:

1. (Rationality) The zeta function

$$Z(X, t) = \exp \left(\sum_{n=1}^{\infty} \frac{\#X(\mathbb{F}_{q^n})}{n} t^n \right)$$

is a rational function which can be expressed as

$$Z(X, t) = \prod_{i=0}^{2d} P_i(t)^{(-1)^{i+1}},$$

where each $P_i(t) = \det(1 - tF \mid H_{\text{ét}}^i(\overline{X}, \mathbb{Q}_\ell))$ is a polynomial with coefficients in $\mathbb{Q}_\ell$.

2. (Functional Equation) The zeta function satisfies a functional equation relating $Z(X, t)$ and $Z(X, q^{-d}t^{-1})$, induced by Poincaré duality on cohomology and the action of F .
3. (Betti Numbers) Each polynomial $P_i(t)$ has degree equal to the i -th Betti number of X (i.e., $\dim_{\mathbb{Q}_\ell} H_{\text{ét}}^i(\overline{X}, \mathbb{Q}_\ell)$).
4. (Riemann Hypothesis) The eigenvalues of the geometric Frobenius F acting on $H_{\text{ét}}^i(\overline{X}, \mathbb{Q}_\ell)$ are algebraic numbers, all of whose complex absolute values equal $q^{i/2}$.

This theorem links the counting of points over finite fields to the action of Frobenius on étale cohomology and provides a cohomological interpretation and proof of the Weil conjectures.

Here is another, more general, formulation of the rationality statement of the Weil conjectures:

Theorem 9.5.2 (Grothendieck, see [?, Rapport 3 Théorème 3.1]; see also [?, Theorem 10.5.1] for a statement and cf. [?, Theorem I.1.1]). Let X_0 be a separated scheme (Definition .3.1) of finite type (Definition C.0.19) over the finite field $\mathbb{F}_q$, and let $K_0 \in D_c^b(X_0, \overline{\mathbb{Q}}_\ell)$ (Definition 9.1.2). Write $X = X_0 \times_{\mathbb{F}_q} \overline{\mathbb{F}}_q$ for the base change, and K for the pullback of K_0 to X .

We have

$$L(X_0, K_0, t) = \prod_{i \in \mathbb{Z}} \det(1 - t \cdot \text{Frob}_q \mid H_c^i(X, K))^{(-1)^{i+1}}.$$

(??) where Frob_q is the geometric Frobenius endomorphism

10. EKEDAHN'S FORMALISM

Deligne [?] defined the ℓ -adic category $D_c^b(X, \mathbb{Z}_\ell)$ (from which the categories $D_c^b(X, \mathbb{Q}_\ell)$ and $D_c^b(X, \overline{\mathbb{Q}}_\ell)$ were produced), intuitively but not precisely thought of as being like a derived category of ℓ -adic sheaves, by taking a 2-limit of derived categories $D_c^b(X, \mathbb{Z}/\ell^n \mathbb{Z})$. In particular, the objects of $D_c^b(X, \mathbb{Z}_\ell)$ are inverse systems of objects coming from various derived categories. Starting from a general ringed topos (S, R) , Ekedahl [?] produced an adic-formalism that is based upon taking literal complexes of sheaves of modules on a ringed topos $(\text{Pro}_{\mathbb{N}}(S), R_\bullet)$ over (S, R) defined on the topos $\text{Pro}_{\mathbb{N}}(S)$ of inverse systems in S , and which is itself equipped with an “adic ring” structure $R_\bullet$. Ekedahl demonstrated that his categories $D_{\text{Ek}}^*(S, R)$ (Definition 10.0.17), under nice circumstances, satisfy nice properties (Theorem 10.0.20) that Deligne’s categories ought to satisfy. Bhatt and Scholze [?] unified Ekedahl’s framework with Deligne’s construction.

Definition 10.0.1 (Categories of Inverse and Direct Sequences of objects in a category). Let $\mathcal{C}$ be a category (Definition .1.1).

1. The *category of inverse sequences* or *Pro-category indexed by $\mathbb{N}$* , denoted by $\text{Pro}_{\mathbb{N}}(\mathcal{C})$ or $\mathcal{C}^{\mathbb{N}}$ (following notation for functor categories (Definition .1.33)), is defined as follows:
 - Objects are sequences $(M_n, p_n)_{n \geq 1}$ where for each $n \geq 1$, $M_n \in \text{Ob}(\mathcal{C})$ and $p_n : M_{n+1} \rightarrow M_n$ is a morphism in $\mathcal{C}$.
 - A morphism $f : (M_n, p_n) \rightarrow (N_n, q_n)$ consists of a collection of morphisms $\{f_n : M_n \rightarrow N_n\}_{n \geq 1}$ in $\mathcal{C}$ satisfying the compatibility condition

$$\forall n \geq 1, \quad q_n \circ f_{n+1} = f_n \circ p_n.$$

Composition and identities are defined componentwise.

The notions of inverse sequences and their morphisms are equivalent to those of towers (Definition 8.4.2) up to indexing.

2. The *category of direct sequences* or *Ind-category indexed by $\mathbb{N}$* , denoted by $\text{Ind}_{\mathbb{N}}(\mathcal{C})$, is defined as follows:
 - Objects are sequences $(M_n, i_n)_{n \geq 1}$ where for each $n \geq 1$, $M_n \in \text{Ob}(\mathcal{C})$ and $i_n : M_n \rightarrow M_{n+1}$ is a morphism in $\mathcal{C}$.
 - A morphism $f : (M_n, i_n) \rightarrow (N_n, j_n)$ consists of a collection of morphisms $\{f_n : M_n \rightarrow N_n\}_{n \geq 1}$ in $\mathcal{C}$ satisfying the compatibility condition

$$\forall n \geq 1, \quad j_n \circ f_n = f_{n+1} \circ i_n.$$

Composition and identities are defined componentwise.

Equivalently, $\text{Pro}_{\mathbb{N}}(\mathcal{C})$ and $\text{Ind}_{\mathbb{N}}(\mathcal{C})$ may be regarded as the functor categories $\text{Fun}(\mathbb{N}^{\text{op}}, \mathcal{C})$ and $\text{Fun}(\mathbb{N}, \mathcal{C})$ respectively, where $\mathbb{N}$ is regarded (Lemma .1.29) as a poset (Definition .1.28) category with objects $1, 2, \dots$ with a unique morphism $n \rightarrow m$ if and only if $n \leq m$.

Context 10.0.2 ([?, Section 1.]). Let (S, R) be a ringed topos and let $m \subseteq R$ be a two-sided ideal. Let $\text{Pro}_{\mathbb{N}}(S)$ be the category of inverse sequences of S (Definition 10.0.1); this is equivalent to the functor category $[\mathbb{N}^{\text{op}}, S]$ (Definition .1.33) and is a topos (Theorem .1.32) itself. Regard it as a ringed topos equipped with $R_{\bullet}$ (Definition 10.0.5).

Context 10.0.3 ([?, Section 1.]). Let S be a topos (Definition .1.31). Write $\pi : \text{Pro}_{\mathbb{N}}(S) \rightarrow S$ (Definition 10.0.1) for the topos morphism given by $\pi_* M_{\bullet} = \varprojlim M_{\bullet}$ and $\pi^* M = \{M, \text{id}\}_{n \geq 1}$, i.e. the system consisting of the objects M at each level and the identity maps between them. For each $n > 0$, further write $i_n : S \rightarrow S^{\mathbb{N}}$ for the topos morphism satisfying $i_n^* M_{\bullet} = M_n$.

Context 10.0.4 ([?, Section 1.]). Assume Context 10.0.2 and Context 10.0.3.

Assume for every n there exists, locally on S , a resolution (Definition .2.7) $F_n^{\bullet} \rightarrow R/m^n$ of R/m^n by a complex of right R -modules which is finitely generated free in each degree and such that modulo essentially zero systems (Definition 10.0.6) $\pi^* F_n^{\bullet} \rightarrow \pi^*(R/m^n)$ is a resolution.

For example, this is satisfied when R is the constant (Definition C.0.10) sheaf of a noetherian, commutative, and local ring whose maximal ideal is m . Moreover, if this assumption is true of (R, m) , then it is also true of $(R[G], m[G])$ where G is a group object in S .

This assumption is assumed throughout [?]

Definition 10.0.5 ([?, Section 1.]). Assume Context 10.0.3. There is a ring object, which we might sometimes denote by $R_\bullet$, given by the projective system

$$\cdots \rightarrow R/m^{n+1} \rightarrow R/m^n \rightarrow \cdots$$

When $\text{Pro}_{\mathbb{N}}(S)$ (Definition 10.0.1) is equipped with $R_\bullet$, π is a morphism of ringed topoi. In particular, for an object M of the ringed topos (Definition .1.31) (S, R) , its pullback π^*M is the inverse system $\{M \otimes_R R_n\}_{n \geq 1}$.

Definition 10.0.6 ([?, Section 1.]). Assume Context 10.0.2

1. We say that an abelian group object $M_\bullet$ of $\text{Pro}_{\mathbb{N}}(S)$ (Definition 10.0.1) is **essentially zero** if there is a covering of the final object (Definition .1.10) of S such that, when $M_\bullet$ is restricted to each element of the covering, there is for all n an $m \geq n$ such that the map $M_m \rightarrow M_n$ in $M_\bullet$ is zero.
2. Let $M^\bullet$ be a complex (Definition .2.1) of abelian group objects of $\text{Pro}_{\mathbb{N}}(S)$. We say that $M^\bullet$ is **essentially zero** if $H^i(M^\bullet)$ (Definition .2.3) is essentially zero for all i .

The category of essentially zero complexes form a triangulated subcategory (Definition B.1.2) of the category of complexes of abelian group objects of $\text{Pro}_{\mathbb{N}}(S)$.

One may say that $M^\bullet$ is **essentially bounded**, **essentially bounded from below**, or **essentially bounded from above**, if $H^m(M^\bullet)$ is essentially zero for all m with $|m| \gg 0$, $-m \gg 0$, and $m \gg 0$ respectively.

The (derived) categories (Definition B.3.1) of essentially bounded, essentially bounded from below, and essentially bounded from above complexes of R -module objects of $\text{Pro}_{\mathbb{N}}(S)$ may be denoted by $D^{eb}(\text{Pro}_{\mathbb{N}}(S), R_\bullet)$, $D^{e+}(\text{Pro}_{\mathbb{N}}(S), R_\bullet)$, and $D^{e-}(\text{Pro}_{\mathbb{N}}(S), R_\bullet)$ respectively. $[?]$ denotes these as $D^{eb}(S^{\mathbb{N}} - R_\bullet)$, etc. Moreover, $[?]$ denotes $D^{eb}(S^{\mathbb{N}} - \text{ab})$ etc. in the case that $R = \mathbb{Z}$.

Notation 10.0.7. Let (S, R) be a ringed topos. Write $D(S, R)$ for the derived category (Definition B.3.1) of the category of R -modules (Definition 7.1.3). Accordingly, $D^+(S, R)$, $D^-(S, R)$, and $D^b(S, R)$ may be used to denote the full subcategories of cohomologically bounded below, cohomologically bounded above, and cohomologically bounded complexes (Definition .2.2) respectively, cf. Definition B.3.1

Definition 10.0.8 ([?, Section 1]). Assume Context 10.0.3.

1. We may say that (S, R) **satisfies Ekedahl's condition A** if the following holds: there exists a class S^{gen} of generators of S (Definition .1.30) and an integer N such that for all $T \in S^{\text{gen}}$, all R/m -modules (Definition 7.1.3) M , and $i > N$, we have $H^i(T, M) = 0$ (Definition 7.1.10).
2. We may say that (S, R) **satisfies Ekedahl's condition B** if the following hold:
 - there is an integer N such that for every $n \geq 1$ and locally on $(\mathcal{C}, J)$, there is a left resolution (Definition .2.7) $F_n^\bullet \rightarrow R/m^n$ of R/m^n by a complex of (right) R -modules of finitely generated free objects such that $\pi^*F_n^\bullet \rightarrow \pi^*(R/m^n)$ is a resolution modulo essentially zero systems (Definition 10.0.6), where $\pi : (\text{Pro}_{\mathbb{N}}(S), R_\bullet) \rightarrow (S, R)$ is considered as a morphism of ringed topoi.
 - m^n/m^{n+1} is, locally on $(\mathcal{C}, J)$, of finite Tor-dimension over R/m . (♠ **TODO: finite Tor dimension**)

Here are sufficient conditions for condition A to be fulfilled:

1. When the topos is that of the small étale site (Definition 4.1.3) $X_{\text{ét}}$, where X is the spectrum of an algebraically closed field, a local field, a global field, or $\mathbb{Z}$, and R is a constant ring (Definition C.0.10) $X_{\text{ét}}$.
2. Whenever condition A holds for $X_{\text{ét}}$ for a scheme (Definition .3.1) X , a constant ring R on $X_{\text{ét}}$, and a two sided ideal $m \subseteq R$, it also holds for $(Y_{\text{ét}}, R, m)$ for any scheme Y of finite type (Definition C.0.19) over X .

Here are sufficient conditions for condition B to be fulfilled:

1. R is a constant ring that is noetherian, commutative (Definition C.0.1), and local (Definition C.0.21) of finite global dimension.
2. Whenever condition A holds for (S, R, m) , it also holds for $(S, R[G], m[G])$ for a group G on S .

Lemma 10.0.9 ([?, Lemma 1.1.]). Assume Context 10.0.2 and Context 10.0.3. Assume Context 10.0.4

Let M be a cohomologically bounded below (Definition .2.2) cochain complex of abelian group objects of $\text{Pro}_{\mathbb{N}}(S)$ (Definition 10.0.1). Then, $R\pi_*M = 0$ (Context 10.0.3) (Definition B.9.7).

Definition 10.0.10 ([?, Between Lemma 1.1. and Lemma 1.3.]). Assume Context 10.0.2.

Let M be a cochain complex of abelian group objects of $\text{Pro}_{\mathbb{N}}(S)$ (Definition 10.0.1). We say that M is *essentially constant* if there exists a complex N of abelian group objects in $\text{Pro}_{\mathbb{N}}(S)$, a complex P of abelian group objects in S , and morphism of complexes (Definition .2.1) $M \xleftarrow{f} N \xrightarrow{g} \pi^*P$ such that the mapping cones (Definition .2.5) of f and g are essentially zero (Definition 10.0.6).

Notation 10.0.11 ([?, After Lemma 1.1]). Assume Context 10.0.2.

Given $M \in D^{e+}(\text{Pro}_{\mathbb{N}}(S), \mathbb{Z}_{\bullet})$ (Definition 10.0.6), [?] lets $\tau(M)$ denote the following:

- If $(\mathcal{C}, J, R)$ satisfies Ekedahl's condition A (Definition 10.0.8), then $\tau(M) = M$, regarded as the constant pro-object (Definition B.5.1).
- Otherwise, $\tau(M)$ is the pro-object (Definition B.5.1)

$$(\cdots \rightarrow \tau_{\geq i-1}M \rightarrow \tau_{\geq i}M \rightarrow \tau_{\geq i+1}M \rightarrow \cdots)$$

(Definition .2.6).

Definition 10.0.12 ([?, After Lemma 1.1.]). Assume Context 10.0.2. let $M \in D^{e+}(\text{Pro}_{\mathbb{N}}(S), \mathbb{Z}_{\bullet})$ (Definition 10.0.6).

In either case, the pro-object $R\pi_*\tau(M)$ obtained by applying $R\pi_*$ (Context 10.0.3) (Definition B.9.7) to $\tau(M)$ (Notation 10.0.11) termwise is essentially constant (Definition 10.0.10) by Lemma 10.0.9. By abuse of language let $R\pi_*M$ denote any limit of this pro-system (Definition .1.18). The functor $R\pi_*$ is a triangulated functor (Definition B.1.3)

$$D^{e+}(\text{Pro}_{\mathbb{N}}(S), R_{\bullet}) \rightarrow D^+(S, R),$$

where $D^+(S, R)$ is the derived category of modules of the ringed topos (S, R) as per Notation 10.0.7.

Context 10.0.13 ([?, Start of Section 2.]). Assume Context 10.0.2 and Context 10.0.3. Assume that Ekedahl's condition A or B holds (Definition 10.0.8). Let

$$* = \begin{cases} - & \text{if Ekedahl's condition A holds} \\ e+ & \text{if Ekedahl's condition B holds} \\ (\text{blank}) & \text{if both Ekedahl's condition A and condition B hold.} \end{cases}$$

In particular, we will consider the category $D^-(\mathrm{Pro}_{\mathbb{N}}, R_{\bullet})$ (Notation 10.0.7), $D^{e+}(\mathrm{Pro}_{\mathbb{N}}, R_{\bullet})$ (Definition 10.0.6), or $D(\mathrm{Pro}_{\mathbb{N}}, R_{\bullet})$ (Notation 10.0.7), depending on which combination of A and B holds.

Definition 10.0.14 ([?, Start of Section 2.]). Assume Context 10.0.13 and let $M \in D^*(\mathrm{Pro}_{\mathbb{N}}, R_{\bullet})$.

Define $\widehat{M} := L\pi^*(R\pi_*M)$ (Context 10.0.3) (Definition B.9.8) (Definition 10.0.10), where

- $R\pi_*$ is given by
 - the literal derived direct image functor of π_* (Context 10.0.3) if Ekedahl's condition A (Definition 10.0.8) holds
 - Definition 10.0.12 if Ekedahl's condition A (Definition 10.0.8) does not hold (so in particular Ekedahl's condition B (Definition 10.0.8) holds)
- $L\pi^*$ is the derived inverse image functor (Definition B.9.8) of π^* .

In any case, $\widehat{-}$ is a functor

$$D^*(\mathrm{Pro}_{\mathbb{N}}, R_{\bullet}) \rightarrow D(\mathrm{Pro}_{\mathbb{N}}, R_{\bullet})$$

where the source is either a literal derived category as per Notation 10.0.7 or a category from Definition 10.0.6, depending on whether Ekedahl's condition A holds, and the target is always a literal derived category.

Moreover, in any case, there is a morphism $\widehat{M} \rightarrow M$ — if Ekedahl's condition A holds, then it is given by the adjunction of (Proposition B.7.4) $L\pi^*$ and $R\pi_*$, and otherwise, it is induced by morphisms $\widehat{M} \rightarrow \tau_{\geq i}M$.

Definition 10.0.15 ([?, Definition 2.1]). Assume Context 10.0.3

Let M, N be objects $D(\mathrm{Pro}_{\mathbb{N}}(S), R_{\bullet})$ (Notation 10.0.7).

1. We may say that M is an ***R-complex*** if $\pi^*(R_1) \otimes_{R_{\bullet}}^L M$ (Context 10.0.3) (Definition B.9.4) is essentially constant (Definition 10.0.10)
2. M is ***negligible*** if $\pi^*(R_1) \otimes_{R_{\bullet}}^L M$ is essentially zero (Definition 10.0.6).
3. M is ***normalised*** if the morphism $\tau(\widehat{M}) \rightarrow \tau(M)$ (Notation 10.0.11) (Definition 10.0.14) is a quasi-isomorphism (Definition .2.4).
4. A morphism $M \rightarrow N$ is ***essentially an isomorphism*** if it has a negligible mapping cone (Definition .2.5).

Proposition 10.0.16 ([?, Proposition 2.2]). Assume Context 10.0.2, Context 10.0.3, Context 10.0.13, and Context 10.0.4

1. M is an R -complex (Definition 10.0.15) if and only if $\tau(\widehat{M}) \rightarrow \tau(M)$ (Notation 10.0.11) (Definition 10.0.14) is essentially an isomorphism (Definition 10.0.15).
2. M is normalised (Definition 10.0.15) if and only if $R_n \otimes_{R_{n+1}}^L i_{n+1}^* M \rightarrow i_n^* M$ (Definition B.9.4) (Context 10.0.3) is an isomorphism for all n .
3. Let $?$ be $-$, $+$, or blank if Ekedahl's condition A, condition B, or conditions A and B hold respectively. Let $M' \in D^?(S, R)$ (Notation 10.0.7). The object $L\pi^* M'$ (Definition B.9.8) (Context 10.0.3) belongs to $D^?(Pro_{\mathbb{N}}, R_{\bullet})$. In particular, $\widehat{M}$ is normalised.

Definition 10.0.17 (cf. [?, Definition 2.5]). Assume Context 10.0.3.

Denote by $D_{\text{Ek}}(S, R)$ or $D_{\text{Ek}}(\mathcal{C}, R)$ the category whose objects are R -complexes (Definition 10.0.15) (and hence in particular objects of $D(Pro_{\mathbb{N}}(S), R_{\bullet})$ (Notation 10.0.7)) and morphisms are the morphisms $\tau(M) \rightarrow \tau(N)$ (Notation 10.0.11) with essential isomorphisms (Definition 10.0.6) inverted. For $*$ $\in \{b, +, -\}$, Denote by $D_{\text{Ek}}^*(S, R)$ or $D_{\text{Ek}}^*(\mathcal{C}, R)$ the full subcategory whose objects are the complexes which are essentially bounded, essentially bounded from below, and essentially bounded from above (Definition 10.0.6) respectively.

[?] denotes these categories by $D(S - R)$, etc.

It is appropriate to call these categories “*Ekedahl's adic categories*”. By abuse of language in conflict with the terminology of Definition 10.0.15, [?] calls the objects of $D_{\text{Ek}}(S, R)$ *R -complexes*.

A priori, there are formal size issues concerning the local smallness of these categories; such issues can be resolved under suitable circumstances. (♠ TODO: probably 2.6, 2.7, 3.6 from the paper)

Definition 10.0.18 ([?, Definition-Lemma 2.6.]). Assume Context 10.0.13. As per Notation 10.0.7, write $D^*(Pro_{\mathbb{N}}(S), R_{\bullet})$ (for $*$ $\in \{+, -, b\}$) and $D(S, R/m^n R)$ respectively for the appropriate derived categories (Definition B.3.1) of modules (Definition 7.1.3) of the ringed topoi $(Pro(S^{\mathbb{N}}), R_{\bullet})$ and $(S, R/m^n R)$

1. The functor

$$R/m^n R \otimes_R^L R\pi_*(-) : D^*(Pro_{\mathbb{N}}(S), R_{\bullet}) \rightarrow D(S, R/m^n R)$$

(Context 10.0.2) ,

$$R\pi_* : D(Pro_{\mathbb{N}}(S), R_{\bullet}) \rightarrow D(S, R)$$

is the right derived direct image (Definition B.9.7) of $\pi : (Pro_{\mathbb{N}}(S), R_{\bullet}) \rightarrow (S, R)$, and

$$R/m^n R \otimes_R^L - : D(S, R) \rightarrow D(S, R/m^n R)$$

is the left derived tensor product (Definition B.9.4) by $R/m^n R$, factors through $D_{\text{Ek}}(S, R)$ (Definition 10.0.17) to yield a functor denoted by

$$R/m^n R \otimes_R^L - : D_{\text{Ek}}(S, R) \rightarrow D(S, R/m^n R)$$

2. The functor

$$(\widehat{-}) : D^*(Pro_{\mathbb{N}}(S), R_{\bullet}) \rightarrow D(Pro_{\mathbb{N}}(S), R_{\bullet})$$

(Definition 10.0.14) factors through $D_{\text{Ek}}(S, R)$ (Definition 10.0.17) to yield a functor $D_{\text{Ek}}(S, R) \rightarrow D(Pro_{\mathbb{N}}(S), R_{\bullet})$ which will also be denoted by $(\widehat{-})$.

Theorem 10.0.19 ([?, Proposition 2.7]). Assume Context 10.0.2, Context 10.0.3, and Context 10.0.4.

Let $* \in \{b, +, -\}$.

1. $R/mR \otimes_R^L (-) : D_{\text{Ek}}(S, R) \rightarrow D(S, R_1)$ (Definition 10.0.18) (Context 10.0.2, Definition 10.0.17, Notation 10.0.7) is conservative. (♠ TODO: conservative functor)
2. Letting $D_{\text{norm}}^*(\text{Pro}_{\mathbb{N}}(S), R_{\bullet}) \subset D^*(\text{Pro}_{\mathbb{N}}(S), R_{\bullet})$ (Definition 10.0.1, Notation 10.0.7) be the subcategory of normalised R -complexes (Definition 10.0.15). The canonical functor

$$D_{\text{norm}}^*(\text{Pro}_{\mathbb{N}}(S), R_{\bullet}) \rightarrow D_{\text{Ek}}^*(S - R)$$

(♠ TODO: What is this canonical functor) and the functor

$$(\widehat{-}) : D_{\text{Ek}}^*(S - R) \rightarrow D_{\text{norm}}^*(\text{Pro}_{\mathbb{N}}(S), R_{\bullet})$$

(Definition 10.0.18) (♠ TODO: how does the normalization functor map into the normalised complex category?) are inverse equivalences of categories (Definition .1.6).

Theorem 10.0.20 ([?, Theorem 6.3]). (♠ TODO: regular ring, scheme) Let S be a regular scheme of dimension (Definition C.0.20) at most 1. Let R be a commutative (Definition C.0.1), local (Definition C.0.21), regular ring with maximal ideal m such that the residue field $R_1 = R/m$ is of positive characteristic invertible in $\mathcal{O}_s$. Write R also for the constant sheaf (Definition C.0.10) of R on $S_{\text{ét}}$ (Definition 4.1.3). Let c be the full subcategory of the category of sheaves of R_1 -modules (Definition 7.1.3) whose objects are constructible (Definition 8.3.1).

Let X, Y be S -schemes

1. The category (♠ TODO: Ekedahl's constructible, bounded category) $D_c^b(X_{\text{ét}}, R)$ and the functor

$$R_1 \otimes_R^L (-) : D_c^b(X_{\text{ét}}, R) \rightarrow D_c^b(X_{\text{ét}}, R_1)$$

is a conservative triangulated functor (♠ TODO: conservative triangulated functor)

2. $D_c^b(X_{\text{ét}}, R)$ has a t -structure (♠ TODO:) whose heart is equivalent to the category of m -adic constructible sheaves and for which every object of $D_c^b(X_{\text{ét}}, R)$ has finite amplitude, i.e. $D_c^b(X_{\text{ét}}, R) = \bigcup_{i \in \mathbb{Z}} D_c^b(X_{\text{ét}}, R)^{\geq i} = \bigcup_{i \in \mathbb{Z}} D_c^b(X_{\text{ét}}, R)^{\leq i}$.
3. (♠ TODO: tensor and RHom)
4. (♠ TODO: pushforward, pullback)
5. (♠ TODO: perverse sheaves)

11. PRO-ÉTALE SHEAVES ON SCHEMES

11.1. Pro-étale topology on a scheme.

Definition 11.1.1 (Diagonal morphism of a morphism of schemes). Let $f : X \rightarrow Y$ be a morphism of schemes (Definition .3.2).

The *diagonal morphism associated to f* is the morphism $\Delta_f : X \rightarrow X \times_Y X$ (Definition C.0.15) which is the diagonal morphism associated to the morphism f in the category of schemes.

In other words, Δ_f is defined as the unique morphism induced by the universal property of the fiber product (Definition C.0.15) making the following diagram commute:

$$\begin{array}{ccccc}
 X, & & \xrightarrow{\text{id}_X} & & X \\
 & \searrow \Delta_f & & \nearrow p_2 & \\
 & X \times_Y X & \xrightarrow{p_2} & X & \\
 & \downarrow p_1 & & \downarrow f & \\
 & X & \xrightarrow{f} & Y &
 \end{array}$$

(Note: The diagram above is a simplified representation of the commutative diagram in the image. The top-left node is X , the top-right node is X , the middle-left node is $X \times_Y X$, the middle-right node is X , the bottom-left node is X , and the bottom-right node is Y . Arrows: $X \xrightarrow{\text{id}_X} X$ (top), $X \xrightarrow{\Delta_f} X \times_Y X$ (diagonal down-right), $X \times_Y X \xrightarrow{p_2} X$ (diagonal up-right), $X \times_Y X \xrightarrow{p_1} X$ (vertical down), $X \xrightarrow{f} Y$ (bottom), $X \xrightarrow{\text{id}_X} X$ (curved arrow from top-left to middle-right), $X \xrightarrow{f} Y$ (curved arrow from middle-left to bottom-right).

where p_1 and p_2 are the natural projections from the fiber product.

In other words, Δ_f is given by the pair of identity morphisms $(\text{id}_X, \text{id}_X)$ over Y :

$$\Delta_f := (\text{id}_X, \text{id}_X) : X \rightarrow X \times_Y X.$$

Definition 11.1.2 (Weakly étale morphism of schemes, [?, Definition 2.2]). A morphism $f : X \rightarrow Y$ of schemes (Definition .3.2) is *weakly étale* if it and its diagonal (Definition 11.1.1) are both flat (Definition 1.1.4).

Definition 11.1.3 (Pro-étale site of a scheme, [?, Definition 1.2]). Let X be a scheme (Definition .3.1). The *pro-étale site* $X_{\text{proét}}$ is the site whose underlying category is the category of weakly étale (Definition 11.1.2) X -schemes and whose covers are covers in the fpqc topology (Definition 4.2.8), i.e. a family $\{Y_i \rightarrow Y\}$ of maps on $X_{\text{proét}}$ is a covering family if any open affine in Y is mapped onto by an open affine in $\coprod_i Y_i$.

11.2. Adically complete derived modules in replete topoi.

Definition 11.2.1 ([?, Definition 3.5.2.]). (♠ TODO: ring on a topos) Let $\mathcal{X}$ be a topos (Definition .1.31), let R be a ring on $\mathcal{X}$, and let $\mathfrak{m} \subseteq R$ be a two-sided ideal (Definition 7.1.6). Let $\widehat{R} = \varprojlim_n R/\mathfrak{m}^n$ be the projective limit (Definition .1.14) of the sheaves $R/\mathfrak{m}^n$.

An object $K \in D(\mathcal{X}, \widehat{R})$ (Notation B.9.1) is called *$\mathfrak{m}$ -adically complete* if the natural map $K \rightarrow R \varprojlim_n K \otimes_R^L R/\mathfrak{m}^n$ (Definition B.5.8) (Definition B.9.4) (♠ TODO: justify the existence of this functor) is an equivalence.

Let $i : D_{\text{comp}}(\mathcal{X}, \widehat{R}) \hookrightarrow D(\mathcal{X}, \widehat{R})$ denote the full subcategory of such complexes.

11.3. Realizing the left-completion of the derived category on the étale site via the pro-étale site.

Definition 11.3.1 ([?, Definition 5.3.1.]). Let X be a scheme. We denote by $D_{\text{cc}}(X_{\text{proét}})$ the full subcategory of $D(X_{\text{proét}})$ (Notation 6.2.2) spanned by complexes whose cohomology sheaves (Definition .2.3)

11.3.1. Comparison to Ekedahl's theory.

Definition 11.3.2 ([?, Definition 5.5.1.]). (♠ TODO: complete ring with respect to ideal) Let R be a commutative (Definition C.0.1) noetherian ring complete for the topology (Definition C.0.6) defined by an ideal (Definition C.0.2) $\mathfrak{m} \subset R$.

We say that $(R, \mathfrak{m})$ satisfy *Bhatt and Scholze's Ekedahl's condition B* if the following holds:

(♠ TODO: regular ideal of a ring) $\mathfrak{m}$ is a regular ideal of R and the $R/\mathfrak{m}$ -module $\mathfrak{m}^n/\mathfrak{m}^{n+1}$ has finite flat dimension bounded independently of n .

When considering R and $\mathfrak{m}$ as constant (Definition C.0.10) sheaves (Definition 4.0.2) on $X_{\text{ét}}$ (Definition 4.1.3), this condition may be slightly stronger than Definition 10.0.8 Ekedahl's condition B.

Definition 11.3.3 ([?, Definition 5.5.3.]). (♠ TODO: complete ring with respect to ideal) Let R be a commutative noetherian ring complete for the topology (Definition C.0.6) defined by an ideal (Definition C.0.2) $\mathfrak{m} \subset R$. Define $D_{\text{Ek}}(X_{\text{proét}}, \widehat{R}) \subset D_{\text{comp}}(X_{\text{proét}}, \widehat{R})$ (Definition 11.1.3) (♠ TODO: D_{comp}) as the full subcategory of complexes K satisfying $K \otimes_{\widehat{R}} R/\mathfrak{m} \in D_{\text{cc}}(X_{\text{proét}})$, i.e. satisfying $H^i(K \otimes_{\widehat{R}} R/\mathfrak{m}) \in \nu^* \text{Ab}(X_{\text{ét}})$ for all i (♠ TODO: D_{cc} , $\text{Ab}(X_{\text{ét}}), \nu$). For $*$ $\in \{+, -, b\}$, define $D_{\text{Ek}}^*(X_{\text{proét}}, \widehat{R}) \subset D_{\text{Ek}}(X_{\text{proét}}, \widehat{R})$ as the full subcategory with corresponding boundedness assumptions.

Proposition 11.3.4 ([?, Proposition 5.5.4.]). (♠ TODO: complete ring with respect to ideal) Let R be a commutative noetherian ring complete for the topology (Definition C.0.6) defined by an ideal (Definition C.0.2) $\mathfrak{m} \subset R$. Let $*$ $= -$ if Ekedahl's condition A (Definition 10.0.8) is satisfied for $(X_{\text{ét}}, R, \mathfrak{m})$, let $*$ $= +$ if Bhatt and Scholze's Ekedahl's condition B (Definition 11.3.2) is satisfied for $(R, \mathfrak{m})$, and let $*$ be blank if both conditions hold.

There is a natural equivalence

$$D_{\text{Ek}}^*(X_{\text{proét}}, \widehat{R}) \cong D_{\text{Ek}}^*(X_{\text{ét}}, R)$$

(Definition 11.3.3) (Definition 10.0.17)

11.4. Comparison to classical ℓ -adic categories.

Theorem 11.4.1. (♠ TODO: the categories) Let ℓ be a prime number. Let E be an algebraic extension of $\mathbb{Q}_{\ell}$. The triangulated category $D_{\text{cons}}(X_{\text{proét}}, E)$ is equivalent to the triangulated category traditionally called $D_c^b(X, E)$

1.1. Categorical definitions.

Definition .1.1 (Category). A *category* $\mathcal{C}$ consists of the following data:

- A class of *objects* denoted $\text{Ob}(\mathcal{C})$.
- For each pair of objects $X, Y \in \text{Ob}(\mathcal{C})$, a class

$$\text{Hom}_{\mathcal{C}}(X, Y)$$

of *morphisms* (also called *arrows* or *homs*). If the category $\mathcal{C}$ is clear, then this *hom-class* is also denoted by $\text{Hom}(X, Y)$. It may also be denoted by $\text{hom}_{\mathcal{C}}(X, Y)$ or $\text{hom}(X, Y)$, especially to distinguish from other types of hom's (e.g. internal hom's)

- For each triple of objects X, Y, Z , a composition law

$$\circ : \text{Hom}_{\mathcal{C}}(Y, Z) \times \text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{\mathcal{C}}(X, Z),$$

denoted $(g, f) \mapsto g \circ f$.

- For each object X , an *identity morphism*

$$\text{id}_X \in \text{Hom}_{\mathcal{C}}(X, X).$$

These data satisfy the following axioms:

- (Associativity) For all morphisms $f \in \text{Hom}_{\mathcal{C}}(X, Y)$, $g \in \text{Hom}_{\mathcal{C}}(Y, Z)$, and $h \in \text{Hom}_{\mathcal{C}}(Z, W)$,

$$h \circ (g \circ f) = (h \circ g) \circ f.$$

- (Identity) For all $f \in \text{Hom}_{\mathcal{C}}(X, Y)$,

$$\text{id}_Y \circ f = f = f \circ \text{id}_X.$$

One often writes $X \in \mathcal{C}$ synonymously with $X \in \text{Ob}(\mathcal{C})$, i.e. to denote that X is an object of $\mathcal{C}$.

We may call a category as above an *ordinary category* to distinguish this notion from the notions of *categories enriched in monoidal categories* or higher/ n -categories. (♠ TODO: define n -categories)

A category as defined above may be called called a *large category* or a *class category* to emphasize that the hom-classes may be proper classes rather than sets (note, however, that the possibility that hom-classes are sets is not excluded for large categories). Accordingly, a *category* may often refer to a locally small category (Definition .1.4), which is a category whose hom-classes are all sets.

Definition .1.2 (Grothendieck Universe). Let U be a set. We say U is a *Grothendieck universe* (or just a *universe*) if the following conditions hold:

1. If $x \in U$ and $y \in x$, then $y \in U$ (transitivity).
2. If $x, y \in U$, then $\{x, y\} \in U$ (closed under pair formation).
3. If $x \in U$, then the power set $\mathcal{P}(x) \in U$.
4. If $I \in U$ and $(x_\alpha)_{\alpha \in I}$ is a family with each $x_\alpha \in U$, then $\bigcup_{\alpha \in I} x_\alpha \in U$.

A set X is called *U -small* or a *U -set* if $X \in U$.

Definition .1.3 (Coarser and finer Grothendieck topologies). Let $\mathcal{C}$ be a category (Definition .1.1) and let τ, τ' be two Grothendieck topologies (Definition A.0.4) on $\mathcal{C}$.

We say that τ is *finer than* τ' (or equivalently, τ' is *coarser than* τ) if for every object $U \in \mathcal{C}$ every τ -covering of U is also a τ' -covering of U .

If τ and τ' are not comparable by inclusion, i.e., neither is finer or coarser than the other, they are said to be *incomparable*.

Definition .1.4 (Locally small category). A (large) category (Definition .1.1) $\mathcal{C}$ is called a *locally small category* if for every pair of objects $X, Y \in \text{Ob}(\mathcal{C})$, the collection $\text{Hom}_{\mathcal{C}}(X, Y)$ of morphisms between them is a (small) *set* (as opposed to a proper class). In other words, each hom-class is a set and may even be called a *hom-set*.

In some contexts, a locally small category may simply be called a *category*, especially when genuinely large categories are not considered.

A category $\mathcal{C}$ is called a *small category* if it is a locally small category and the class $\text{Ob}(\mathcal{C})$ of objects is a set.

Given a universe (Definition .1.2) U , we can define the notion of a *U -locally small category* and of a *U -small category* similarly [?, Exposé I, Définition 1.1.]. More explicitly,

1. a U -locally small category is a category such that for every pair of objects $X, Y \in \text{Ob}(\mathcal{C})$, the collection $\text{Hom}_{\mathcal{C}}(X, Y)$ of morphisms between them is a U -set.
2. a U -small category is a category such that $\text{Ob}(\mathcal{C})$ is a U -set and for every pair of objects $X, Y \in \text{Ob}(\mathcal{C})$, the collection $\text{Hom}_{\mathcal{C}}(X, Y)$ of morphisms between them is a U -set; in particular the collection of all objects and morphisms in a U -small category is a U -set.

A *U -category* refers to a U -locally small category.

Remark .1.5. Many “concrete” categories considered in “classical mathematics” or outside of more “abstract” category theory tend to be locally small. For example, the categories of sets, groups, R -modules, vector spaces, topological spaces, schemes, manifolds, sheaves on “small enough” sites are all locally small.

Definition .1.6. An *equivalence of categories* between two (large) categories (Definition .1.1) $\mathcal{C}$ and $\mathcal{D}$ consists of a pair of functors

$$F : \mathcal{C} \rightarrow \mathcal{D} \quad \text{and} \quad G : \mathcal{D} \rightarrow \mathcal{C}$$

together with natural isomorphisms

$$\eta : \text{Id}_{\mathcal{C}} \xrightarrow{\sim} G \circ F \quad \text{and} \quad \epsilon : F \circ G \xrightarrow{\sim} \text{Id}_{\mathcal{D}}.$$

Such functors F and G may be called *(natural) inverses of each other*.

When $\mathcal{C}$ and $\mathcal{D}$ are locally small categories (Definition .1.4), F is an equivalence of categories if and only if F is fully faithful and essentially surjective

Lemma .1.7. Let $\mathcal{C}$ be a small category (Definition .1.4) (resp. U -small category where U is some universe (Definition .1.2)) and let $\mathcal{A}$ be a locally small (Definition .1.4) category (resp. U -locally small category). The presheaf category $\text{PreShv}(\mathcal{C}, \mathcal{A})$ (Definition 4.0.1) is locally small (resp. U -locally small).

Proof. A morphism $\mathcal{F} \rightarrow \mathcal{G}$ in $\text{PreShv}(\mathcal{C}, \mathcal{A})$ is a natural transformation of the functors $\mathcal{F}, \mathcal{G} : \mathcal{C}^{\text{op}} \rightarrow \mathcal{A}$. Such a natural transformation is encoded by a family $(\eta_C)_C$ of morphisms (satisfying certain conditions) $\eta_C : \mathcal{F}(C) \rightarrow \mathcal{G}(C)$ in $\mathcal{A}$ over objects C of $\mathcal{C}^{\text{op}}$. The product $\prod_{C \in \text{Ob } \mathcal{C}^{\text{op}}} \text{Hom}_{\mathcal{A}}(\mathcal{F}(C), \mathcal{G}(C))$ is a product of (U -small) sets indexed by a (U -small) set, and the collection of natural transformations is a subset of this set. Therefore, $\text{Hom}_{\text{PreShv}(\mathcal{C}, \mathcal{A})}(\mathcal{F}, \mathcal{G})$ is a (U -small) set. $\square$

Definition .1.8. Let $\mathcal{C}$ and $\mathcal{D}$ be categories (Definition .1.1). Let $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ be functors.

An **adjunction between F and G** consists of two natural transformations: $\eta : \text{Id}_{\mathcal{C}} \Rightarrow GF$ (the **unit**), and $\varepsilon : FG \Rightarrow \text{Id}_{\mathcal{D}}$ (the **counit**)

These must satisfy the triangle identities: For every object $X \in \mathcal{C}$ and $Y \in \mathcal{D}$,

$$\varepsilon_{FX} \circ F(\eta_X) = \text{id}_{FX}$$

$$G(\varepsilon_Y) \circ \eta_{GY} = \text{id}_{GY}.$$

In diagrammatic form, the triangle identities assert that the following are commutative diagrams:

$$\begin{array}{ccc} F(X) & \xrightarrow{F(\eta_X)} & FGF(X) \\ & \searrow \text{id}_{F(X)} & \downarrow \varepsilon_{F(X)} \\ & & F(X) \end{array} \quad \begin{array}{ccc} G(Y) & \xrightarrow{\eta_{G(Y)}} & GFG(Y) \\ & \searrow \text{id}_{G(Y)} & \downarrow G(\varepsilon_Y) \\ & & G(Y) \end{array}$$

We say that F is a **left adjoint to G** and G is a **right adjoint to F** (written $F \dashv G$).

In the case that $\mathcal{C}$ and $\mathcal{D}$ are locally small (Definition .1.4) categories (or U -locally small categories if a universe (Definition .1.2) U is available), we have an adjunction $F \dashv G$ if and only if for every object X in $\mathcal{C}$ and Y in $\mathcal{D}$ there is a natural isomorphism

$$\text{Hom}_{\mathcal{D}}(F(X), Y) \cong \text{Hom}_{\mathcal{C}}(X, G(Y))$$

that is natural in both X and Y . More precisely, there are two functors

$$\mathcal{C}^{\text{op}} \times \mathcal{D} \rightarrow \mathbf{Sets}$$

given by

$$(X, Y) \mapsto \text{Hom}_{\mathcal{D}}(F(X), Y)$$

$$(X, Y) \mapsto \text{Hom}_{\mathcal{C}}(X, G(Y))$$

respectively, and there is a natural isomorphism between them.

In this case, the **unit of the adjunction** is the natural transformation $\eta : \text{Id}_{\mathcal{C}} \Rightarrow GF$ such that,

1. for every $X \in \mathcal{C}$, the morphism $\eta_X : X \rightarrow GF(X)$ (each called a **unit morphism**) in $\mathcal{C}$ is obtained as the image of $\text{id}_{F(X)}$ via the adjoint isomorphism

$$\text{Hom}_{\mathcal{D}}(F(X), F(X)) \cong \text{Hom}_{\mathcal{C}}(X, GF(X)).$$

2. for every $Y \in \mathcal{D}$, the morphism $\varepsilon_Y : FG(Y) \rightarrow Y$ (each called a **counit morphism**) in $\mathcal{D}$ is obtained as the image of $\text{id}_{G(Y)}$ via the adjoint isomorphism

$$\text{Hom}_{\mathcal{C}}(G(Y), G(Y)) \cong \text{Hom}_{\mathcal{D}}(FG(Y), Y).$$

Definition .1.9. Let (X, τ_X) be a topological space. The **small site associated to X** or **the site of open covers of X** or **the canonical site on $\text{Open } X$** is the category $\text{Open}(X)$ of open subsets of X with inclusion morphisms, equipped with the canonical Grothendieck topology (Definition A.0.4) generated by (Definition 4.2.1) the Grothendieck pretopology (Definition 4.2.2) whose covering families $\{U_i \rightarrow U\}_{i \in I}$, for $U \in \text{Open}(X)$ are families of morphisms in $\text{Open}(X)$ such that $\bigcup_{i \in I} U_i = U$. In other words, $\{U_i \rightarrow U\}_{i \in I}$ is a covering for the pretopology if it is an open coverings.

Definition .1.10. Let $\mathcal{C}$ be a (large) category (Definition .1.1).

1. An object $I \in \mathcal{C}$ is called an **initial object** if for every object $X \in \mathcal{C}$ there exists a unique morphism

$$I \rightarrow X.$$

Equivalently, an initial object is a limit (Definition .1.13) of the empty diagram (Definition .1.33), if such a limit exists.

2. An object $F \in \mathcal{C}$ is called a **final object** (or **terminal object**) if for every object $X \in \mathcal{C}$ there exists a unique morphism

$$X \rightarrow F.$$

Equivalently, a final object is a colimit (Definition .1.13) of the empty diagram (Definition .1.33), if such a colimit exists.

3. An object $Z \in \mathcal{C}$ is called a **zero object** if Z is both initial and final in $\mathcal{C}$. In particular, for every object $X \in \mathcal{C}$ there exist unique morphisms

$$Z \rightarrow X \quad \text{and} \quad X \rightarrow Z.$$

In particular, if initial/final/zero objects exist in a category, then they are unique up to unique isomorphism.

Definition .1.11 (connected category). A **non-empty category $\mathcal{I}$ is connected** if, for any two objects $i, j \in \mathcal{I}$, there exists a finite sequence of objects $x_0 = i, x_1, \dots, x_n = j$ in $\mathcal{I}$ and morphisms $f_k : x_{k-1} \rightarrow x_k$ or $f_k : x_k \rightarrow x_{k-1}$ for each $k = 1, \dots, n$. That is, any two objects can be joined by a zigzag of morphisms. (♠ TODO: AI generated, verify)

Definition .1.12 (Filtered category). 1. A **filtered category** is a (nonempty, large) category $\mathcal{I}$ satisfying the following conditions:

- For every finite collection of objects $i_1, i_2, \dots, i_n$ in $\mathcal{I}$, there exists an object j and morphisms

$$\phi_k : i_k \rightarrow j, \quad \text{for each } k = 1, \dots, n.$$

- For every pair of morphisms $f, g : i \rightarrow j$ in $\mathcal{I}$, there exists an object k and a morphism

$$h : j \rightarrow k$$

that satisfies

$$h \circ f = h \circ g.$$

In other words, $\mathcal{I}$ is nonempty, any finite diagram of objects admits a cocone (Definition .1.13), and any pair of parallel morphisms become equal after post-composition with an appropriate morphism.

2. Dually, a **Cofiltered category** is a category whose opposite category is filtered. More explicitly, A cofiltered category is a (nonempty, large) category $\mathcal{I}$ satisfying the following conditions:

- For every finite collection of objects $i_1, i_2, \dots, i_n$ in $\mathcal{I}$, there exists an object j and morphisms

$$\phi_k : j \rightarrow i_k, \quad \text{for each } k = 1, \dots, n.$$




FIGURE 1. *

Condition 1: Upper Bound

- For every pair of morphisms $f, g : j \rightarrow i$ in $\mathcal{I}$, there exists an object k and a morphism

$$h : k \rightarrow j$$

that satisfies

$$f \circ h = g \circ h.$$




FIGURE 3. *

Condition 1: Lower Bound

In other words, $\mathcal{I}$ is nonempty, any finite diagram of objects admits a cone, and any pair of parallel morphisms become equal after pre-composition with an appropriate morphism.

Definition .1.13 (Cones, limits and colimits in a category). Let $\mathcal{C}$ be a (large) category (Definition .1.1), let I be a (large) category, and let $D : I \rightarrow \mathcal{C}$ be a diagram (Definition .1.33) (Definition .1.33).

1. A **cone to the diagram D** is an object $L \in \mathcal{C}$ together with a family of morphisms

$$\{\pi_i : L \rightarrow D(i)\}_{i \in I}$$

such that for every morphism $f : i \rightarrow j$ in I , the diagram

$$\begin{array}{ccc} & L & \\ \pi_i \swarrow & & \searrow \pi_j \\ D(i) & \xrightarrow{D(f)} & D(j) \end{array}$$

$$i \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} j \dashrightarrow^h k$$

FIGURE 2. *

Condition 2: Coequalizing map

$$k \dashrightarrow^h j \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} i$$

FIGURE 4. *


Condition 2: Equalizing map

commutes, i.e. $D(f) \circ \pi_i = \pi_j$.

2. A cone $(L, \{\pi_i\})$ is called a **limit of D** if it satisfies the following “universal property”: for any cone $(C, \{f_i\})$ over D , there exists a *unique* morphism $u : C \rightarrow L$ such that

$$\pi_i \circ u = f_i \quad \text{for all } i \in I.$$

Visually, the following diagrams commute every morphism $f : i \rightarrow j$ in I :




If such a cone exists, then the object L is necessarily unique up to unique isomorphism by the universal property. In this case, L is denoted by $\lim_{i \in I} D$ or **lim D** .

3. A **cocone from the diagram D** is an object $C \in \mathcal{C}$ together with a family of morphisms

$$\{\iota_i : D(i) \rightarrow C\}_{i \in I}$$

such that for every morphism $f : i \rightarrow j$ in I , the diagram




commutes, i.e. $\iota_j \circ D(f) = \iota_i$.

4. A cocone $(L, \{\iota_i\})$ is called a **colimit of D** if it satisfies the following “universal property”: for any cocone $(C, \{g_i\})$ under D , there exists a *unique* morphism $u : L \rightarrow C$ such that

$$u \circ \iota_i = g_i \quad \text{for all } i \in I.$$

Visually, the following diagrams commute every morphism $f : i \rightarrow j$ in I :



If such a cocone exists, then the object L is necessarily unique up to unique isomorphism by the universal property. In this case, L is denoted by $\text{colim}_{i \in I} D$ or **colim D** .

A limit/colimit is called **fnite** (resp. **small**) if the index category I is finite (resp. small).

Some authors use the terms *projective limit* or *inverse limit* to refer to what is defined here as a limit. Similarly, the terms *inductive limit* or *direct limit* are sometimes used to mean a colimit. However, these phrases can have more specific meanings to other authors: a *projective* or *inverse limit* may refer to a limit over a diagram indexed by a codirected poset (Definition .1.28). Likewise, an *inductive* or *direct limit* may refer to a colimit over a directed poset (Definition .1.28) (see Definition .1.14).

Thus, while the terms are sometimes used interchangeably with “limit” and “colimit,” they may also emphasize particular indexing shapes and directions, distinguishing them from general limits and colimits taken over arbitrary small categories.

Definition .1.14 (Special cases of limits). Let $\mathcal{C}$ be a (large) category. Let I be a (large) category. Let $I \rightarrow \mathcal{C}$ be a diagram (Definition .1.33)/system.

- Suppose that the system is a cofiltered system (Definition .1.18), i.e. I is a cofiltered category. A limit (Definition .1.13) of this diagram is often denoted by

$$\varprojlim_{i \in I} D(i)$$

and may be called a *cofiltered (inverse/projective) limit*. In case that the system is more specifically an inverse/projective system (Definition .1.18), i.e. I is a cofiltered poset (Definition .1.28), the preferred term for such a limit is *inverse/projective limit*.

- Suppose that the system is a filtered system, i.e. I is a filtered category. A colimit of this diagram is often denoted by

$$\varinjlim_{i \in I} D(i)$$

and may be called a *filtered colimit* or a *direct/inductive/injective limit*. In case that the system is more specifically a direct/inductive system, i.e. I is a filtered poset (Definition .1.28), the preferred term for such a limit is *direct/inductive limit*.

Definition .1.15 (Ind-category). Let $\mathcal{C}$ be a locally small category (Definition .1.4).

1. The *Ind-category of $\mathcal{C}$* , denoted $\text{Ind}(\mathcal{C})$, is defined as follows:
 - Objects of $\text{Ind}(\mathcal{C})$ are formal filtered colimits (Definition .1.14) of objects in $\mathcal{C}$. More precisely, an object is given by a filtered (Definition .1.12) small category I and a functor

$$X : I \rightarrow \mathcal{C}.$$

- Morphisms between objects $X : I \rightarrow \mathcal{C}$ and $Y : J \rightarrow \mathcal{C}$ are defined by

$$\text{Hom}_{\text{Ind}(\mathcal{C})}(X, Y) := \varprojlim_{i \in I} \varinjlim_{j \in J} \text{Hom}_{\mathcal{C}}(X_i, Y_j),$$

(Definition .1.14) where X_i and Y_j denote the images of $i \in I$ and $j \in J$ under X and Y , respectively.

The composition of morphisms is induced naturally from composition in $\mathcal{C}$. Hence, $\text{Ind}(\mathcal{C})$ is the completion of $\mathcal{C}$ under filtered colimits. Objects of $\text{Ind}(\mathcal{C})$ are called *Ind-objects of $\mathcal{C}$* .

2. The *Pro-category of $\mathcal{C}$* , denoted $\text{Pro}(\mathcal{C})$, is defined as follows:

- Objects of $\text{Pro}(\mathcal{C})$ are formal cofiltered limits (Definition .1.14) of objects in $\mathcal{C}$. More precisely, an object is given by a cofiltered small category I and a functor

$$X : I \rightarrow \mathcal{C}.$$

- Morphisms between objects $X : I \rightarrow \mathcal{C}$ and $Y : J \rightarrow \mathcal{C}$ are defined by

$$\text{Hom}_{\text{Pro}(\mathcal{C})}(X, Y) := \varinjlim_{j \in J} \varprojlim_{i \in I} \text{Hom}_{\mathcal{C}}(X_i, Y_j),$$

where X_i and Y_j denote the images of $i \in I$ and $j \in J$ under X and Y , respectively. The composition of morphisms is induced naturally from composition in $\mathcal{C}$.

Hence, $\text{Pro}(\mathcal{C})$ is the completion of $\mathcal{C}$ under cofiltered limits. Objects of $\text{Pro}(\mathcal{C})$ are called *Pro-objects of $\mathcal{C}$* .

Since **Sets** has all limits and colimits (♠ **TODO:**) and hence has all projective and inductive limits and since $\mathcal{C}$ is locally small, $\text{Ind}(\mathcal{C})$ and $\text{Pro}(\mathcal{C})$ are locally small.

The *canonical embedding of a locally small category $\mathcal{C}$ into its Ind-category or Pro-category* is defined by treating each object as a "constant" diagram. Let $\mathbf{1}$ be the category with a single object $*$ and its identity morphism.

1. The Embedding $\iota_{\text{Ind}} : \mathcal{C} \rightarrow \text{Ind}(\mathcal{C})$:

- Objects: An object $C \in \mathcal{C}$ is mapped to the functor $X : \mathbf{1} \rightarrow \mathcal{C}$ sending $*$ to C . This represents C as a formal filtered colimit over the trivial index category.
- Morphisms: Given objects X (representing C) and Y (representing D), the Hom-set in the Ind-category simplifies:

$$\text{Hom}_{\text{Ind}(\mathcal{C})}(X, Y) \cong \varinjlim_{i \in \mathbf{1}} \varprojlim_{j \in \mathbf{1}} \text{Hom}_{\mathcal{C}}(X_i, Y_j) \cong \text{Hom}_{\mathcal{C}}(C, D).$$

2. The Embedding $\iota_{\text{Pro}} : \mathcal{C} \rightarrow \text{Pro}(\mathcal{C})$:

- Objects: An object $C \in \mathcal{C}$ is mapped to the functor $X : \mathbf{1} \rightarrow \mathcal{C}$ sending $*$ to C , representing C as a formal cofiltered limit.
- Morphisms: Similarly, the Pro-morphism definition reduces to:

$$\text{Hom}_{\text{Pro}(\mathcal{C})}(X, Y) \cong \varinjlim_{j \in \mathbf{1}} \varprojlim_{i \in \mathbf{1}} \text{Hom}_{\mathcal{C}}(X_i, Y_j) \cong \text{Hom}_{\mathcal{C}}(C, D).$$

In both cases, the functor ι is fully faithful, allowing us to identify $\mathcal{C}$ with a full subcategory of its respective completion. Given an object C of $\mathcal{C}$, the pro/ind-object $\iota(C)$ may be called the *constant pro/ind-object of C* .

Definition .1.16 (Product in a category). Let $\mathcal{C}$ be a category and let $\{X_i\}_{i \in I}$ be a family of objects in $\mathcal{C}$ indexed by a class I .

1. A *product of the family $\{X_i\}$* is an object P of $\mathcal{C}$ together with a “universal” family of morphisms

$$\pi_i : P \rightarrow X_i, \quad \text{for each } i \in I.$$

More precisely, for any object Y and any family of morphisms $\{f_i : Y \rightarrow X_i\}_{i \in I}$, there exists a unique morphism

$$f : Y \rightarrow P$$

making the following diagram commute for all $i \in I$, i.e. $\pi_i \circ f = f_i$:

$$\begin{array}{ccc} Y & & \\ \downarrow \exists! f & \searrow f_i & \\ \prod X_i & \xrightarrow{\pi_i} & X_i \end{array}$$

Such a product is often denoted by $\prod_{i \in I} X_i$. If $\prod_{i \in I} X_i$ exists in $\mathcal{C}$, then it is unique up to unique isomorphism by the universal property described above.

Equivalently, the product $\prod_{i \in I} X_i$ is the limit (Definition .1.13) of the diagram (Definition .1.33) $I \rightarrow \mathcal{C}, i \mapsto X_i$, where I is made into a category whose objects are the members of I and whose morphisms are just the identity morphisms.

2. A **coproduct** (or synonymously **direct sum**) of the family $\{X_i\}$ is an object C of $\mathcal{C}$ together with a “universal” family of morphisms

$$\iota_i : X_i \rightarrow C, \quad \text{for each } i \in I.$$

More precisely, for any object Y and any family of morphisms $\{g_i : X_i \rightarrow Y\}_{i \in I}$, there exists a unique morphism

$$g : C \rightarrow Y$$

making the following diagram commute for all $i \in I$, i.e. $g \circ \iota_i = g_i$:

$$\begin{array}{ccc} X_i & \xrightarrow{\iota_i} & \coprod X_i \\ & \searrow g_i & \downarrow \exists! g \\ & & Y \end{array}$$

Such a coproduct is often denoted by $\coprod_{i \in I} X_i$ or $\oplus_{i \in I} X_i$. If $\coprod_{i \in I} X_i$ exists in $\mathcal{C}$, then it is unique up to unique isomorphism by the universal property described above.

Equivalently, the coproduct $\coprod_{i \in I} X_i$ is the colimit (Definition .1.13) of the diagram (Definition .1.33) $I \rightarrow \mathcal{C}, i \mapsto X_i$, where I is made into a category whose objects are the members of I and whose morphisms are just the identity morphisms.

Definition .1.17 (Equalizer in a category). Let $\mathcal{C}$ be a (large) category (Definition .1.1) and let $f, g : X \rightarrow Y$ be morphisms in $\mathcal{C}$.

1. An **equalizer of f and g** is an object E together with a morphism

$$e : E \rightarrow X$$

such that

$$f \circ e = g \circ e$$

and for any object Z with morphism $z : Z \rightarrow X$ satisfying

$$f \circ z = g \circ z,$$

there exists a unique morphism $u : Z \rightarrow E$ making the diagram commute:

$$e \circ u = z.$$

$$\begin{array}{ccccc}
Z & & & & \\
\downarrow \exists! u & \searrow z & & & \\
E & \xrightarrow{e} & X & \xrightleftharpoons[f]{g} & Y
\end{array}$$

If such an equalizer of f and g exists, then we say that the following **equalizer diagram is exact**:

$$E \xrightarrow{e} X \xrightleftharpoons[f]{g} Y$$

2. A **coequalizer of f and g** is an object Q together with a morphism

$$q : Y \rightarrow Q$$

such that

$$q \circ f = q \circ g$$

and for any object Z with morphism $w : Y \rightarrow Z$ satisfying

$$w \circ f = w \circ g,$$

there exists a unique morphism $v : Q \rightarrow Z$ making the diagram commute:

$$\begin{array}{ccccc}
X & \xrightleftharpoons[f]{g} & Y & \xrightarrow{q} & Q \\
& & \searrow w & & \downarrow \exists! v \\
& & & & Z
\end{array}$$

If such a coequalizer of f and g exists, then we say that the following **coequalizer diagram is exact**:

$$X \xrightleftharpoons[f]{g} Y \xrightarrow{q} Q$$

Definition .1.18 (Systems in a category). Let $\mathcal{C}$ be a (large) category. Let I be a (large) category.

1. A diagram/system (Definition .1.33) $I \rightarrow \mathcal{C}$ is called **filtered** (resp. **cofiltered**) if I is a filtered (Definition .1.12) (resp. cofiltered (Definition .1.12)) category.
2. A diagram/system $I \rightarrow \mathcal{C}$ is called **directed** (resp. **codirected**) if I is small and thing, i.e. is regardable/comes from (Lemma .1.29) a directed (resp. codirected) partially ordered set (Definition .1.28). A **direct system** or **inductive system** is synonymous for a directed system and a **inverse system** or **projective system** is synonymous for a codirected system.

One might also speak of a **filtered direct/inductive system** synonymously for a filtered system to emphasize that the indexing category is a general filtered category, rather than a directed poset.

Definition .1.19 (Cofinal and final sub-posets). Let $(I, \leq)$ be a partially ordered set (Definition .1.28), and let $J \subseteq I$ be a sub-poset. Let $(K, \leq)$ be a partially ordered set (Definition .1.28), and let $L \subseteq K$ be a sub-poset.

1. J is **cofinal in I** if for every $i \in I$, there exists $j \in J$ such that $i \leq j$.
2. L is **final in K** if for every $k \in K$, there exists $\ell \in L$ such that $\ell \leq k$.

Definition .1.20 (Cofinal and final subsystems of direct/inverse systems). (♠ TODO: AI generated, verify) Let $\mathcal{C}$ be a category (Definition .1.1).

1. Let $(I, \leq)$ be a directed partially ordered set (Definition .1.28), and let $(A_i)_{i \in I}$ be a direct system (Definition .1.18) in $\mathcal{C}$ indexed by I . Let $J \subseteq I$ be a sub-poset that is cofinal in I (Definition .1.19). The restriction of the direct system to J , denoted $(A_j)_{j \in J}$, is called a **cofinal subsystem** of $(A_i)_{i \in I}$.
2. Let $(K, \leq)$ be a codirected partially ordered set (Definition .1.28), and let $(B_k)_{k \in K}$ be an inverse system (Definition .1.18) in $\mathcal{C}$ indexed by K . Let $L \subseteq K$ be a sub-poset that is final in K (Definition .1.19). The restriction of the inverse system to L , denoted $(B_\ell)_{\ell \in L}$, is called a **final subsystem** of $(B_k)_{k \in K}$.

Theorem .1.21 (Cofinal and final subsystems suffice for limits). Let $\mathcal{C}$ be a category (Definition .1.1).

1. Let $(I, \leq)$ be a directed partially ordered set (Definition .1.28), and let $J \subseteq I$ be a sub-poset. Let $(A_i)_{i \in I}$ be a direct system (Definition .1.18) in $\mathcal{C}$ indexed by I . Suppose that J is cofinal in I (Definition .1.19). Let $(A_j)_{j \in J}$ denote the restriction of the direct system to J . Suppose that $\varinjlim_{i \in I} A_i$ exists in $\mathcal{C}$. Then $\varinjlim_{j \in J} A_j$ also exists in $\mathcal{C}$ and the canonical morphism

$$(D) \quad \varinjlim_{j \in J} A_j \rightarrow \varinjlim_{i \in I} A_i$$

(Definition .1.14) induced by the inclusions $J \hookrightarrow I$ and the transition morphisms of the direct system is an isomorphism.

2. Let $(K, \leq)$ be a codirected partially ordered set (Definition .1.28), and let $L \subseteq K$ be a sub-poset. Let $(B_k)_{k \in K}$ be an inverse system (Definition .1.18) in $\mathcal{C}$ indexed by K . Suppose that L is final in K (Definition .1.19). Let $(B_\ell)_{\ell \in L}$ denote the restriction of the inverse system to L . Suppose that $\varprojlim_{k \in K} B_k$ exists in $\mathcal{C}$. Then $\varprojlim_{\ell \in L} B_\ell$ also exists in $\mathcal{C}$ and the canonical morphism

$$(E) \quad \varprojlim_{k \in K} B_k \rightarrow \varprojlim_{\ell \in L} B_\ell$$

(Definition .1.14) induced by the inclusions $L \hookrightarrow K$ and the transition morphisms of the inverse system is an isomorphism.

Proof. 1. Suppose that $\varinjlim_{i \in I} A_i$ exists in $\mathcal{C}$. We show that it satisfies the universal property of $\varinjlim_{j \in J} A_j$, i.e. that $\varinjlim_{i \in I} A_i$ is a cocone (Definition .1.13) of $(A_j)_{j \in J}$ and that for any cocone D of $(A_j)_{j \in J}$ there is a unique morphism $\varinjlim_{j \in J} A_j \rightarrow D$ compatible with the structures of these objects as cocones of $(A_j)_{j \in J}$. For every $j \in J$,

there is a canonical map A_j to $\varinjlim_{i \in I} A_i$. Moreover, these canonical maps over $j \in J$ admit $\varinjlim_{i \in I} A_i$ as a cocone (Definition .1.13) of $(A_j)_{j \in J}$ because $\varinjlim_{i \in I} A_i$ is a cocone of $(A_i)_{i \in I}$.

Suppose that $D \in \mathcal{C}$ is a cocone of $(A_j)_{j \in J}$, so equivalently it is equipped with morphisms $\pi_j : A_j \rightarrow D$ for all $j \in J$ such that for every morphism $f : j_1 \rightarrow j_2$ in J , we have that

$$\begin{array}{ccc} A_{j_1} & & \\ \downarrow A_f & \searrow \pi_{j_1} & \\ & D & \\ \uparrow \pi_{j_2} & \nearrow & \\ A_{j_2} & & \end{array}$$

commutes. To first show the existence of the desired map $\varinjlim_{i \in I} A_i \rightarrow D$, we extend this cocone D of $(A_j)_{j \in J}$ to a cocone of $(A_i)_{i \in I}$. Given $i \in I$, define $\pi_i : A_i \rightarrow D$ as follows: by the cofinality of J in I , there exists some $j \in J$ such that $i \leq j$. Writing $A_{i \rightarrow j}$ for morphism in $\mathcal{C}$ assigned to $i \rightarrow j$ by the diagram A , let

$$(F) \quad \pi_i = \pi_j \circ A_{i \rightarrow j}.$$

We show that this is well defined, i.e. is independent of the choice of $j \in J$. To this end, suppose that j_1, j_2 are objects of J such that $i \leq j_1, j_2$. Since I is a directed poset, there exists some $j_3 \in J$ such that $j_1, j_2 \leq j_3$. The following diagram commutes:

$$\begin{array}{ccccc} & & A_{j_1} & & \\ & \nearrow & \searrow \pi_{j_1} & \nearrow & \\ A_i & & & & D \\ & \searrow & \nearrow \pi_{j_3} & \searrow & \\ & & A_{j_3} & & \\ & \nearrow & \searrow \pi_{j_2} & \nearrow & \\ & & A_{j_2} & & \end{array}$$

Therefore, $\pi_j \circ A_{i \rightarrow j_1}$ and $\pi_j \circ A_{i \rightarrow j_2}$ both equal any of the maps from A_i to D above, so π_i is indeed well defined.

We show that D equipped with the maps $\pi_i : A_i \rightarrow D$ is indeed a cocone of $(A_i)_{i \in I}$. Suppose that $i_1 \rightarrow i_2$ is an arrow in I . We show that

$$\begin{array}{ccc} A_{i_1} & & \\ \downarrow A_{i_1 \rightarrow i_2} & \searrow \pi_{i_1} & \\ & D & \\ \uparrow \pi_{i_2} & \nearrow & \\ A_{i_2} & & \end{array}$$

commutes. Let $j_1, j_2 \in J$ satisfy $i_1 \leq j_1$ and $i_2 \leq j_2$. By the directedness of I , choose $i \in I$ such that $j_1, j_2 \leq i$ and by the cofinality of J in I , choose $j \in I$ such that $i \leq j$. In particular, we have that $i_1, i_2 \leq j$. By construct, $\pi_{i_1} = \pi_j \circ A_{i_1 \rightarrow j}$ and

$\pi_{i_2} = \pi_j \circ A_{i_2 \rightarrow j}$. Therefore,


$$\pi_{i_2} \circ A_{i_1 \rightarrow i_2} = \pi_j \circ A_{i_2 \rightarrow j} \circ A_{i_1 \rightarrow i_2} = \pi_j \circ A_{i_1 \rightarrow j} = \pi_{i_1}$$

as desired.

As D equipped with the π_i is a cocone of $(A_i)_{i \in I}$, the universal property of $\varinjlim_{i \in I} A_i$ means that there is a canonical morphism

$$\varinjlim_{i \in I} A_i \rightarrow D.$$

We have so far thus shown that $\varinjlim_{i \in I} A_i$ is a cocone of $(A_j)_{j \in J}$ and that for every cocone D of $(A_j)_{j \in J}$, there is a canonical morphism $\varinjlim_{i \in I} A_i \rightarrow D$ such that, for every morphism $j \rightarrow j'$ in J ,



commutes.

We show that the morphism is unique, i.e. that the desired map $\varinjlim_{i \in I} A_i \rightarrow D$ is unique. To this end, suppose that $\varinjlim_{i \in I} A_i \rightarrow D$ is a morphism making the above diagram commute for all $j \rightarrow j'$ in J . To give a morphism $\varinjlim_{i \in I} A_i \rightarrow D$ is equivalent to giving D as a cocone of A_i (♠ TODO: write the notion that a morphism from an injective limit to an object is equivalent to giving the object as a cocone of the system, assuming that the injective limit exists). Write $\pi_i : A_i \rightarrow D$ for the morphisms of this cocone. For any $j_0 \in J$ such that $i \leq j_0$ in I , we must have that $\pi_i = \pi_{j_0} \circ A_{i \rightarrow j_0}$ where $\pi_{j_0} : A_{j_0} \rightarrow D$ here is written as a morphism for D as a cocone of $(A_i)_{i \in I}$. On the other hand, π_{j_0} as a morphism for D as a cocone of $(A_i)_{i \in I}$ coincides with π_{j_0} as a morphism for D as a cocone of $(A_j)_{j \in J}$ because the above diagram has to commute and because the map $A_{j_0} \rightarrow A_{j_0}$ where the first copy is from $(A_j)_{j \in J}$ and the second copy is from $(A_i)_{i \in I}$ is the identity map. Therefore, π_i must be constructed as in (F), so the morphism is indeed unique as desired. We have thus shown that the desired morphism both exists and is unique as desired.

2. This is dual to the previous part.

□

Definition .1.22 (cofinal and final functors). Let $\mathcal{I}$ and $\mathcal{J}$ be categories (Definition .1.1), and let $u : \mathcal{I} \rightarrow \mathcal{J}$ be a functor.

1. A **cofinal functor** is a functor $u : \mathcal{I} \rightarrow \mathcal{J}$ such that, for every object $j \in \mathcal{J}$, the comma category $j \downarrow u$ is non-empty and a connected category (Definition .1.11).
2. A **final functor** is a functor $u : \mathcal{I} \rightarrow \mathcal{J}$ such that, for every object $j \in \mathcal{J}$, the comma category $u \downarrow j$ is non-empty and a connected category (Definition .1.11).

(♠ TODO: AI generated, verify)

(♠ TODO: cofinal and final functors induce isomorphisms of direct/inverse limits and guarantee existence of direct/inverse limits)

Definition .1.23 (action of a group on an object in a category). Let G be a group, let $\mathcal{C}$ be a category (Definition .1.1), and let c be an object of $\mathcal{C}$. An **action of G on c in $\mathcal{C}$** is a family $(\alpha(g))_{g \in G}$ where, for each $g \in G$, the morphism $\alpha(g) : c \rightarrow c$ is an automorphism of c such that

$$\alpha(e) = \text{id}_c \quad \text{and} \quad \alpha(gh) = \alpha(g) \circ \alpha(h) \quad \text{for all } g, h \in G,$$

where e denotes the identity element of G . If $\mathcal{C}$ is a locally small category (Definition .1.4), an action of G on c is equivalently given by a group homomorphism $\alpha : G \rightarrow \text{Aut}_{\mathcal{C}}(c)$ to the automorphism group of c .

Definition .1.24 (equivariant morphism between objects with G -action in $\mathcal{C}$). Let G be a group, let $\mathcal{C}$ be a category (Definition .1.1), and let (c, α) and (d, β) be objects of $\mathcal{C}$ equipped with an action of G (Definition .1.23). An **equivariant morphism from (c, α) to (d, β) in $\mathcal{C}$** is a morphism $f : c \rightarrow d$ in $\mathcal{C}$ such that the following diagram commutes for every $g \in G$:

$$\begin{array}{ccc} c & \xrightarrow{\alpha(g)} & c \\ f \downarrow & & \downarrow f \\ d & \xrightarrow{\beta(g)} & d \end{array}$$

Equivalently, $f \circ \alpha(g) = \beta(g) \circ f$ for all $g \in G$.

Definition .1.25 (category of objects with G -action in $\mathcal{C}$). Let G be a group and let $\mathcal{C}$ be a category (Definition .1.1). The **category of objects with G -action in $\mathcal{C}$** , denoted by $G\text{-}\mathcal{C}$, is defined as follows:

1. An object is a pair (c, α) where c is an object of $\mathcal{C}$ and α is an action of G on c in $\mathcal{C}$ (Definition .1.23).
2. A morphism from (c, α) to (d, β) is an equivariant morphism from (c, α) to (d, β) in $\mathcal{C}$ (Definition .1.24).

Definition .1.26. A category $\mathcal{C}$ is called **essentially small** if it is equivalent (Definition .1.6) to a small category (Definition .1.4), i.e., there exists a small category $\mathcal{D}$ and an equivalence of categories

$$F : \mathcal{D} \rightarrow \mathcal{C}.$$

Note that an essentially small category is necessarily locally small (Definition .1.4).

Definition .1.27 (Monomorphism and Epimorphism in Categories). Let $\mathcal{C}$ be a category (Definition .1.1). For objects $A, B \in \mathcal{C}$, let $f : A \rightarrow B$ be a morphism in $\mathcal{C}$.

- The morphism f is called a **monomorphism** (or a **monic morphism**) if for every object X and every pair of morphisms $g_1, g_2 : X \rightarrow A$, the equality $f \circ g_1 = f \circ g_2$ implies $g_1 = g_2$.

- The morphism f is called an **epimorphism** (or an **epic morphism**) if for every object Y and every pair of morphisms $h_1, h_2 : B \rightarrow Y$, the equality $h_1 \circ f = h_2 \circ f$ implies $h_1 = h_2$.

Definition .1.28 (Partially ordered set). 1. A **partially ordered set** (or **poset**), or **ordered set** is a pair $(P, \leq)$ where P is a set and

$$\leq : P \times P \rightarrow \{\text{true}, \text{false}\}$$

is a binary relation on P satisfying the following axioms for all $a, b, c \in P$:

- **Reflexivity**: $a \leq a$,
- **Antisymmetry**: if $a \leq b$ and $b \leq a$, then $a = b$,
- **Transitivity**: if $a \leq b$ and $b \leq c$, then $a \leq c$.

The relation $\leq$ is called an **order** or a **partial order**

2. A partially ordered set $(P, \leq)$ is called a **directed partially ordered set** if for every pair $a, b \in P$, there exists $c \in P$ such that

$$a \leq c \quad \text{and} \quad b \leq c.$$

3. A partially ordered set $(P, \leq)$ is called a **codirected partially ordered set** (or **downward directed poset**) if for every pair $a, b \in P$, there exists $d \in P$ such that

$$d \leq a \quad \text{and} \quad d \leq b.$$

Lemma .1.29. Let $(P, \leq)$ be a nonempty poset (Definition .1.28).

1. Regarding P as a category whose objects are the elements of P and such that there is a unique arrow $a \rightarrow b$ if and only if $a \leq b$, the category is filtered.
2. Every nonempty small (Definition .1.4) thin filtered category (Definition .1.12) corresponds to a poset in this way.
3. Moreover, the poset P is directed (Definition .1.28) if and only if the category is filtered. The poset P is codirected (Definition .1.28) if and only if the category is cofiltered.

Definition .1.30 (Generator of a category). Let $\mathcal{C}$ be a category (Definition .1.1).

1. An object $G \in \mathcal{C}$ is called a **generator** (or **separator**) if for every pair of distinct morphisms $f, g : X \rightarrow Y$ in $\mathcal{C}$, there exists a morphism $h : G \rightarrow X$ such that

$$f \circ h \neq g \circ h.$$

In case that $\mathcal{C}$ is locally small (Definition .1.4), this is equivalent to the condition that the representable functor (Definition 4.4.3)

$$\text{Hom}_{\mathcal{C}}(G, -) : \mathcal{C} \rightarrow \mathbf{Set}$$

is faithful, which in turn is equivalent to the condition that for every object $X \in \mathcal{C}$, there exists an epimorphism

$$\bigoplus_{i \in I} G \twoheadrightarrow X$$

for some indexing set I , where $\bigoplus$ denotes the coproduct (Definition .1.16) in $\mathcal{C}$.

2. A family $\{G_i\}_{i \in I}$ is called a **generating family** if for every pair of distinct morphisms $f, g : X \rightarrow Y$ in $\mathcal{C}$, there exists some index $i \in I$ and a morphism $h : G_i \rightarrow X$ such that

$$f \circ h \neq g \circ h.$$

In case $\mathcal{C}$ is locally small, this is equivalent to the condition that the collection of representable functors

$$\{\mathrm{Hom}_{\mathcal{C}}(G_i, -) : \mathcal{C} \rightarrow \mathbf{Set}\}_{i \in I}$$

is jointly faithful, which in turn is equivalent to the condition that for every object $X \in \mathcal{C}$, there exists a family of objects $\{G_i\}_{i \in J}$ from the generating set indexed by some set J , and an epimorphism

$$\bigoplus_{i \in J} G_i \twoheadrightarrow X.$$

Definition .1.31 (Topos). There are multiple notions of a topos depending on the context (geometric vs. logical).

1. A **Grothendieck topos** (or **sheaf topos**) is a category (Definition .1.1) equivalent (Definition .1.6) to the category of sheaves (Definition 4.0.2) of sets on a small site. That is, there exists a small site $(\mathcal{C}, J)$ such that the category is equivalent to $\mathrm{Sh}(\mathcal{C}, J, \mathbf{Sets})$.

If a Grothendieck universe (Definition .1.2) $\mathcal{U}$ is available, then a **$\mathcal{U}$ -(Grothendieck/sheaf) topos** is a category equivalent to (Definition .1.6) the category of sheaves of $\mathcal{U}$ -sets (Definition .1.2) on a $\mathcal{U}$ -small site (Definition A.0.5) $(\mathcal{C}, J)$ [?, Exposé IV Définition 1.1].

2. An **elementary topos** is a category which has all finite limits (Definition .1.13), is cartesian closed, and has a subobject classifier.

Unless otherwise specified, a **topos** refers to a Grothendieck/sheaf topos. The plural for topos is **topoi**. *Remark:* Every Grothendieck topos is an elementary topos, but the converse is not true (e.g., the category of finite sets is an elementary topos but not a Grothendieck topos).

Theorem .1.32 (Functor Category of a Grothendieck Topos is a Topos). Let I be a small category (Definition .1.4), and let $\mathcal{S}$ be a Grothendieck topos (Definition .1.31). Then the functor category $\mathrm{Fun}(I, \mathcal{S})$ (Definition .1.33) is a Grothendieck topos.

Definition .1.33 (Diagram in a category and category of diagrams). Let $\mathcal{C}$ be a (large) category (Definition .1.1), and let I be a (large) category (Definition .1.1).

1. A **diagram of shape I in $\mathcal{C}$** is a functor $D : I \rightarrow \mathcal{C}$. We often denote such a diagram by the family $\{D(i)\}_{i \in \mathrm{Ob}(I)}$ with transition maps given by the functorial image of morphisms in I .

A diagram is also synonymously called a **system**. Moreover, the category I is called the **index category** or the **indexing category of the diagram D** .

2. Given two diagrams $D, E : I \rightarrow \mathcal{C}$, a **morphism of diagrams** is simply a natural transformation $D \Rightarrow E$ of the functors D and E .
3. The **category of I -shaped diagrams in $\mathcal{C}$** or simply **diagram category (of I -shaped diagrams in $\mathcal{C}$)**, often denoted $\mathcal{C}^I$, $[I, \mathcal{C}]$, or $\mathrm{Fun}(I, \mathcal{C})$, is the (large) category whose

objects are functors $I \rightarrow \mathcal{C}$ (that is, diagrams of shape I in $\mathcal{C}$) and whose morphisms are natural transformations between such functors. The category $\mathcal{C}^I$ is also called the *functor category of functors $I \rightarrow \mathcal{C}$* . Equivalently, the functor category $\mathcal{C}^I$ is the category $\text{PreShv}(I^{\text{op}}, \mathcal{C})$ of presheaves (Definition 4.0.1) on I^{op} with values in $\mathcal{C}$ and hence notations for presheaf categories are applicable as notations for functor categories.

If $\mathcal{C}$ is locally small (Definition .1.4) and I is small, then $\mathcal{C}^I$ is locally small by Lemma .1.7.

.1.1. Abelian categories.

Definition .1.34 (Abelian category). Let $\mathcal{A}$ be a category. The category $\mathcal{A}$ is an *abelian category* if:

- $\mathcal{A}$ is an additive category.
- Every morphism $f : A \rightarrow B$ has a kernel $\ker(f)$ and a cokernel $\text{coker}(f)$.
- For every morphism $f : A \rightarrow B$, the canonical morphism $\text{coim}(f) \rightarrow \text{im}(f)$ is an isomorphism, where

$$\text{coim}(f) = \text{coker}(\ker(f) \rightarrow A), \quad \text{im}(f) = \ker(B \rightarrow \text{coker}(f)).$$

(♠ TODO: I think I need to re-check this definition) (♠ TODO: coimage)

.

It is also worth considering Grothendieck's additional axioms for abelian categories.

Proposition .1.35. The following are examples of abelian categories (Definition .1.34):

1. The category of R - S bimodules where R, S are (not necessarily commutative) rings (Definition C.0.8) (??).
2. The category $\mathbf{Ab}$ of abelian groups and group homomorphisms is abelian.
3. The category Vect_k of vector spaces over a field k and k -linear maps is abelian.
4. More generally, if R is a noetherian ring, then the category of finitely generated R -modules is abelian.
5. For a ringed space $(X, \mathcal{O}_X)$, the category of $\mathcal{O}_X$ -modules (Definition 7.1.3) is abelian.
(♠ TODO: a quasi-coherent sheaf on a locally ringed space)
6. If X is a scheme (Definition .3.1) (or more generally a locally ringed space), the category of quasi-coherent sheaves on X is abelian.
7. For any essentially small category (Definition .1.26) $\mathcal{C}$ and any abelian category $\mathcal{A}$, the functor category $[\mathcal{C}, \mathcal{A}]$ (Definition .1.33) and the category $\text{PreShv}(\mathcal{C}, \mathcal{A})$ of presheaves (Definition 4.0.1) are abelian. (♠ TODO: apparently, the essentially smallness condition is removable, provided that the sheafification functor exists. However, the essentially small assumption is needed to show that the category of sheaves of $\mathcal{O}$ -modules is a Grothendieck abelian category. Verify all this. Moreover, when working with a big site of a scheme, one typically fixes a universe or work relative to a cardinal cutoff to treat it as essentially small)
8. For any site (Definition A.0.4) $(\mathcal{C}, J)$ on an essentially small category (Definition .1.26) $\mathcal{C}$ and any abelian category $\mathcal{A}$, the category $\text{Shv}(\mathcal{C}, \mathcal{A})$ of sheaves (Definition 4.0.2) is abelian.

9. For any site (Definition A.0.4) $(\mathcal{C}, J)$ on an essentially small category (Definition .1.26) $\mathcal{C}$ and a sheaf of rings (Definition 4.0.2) $\mathcal{O}$ on $\mathcal{C}$, the category $\mathbf{Mod}(\mathcal{O})$ of $\mathcal{O}$ -modules (Definition 7.1.3) is an abelian category.

Definition .1.36. Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be an additive functor between abelian categories (Definition .1.34).

1. F is called *left exact* if it preserves all finite limits (Definition .1.13); since F is an additive functor between abelian categories, this is equivalent to saying that F preserves kernels⁶

Equivalently, for every left exact sequence in $\mathcal{A}$

$$0 \rightarrow A' \xrightarrow{f} A \xrightarrow{g} A''$$

the sequence

$$0 \rightarrow F(A') \xrightarrow{F(f)} F(A) \xrightarrow{F(g)} F(A'')$$

is exact at $F(A')$ and $F(A)$ (i.e., F preserves monomorphisms (Definition .1.27) and exactness at the first two terms).

2. Dually, F is called *right exact* if it preserves all finite colimits (Definition .1.13); since F is an additive functor between abelian categories, this is equivalent to saying that F preserves cokernels.

Equivalently, for every right exact sequence in $\mathcal{A}$

$$A' \xrightarrow{f} A \xrightarrow{g} A'' \rightarrow 0,$$

the sequence

$$F(A') \xrightarrow{F(f)} F(A) \xrightarrow{F(g)} F(A'') \rightarrow 0$$

is exact at $F(A)$ and $F(A'')$ (i.e., F preserves epimorphisms (Definition .1.27) and exactness at the last two terms).

3. F is called *exact* if it is both left and right exact.

Definition .1.37. Let $\mathcal{A}$ be an abelian category (Definition .1.34).

1. $\mathcal{A}$ is said to *have enough injectives* if for every object A in $\mathcal{A}$, there is an monomorphism (Definition .1.27) $A \rightarrow I$ with I an injective object of $\mathcal{A}$.
2. $\mathcal{A}$ is said to *have enough projectives* if for every object A in $\mathcal{A}$, there is a epimorphism (Definition .1.27) $P \rightarrow A$ with P a projective object of $\mathcal{A}$.

Lemma .1.38 (cf. [?, Lemma 2.2.5, Lemma 2.3.6]). Let $\mathcal{A}$ be an abelian category (Definition .1.34) and let $\mathcal{X}$ be a class of objects in $\mathcal{A}$.

⁶To see this equivalence, note that

- finite products (Definition .1.16) coincide with finite direct sums in abelian categories,
- additive functors preserve finite direct sums by definition,
- kernels in a pointed category are equalizers (Definition .1.17), and all equalizers in abelian categories are kernels,
- if all finite products and equalizers exist in a general category, then all finite limits exist in the category (??)

1. If $\mathcal{A}$ has enough objects of class $\mathcal{X}$ on the right, then for every object $A \in \mathcal{A}$ there exists an $\mathcal{X}$ -right resolution of A (Definition .2.7).
2. If $\mathcal{A}$ has enough objects of class $\mathcal{X}$ on the left, then for every object $A \in \mathcal{A}$ there exists an $\mathcal{X}$ -left resolution of A (Definition .2.7).

Note that this is a special case of ?? obtained by letting the complex $M^\bullet$ be the complex such that

$$M^i = \begin{cases} A & \text{if } i = 0 \\ 0 & \text{otherwise} \end{cases}.$$

In particular,

- If $\mathcal{A}$ has enough injective objects (Definition .1.37), then for every object $A \in \mathcal{A}$ there exists an injective resolution of A (Definition .2.7).
- If $\mathcal{A}$ has enough projective objects (Definition .1.37), then for every object $A \in \mathcal{A}$ there exists a projective resolution of A (Definition .2.7).
- If $F : \mathcal{A} \rightarrow \mathcal{B}$ is a left (resp. right) exact functor (Definition .1.36) between abelian categories and $\mathcal{A}$ has enough F -acyclic objects on the right (resp. left), then for every object $A \in \mathcal{A}$, there exists an right (resp. left) F -acyclic resolution of A .

Proof. 1. Let $A \in \mathcal{A}$ be an object. Since $\mathcal{A}$ has enough objects of class $\mathcal{X}$ of the right, there is an object X_0 of $\mathcal{X}$ and a monomorphism $\varepsilon_0 : A \rightarrow X_0$. Let $A_0 = \text{coker } \varepsilon_0$. Inductively, given an object A_{n-1} of $\mathcal{A}$, choose an object X_n of $\mathcal{X}$ and a monomorphism $\varepsilon_n : A_{n-1} \hookrightarrow X_n$. Let $A_n = \text{coker } \varepsilon_n$. In particular, there is a surjection $X_n \twoheadrightarrow A_n$. Let d_n be the composition

$$X_{n-1} \twoheadrightarrow A_{n-1} \xrightarrow{\varepsilon_n} X_n.$$

The chain complex

$$0 \rightarrow A \xrightarrow{\varepsilon_0} X_0 \xrightarrow{d_0} X_1 \xrightarrow{d_1} \dots$$

is thus an $\mathcal{X}$ -right resolution of A .

2. This is simply the dual statement of the next statement.

□

Definition .1.39. (♠ TODO: I think that the definition of derived categories might be doable for more general kinds of resolutions? Perhaps it is that if I have a right exact functor F , then $L^i F$ can be computed with resolutions of F -acyclic objects?) (♠ TODO: Apparently, left/right derived functors may be defined for functors that are additive and preserve finite coproducts, and not necessarily right/left exact; the exactness condition ensures that the zeroth derived functor agrees with F .) Let $\mathcal{A}$ and $\mathcal{B}$ be abelian categories (Definition .1.34), and let

$$F : \mathcal{A} \rightarrow \mathcal{B}$$

be an additive functor.

1. Suppose that the functor F is right exact (Definition .1.36) and suppose that $A \in \mathcal{A}$ is an object for which a projective resolution (Definition .2.7)

$$\cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow A \rightarrow 0$$

exists in $\mathcal{A}$. We define the *left derived object* $L_n F A \in \mathcal{B}$ by applying F to obtain a complex

$$\cdots \rightarrow F(P_2) \rightarrow F(P_1) \rightarrow F(P_0) \rightarrow 0$$

and letting $L_n F(A)$ be the n -th homology object (Definition .2.3) of this complex in $\mathcal{B}$:

$$L_n F(A) := H_n(F(P_\bullet)).$$

The object $L_n F(A)$ is independent of the choice of projective resolution up to natural isomorphism (Proposition .1.40).

By convention, set $L_n F = 0$ for $n < 0$.

The *higher left derived objects* refer to the object $L_n F(A)$ for $n > 0$.

2. Suppose that the functor F is right exact (Definition .1.36) and that $\mathcal{A}$ has enough projectives (Definition .1.37). The *left derived functors* refer to the family of functors

$$L_n F : \mathcal{A} \rightarrow \mathcal{B}, \quad A \mapsto L_n F(A).$$

The *higher left derived functors* refer to the functors $L_n F$ for $n > 0$.

3. Suppose that the functor F is left exact (Definition .1.36) and suppose that $A \in \mathcal{A}$ is an object for which a injective resolution (Definition .2.7)

$$0 \rightarrow A \rightarrow I^0 \rightarrow I^1 \rightarrow I^2 \rightarrow \cdots$$

exists in $\mathcal{A}$. We define the *right derived object* $R_n F A \in \mathcal{B}$, also often denoted by $R^n F A$, by applying F to obtain a complex

$$0 \rightarrow F(I^0) \rightarrow F(I^1) \rightarrow F(I^2) \rightarrow \cdots.$$

and letting $R_n F(A)$ be the n -th cohomology object (Definition .2.3) of this complex in $\mathcal{B}$:

$$R_n F(A) := H^n(F(I_\bullet)).$$

The object $R_n F(A)$ is independent of the choice of injective resolution up to natural isomorphism (Proposition .1.40).

By convention, set $R_n F = 0$ for $n < 0$.

The *higher right derived objects* refer to the object $R_n F(A)$ for $n > 0$.

4. Suppose that the functor F is left exact (Definition .1.36) and that $\mathcal{A}$ has enough injectives (Definition .1.37). The *right derived functors* refer to the family of functors

$$R_n F : \mathcal{A} \rightarrow \mathcal{B}, \quad A \mapsto R_n F(A).$$

The right derived functors are also often denoted by $R^n F$. The *higher right derived functors* refer to the functors $R_n F$ for $n > 0$.

Proposition .1.40 (cf.[?, Lemma 2.4.1]). Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be an additive functor between abelian categories (Definition .1.34). Let A be an object of $\mathcal{A}$.

1. Suppose that F is right exact (Definition .1.36), and suppose that a projective resolution (Definition .2.7)

$$\cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow A \rightarrow 0$$

of A exists in $\mathcal{A}$. Let

$$\cdots \rightarrow Q_2 \rightarrow Q_1 \rightarrow Q_0 \rightarrow A \rightarrow 0$$

be any projective resolution of A in $\mathcal{A}$. For all n , there are natural isomorphisms

$$H_n(F(P_\bullet)) \cong H_n(F(Q_\bullet)).$$

In other words, the left derived objects $L_n F(A)$ (Definition .1.39) is well defined.

2. Suppose that F is left exact (Definition .1.36), and suppose that a injective resolution (Definition .2.7)

$$0 \rightarrow A \rightarrow I^0 \rightarrow I^1 \rightarrow I^2 \rightarrow \cdots$$

of A exists in $\mathcal{A}$. Let

$$0 \rightarrow A \rightarrow Q^0 \rightarrow Q^1 \rightarrow Q^2 \rightarrow \cdots$$

be any injective resolution of A in $\mathcal{A}$. For all n , there are natural isomorphisms

$$H_n(F(I^\bullet)) \cong H_n(F(Q^\bullet)).$$

In other words, the right derived objects $R_n F(A)$ (Definition .1.39) is well defined.

- Proof.* 1. By Lemma .1.41, there is a lift $f : P_\bullet \rightarrow Q_\bullet$ of the identity map $A \rightarrow A$ unique up to chain homotopy. There are then induced natural maps $H_n(F(f)) : H_n(F(P_\bullet)) \rightarrow H_n(F(Q_\bullet))$ (??). There is also a lift $f' : Q_\bullet \rightarrow P_\bullet$ of the identity map $A \rightarrow A$ unique up to chain homotopy, and this also induces natural maps $H_n(F(f')) : H_n(F(Q_\bullet)) \rightarrow H_n(F(P_\bullet))$. The chain maps f and f' are in fact chain homotopy inverses because Lemma .1.41 also implies that any lifts $P_\bullet \rightarrow P_\bullet$ and $Q_\bullet \rightarrow Q_\bullet$ of the identity map $A \rightarrow A$ are chain homotopic to the identity chain maps. Therefore, $H_n(F(f))$ and $H_n(F(f'))$ are inverses of each other as morphisms in $\mathcal{B}$ by ??.
2. This is dual to the previous part.

□

Lemma .1.41 (cf. [?, Porism 2.2.7, Comparison Theorem 2.2.6, Comparison Theorem 2.3.7]). Let $\mathcal{A}$ be an abelian category (Definition .1.34).

1. Let

$$\cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0$$

be a chain complex (Definition .2.1) with P_i projective. For every left resolution (Definition .2.7) $Q_\bullet \rightarrow N$ of an object N , every map $M \rightarrow N$ lifts to a complex map (Definition .2.1) $P_\bullet \rightarrow Q_\bullet$ unique up to chain homotopy.

2. Let

$$0 \rightarrow M \rightarrow I^0 \rightarrow I^1 \rightarrow I^2 \rightarrow \cdots$$

be a (co)chain complex (Definition .2.1) with I^i injective. For every right resolution (Definition .2.7) $N \rightarrow Q^\bullet$ of an object N , every map $N \rightarrow M$ lifts to a complex map (Definition .2.1) $Q^\bullet \rightarrow I^\bullet$ unique up to chain homotopy.

Proof. 1. The map $P_0 \rightarrow M \rightarrow N$ lifts to a map $P_0 \rightarrow Q_0$ because P_0 is projective and $Q_0 \rightarrow N$ is an epimorphism. Inductively suppose that there are morphisms $P_i \rightarrow Q_i$ for $0 \leq i \leq n$, where $n \geq 0$ that make

$$\begin{array}{ccccccccc} P_n & \longrightarrow & P_{n-1} & \longrightarrow & \cdots & \longrightarrow & P_0 & \longrightarrow & M & \longrightarrow & 0 \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \\ Q_n & \longrightarrow & Q_{n-1} & \longrightarrow & \cdots & \longrightarrow & Q_0 & \longrightarrow & N & \longrightarrow & 0 \end{array}$$

into a commuting diagram are established. The morphism $Q_n \rightarrow Q_{n-1}$ (where we let $Q_{-1} = N$ and $P_{-1} = M$ here in case that $n = 0$) acts as 0 when restricted to $\mathfrak{J} := \text{im}(P_{n+1} \rightarrow P_n \rightarrow Q_n)$ because the composition

$$P_{n+1} \rightarrow P_n \rightarrow Q_n \rightarrow Q_{n-1}$$

equals the composition

$$P_{n+1} \rightarrow P_n \rightarrow P_{n-1} \rightarrow Q_{n-1}.$$

In other words, $\mathfrak{J}$ is a subobject of $\ker(Q_n \rightarrow Q_{n-1})$, which is isomorphic to $\text{im}(Q_{n+1} \rightarrow Q_n)$ by the acyclicity of the sequence of the Q_i 's. Therefore, we have a map $P_{n+1} \twoheadrightarrow \mathfrak{J} \hookrightarrow \text{im}(Q_{n+1} \rightarrow Q_n)$ along with an epimorphism $Q_{n+1} \twoheadrightarrow \text{im}(Q_{n+1} \rightarrow Q_n)$. Since P_{n+1} is projective, the former map lifts to a map $P_{n+1} \rightarrow Q_{n+1}$ in a way that is compatible with the latter, i.e. the following commutes:

$$\begin{array}{ccc} P_{n+1} & & \\ \downarrow \text{dotted} & \searrow & \\ Q_{n+1} & \longrightarrow & \text{im}(Q_{n+1} \rightarrow Q_n). \end{array}$$

By induction, this shows that $M \rightarrow N$ lifts to a morphism $P_\bullet \rightarrow Q_\bullet$ of complexes.

We show that the morphism of complexes is unique up to chain homotopy, i.e. if $f_1, f_2 : P_\bullet \rightarrow Q_\bullet$ are two morphisms of complexes, then $h := f_1 - f_2$ is null homotopic. We construct a chain contraction $\{s_n : P_n \rightarrow Q_{n+1}\}$ of h by induction on n . If $n < 0$, then set $s_n = 0$. If $n = 0$, note that the composition $P_0 \xrightarrow{h_0} Q_0 \rightarrow N$ equals the composition $P_0 \rightarrow M \xrightarrow{0} N$, so $\text{im}(h_0)$ is a subobject of $\ker(Q_0 \rightarrow N) \cong \text{im}(Q_1 \rightarrow Q_0)$. The projectivity of P_0 thus yields a lift $s_0 : P_0 \rightarrow Q_1$ such that h_0 equals the composition $P_0 \xrightarrow{s_0} Q_1 \xrightarrow{d} Q_0$:

$$\begin{array}{ccc} & P_0 & \\ & \downarrow h_0 & \\ Q_1 & \xrightarrow{d} & Q_0 \end{array} \quad \begin{array}{c} \nearrow \text{dotted } s_0 \\ \nwarrow \end{array}$$

Note moreover that $h_0 = ds_0 + s_{-1}d$ because $s_{-1} = 0$. Inductively suppose that we have maps s_i for $i \leq n$ such that $h_n = ds_n + s_{n-1}d$ or equivalently that $ds_n = h_n - s_{n-1}d$. Consider the map $h_{n+1} - s_nd : P_{n+1} \rightarrow Q_{n+1}$. Compute

$$d(h_{n+1} - s_nd) = dh_{n+1} - ds_nd = dh_{n+1} - (h_n - s_{n-1}d)d = (dh_{n+1} - h_nd) + s_{n-1}dd = 0$$

Therefore, $\text{im}(h_{n+1} - s_n d)$ is a subobject of $\ker(Q_{n+1} \rightarrow Q_n) \cong \text{im}(Q_{n+2} \rightarrow Q_{n+1})$, which is in turn a quotient of Q_{n+2} . Since P_{n+1} is projective, there is a morphism $s_{n+1} : P_{n+1} \rightarrow Q_{n+2}$ such that $ds_{n+1} = h_{n+1} - s_n d$.

$$\begin{array}{ccc} & & P_{n+1} \\ & \nearrow s_{n+1} & \downarrow h_n - s_{n-1}d = ds_n \\ Q_{n+2} & \xrightarrow{d} & \text{im}(Q_{n+2} \rightarrow Q_{n+1}) \cong \ker(Q_{n+1} \rightarrow Q_n) \end{array}$$

The s_n thus form a chain contraction as needed.

2. This is simply dual to the previous part.

□

2. Homological algebra.

Definition .2.1 (Chain complex in a preadditive category). Let $\mathcal{A}$ be a preadditive category and let I be a totally ordered set (typically $\mathbb{Z}$, but $I \subseteq \mathbb{Z}$ is also allowed).

1. A **chain complex** (or **homological (chain) complex**) $(K_\bullet, d_\bullet)$ in $\mathcal{A}$ indexed by I is the homological convention for sequences with decreasing degrees. It consists of:
 - Objects $\{K_i\}_{i \in I}$ in $\mathcal{A}$, called the **terms in degree i** ,
 - Morphisms $d_i : K_i \rightarrow K_{i-1}$ in $\mathcal{A}$, called the **boundary maps** or **differentials in degree i** ,

such that for every $i \in I$, $d_{i-1} \circ d_i = 0$. That is,

$$K_\bullet : \cdots \xrightarrow{d_{i+1}} K_i \xrightarrow{d_i} K_{i-1} \xrightarrow{d_{i-1}} K_{i-2} \rightarrow \cdots$$

with $d_{i-1}d_i = 0$ for all i . We typically use the notation $K_\bullet = (K_i, d_i)_{i \in I}$.

2. Dually, a **cochain complex** (or **cohomological complex**) $(K^\bullet, d^\bullet)$ in $\mathcal{A}$ follows the cohomological convention with increasing degrees. It consists of objects $\{K^i\}_{i \in I}$ and **coboundary maps** $d^i : K^i \rightarrow K^{i+1}$ such that $d^{i+1} \circ d^i = 0$:

$$K^\bullet : \cdots \xrightarrow{d^{i-1}} K^i \xrightarrow{d^i} K^{i+1} \xrightarrow{d^{i+1}} K^{i+2} \rightarrow \cdots$$

We typically use the notation $K^\bullet = (K^i, d^i)_{i \in I}$.

3. Let $K_\bullet = (K_i, d_i^K)$ and $L_\bullet = (L_i, d_i^L)$ be chain complexes (Definition .2.1) in $\mathcal{A}$ indexed by the same set I . A **morphism of chain complexes** (or **chain map**)

$$f_\bullet : K_\bullet \rightarrow L_\bullet$$

consists of morphisms $f_i : K_i \rightarrow L_i$ for all $i \in I$, such that for every $i \in I$, the following diagram commutes:

$$\begin{array}{ccc} K_i & \xrightarrow{d_i^K} & K_{i-1} \\ \downarrow f_i & & \downarrow f_{i-1} \\ L_i & \xrightarrow{d_i^L} & L_{i-1} \end{array}$$

i.e., $d_i^L \circ f_i = f_{i-1} \circ d_i^K$.

A **morphism of cochain complexes** $f^\bullet : K^\bullet \rightarrow L^\bullet$ is defined similarly, satisfying the commutativity condition $d_L^i \circ f^i = f^{i+1} \circ d_K^i$.

The collection of these objects and morphisms forms a category. Notation for these categories is as follows:

- $\mathbf{Ch}(\mathcal{A})$ or $\mathbf{Ch}(\mathcal{A})$ is often used as a general term.
- To be explicit about the indexing convention, one uses $\mathbf{Ch}_\bullet(\mathcal{A})$ for chain complexes and $\mathbf{Ch}^\bullet(\mathcal{A})$ (or sometimes $\mathbf{CoCh}(\mathcal{A})$) for cochain complexes.
- The set of chain maps between two complexes is denoted by $\mathbf{Hom}_{\mathbf{Ch}(\mathcal{A})}(K_\bullet, L_\bullet)$; it is an abelian group under pointwise addition $(f + g)_i = f_i + g_i$.

Notations used to write chain complexes are *homological conventions* whereas notations used to write cochain complexes are *cohomological conventions*. In homological conventions, indices are written as subscripts and decrease when “following the differential maps”. In cohomological conventions, indices are written as superscripts and increase when “following the differential maps”. The theories of chain complexes and of cochain complexes are not genuinely different, however, as the conventions are essentially notational — starting from a chain complex $(K_\bullet, d_\bullet)$, one can obtain a cochain complex $(K^\bullet, d^\bullet)$ simply by defining $K^i = K_{-i}$ and d^i as d_{-i} and vice versa. As such, cochain complexes may be simply called *chain complexes* when careful distinctions between the conventions are unnecessary. Homological and cohomological notational conventions persist throughout homological algebra, see e.g. spectral sequences, double complexes, etc, but again the distinction between such conventions are merely notational.

Definition .2.2 (Boundedness conditions on chain complexes). Let $\mathcal{A}$ be an additive category.

Cohomological convention: Let $K^\bullet = (K^i, d^i)_{i \in \mathbb{Z}}$ be a cohomologically indexed chain complex in $\mathcal{A}$.

- $K^\bullet$ is *bounded above* if there exists $n \in \mathbb{Z}$ such that $K^i = 0$ for all $i > n$.
- $K^\bullet$ is *bounded below* if there exists $m \in \mathbb{Z}$ such that $K^i = 0$ for all $i < m$.
- $K^\bullet$ is *bounded* if it both bounded above and below, or equivalently if $K^i = 0$ for all but finitely many $i \in \mathbb{Z}$.
- Assuming that $\mathcal{A}$ is an abelian category (Definition .1.34),
 - $K^\bullet$ is *cohomologically bounded above* if there exists $n \in \mathbb{Z}$ such that $H^i(K^\bullet) = 0$ for all $i > n$.
 - $K^\bullet$ is *cohomologically bounded below* if there exists $m \in \mathbb{Z}$ such that $H^i(K^\bullet) = 0$ for all $i < m$.
 - $K^\bullet$ is *cohomologically bounded* if it is both cohomologically bounded above and below, or equivalently if the cohomology objects (Definition .2.3) $H^i(K^\bullet)$ vanish for all but finitely many $i \in \mathbb{Z}$.

Homological convention: Let $K_\bullet = (K_i, d_i)_{i \in \mathbb{Z}}$ be a homologically indexed chain complex in $\mathcal{A}$.

- $K_\bullet$ is *bounded above* if there exists $n \in \mathbb{Z}$ such that $K_i = 0$ for all $i > n$.
- $K_\bullet$ is *bounded below* if there exists $m \in \mathbb{Z}$ such that $K_i = 0$ for all $i < m$.

- $K_\bullet$ is **bounded** if it both bounded above and below, or equivalently if $K_i = 0$ for all but finitely many $i \in \mathbb{Z}$.
- Assuming that $\mathcal{A}$ is an abelian category (Definition .1.34),
 - $K_\bullet$ is **homologically bounded above** if there exists $n \in \mathbb{Z}$ such that $H_i(K_\bullet) = 0$ for all $i > n$.
 - $K_\bullet$ is **homologically bounded below** if there exists $m \in \mathbb{Z}$ such that $H_i(K_\bullet) = 0$ for all $i < m$.
 - $K_\bullet$ is **homologically bounded** if it is both homologically bounded above and below, or equivalently if the homology objects (Definition .2.3) $H_i(K_\bullet)$ vanish for all but finitely many $i \in \mathbb{Z}$.

Definition .2.3 (Chain complexes and their (co)homology objects). Let $\mathcal{A}$ be an abelian category (Definition .1.34).

- For a cochain complex $K^\bullet$, its **cohomology object in degree i** is defined as the quotient of the object of i -cocycles by the object of i -coboundaries:

$$H^i(K^\bullet) := Z^i(K)/B^i(K) = \ker(d^i)/\operatorname{im}(d^{i-1}).$$

In fact, each H^i is a functor $H^i : \mathbf{Ch}^\bullet(\mathcal{A}) \rightarrow \mathcal{A}$ (Definition .2.1); given a morphism $f^\bullet : K^\bullet \rightarrow L^\bullet$ of cochain complexes, it is thus customary to write $H^i(f^\bullet) : H^i(K^\bullet) \rightarrow H^i(L^\bullet)$ for the induced morphism on the i th cohomology objects.

- For a chain complex (Definition .2.1) $K_\bullet$, its **homology object in degree i** is defined as the quotient of the object of i -cycles by the object of i -boundaries:

$$H_i(K_\bullet) := Z_i(K)/B_i(K) = \ker(d_i)/\operatorname{im}(d_{i+1}).$$

In fact, each H_i is a functor $H_i : \mathbf{Ch}_\bullet(\mathcal{A}) \rightarrow \mathcal{A}$ (Definition .2.1); given a morphism $f_\bullet : K_\bullet \rightarrow L_\bullet$ of chain complexes, it is thus customary to write $H_i(f_\bullet) : H_i(K_\bullet) \rightarrow H_i(L_\bullet)$ for the induced morphism on the i th homology objects.

Definition .2.4 (Quasi-isomorphism). Let $\mathcal{A}$ be an abelian category (Definition .1.34), and let

$$f_\bullet : (C_\bullet, d_\bullet^C) \rightarrow (D_\bullet, d_\bullet^D)$$

be a chain map between complexes (Definition .2.1) in $\mathcal{A}$.

The morphism $f_\bullet$ is called a **quasi-isomorphism** if it induces isomorphisms on all cohomology objects, i.e., for every integer n , the induced morphism on homology (Definition .2.3) (or cohomology, depending on the convention)

$$H_n(f_\bullet) : H_n(C_\bullet) \rightarrow H_n(D_\bullet)$$

is an isomorphism in $\mathcal{A}$.

Note that all of these notions are applicable to the cohomological convention as well (Definition .2.1).

Definition .2.5. Let $\mathcal{A}$ be an additive category

1. Let $f : (C_\bullet, d_\bullet^C) \rightarrow (D_\bullet, d_\bullet^D)$ be a morphism of chain complexes (Definition .2.1) in $\mathcal{A}$. The **mapping cone of f** , denoted $\operatorname{Cone}(f)$, is the chain complex defined by:

- Objects: For each n ,

$$\text{Cone}(f)_n = D_n \oplus C_{n-1}.$$

- Differential: For each n , define

$$d_n^{\text{Cone}(f)} : \text{Cone}(f)_n \rightarrow \text{Cone}(f)_{n-1}$$

by the matrix morphism

$$d_n^{\text{Cone}(f)} = \begin{pmatrix} d_n^D & f_{n-1} \\ 0 & -d_{n-1}^C \end{pmatrix} : D_n \oplus C_{n-1} \rightarrow D_{n-1} \oplus C_{n-2}.$$

This construction defines a chain complex, i.e., $d_{n-1}^{\text{Cone}(f)} \circ d_n^{\text{Cone}(f)} = 0$.

2. Dually, let $g : (C^\bullet, d_C^\bullet) \rightarrow (D^\bullet, d_D^\bullet)$ be a morphism of cochain complexes (Definition .2.1) in $\mathcal{A}$.

The *mapping cone of g* , denoted $\text{Cone}(g)$, is the cochain complex (Definition .2.1) defined by:

- Objects: For each n ,

$$\text{Cone}(g)^n = D^n \oplus C^{n+1}.$$

- Differential: For each n , define

$$d_{\text{Cone}(g)}^n : \text{Cone}(g)^n \rightarrow \text{Cone}(g)^{n+1}$$

by the matrix morphism

$$d_{\text{Cone}(g)}^n = \begin{pmatrix} d_D^n & g^{n+1} \\ 0 & -d_C^{n+1} \end{pmatrix} : D^n \oplus C^{n+1} \rightarrow D^{n+1} \oplus C^{n+2}.$$

This construction defines a cochain complex, i.e., $d_{\text{Cone}(g)}^{n+1} \circ d_{\text{Cone}(g)}^n = 0$.

Given a chain complex $(C_\bullet, d_\bullet)$ (or a cochain complex), its *mapping cone* $\text{Cone}(C_\bullet)$ simply refers to the mapping cone of the identity morphism $\text{id}_{C_\bullet} : C_\bullet \rightarrow C_\bullet$.

Definition .2.6. Let $\mathcal{A}$ be an abelian category (Definition .1.34).

1. The *canonical truncations* of a chain complex (Definition .2.1) $A_\bullet$ in $\mathbf{Ch}(\mathcal{A})$ (Definition .2.1) are defined by

$$(\tau_{\geq n} A)_i = \begin{cases} A_i, & i > n, \\ \ker(d_n : A_n \rightarrow A_{n-1}), & i = n, \\ 0, & i < n, \end{cases} \quad (\tau_{\leq n} A)_i = \begin{cases} 0, & i > n, \\ \text{coker}(d_{n+1} : A_{n+1} \rightarrow A_n), & i = n, \\ A_i, & i < n. \end{cases}$$

The differentials are the restrictions and/or quotient maps induced from $A_\bullet$. In particular,

$$H_i(\tau_{\geq n} A_\bullet) = \begin{cases} H_i(A_\bullet), & i \geq n, \\ 0, & i < n, \end{cases} \quad \text{and} \quad H_i(\tau_{\leq n} A_\bullet) = \begin{cases} H_i(A_\bullet), & i \leq n, \\ 0, & i > n. \end{cases}$$

The assignments $A_\bullet \mapsto \tau_{\geq n} A_\bullet$ and $A_\bullet \mapsto \tau_{\leq n} A_\bullet$ extend to endofunctors

$$\tau_{\geq n}, \tau_{\leq n} : \mathbf{Ch}(\mathcal{A}) \rightarrow \mathbf{Ch}(\mathcal{A}),$$

called the *truncation functors*. They are natural in both $A_\bullet$ and n , and fit into canonical morphisms of complexes

$$\tau_{\geq n} A_\bullet \longrightarrow A_\bullet \longrightarrow \tau_{\leq n} A_\bullet.$$

2. Similarly, let $A^\bullet$ be a cochain complex in $\mathbf{Ch}(\mathcal{A})$, i.e.

$$\cdots \xrightarrow{d^{n-2}} A^{n-1} \xrightarrow{d^{n-1}} A^n \xrightarrow{d^n} A^{n+1} \xrightarrow{d^{n+1}} \cdots$$

with $d^{n+1} \circ d^n = 0$ for all $n \in \mathbb{Z}$. The *canonical truncations of $A^\bullet$* are defined by

$$(\tau_{\leq n} A)^i = \begin{cases} A^i, & i < n, \\ \ker(d^n : A^n \rightarrow A^{n+1}), & i = n, \\ 0, & i > n, \end{cases} \quad (\tau_{\geq n} A)^i = \begin{cases} 0, & i < n, \\ \operatorname{coker}(d^{n-1} : A^{n-1} \rightarrow A^n), & i = n, \\ A^i, & i > n. \end{cases}$$

The differentials are the restrictions or quotient maps induced by those of $A^\bullet$. These truncations satisfy

$$H^i(\tau_{\leq n} A^\bullet) = \begin{cases} H^i(A^\bullet), & i \leq n, \\ 0, & i > n, \end{cases} \quad \text{and} \quad H^i(\tau_{\geq n} A^\bullet) = \begin{cases} 0, & i < n, \\ H^i(A^\bullet), & i \geq n. \end{cases}$$

They also extend to endofunctors

$$\tau_{\leq n}, \tau_{\geq n} : \mathbf{Ch}(\mathcal{A}) \rightarrow \mathbf{Ch}(\mathcal{A}),$$

natural in both $A^\bullet$ and n , fitting into canonical morphisms of cochain complexes

$$\tau_{\leq n} A^\bullet \longrightarrow A^\bullet \longrightarrow \tau_{\geq n} A^\bullet.$$

Definition .2.7. Let $\mathcal{A}$ be an abelian category (Definition .1.34) and let $\mathcal{X}$ be a class of objects in $\mathcal{A}$. Let M be an object of $\mathcal{A}$.

1. A *right resolution of M* is a cochain complex (Definition .2.1) $I^\bullet$ with $I^i = 0$ for $i < 0$ and a map $M \rightarrow I^0$ such that the augmented complex

$$0 \rightarrow M \rightarrow I^0 \rightarrow I^1 \rightarrow I^2 \rightarrow \cdots$$

is exact.

2. A *left resolution of M* is a chain complex (Definition .2.1) $P_\bullet$ with $P_i = 0$ for $i < 0$ and a map $P_0 \rightarrow M$ such that the augmented complex

$$\cdots P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0$$

is exact.

3. An *$\mathcal{X}$ -left resolution* of an object $M \in \mathcal{A}$ a left resolution (Definition .2.7) by objects of $\mathcal{X}$, i.e. an exact complex

$$\cdots \rightarrow X_2 \rightarrow X_1 \rightarrow X_0 \rightarrow M \rightarrow 0$$

with each $X_i \in \mathcal{X}$.

4. An *$\mathcal{X}$ -right resolution* of an object $M \in \mathcal{A}$ a right resolution (Definition .2.7) by objects of $\mathcal{X}$, i.e. an exact complex

$$0 \rightarrow M \rightarrow X^0 \rightarrow X^1 \rightarrow X^2 \rightarrow \cdots$$

with each $X_i \in \mathcal{X}$.

5. A **projective resolution of M** is a left resolution $P^\bullet$ for which the objects P^i are all projective.
6. An **injective resolution of M** is a right resolution $I^\bullet$ for which the objects I^i are all injective.

The notion of right/left resolutions above may be regarded as special cases of the notions of right/left resolutions of Definition B.6.1 where $\mathcal{A}$ is embedded into the category $\mathbf{Ch}(\mathcal{A})$ by identifying an object A of $\mathcal{A}$ with the (co)chain complex concentrated in degree 0 associated to A — the right/left resolutions as defined above are then right/left resolutions of $A^\bullet$ by cochain/chain complexes $K^\bullet$ whose negative degree components are 0.

.3. Schemes.

Definition .3.1 (Scheme). A **scheme** is a locally ringed space $(X, \mathcal{O}_X)$ that admits an open cover $\{U_i\}_{i \in I}$ such that each $(U_i, \mathcal{O}_X|_{U_i})$ (??) is isomorphic (as a locally ringed space) to an affine scheme $(\mathrm{Spec}(A_i), \mathcal{O}_{\mathrm{Spec}(A_i)})$ for some commutative ring (Definition C.0.1) A_i . In other words, a scheme is a locally ringed space locally isomorphic to affine schemes.

As is the case with ringed spaces in general, the sheaf $\mathcal{O}_X$ may be suppressed from the notation and only X may be used to denote a ringed space. Similarly, the sheaf $\mathcal{O}_X$ is called the **structure sheaf of X** .

Definition .3.2 (Morphism of schemes). Let $(X, \mathcal{O}_X)$ and $(Y, \mathcal{O}_Y)$ be schemes (Definition .3.1). A **morphism of schemes** is a morphism as locally ringed spaces.

In particular, there is a category (Definition .1.1), often denoted by **Sch**, **Sch** etc., whose objects are schemes and whose morphisms are morphisms of schemes.

Definition .3.3 (Scheme over a scheme). Let $(S, \mathcal{O}_S)$ be a scheme. A **scheme over S** (or an **S -scheme**) is a scheme $(X, \mathcal{O}_X)$ together with a morphism of schemes

$$\pi : (X, \mathcal{O}_X) \rightarrow (S, \mathcal{O}_S).$$

This morphism π is called the **structure morphism of the scheme X over S** .

If $S = \mathrm{Spec}(R)$ is an affine scheme for a commutative ring R , then an S -scheme is synonymously called an **R -scheme** or a **scheme over R** .

Let $(S, \mathcal{O}_S)$ be a scheme, and let $(X, \mathcal{O}_X)$ and $(Y, \mathcal{O}_Y)$ be schemes over S with structure morphisms

$$\pi_X : X \rightarrow S, \quad \pi_Y : Y \rightarrow S.$$

A **morphism of S -schemes** (or synonymously a **S -scheme morphism**) is a morphism of schemes (Definition .3.2)

$$(f, f^\#) : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$$

such that the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \pi_X \downarrow & & \downarrow \pi_Y \\ S & = & S \end{array}$$

In other words,

$$\pi_Y \circ f = \pi_X.$$

Given a fixed scheme S , there is a category, often denoted by $\mathbf{Sch}_S$, $\mathbf{Sch}/_S$, $\mathbf{Sch}/S$, $\mathbf{Sch}_S$, $\mathbf{Sch}/_S$, $\mathbf{Sch}/S$ etc. whose objects are schemes T over S and whose morphisms $T_1 \rightarrow T_2$ are morphisms of schemes over S . If $S = \operatorname{Spec} R$ for some commutative ring R , then we may instead write $\mathbf{Sch}_R$ to denote $\mathbf{Sch}_{\operatorname{Spec} R}$, etc. It is noteworthy that $\mathbf{Sch}_{\mathbb{Z}}$ coincides with the category $\mathbf{Sch}$ (Definition .3.2) of all schemes. In other words, a $\mathbb{Z}$ -scheme can be identified simply with a scheme.

Equivalently, the category $\mathbf{Sch}/_S$ is the category of schemes over S in the sense of Definition A.0.6.

Definition .3.4. Let $f : X \rightarrow Y$ be a morphism of schemes (Definition .3.2). We say that f is an *affine morphism* and that X is an *affine Y -scheme* if for every affine open $V = \operatorname{Spec} B \subseteq Y$, the preimage $U = f^{-1}(V)$ is an affine scheme.

Definition .3.5. Let $f : X \rightarrow Y$ be a morphism of schemes (Definition .3.2). (♠ **TODO: locally of finite type**) We say that f is a *finite type morphism* if the following equivalent conditions hold:

1. for every affine open $\operatorname{Spec} B \subseteq Y$, the inverse image $f^{-1}(\operatorname{Spec} B)$ is quasi-compact and has a covering $f^{-1}(\operatorname{Spec} B) = \bigcup_j \operatorname{Spec} A_j$ by affine opens such that each A_j is a *finitely generated B algebra*.
2. there exists an affine open covering $Y = \bigcup_i \operatorname{Spec} B_i$ of Y such that the inverse image $f^{-1}(\operatorname{Spec} B_i)$ has a finite cover $\{\operatorname{Spec} A_{ij}\}_j$ by affine opens where each A_{ij} is a finitely generated B_i algebra.
3. f is quasi-compact and locally of finite type.

When X is equipped with a finite type morphism $f : X \rightarrow Y$, we say that X is a *finite type scheme over Y* or a *finite type Y -scheme* or a *Y -scheme of finite type* (Definition .3.3), etc.

Definition .3.6 (Finitely presented algebra over a ring). Let R be a (not necessarily commutative) ring (Definition C.0.8). An R -algebra A is said to be *finitely presented* if there exists an integer $n \geq 0$ and a surjective R -algebra homomorphism

$$\varphi : R\langle x_1, \dots, x_n \rangle \twoheadrightarrow A$$

where $R\langle x_1, \dots, x_n \rangle$ is the free R -algebra on n generators, such that the kernel $\ker(\varphi)$ is a finitely generated two-sided ideal (Definition C.0.2) of $R\langle x_1, \dots, x_n \rangle$.

In other words, A admits a presentation as

$$A \cong R\langle x_1, \dots, x_n \rangle / I,$$

where I is a finitely generated two-sided ideal.

If R and A are commutative rings, this recovers the usual definition of a finitely presented commutative R -algebra by replacing $R\langle x_1, \dots, x_n \rangle$ with the polynomial ring $R[x_1, \dots, x_n]$ and I a finitely generated ideal.

Definition .3.7. (♠ TODO: TODO: define a (commutative) group scheme) (♠ TODO: TODO: define the multiplicative group scheme) Let S be a scheme. The *additive group scheme over S* is the group scheme $\mathbb{G}_a = \mathbb{G}_{a,S}$ over S defined by

$$\mathbb{G}_{a,S} = \operatorname{Spec} \mathcal{O}_S[T]$$

with group structure morphisms:

- **Comultiplication (Addition):** The morphism $\Delta : \mathbb{G}_{a,S} \rightarrow \mathbb{G}_{a,S} \times_S \mathbb{G}_{a,S}$ corresponding to the ring homomorphism $\mathcal{O}_S[T] \rightarrow \mathcal{O}_S[T] \otimes_{\mathcal{O}_S} \mathcal{O}_S[T]$ given by $T \mapsto T \otimes 1 + 1 \otimes T$.
- **Counit (Identity):** The morphism $\varepsilon : \mathbb{G}_{a,S} \rightarrow S$ corresponding to $T \mapsto 0$.
- **Coinverse (Inversion):** The morphism $\iota : \mathbb{G}_{a,S} \rightarrow \mathbb{G}_{a,S}$ corresponding to $T \mapsto -T$.

Thus, $(\mathbb{G}_{a,S}, \Delta, \varepsilon, \iota)$ is a commutative group scheme over S . Note that we may speak of the *additive group scheme over a ring R* as the additive group scheme over $\operatorname{Spec} R$.

Definition .3.8 (Dimension of a Scheme). Let X be a scheme (Definition .3.1) with underlying topological space $|X|$.

- The *dimension at a point $x \in |X|$* , denoted $\dim_x(X)$, is the Krull dimension of the local ring (Definition C.0.21) $\mathcal{O}_{X,x}$ (Definition C.0.23), which is local by definition. This is the supremum of the lengths n of chains of prime ideals

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n \subseteq \mathcal{O}_{X,x}.$$

- The *dimension of the scheme X* is defined as

$$\dim(X) := \sup_{x \in |X|} \dim_x(X).$$

Equivalently, it is the supremum of the lengths of chains of distinct irreducible closed subsets of $|X|$ ordered by inclusion. It is also equivalent to the dimension of the underlying topological space $|X|$.

APPENDIX A. PRESHEAVES AND SHEAVES

Definition A.0.1 ([?, Exposé I Définition 4.1]). Let C be a (large) category (Definition .1.1).

1. A *sieve S on the category C* is a full subcategory (Definition 8.3.2) D of C such that for any object U of C there exists an object V of (♠ TODO: correctly parse the definition)
2. A *sieve S on an object $U \in \operatorname{Ob}(C)$* is a collection of morphisms in C with codomain U that is closed under precomposition by any compatible morphism in C . In other words, S is a sieve if for every $f : V \rightarrow U$ in S and morphism $g : W \rightarrow V$ in C , the composition $f \circ g : W \rightarrow U$ is also in S .

Given a morphism $f : V \rightarrow U$ in a sieve S , we also say that *f factors through U* .

Definition A.0.2. Let C be a category, let $U \in \operatorname{Ob}(C)$, and let S be a sieve on U (Definition A.0.1). For a morphism $f : V \rightarrow U$ in C , the *pullback sieve f^*S* (or *basechange sieve $S \times_U V$*) on V is defined by

$$f^*S = \{g : W \rightarrow V \mid f \circ g \in S\}.$$

In other words, f^*S consists of all morphisms into V whose composite with f belongs to the sieve S on U . It is indeed a sieve.

Definition A.0.3. Let $\mathcal{C}$ be a category (Definition .1.1) and $U \in \mathcal{C}$ an object. Let $\mathcal{S} = \{f_i : U_i \rightarrow U\}_{i \in I}$ be a family of morphisms with codomain U .

The *sieve generated by $\mathcal{S}$* , denoted $\langle \mathcal{S} \rangle$ or $\langle \mathcal{S} \rangle$, is the smallest sieve on U (Definition A.0.1) containing all the morphisms in $\mathcal{S}$.

Explicitly, a morphism $h : V \rightarrow U$ belongs to the generated sieve if and only if h factors through some morphism in $\mathcal{S}$. That is, there exists an index $i \in I$ and a morphism $g : V \rightarrow U_i$ such that

$$h = f_i \circ g.$$

Definition A.0.4 (Grothendieck topology). Let $\mathcal{U}$ be a universe (Definition .1.2).

1. (See [?, Exposé II, Définition 1.1]) Let $\mathcal{C}$ be a category (Definition .1.1). A *Grothendieck topology on $\mathcal{C}$* assigns to each object U of $\mathcal{C}$ a collection $J(U)$ of sieves (Definition A.0.1) $\{U_i \rightarrow U\}_{i \in I}$, each called a *covering sieve of U* , satisfying:
 - (a) (Stability under “base change”): If $S \in J(U)$ is a covering sieve of an object U , and $f : V \rightarrow U$ is any morphism in $\mathcal{C}$, then the pullback sieve (Definition A.0.2) f^*S is a covering sieve of U .
 - (b) (Local character condition) If S is a sieve on U , and if there exists a covering sieve $R \in J(U)$ such that for all $f : V \rightarrow U$ in R the pullback sieve (Definition A.0.2) f^*S is in $J(V)$, then $S \in J(U)$.
 - (c) The maximal sieve is a covering sieve.

Some will refer to a Grothendieck topology as simply a *topology*, not to be confused with the related, but less general, notion of a topology on a set (Definition C.0.6).

2. (See [?, Exposé II, 1.1.5]) A *site* is a category $\mathcal{C}$ equipped with a Grothendieck topology. When we are working with a Grothendieck pretopology (Definition 4.2.2) K on a category $\mathcal{C}$, we may regard $\mathcal{C}$ as a site by equipping it with the Grothendieck topology generated by (Definition 4.2.1) K . In this case, we may call $\mathcal{C}$ equipped with this Grothendieck topology a *site with a pretopology*. Note that the Stacks project notion of “site” is that of a small (Definition .1.4) site (i.e. a site whose underlying category is small) with a pretopology, cf. [?, Tag 00V0] for a discussion for this choice, [?, Tag 00VH] for the definition and [?, Tags 0005 and 0015] for the Stacks project’s convention of “categories” (what we call small categories (Definition .1.4)) and “big categories” (what we call locally small categories (Definition .1.4)). (♠ TODO: talk more about how, at least when the axiom of universes is assumed, topoi coming from general Grothendieck topology are equivalent to those on pretopologies).
3. (See [?, Exposé II, Définition 1.2]) Let $(\mathcal{C}, J)$ be a site. A family of morphisms $(U_i \rightarrow U)_{i \in I}$ is called a *covering family of U (with respect to the site/topology)* or a *cover of U (with respect to the site/topology)* if the sieve generated by (Definition A.0.3) the family is a covering sieve of U .
4. (See [?, Exposé II, Définition 3.0.1]) Let $(\mathcal{C}, J)$ be a site (Definition A.0.4), where J is a Grothendieck topology on $\mathcal{C}$.

A family G of objects $\mathcal{C}$ is called a *topologically generating family of the site/topology* or a *generating family/collection of the site/topology* if for every object $X \in \mathcal{C}$, there is a covering family $\{X_\alpha \rightarrow X\}_{\alpha \in A}$ of X such that every X_α is a member of G .

Equivalently, the Grothendieck topology J is the smallest Grothendieck topology containing all covers of the U_i . Also equivalently, for any $S \in J(X)$, the sieve S contains a covering family $\{V_i \rightarrow X\}$ such that each morphism $V_i \rightarrow X$ factors through some member of G . (♠ TODO: Verify that these claimed equivalences are indeed equivalences)

- Definition A.0.5.** 1. An *small site* is a site (Definition A.0.4) whose underlying category is small (Definition .1.4).
 2. A *locally small site* is a site whose underlying category is locally small (Definition .1.4).
 3. Given a Grothendieck universe (Definition .1.2) $\mathcal{U}$, a *$\mathcal{U}$ -site* is a site (Definition A.0.4) whose underlying category (Definition .1.1) is a $\mathcal{U}$ -(locally small) category (Definition .1.4) and that has a $\mathcal{U}$ -small topologically generating family (Definition A.0.4); a Grothendieck topology making the site a $\mathcal{U}$ -site is called a *$\mathcal{U}$ -topology*.

A $\mathcal{U}$ -site is called *$\mathcal{U}$ -small* if its underlying category is $\mathcal{U}$ -small (Definition .1.4).

Definition A.0.6 (Category of objects over a fixed object). Let $\mathcal{C}$ be a category (Definition .1.1) and let $X \in \text{Ob}(\mathcal{C})$ be a fixed object.

- The *category of objects over X* (or synonymously the *slice category of X in $\mathcal{C}$* or the *over category of X in $\mathcal{C}$*), commonly denoted $\mathcal{C}/X$, $\mathcal{C}_{/X}$, or $(\mathcal{C} \downarrow X)$ is the category defined as follows:
 - An object of $\mathcal{C}/X$ is a morphism $f: A \rightarrow X$ in $\mathcal{C}$, where $A \in \text{Ob}(\mathcal{C})$.
 - A morphism from $f: A \rightarrow X$ to $g: B \rightarrow X$ in $\mathcal{C}/X$ is a morphism $h: A \rightarrow B$ in $\mathcal{C}$ such that the following diagram commutes:

$$\begin{array}{ccc} A & \xrightarrow{h} & B \\ & \searrow f & \downarrow g \\ & & X \end{array}$$

i.e. such that $g \circ h = f$.

- The identity morphisms and composition in $\mathcal{C}/X$ are inherited from $\mathcal{C}$.
- The *category of objects under X* (or synonymously the *coslice category of X in $\mathcal{C}$* or the *under category of X in $\mathcal{C}$*), commonly denoted $X/\mathcal{C}$, $X \backslash \mathcal{C}$, $\mathcal{C}_{X/}$, or $(X \downarrow \mathcal{C})$, is the category defined as follows:
 - An object of $X/\mathcal{C}$ is a morphism $f: X \rightarrow A$ in $\mathcal{C}$, where $A \in \text{Ob}(\mathcal{C})$.
 - A morphism from $f: X \rightarrow A$ to $g: X \rightarrow B$ in $X/\mathcal{C}$ is a morphism $h: A \rightarrow B$ in $\mathcal{C}$ such that the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{f} & A \\ & \searrow g & \downarrow h \\ & & B \end{array}$$

i.e. such that $h \circ f = g$.

- The identity morphisms and composition in $X/\mathcal{C}$ are inherited from $\mathcal{C}$.

Lemma A.0.7. Let $\mathcal{C}$ be a category (Definition .1.1) and let $X \in \text{Ob}(\mathcal{C})$ be a fixed object. The slice category $\mathcal{C}/X$ (Definition A.0.6) has X as its final object (Definition .1.10).

Proof. This is clear. $\square$

Definition A.0.8 (Slice site). Let $(\mathcal{C}, \tau)$ be a site (Definition A.0.4), where τ is a Grothendieck topology on the (locally small or U -locally small (Definition .1.4), if a universe (Definition .1.2) U is available) category $\mathcal{C}$. For a fixed object X in $\mathcal{C}$, the *slice site* (or the *over site*, the *site on the slice category $\mathcal{C}/X$* , the *site induced on the over category $\mathcal{C}/X$* , the *localization of the site $\mathcal{C}$ at the object X* , etc.) $(\mathcal{C}/X, \tau/X)$ is the site whose underlying category is the slice category $\mathcal{C}/X$ (Definition A.0.6), and whose Grothendieck topology τ/X (also denoted by notations such as $\tau|_X$ or τ/X) is defined by declaring a family of morphisms $\{f_i : Y_i \rightarrow Y\}$ in $\mathcal{C}/X$ to be a covering if and only if the family $\{f_i : Y_i \rightarrow Y\}$ is a covering in $(\mathcal{C}, \tau)$.

APPENDIX B. DERIVED CATEGORIES

B.1. Homotopy category of chain complexes of an additive category. We will define homotopy categories and derived categories using the cohomological convention (Definition .2.1)

Definition B.1.1 (Homotopy category $K(\mathcal{A})$). Let $\mathcal{A}$ be an additive category. (♠ **TODO:** describe the distinguished triangles of this category)

1. The *homotopy category* $K(\mathcal{A})$ has as objects the (cochain) complexes over $\mathcal{A}$, and as morphisms the cochain maps (Definition .2.1) modulo cochain homotopy equivalence. More explicitly,

$$\text{Hom}_{K(\mathcal{A})}(C^\bullet, D^\bullet) := \frac{\{\text{cochain maps } C^\bullet \rightarrow D^\bullet\}}{\{\text{cochain homotopies}\}},$$

where morphisms differing by a cochain homotopy are identified.

2. Let

$$F : \mathcal{A} \rightarrow \mathcal{B}$$

be an additive functor of additive categories.

The functor F naturally induces a functor between the homotopy categories of complexes,

$$K(F) : K(\mathcal{A}) \rightarrow K(\mathcal{B}),$$

by applying F degreewise to complexes and cochain maps. That is, for a complex $(C^\bullet, d^\bullet)$ in $\mathcal{A}$, define

$$K(F)(C^\bullet) := (F(C^\bullet), F(d^\bullet)),$$

and for a cochain map $f^\bullet : C^\bullet \rightarrow D^\bullet$, define

$$K(F)(f^\bullet) := (F(f^\bullet)).$$

The functor $K(F)$ respects cochain homotopies, thus it is well-defined on the homotopy category.

3. (See [?, Catégories dérivées Chapitre 1 2-3] for notation) We write $K^+(\mathcal{A})$, $K^-(\mathcal{A})$, and $K^b(\mathcal{A})$ for the full subcategories of bounded below, bounded above, and bounded (Definition .2.2) complexes respectively.
4. (See [?, Catégories dérivées Chapitre 1 2-3] for notation) If $\mathcal{A}$ is abelian, we may also write $K^{\infty,+}(\mathcal{A})$, $K^{\infty,-}(\mathcal{A})$, $K^{\infty,b}(\mathcal{A})$, and $K^{\infty,\emptyset}(\mathcal{A})$ for the full subcategories of cohomologically bounded below, cohomologically bounded above, cohomologically bounded (Definition .2.2), and acyclic complexes respectively. We may further write $K^{?,??}(\mathcal{A})$ for the full subcategory where $? \in \{+, -, b\}$ indicates the boundedness of the complex and $?? \in \{+, -, b, (\text{blank})\}$ indicates the cohomological boundedness of the complex.

The various categories here are (Proposition B.1.6) triangulated categories.

Definition B.1.2 (Triangulated category). A *triangulated category* is a triple $(\mathcal{T}, [1], \Delta)$ where:

- $\mathcal{T}$ is an additive category,
- $[1] : \mathcal{T} \rightarrow \mathcal{T}$ (also denoted by notations such as Σ or T) is an additive auto-equivalence called the *shift* or *suspension functor*,
- Δ is a class of *distinguished triangles* of the form

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} X[1]$$

with objects $X, Y, Z \in \mathcal{T}$ and morphisms f, g, h in $\mathcal{T}$,

satisfying the following axioms:

- (TR1) The class Δ is closed under isomorphisms of triangles and contains the triangles isomorphic to

$$A \xrightarrow{\text{id}} A \rightarrow 0 \rightarrow A[1].$$

Moreover, for every morphism $f : X \rightarrow Y$ in $\mathcal{T}$, there exists a distinguished triangle

$$X \xrightarrow{f} Y \rightarrow Z \rightarrow X[1].$$

- (TR2) (Rotation) If

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} X[1]$$

is a distinguished triangle, then so are

$$Y \xrightarrow{g} Z \xrightarrow{h} X[1] \xrightarrow{-f[1]} Y[1]$$

and

$$Z[-1] \xrightarrow{-h[-1]} X \xrightarrow{f} Y \xrightarrow{g} Z.$$

- (TR3) (Octahedral axiom) Given morphisms $X \xrightarrow{f} Y$ and $Y \xrightarrow{g} Z$ in $\mathcal{T}$, there exist distinguished triangles

$$\begin{aligned} X &\xrightarrow{f} Y \xrightarrow{u} C(f) \xrightarrow{v} X[1], \\ Y &\xrightarrow{g} Z \xrightarrow{w} C(g) \xrightarrow{x} Y[1], \\ X &\xrightarrow{g \circ f} Z \xrightarrow{y} C(g \circ f) \xrightarrow{z} X[1], \end{aligned}$$

and morphisms

$$s : C(f) \rightarrow C(g \circ f), \quad t : C(g \circ f) \rightarrow C(g)$$

such that the following diagram commutes and its rows and columns are distinguished triangles:

$$\begin{array}{ccccccc}
X & \xrightarrow{f} & Y & \xrightarrow{u} & C(f) & \xrightarrow{v} & X[1] \\
\parallel & & \downarrow g & & \downarrow s & & \parallel \\
X & \xrightarrow{g \circ f} & Z & \xrightarrow{y} & C(g \circ f) & \xrightarrow{z} & X[1] \\
& & \downarrow w & & \downarrow t & & \\
& & C(g) & \xlongequal{\quad} & C(g) & & \\
& & \downarrow x & & \downarrow & & \\
& & Y[1] & \xrightarrow{f[1]} & X[1][1] & &
\end{array}$$

In particular,

- The compositions $s \circ u = y$, $t \circ s = w$, and $x \circ t = v[1]$ hold,
- All rows and columns form distinguished triangles.

(TR4) The class Δ is closed under the shift functor $[1]$: if

$$X \rightarrow Y \rightarrow Z \rightarrow X[1]$$

is distinguished, then so is

$$X[1] \rightarrow Y[1] \rightarrow Z[1] \rightarrow X[2].$$

Definition B.1.3. A functor $T : D_1 \rightarrow D_2$ between triangulated categories (Definition B.1.2) is called *exact* if it is additive, is translation preserving, and transforms distinguished triangles (Definition B.1.2) to distinguished triangles. Oftentimes, a *morphism between triangulated categories* or a *triangulated functor* refers to an exact functor between triangulated categories.

Definition B.1.4. In the context of a triangulated category (Definition B.1.2) $\mathcal{D}$, a multiplicative system (Definition B.2.1) S is *compatible with the triangulation* if:

1. For any $s \in S$, $T(s) \in S$ where T is the translation functor.
2. For any commutative square between the first two steps of two distinguished triangles where the vertical arrows are in S , there exists a morphism in S between the third objects that makes the whole diagram of triangles commute.

Definition B.1.5. Let I be a finite set and $\varepsilon : I \rightarrow \{1, -1\}$ a variance function. Let $(\mathcal{A}_i)_{i \in I}$ and $\mathcal{A}$ be triangulated categories (Definition B.1.2). A multifunctor $F : \prod_{i \in I} \mathcal{A}_i \rightarrow \mathcal{A}$ is an *exact multifunctor of variance ε* if for every index $k \in I$, the partial functor

$$F_k : \mathcal{A}_k \rightarrow \mathcal{A}$$

(obtained by fixing all objects A_j for $j \neq k$) satisfies the following:

1. If $\varepsilon(k) = 1$, F_k is a covariant exact functor: it maps any distinguished triangle $X \rightarrow Y \rightarrow Z \rightarrow T_k X$ in $\mathcal{A}_k$ to a distinguished triangle

$$F_k(X) \rightarrow F_k(Y) \rightarrow F_k(Z) \rightarrow TF_k(X)$$

in $\mathcal{A}$.

2. If $\varepsilon(k) = -1$, F_k is a contravariant exact functor: it maps any distinguished triangle $X \rightarrow Y \rightarrow Z \rightarrow T_k X$ in $\mathcal{A}_k$ to a distinguished triangle

$$F_k(Z) \rightarrow F_k(Y) \rightarrow F_k(X) \rightarrow F_k(T_k^{-1}X)$$

in $\mathcal{A}$, using the isomorphism $F_k(T_k^{-1}X) \cong TF_k(X)$.

Furthermore, F must be a graded functor equipped with natural isomorphisms $\varphi_i : T \circ F \xrightarrow{\sim} F \circ T_i^{\varepsilon_i}$ for each $i \in I$. These must satisfy the anticommutativity condition: for any pair of distinct indices $i, j \in I$ ($i \neq j$), the following diagram of natural transformations commutes with a factor of -1 :

$$(G) \quad \begin{array}{ccc} T^2 F & \xrightarrow{T\varphi_i} & T F T_i^{\varepsilon_i} \\ \downarrow T\varphi_j & & \downarrow \varphi_j T_i^{\varepsilon_i} \\ T F T_j^{\varepsilon_j} & \xrightarrow{-\varphi_i T_j^{\varepsilon_j}} & F T_i^{\varepsilon_i} T_j^{\varepsilon_j} \end{array}$$

Equivalently, the following square is anticommutative:

$$(H) \quad (\varphi_j T_i^{\varepsilon_i}) \circ (T\varphi_i) = -(\varphi_i T_j^{\varepsilon_j}) \circ (T\varphi_j)$$

as morphisms of functors $T^2 F \rightarrow F T_i^{\varepsilon_i} T_j^{\varepsilon_j}$.

In particular in the case that $\varepsilon(i) = 1$ for all i , we are reduced to the following definition: Let $(\mathcal{A}_i)_{1 \leq i \leq n}$ and $\mathcal{A}$ be triangulated categories (Definition B.1.2). A multifunctor $F : \prod \mathcal{A}_i \rightarrow \mathcal{A}$ is called an *exact covariant multifunctor* if it is covariant in each argument and if for every index $i \in \{1, \dots, n\}$ and every fixed collection of objects $(A_j)_{j \neq i}$, the partial functor $F(\dots, A_{i-1}, -, A_{i+1}, \dots) : \mathcal{A}_i \rightarrow \mathcal{A}$ is an exact functor (Definition B.1.3) of triangulated categories.

Proposition B.1.6. Let $\mathcal{A}$ be an additive category.

1. The homotopy category $K(\mathcal{A})$ (Definition B.1.1) is a triangulated category (Definition B.1.2). (♠ TODO: shift of chain complexes, describe what the distinguished triangles are)
2. The subcategory $K^?(\mathcal{A})$ is a triangulated subcategory of $K(\mathcal{A})$ for $? \in \{+, -, b\}$.
3. Let $\mathcal{A}$ be an abelian category. The subcategory $K^{?,??}(\mathcal{A})$ is a triangulated subcategory of $K(\mathcal{A})$ for $? \in \{\infty, +, -, b\}$ and $?? \in \{+, -, b, (\text{blank})\}$.

(♠ TODO: How homotopy categories of chain complexes relate to topological homotopy categories)

Theorem B.1.7 (Exact functor induces a functor on homotopy categories, cf. [?, Exposé XVII, Proposition 1.1.7.]). Let $F : \prod_{i=1}^n \mathcal{A}_i \rightarrow \mathcal{B}$ be an additive multifunctor (Definition C.0.13) of abelian categories (Definition .1.34). F induces exact multifunctors (Definition B.1.5)⁷ between the corresponding homotopy categories (Definition B.1.1):

$$\begin{aligned} K(\mathcal{A}) &\rightarrow K(\mathcal{B}) \\ K^?(\prod \mathcal{A}_i) &\rightarrow K^?(\mathcal{B}) \\ K^{?,??}(\prod \mathcal{A}_i) &\rightarrow K^{?,??}(\mathcal{B}) \end{aligned}$$

(Definition B.1.1). We write these functors by $K(F)$.

B.2. Multiplicative systems in categories.

Definition B.2.1 (Multiplicative system). (♠ TODO: Write a precise definition) Let $\mathcal{C}$ be a category. A *multiplicative system S in $\mathcal{C}$* consists of a collection of morphisms in $\mathcal{C}$ satisfying:

- S contains all identity morphisms,
- S is closed under composition,
- (Ore conditions) For any morphism f in $\mathcal{C}$ and any morphism $s \in S$ with suitable domain and codomain, there exist morphisms to form commutative squares allowing localization.

The precise Ore condition varies depending on context but guarantees the localization exists.

Definition B.2.2 (Localization by a multiplicative system). Given a category $\mathcal{C}$ and a multiplicative system (Definition B.2.1) $S \subseteq \text{Mor}(\mathcal{C})$, the *localization $\mathcal{C}[S^{-1}]$* is a category equipped with a functor

$$Q : \mathcal{C} \rightarrow \mathcal{C}[S^{-1}]$$

that sends every morphism in S to an isomorphism satisfying the universal property: any functor from $\mathcal{C}$ sending the morphisms in S to isomorphisms factors uniquely through Q .

Definition B.2.3. A multiplicative system (Definition B.2.1) S in a category (Definition .1.1) $\mathcal{C}$ is called *saturated* if it satisfies the following condition: any morphism f in $\mathcal{C}$ whose image $Q(f)$ in the localized category (Definition B.2.2) $\mathcal{C}(S^{-1})$ is an isomorphism belongs to S .

Proposition B.2.4. Let $\mathcal{C}$ be a category (Definition .1.1) and S a multiplicative system (Definition B.2.1). There exists a unique saturated multiplicative system (Definition B.2.3) $\bar{S}$ containing S such that the canonical functors $\mathcal{C}(S^{-1}) \rightarrow \mathcal{C}(\bar{S}^{-1})$ define an equivalence of categories (Definition .1.6). $\bar{S}$ consists of all morphisms f such that there exist morphisms g, h with $g \circ f \in S$ and $f \circ h \in S$.

⁷wlog covariant in each variable, but this can be made multi-variant by taking opposite categories

If F is contravariant in the j -th variable, the definitions are adjusted by replacing $\mathcal{A}_j$ with its opposite category $\mathcal{A}_j^{\text{op}}$, thereby interchanging K^+ with K^- and RF with LF for that specific variable.

B.3. Derived categories of abelian categories. (♠ TODO: describe the objects and morphisms in a derived category)

Definition B.3.1 (Derived category). Let $\mathcal{A}$ be an abelian category. Consider the class S of quasi-isomorphisms (Definition .2.4) in the homotopy category $K(\mathcal{A})$ (Definition B.1.1) (Definition B.1.1).

The *derived category* $D(\mathcal{A})$ is the localization (Definition B.2.2) of the homotopy category at the multiplicative system (Definition B.2.1) S :

$$D(\mathcal{A}) := K(\mathcal{A})[S^{-1}],$$

We often write the localization functor (Definition B.2.2) $K(\mathcal{A}) \rightarrow D(\mathcal{A})$ by q or Q .

- We often write $D^+(\mathcal{A})$, $D^-(\mathcal{A})$, and $D^b(\mathcal{A})$ for the full subcategories of cohomologically bounded below, cohomologically bounded above, and cohomologically bounded (Definition .2.2) complexes respectively.

In general, $D(\mathcal{A})$ (and the aforementioned subcategories) may only exist as a large category (Definition .1.1), rather than a locally small category. When $\mathcal{A}$ is a Grothendieck abelian category, however, $D(\mathcal{A})$ is a locally small category (see [?, Tag 09PA]).

Convention B.3.2. Given an abelian category (Definition B.3.1) $\mathcal{A}$, any object M of $\mathcal{A}$ may be identified with an object of the homotopy category $K(\mathcal{A})$ (Definition B.1.1) and the derived category $D(\mathcal{A})$ (Definition B.3.1) — simply take the chain complex concentrated at degree 0 with the object M . In fact, this derived object can often be regarded as an object of “nice” enough subcategories of $D(\mathcal{A})$, such as the bounded derived categories $D^+(\mathcal{A})$, $D^-(\mathcal{A})$, $D^b(\mathcal{A})$.

(♠ TODO: State how a derived category is a triangulated category)

Definition B.3.3. Let $\mathcal{A}$ be an additive category. For $? \in \{+, -, b, (\text{blank})\}$, write $K^?(\mathcal{I})$ (resp. $K^?(\mathcal{P})$) for the full subcategory of $K^?(\mathcal{A})$ (Definition B.1.1) whose objects are appropriately bounded (Definition .2.2) complexes of injectives (resp. projectives).

Even though derived categories of abelian categories may not be guaranteed to be locally small, the following gives sufficient conditions for the appropriately (cohomologically) bounded derived categories to be locally small. The following also shows that appropriately bounded complexes under these circumstances can be represented in the derived category by a complex of injective or projective objects; we might think of such complexes of injectives or projectives as acting like an injective/projective resolution.

Theorem B.3.4 (see e.g. [?, Theorem 10.4.8]). Let $\mathcal{A}$ be an abelian category.

1. If $\mathcal{A}$ has enough injectives (Definition .1.37), then the category $D^+(\mathcal{A})$ (Definition B.3.1) is equivalent to the category $K^+(\mathcal{I})$ (Definition B.3.3), which is locally small (Definition .1.4). In fact, an equivalence is given by

$$K^+(\mathcal{I}) \xrightarrow{q} D^+(\mathcal{I}) \hookrightarrow D^+(\mathcal{A}).$$

2. If $\mathcal{A}$ has enough projectives (Definition .1.37), then the category $D^-(\mathcal{A})$ (Definition B.3.1) is equivalent to the category $K^-(\mathcal{P})$ (Definition B.3.3), which is locally small (Definition .1.4).

$$K^-(\mathcal{P}) \xrightarrow{q} D^-(\mathcal{P}) \hookrightarrow D^-(\mathcal{A}).$$

B.4. Total derived functors. The following subsection mostly takes the point of view of [?, Chapter 10]

Definition B.4.1 (cf. [?, Definition 10.5.1]). Let $\mathcal{A}$ and $\mathcal{B}$ be abelian categories. Let $F : K(\mathcal{A}) \rightarrow K(\mathcal{B})$ (Definition B.1.1) be an exact functor (Definition B.1.3) of triangulated categories. Write $q_{\mathcal{A}} : K(\mathcal{A}) \rightarrow D(\mathcal{A})$ and $q_{\mathcal{B}} : K(\mathcal{B}) \rightarrow D(\mathcal{B})$ for the localization functors (Definition B.3.1),

1. A *(total) right derived functor on F on $K(\mathcal{A})$* is an exact functor $RF : D(\mathcal{A}) \rightarrow D(\mathcal{B})$ together with a natural transformation

$$\xi : q_{\mathcal{B}}F \Rightarrow (RF)q_{\mathcal{A}},$$

which is universal in the sense that if $G : D(\mathcal{A}) \rightarrow D(\mathcal{B})$ is another morphism equipped with a natural transformation $\zeta : q_{\mathcal{B}}F \Rightarrow Gq_{\mathcal{A}}$, then there exists a unique natural transformation $\eta : RF \Rightarrow G$ so that $\zeta_A = \eta_{q_{\mathcal{A}}A} \circ \xi_A$ for all A in $K(\mathcal{A})$.

2. Dually, a *(total) left derived functor on F on $K(\mathcal{A})$* is an exact functor $LF : D(\mathcal{A}) \rightarrow D(\mathcal{B})$ together with a natural transformation

$$\xi : (LF)q_{\mathcal{A}} \Rightarrow q_{\mathcal{B}}F,$$

which is universal in the sense that if $G : D(\mathcal{A}) \rightarrow D(\mathcal{B})$ is another exact functor equipped with a natural transformation $\zeta : Gq_{\mathcal{A}} \Rightarrow q_{\mathcal{B}}F$, then there exists a unique natural transformation $\eta : G \Rightarrow LF$ such that

$$\zeta_A = \xi_A \circ \eta_{q_{\mathcal{A}}A}$$

for all A in $K(\mathcal{A})$.

The universal properties guarantee that if the total derived functors exist, then they are unique up to natural isomorphism, and that

These definitions/ideas may be extended to exact functors $K^?(\mathcal{A}) \rightarrow K(\mathcal{B})$ for $? \in \{+, -, b\}$ and to exact functors $K^{?,??}(\mathcal{A}) \rightarrow K(\mathcal{B})$ for $? \in \{\infty, +, -, b\}$ and $?? \in \{+, -, b, (\text{blank})\}$ (Definition B.1.1)

See Definition B.5.4 for a more general treatment of derived functors

In particular, the above definitions are specializations of those of Definition B.5.8, which fit into the framework of Definition B.5.4.

Proposition B.4.2. See e.g. [?, 10.5.2] Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be an exact functor (Definition .1.36) of abelian categories (Definition .1.34). There is an induced functor

$$D(\mathcal{A}) \rightarrow D(\mathcal{B})$$

(Definition B.3.1) also denoted by $\mathbf{F}$ by abuse of notation.

This functor is constructible in the “simple” way —

- For an object M of $D(\mathcal{A})$, take a chain complex K representing it. The object FM of $D(\mathcal{B})$ is the object represented by the chain complex FK .
- For a morphism $f : M \rightarrow N$ in $D(\mathcal{A})$, take a morphism $\tilde{f} : K \rightarrow L$ representing it. The morphism $Ff : FM \rightarrow FN$ is the morphism represented by $F\tilde{f} : FK \rightarrow FL$.

Due to the exactness of F , these do not depend on the choices of representatives.

The below shows that right and left total derived functors exist for nice enough abelian categories on sufficiently (cohomologically) bounded objects and can be computed using injective/projective resolutions.

Theorem B.4.3 (See e.g. [?, 10.5.6]). Let $\mathcal{A}, \mathcal{B}$ be abelian categories. Let $F : K^+(\mathcal{A}) \rightarrow K(\mathcal{B})$ (Definition B.1.1) be a morphism of triangulated categories (Definition B.1.3).

1. If $\mathcal{A}$ has enough injectives (Theorem B.1.7), then the right derived functor RF (Definition B.4.1) exists (as a functor $D^+(\mathcal{A}) \rightarrow D(\mathcal{B})$ (Definition B.3.1)), and if I is a bounded below (Definition .2.2) complex of injectives, then

$$RF(I) \cong qF(I)$$

where $q : K(\mathcal{B}) \rightarrow D(\mathcal{B})$ is the localization functor (Definition B.3.1).

2. Dually, if $\mathcal{A}$ has enough projectives, then the left derived functor LF exists (as a functor $D^-(\mathcal{A}) \rightarrow D(\mathcal{B})$), and if I is a bounded above complex of projectives, then

$$LF(P) \cong qF(P)$$

Corollary B.4.4. Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be an additive functor between abelian categories (Definition .1.34). Write $q : K(\mathcal{B}) \rightarrow D(\mathcal{B})$ (Definition B.1.1 Definition B.3.1) for the localization functor (Definition B.2.2).

1. Assuming that $\mathcal{A}$ has enough injectives (Definition .1.37), there exists a right derived functor (Definition B.4.1) $RF : D^+(\mathcal{A}) \rightarrow D(\mathcal{B})$ and if $I \in K^+(\mathcal{I})$ (Definition B.3.3) is a bounded below complex of injectives, then

$$RF(I) \cong q(K(F))(I).$$

where $K(F) : K^+(\mathcal{A}) \rightarrow K(\mathcal{B})$ is the exact functor (Definition B.1.3) induced by F (Theorem B.1.7).

If F is additionally left exact (Definition .1.36), then for all objects A of $\mathcal{A}$ we have

$$H^i(RF(A)) \cong R^iF(A)$$

(Definition .1.39) for all integers i where we regard A as the chain complex (Definition .2.1) object of $D^+(\mathcal{A})$ with A concentrated in degree 0 (Convention B.3.2).

2. Assuming that $\mathcal{A}$ has enough projectives (Definition .1.37), there exists a left derived functor (Definition B.4.1) $LF : D^-(\mathcal{A}) \rightarrow D(\mathcal{B})$ and if $P \in K^-(\mathcal{P})$ (Definition B.3.3) is a bounded above complex of projectives, then

$$LF(P) \cong q(K(F))(P).$$

where $K(F) : K^-(\mathcal{A}) \rightarrow K(\mathcal{B})$ is the exact functor (Definition B.1.3) induced by F (Theorem B.1.7).

If F is additionally right exact (Definition .1.36), then for all objects A of $\mathcal{A}$ we have

$$H^{-i}(LF(A)) \cong L_i F(A)$$

(Definition .1.39) for all integers i where we regard A as the chain complex (Definition .2.1) object of $D^+(\mathcal{A})$ with A concentrated in degree 0 (Convention B.3.2).

Proof. Apply Theorem B.4.3 to the exact functor $K(F)$ to obtain the isomorphisms

$$RF(I) \cong q(K(F))(I)$$

$$LF(P) \cong q(K(F))(P).$$

If F is left exact and A is an object of $\mathcal{A}$, then the right derived functors $R^i F(A)$ (Definition .1.39) are defined. Recall that they are defined first by taking an injective resolution (Definition .2.7)

$$0 \rightarrow A \rightarrow I^0 \rightarrow I^1 \rightarrow \dots$$

Since the injective resolution is acyclic, The object A and the complex $0 \rightarrow I^0 \rightarrow I^1 \rightarrow \dots$ are equivalent as objects of $D^+(\mathcal{A})$ as an object of $D^+(\mathcal{A})$ are equivalent objects. Moreover, we have the isomorphism

$$RF(I) \cong q(K(F))(I).$$

In fact, the i th cohomology of the right hand side object precisely describes $R^i F(A)$ by definition. Therefore,

$$H^i(RF(A)) \cong R^i F(A)$$

as desired.

The analogous statement for $L_i F(A)$ holds similarly. □

(♠ TODO: define hyper-derived functors)

Corollary B.4.5 (See e.g. [?, Corollary 10.5.7, Remark 10.5.8]). Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be an additive functor between abelian categories (Definition .1.34).

1. If $\mathcal{A}$ has enough injectives (Definition .1.37), the hyper-derived functors $\mathbb{R}^i F(X)$ are the cohomology of $\mathbf{R}F(X) : \mathbb{R}^i F(X) \cong H^i \mathbf{R}F(X)$ for all i .
2. If $\mathcal{A}$ has enough projectives (Definition .1.37), the hyper-derived functors $\mathbb{L}_i F(X)$ are the cohomology of $\mathbf{L}F(X) : \mathbb{L}_i F(X) \cong H^{-i} \mathbf{L}F(X)$ for all i .

B.5. General derived functors for exact multifunctors between triangulated categories. The following subsection mostly takes the point of view of [?, Exposé XVII, 1.2]

Definition B.5.1 (Ind-category). Let $\mathcal{C}$ be a locally small category (Definition .1.4).

1. The *Ind-category of $\mathcal{C}$* , denoted $\mathbf{Ind}(\mathcal{C})$, is defined as follows:
 - Objects of $\mathbf{Ind}(\mathcal{C})$ are formal filtered colimits (Definition .1.14) of objects in $\mathcal{C}$. More precisely, an object is given by a filtered (Definition .1.12) small category I and a functor

$$X : I \rightarrow \mathcal{C}.$$

- Morphisms between objects $X : I \rightarrow \mathcal{C}$ and $Y : J \rightarrow \mathcal{C}$ are defined by

$$\mathrm{Hom}_{\mathrm{Ind}(\mathcal{C})}(X, Y) := \varprojlim_{i \in I} \varinjlim_{j \in J} \mathrm{Hom}_{\mathcal{C}}(X_i, Y_j),$$

(Definition .1.14) where X_i and Y_j denote the images of $i \in I$ and $j \in J$ under X and Y , respectively.

The composition of morphisms is induced naturally from composition in $\mathcal{C}$. Hence, $\mathrm{Ind}(\mathcal{C})$ is the completion of $\mathcal{C}$ under filtered colimits. Objects of $\mathrm{Ind}(\mathcal{C})$ are called *Ind-objects of $\mathcal{C}$* .

2. The *Pro-category of $\mathcal{C}$* , denoted $\mathrm{Pro}(\mathcal{C})$, is defined as follows:

- Objects of $\mathrm{Pro}(\mathcal{C})$ are formal cofiltered limits (Definition .1.14) of objects in $\mathcal{C}$. More precisely, an object is given by a cofiltered small category I and a functor

$$X : I \rightarrow \mathcal{C}.$$

- Morphisms between objects $X : I \rightarrow \mathcal{C}$ and $Y : J \rightarrow \mathcal{C}$ are defined by

$$\mathrm{Hom}_{\mathrm{Pro}(\mathcal{C})}(X, Y) := \varinjlim_{j \in J} \varprojlim_{i \in I} \mathrm{Hom}_{\mathcal{C}}(X_i, Y_j),$$

where X_i and Y_j denote the images of $i \in I$ and $j \in J$ under X and Y , respectively.

The composition of morphisms is induced naturally from composition in $\mathcal{C}$.

Hence, $\mathrm{Pro}(\mathcal{C})$ is the completion of $\mathcal{C}$ under cofiltered limits. Objects of $\mathrm{Pro}(\mathcal{C})$ are called *Pro-objects of $\mathcal{C}$* .

Since **Sets** has all limits and colimits (♠ TODO:) and hence has all projective and inductive limits and since $\mathcal{C}$ is locally small, $\mathrm{Ind}(\mathcal{C})$ and $\mathrm{Pro}(\mathcal{C})$ are locally small.

The *canonical embedding of a locally small category $\mathcal{C}$ into its Ind-category or Pro-category* is defined by treating each object as a "constant" diagram. Let $\mathbf{1}$ be the category with a single object $*$ and its identity morphism.

1. The Embedding $\iota_{\mathrm{Ind}} : \mathcal{C} \rightarrow \mathrm{Ind}(\mathcal{C})$:

- Objects: An object $C \in \mathcal{C}$ is mapped to the functor $X : \mathbf{1} \rightarrow \mathcal{C}$ sending $*$ to C . This represents C as a formal filtered colimit over the trivial index category.
- Morphisms: Given objects X (representing C) and Y (representing D), the Hom-set in the Ind-category simplifies:

$$\mathrm{Hom}_{\mathrm{Ind}(\mathcal{C})}(X, Y) \cong \varprojlim_{i \in \mathbf{1}} \varinjlim_{j \in \mathbf{1}} \mathrm{Hom}_{\mathcal{C}}(X_i, Y_j) \cong \mathrm{Hom}_{\mathcal{C}}(C, D).$$

2. The Embedding $\iota_{\mathrm{Pro}} : \mathcal{C} \rightarrow \mathrm{Pro}(\mathcal{C})$:

- Objects: An object $C \in \mathcal{C}$ is mapped to the functor $X : \mathbf{1} \rightarrow \mathcal{C}$ sending $*$ to C , representing C as a formal cofiltered limit.
- Morphisms: Similarly, the Pro-morphism definition reduces to:

$$\mathrm{Hom}_{\mathrm{Pro}(\mathcal{C})}(X, Y) \cong \varinjlim_{j \in \mathbf{1}} \varprojlim_{i \in \mathbf{1}} \mathrm{Hom}_{\mathcal{C}}(X_i, Y_j) \cong \mathrm{Hom}_{\mathcal{C}}(C, D).$$

In both cases, the functor ι is fully faithful, allowing us to identify $\mathcal{C}$ with a full subcategory of its respective completion. Given an object C of $\mathcal{C}$, the pro/ind-object $\iota(C)$ may be called the *constant pro/ind-object of C* .

Definition B.5.2. Let $\mathcal{C}$ be a locally small category (Lemma .1.7).

1. An Ind-object (Definition B.5.1) $X = \varinjlim X_i$ is said to be *essentially constant* if it is isomorphic in $\text{Ind}(\mathcal{C})$ to an object in the image of the canonical fully faithful functor $h : \mathcal{C} \rightarrow \text{Ind}(\mathcal{C})$, which maps an object $A \in \text{ob}(\mathcal{C})$ to the Ind-object represented by the constant diagram.
2. A Pro-object (Definition B.5.1) $X = \varprojlim X_i$ is said to be *essentially constant* if it is isomorphic in $\text{Pro}(\mathcal{C})$ to an object in the image of the canonical fully faithful functor $h : \mathcal{C} \rightarrow \text{Pro}(\mathcal{C})$, which maps an object $A \in \text{ob}(\mathcal{C})$ to the Pro-object represented by the constant diagram.

In either definition, the object A may be referred to as the *value/limit/constant value of the Ind/Pro object* or the *object representing the Ind/Pro object*.

Proposition B.5.3. Let $\mathcal{C}$ be a locally small category (Lemma .1.7).

1. An Ind-object (Definition B.5.1) $X = \varinjlim_{i \in I} X_i$ (where I is a filtered (Definition .1.12) index category) is essentially constant (Definition B.5.2) and isomorphic to $A \in \text{ob}(\mathcal{C})$ if and only if there exists a family of morphisms $(f_i : X_i \rightarrow A)_{i \in I}$ such that for every morphism $u : i \rightarrow j$ in I , the diagram

$$\begin{array}{ccc} X_i & \xrightarrow{X(u)} & X_j \\ & \searrow f_i & \downarrow f_j \\ & & A \end{array}$$

commutes, and the induced morphism $\varinjlim X_i \rightarrow A$ (Definition .1.14) is an isomorphism in $\text{Ind}(\mathcal{C})$.

2. A Pro-object (Definition B.5.1) $X = \varprojlim_{i \in I} X_i$ (where I is a filtered (Definition .1.12) index category) is essentially constant (Definition B.5.2) and isomorphic to $A \in \text{ob}(\mathcal{C})$ if and only if there exists a family of morphisms $(g_i : A \rightarrow X_i)_{i \in I}$ such that for every morphism $u : i \rightarrow j$ in I , the diagram

$$\begin{array}{ccc} A & \xrightarrow{g_i} & X_i \\ & \searrow g_j & \downarrow X(u) \\ & & X_j \end{array}$$

commutes, and the induced morphism $A \rightarrow \varprojlim X_i$ (Definition .1.14) is an isomorphism in $\text{Pro}(\mathcal{C})$.

Definition B.5.4 (Mixed-Variance Derived Multifunctors, cf. [?, Exposé XVII, Définition 1.2.1., 1.2.5.]). Let $(\mathcal{A}_i)_{1 \leq i \leq n}$ and $(\mathcal{B}_j)_{1 \leq j \leq m}$ be families of triangulated categories (Definition B.1.2), and let $\mathcal{C}$ be a triangulated category. Let

$$F : \prod_{i=1}^n \mathcal{A}_i \times \prod_{j=1}^m \mathcal{B}_j \rightarrow \mathcal{C}$$

be an exact multifunctor (Definition B.1.5), covariant in $\mathcal{A}_i$ and contravariant in $\mathcal{B}_j$. Let S_i, T_j, U be saturated multiplicative systems (Definition B.2.3) in $\mathcal{A}_i, \mathcal{B}_j, \mathcal{C}$ respectively

assumed compatible with the triangulated structure (Definition B.1.4) of its respective category. Write

$$\begin{aligned} Q_{A_i} &: \mathcal{A}_i \rightarrow \mathcal{A}_i(S_i^{-1}) \\ Q_{B_j} &: \mathcal{B}_j \rightarrow \mathcal{B}_j(T_j^{-1}) \\ Q_C &: \mathcal{C} \rightarrow \mathcal{C}(U^{-1}) \end{aligned}$$

for the localization functors (Definition B.2.2).

1. The *right derived functor* RF of F relative to (S_i, T_j) and U is the functor

$$RF : \prod_{i=1}^n \mathcal{A}_i(S_i^{-1}) \times \prod_{j=1}^m \mathcal{B}_j(T_j^{-1}) \rightarrow \text{Ind}(\mathcal{C}(U^{-1}))$$

which makes the following diagram commutative

$$\begin{array}{ccc} \prod_{i=1}^n \mathcal{A}_i \times \prod_{j=1}^m \mathcal{B}_j & \xrightarrow{F} & \mathcal{C} \\ \downarrow \prod Q_{A_i} \times \prod Q_{B_j} & & \downarrow \bar{Q}_C \\ \prod_{i=1}^n \mathcal{A}_i(S_i^{-1}) \times \prod_{j=1}^m \mathcal{B}_j(T_j^{-1}) & \xrightarrow{RF} & \text{Ind}(\mathcal{C}(U^{-1})) \end{array}$$

where $\bar{Q}_C$ is the composition

$$\mathcal{C} \xrightarrow{Q_C} \mathcal{C}(U^{-1}) \xrightarrow{\iota_{\text{Ind}}} \text{Ind}(\mathcal{C}(U^{-1}))$$

of Q_C and the canonical embedding (Definition B.5.1) ι_{Ind} .

2. The *left derived functor* LF of F relative to (S_i, T_j) and U is the functor

$$LF : \prod_{i=1}^n \mathcal{A}_i(S_i^{-1}) \times \prod_{j=1}^m \mathcal{B}_j(T_j^{-1}) \rightarrow \text{Pro}(\mathcal{C}(U^{-1}))$$

which makes the following diagram commutative:

$$\begin{array}{ccc} \prod_{i=1}^n \mathcal{A}_i \times \prod_{j=1}^m \mathcal{B}_j & \xrightarrow{F} & \mathcal{C} \\ \downarrow \prod Q_{A_i} \times \prod Q_{B_j} & & \downarrow \bar{Q}_C \\ \prod_{i=1}^n \mathcal{A}_i(S_i^{-1}) \times \prod_{j=1}^m \mathcal{B}_j(T_j^{-1}) & \xrightarrow{LF} & \text{Pro}(\mathcal{C}(U^{-1})) \end{array}$$

where $\bar{Q}_C$ is the composition

$$\mathcal{C} \xrightarrow{Q_C} \mathcal{C}(U^{-1}) \xrightarrow{\iota_{\text{Pro}}} \text{Pro}(\mathcal{C}(U^{-1}))$$

of Q_C and the canonical embedding (Definition B.5.1) ι_{Pro} .

Definition B.5.5 (Derivability and Adapted Families). Let $(\mathcal{A}_i)_{1 \leq i \leq n}$ and $(\mathcal{B}_j)_{1 \leq j \leq m}$ be families of triangulated categories (Definition B.1.2), and let $\mathcal{C}$ be a triangulated category. Let

$$F : \prod_{i=1}^n \mathcal{A}_i \times \prod_{j=1}^m \mathcal{B}_j \rightarrow \mathcal{C}$$

be an exact multifunctor (Definition B.1.5), covariant in $\mathcal{A}_i$ and contravariant in $\mathcal{B}_j$. Let S_i, T_j, U be saturated multiplicative systems (Definition B.2.3) in $\mathcal{A}_i, \mathcal{B}_j, \mathcal{C}$ respectively

assumed compatible with the triangulated structure (Definition B.1.4) of its respective category. Write

$$\begin{aligned} Q_{A_i} : \mathcal{A}_i &\rightarrow \mathcal{A}_i(S_i^{-1}) \\ Q_{B_j} : \mathcal{B}_j &\rightarrow \mathcal{B}_j(T_j^{-1}) \\ Q_C : \mathcal{C} &\rightarrow \mathcal{C}(U^{-1}) \end{aligned}$$

for the localization functors (Definition B.2.2). Write ι for the appropriate canonical embedding functors of the localized categories into their Ind or Pro categories (Definition B.5.1).

1. Let RF be the right derived functor (Definition B.5.4) of F relative to the multiplicative systems.
 - (a) RF is *defined at a family* (A_i, B_j) if the Ind-object (Definition B.5.1) $RF(A_i, B_j)$ is essentially constant (Definition B.5.2), i.e. isomorphic to an object of $\mathcal{C}(U^{-1})$. This object of $\mathcal{C}(U^{-1})$ is the *value of RF at (A_i, B_j)* .
 - (b) F is *right derivable* if RF is defined at all objects. In this case, we view RF as a functor into $\mathcal{C}(U^{-1})$.
 - (c) A family (A_i, B_j) is *adapted for RF* (or *deployed for RF* (*déployée pour RF* in French) or *unfolded for RF* , cf. [?]) if the natural morphism $(\iota \circ Q_C \circ F)(A_i, B_j) \rightarrow RF(Q_{A_i} A_i, Q_{B_j} B_j)$ is an isomorphism in $\text{Ind}(\mathcal{C}(U^{-1}))$, i.e. the Ind-object $RF(A_i)$ of $\mathcal{A}(S^{-1})$ is essentially constant (Definition B.5.2) with value $F((A_i), (B_j))$.
2. Let LF be the left derived functor (Definition B.5.4) of F relative to the multiplicative systems.
 - (a) LF is *defined at a family* (A_i, B_j) if the Pro-object (Definition B.5.1) $LF(A_i, B_j)$ is essentially constant (Definition B.5.2), i.e. isomorphic to an object of $\mathcal{C}(U^{-1})$. This object of $\mathcal{C}(U^{-1})$ is the *value of LF at (A_i, B_j)* .
 - (b) F is *left derivable* if LF is defined at all objects. In this case, we view LF as a functor into $\mathcal{C}(U^{-1})$.
 - (c) A family (A_i, B_j) is *adapted for LF* (or *deployed for LF* (*déployée pour LF* in French) *unfolded for LF* , cf. [?]) if the natural morphism $LF(Q_{A_i} A_i, Q_{B_j} B_j) \rightarrow (\iota \circ Q_C \circ F)(A_i, B_j)$ is an isomorphism in $\text{Pro}(\mathcal{C}(U^{-1}))$, i.e. the Pro-object $LF(A_i)$ of $\mathcal{A}(S^{-1})$ is essentially constant (Definition B.5.2) with value $F((A_i), (B_j))$.

In particular, note that if a family (A_i, B_j) is adapted for RF resp. LF , then RF resp. LF is defined at (A_i, B_j) .

Proposition B.5.6 (cf. [?]). Let $(\mathcal{A}_i)_{1 \leq i \leq n}$ and $(\mathcal{B}_j)_{1 \leq j \leq m}$ be families of triangulated categories (Definition B.1.2), and let $\mathcal{C}$ be a triangulated category. Let

$$F : \prod_{i=1}^n \mathcal{A}_i \times \prod_{j=1}^m \mathcal{B}_j \rightarrow \mathcal{C}$$

be an exact multifunctor (Definition B.1.5), covariant in $\mathcal{A}_i$ and contravariant in $\mathcal{B}_j$. Let S_i, T_j, U be saturated multiplicative systems (Definition B.2.3) in $\mathcal{A}_i, \mathcal{B}_j, \mathcal{C}$ respectively assumed compatible with the triangulated structure (Definition B.1.4) of its respective category.

A family (A_i, B_j) is adapted (or deployed/unfolded) (Definition B.5.5) for RF (Definition B.5.4) if and only if the following holds: For every family of morphisms $s_i : A_i \rightarrow X_i$ in S_i and $t_j : Y_j \rightarrow B_j$ in T_j , there exist morphisms $\sigma_i : X_i \rightarrow \mathcal{X}_i$ in S_i and $\tau_j : \mathcal{Y}_j \rightarrow Y_j$ in T_j such that the induced morphism in $\mathcal{C}$

$$F(\sigma_i \circ s_i, t_j \circ \tau_j) : F(A_i, B_j) \rightarrow F(\mathcal{X}_i, \mathcal{Y}_j)$$

is an isomorphism in the localized category $\mathcal{C}(U^{-1})$ (i.e., its image under Q_C is an isomorphism, or equivalently, it is an element of U).

See also Proposition B.7.1

Proof. (♠ TODO: verify this proof) Let $\mathcal{D} = \prod_i \mathcal{A}_i \times \prod_j \mathcal{B}_j^{op}$ and let S be the product multiplicative system $\prod_i S_i \times \prod_j T_j^{op}$. The right derived functor RF evaluated at an object $\mathbf{A} = (A_i, B_j)$ is defined as the Ind-object in $\mathcal{C}(U^{-1})$ represented by the diagram:

$$RF(\mathbf{A}) = \varinjlim_{\mathbf{A} \rightarrow \mathbf{X} \in S} Q_C(F(\mathbf{X}))$$

By definition, $\mathbf{A}$ is *adapted* (deployed) for RF if the natural morphism from the constant Ind-object to the system is an isomorphism:

$$\eta : \iota(Q_C(F(\mathbf{A}))) \xrightarrow{\cong} RF(Q_{\mathcal{D}}(\mathbf{A})) \quad \text{in } \text{Ind}(\mathcal{C}(U^{-1}))$$

($\Rightarrow$) Suppose $\mathbf{A}$ is adapted. An isomorphism in an Ind-category $\text{Ind}(\mathcal{M})$ between a constant object M and a filtered system $\{N_k\}_{k \in K}$ implies that for every index $k \in K$ (representing a resolution $\mathbf{A} \rightarrow \mathbf{X}$), there exists a morphism $k \rightarrow \ell$ in the filtering category (representing a further resolution $\mathbf{X} \rightarrow \mathcal{X}$) such that the composite map $M \rightarrow N_\ell$ is an isomorphism in $\mathcal{M}$.

Translating this to our categories: for any family of morphisms $s_i : A_i \rightarrow X_i$ and $t_j : Y_j \rightarrow B_j$ (defining the resolution $\mathbf{A} \rightarrow \mathbf{X}$), there must exist a further resolution $\mathbf{X} \rightarrow \mathcal{X}$ (defined by $\sigma_i : X_i \rightarrow \mathcal{X}_i$ and $\tau_j : \mathcal{Y}_j \rightarrow Y_j$) such that the induced morphism $Q_C(F(\mathbf{A})) \rightarrow Q_C(F(\mathcal{X}))$ is an isomorphism in $\mathcal{C}(U^{-1})$. This is exactly the condition stated.

($\Leftarrow$) Conversely, assume the condition holds. Let J be the category of resolutions of $\mathbf{A}$. The condition implies that the constant object $Q_C(F(\mathbf{A}))$ is *cofinal* in the system defining $RF(\mathbf{A})$.

Specifically, the assumption guarantees that for every $\mathbf{X} \in J$, there is a $\mathcal{X} \in J$ further in the diagram such that $Q_C(F(\mathbf{A}))$ and $Q_C(F(\mathcal{X}))$ are isomorphic via the transition maps of the system. In the category of Ind-objects, if a filtered system contains a cofinal subsystem of objects isomorphic to a constant object M , then the Ind-limit of that system is isomorphic to M . Thus, $RF(\mathbf{A}) \cong \iota(Q_C(F(\mathbf{A})))$, which means $\mathbf{A}$ is adapted. $\square$

Proposition B.5.7 (cf. [?]). Let $(\mathcal{A}_i)_{1 \leq i \leq n}$ and $(\mathcal{B}_j)_{1 \leq j \leq m}$ be families of abelian categories, and let $\mathcal{C}$ be an abelian category. Let

$$F : \prod_{i=1}^n \mathcal{A}_i \times \prod_{j=1}^m \mathcal{B}_j \rightarrow \mathcal{C}$$

be an additive multifunctor, covariant in $K(\mathcal{A}_i)$ and contravariant in $K(\mathcal{B}_j)$.

A family of complexes $(A_i^\bullet, B_j^\bullet)$ is adapted (or deployed/unfolded) (Definition B.5.5) for the right derived functor (Definition B.9.14) RF if and only if the following holds: For every family of quasi-isomorphisms (Definition .2.4) $s_i : A_i^\bullet \rightarrow X_i^\bullet$ and $t_j : Y_j^\bullet \rightarrow B_j^\bullet$, there exist quasi-isomorphisms $\sigma_i : X_i^\bullet \rightarrow \mathcal{X}_i^\bullet$ and $\tau_j : \mathcal{Y}_j^\bullet \rightarrow Y_j^\bullet$ such that the induced morphism in $K(\mathcal{C})$

$$F(\sigma_i \circ s_i, t_j \circ \tau_j) : F(A_i^\bullet, B_j^\bullet) \rightarrow F(\mathcal{X}_i^\bullet, \mathcal{Y}_j^\bullet)$$

is a quasi-isomorphism (i.e., an isomorphism in the derived category $D(\mathcal{C})$).

Proof. This follows from Proposition B.5.6. □

B.5.1. *Derived multifunctors of additive multifunctors between abelian categories.*

Definition B.5.8. Let $F : \prod_{i=1}^n \mathcal{A}_i \rightarrow \mathcal{B}$ be an additive multifunctor (Definition C.0.13) of abelian categories (Definition .1.34).

The *right (resp. left) derived multifunctors of F* are the derived functors (Definition B.5.4) RF (resp. LF) of the induced multifunctors (Theorem B.1.7)

- $K(\prod \mathcal{A}_i) \rightarrow K(\mathcal{B})$
- $K^+(\prod \mathcal{A}_i) \rightarrow K^+(\mathcal{B})$
- $K^-(\prod \mathcal{A}_i) \rightarrow K^-(\mathcal{B})$

(Recall Proposition B.1.6) relative to the systems (Definition B.2.3) of quasi-isomorphisms (Definition .2.4) in the respective homotopy categories (Definition B.1.1). In particular, the functor RF a priori may refer to one of the functors

$$D^+(\prod \mathcal{A}_i) \rightarrow \text{Ind } D^+(\mathcal{B})$$

$$D^-(\prod \mathcal{A}_i) \rightarrow \text{Ind } D^-(\mathcal{B})$$

$$D(\prod \mathcal{A}_i) \rightarrow \text{Ind } D(\mathcal{B})$$

and the functor LF refers to one of the functors

$$D^+(\prod \mathcal{A}_i) \rightarrow \text{Pro } D^+(\mathcal{B})$$

$$D^-(\prod \mathcal{A}_i) \rightarrow \text{Pro } D^-(\mathcal{B})$$

$$D(\prod \mathcal{A}_i) \rightarrow \text{Pro } D(\mathcal{B})$$

(Definition B.3.1) (Definition B.5.1).

Definition B.5.9 ([?, Definition 1.2.3]). Let $F : \prod_{i=1}^n \mathcal{A}_i \rightarrow \mathcal{B}$ be an additive multifunctor (Definition C.0.13) of abelian categories (Definition .1.34).

1. An object $(A_1, \dots, A_n)$ with $A_i \in \text{ob}(\mathcal{A}_i)$ is *right acyclic for F* or *acyclic for RF* if the complex concentrated in degree 0 associated to (A_i) is adapted for (Definition B.5.5) RF .

2. Similarly, (A_i) is *left acyclic for F* or *acyclic for LF* if the complex concentrated in degree 0 associated to (A_i) is adapted for (Definition B.5.5) LF .

In the case that $n = 1$ and $A_1 \in \mathcal{A}_1$ has a projective resolution (Definition .2.7) (resp. injective resolution), the notion of left acyclic for F (resp. right acyclic for F) is equivalent to the notion of F -acyclicity. (♠ TODO: justify)

Proposition B.5.10 (cf. [?, Exposé XVII, Proposition 1.2.4.]). Let $F : \prod_{i=1}^n \mathcal{A}_i \rightarrow \mathcal{B}$ be an additive multifunctor (Definition C.0.13) of abelian categories (Definition .1.34).

1. The following diagram of categories is commutative for the right derived multifunctor:

$$\begin{array}{ccc} D^+(\prod \mathcal{A}_i) & \longrightarrow & D(\prod \mathcal{A}_i) \\ \downarrow RF & & \downarrow RF \\ \text{Ind } D^+(\mathcal{B}) & \longrightarrow & \text{Ind } D(\mathcal{B}) \end{array}$$

(Definition B.3.1) (??) (Definition B.5.8) where the horizontal arrows are the natural functors induced by inclusions. If RF is everywhere defined on $D(\prod \mathcal{A}_i)$, its restriction to $D^+(\prod \mathcal{A}_i)$ factors through $D^+(\mathcal{B})$ [?, Proposition 1.2.4].

2. Dually, the following diagram is commutative for the left derived multifunctor:

$$\begin{array}{ccc} D^-(\prod \mathcal{A}_i) & \longrightarrow & D(\prod \mathcal{A}_i) \\ \downarrow LF & & \downarrow LF \\ \text{Pro } D^-(\mathcal{B}) & \longrightarrow & \text{Pro } D(\mathcal{B}) \end{array}$$

If LF is everywhere defined on $D(\prod \mathcal{A}_i)$, its restriction to $D^-(\prod \mathcal{A}_i)$ factors through $D^-(\mathcal{B})$.

Theorem B.5.11 ([?, Exposé XVII, Proposition 1.2.7.]). (♠ TODO:) Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be an additive functor between abelian categories (Definition .1.34).

If $\mathcal{A}$ has enough objects that are acyclic for (Definition B.5.9) LF (resp. for RF), the derived functor (Definition B.9.14) $LF : D^-(\mathcal{A}) \rightarrow D^-(\mathcal{B})$ (resp. $RF : D^+(\mathcal{A}) \rightarrow D^+(\mathcal{B})$) is left (resp. right) derivable (Definition B.5.5) and every homologically upper bounded (Definition .2.2) (resp. homologically lower bounded) complex of acyclics for LF (resp. for RF) is adapted for LF (resp. for RF) (Definition B.5.5).

(♠ TODO:

- **Address the Biadditive/Multifunctor Gap**

- Add a Lemma or Remark following Definition 5.5.5 clarifying that for a biadditive functor $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$, the condition of a pair of complexes $(M^\bullet, N^\bullet)$ being adapted (deployed) is what allows the representation of LF via the Total Complex construction.
- Explicitly define the Total Complex functor $\text{Tot} : K(\mathcal{A}) \times K(\mathcal{B}) \rightarrow K(\mathcal{C})$ used to induce the multifunctor in Definition 5.5.6. Specify the sign convention for the total differential: $d(x \otimes y) = d_x(x) \otimes y + (-1)^p x \otimes d_y(y)$ for x of degree p .

- **Formulate the Balancing Results**

- Add a Proposition (The Balancing Lemma) stating that if $\mathcal{P} \subset \mathcal{A}$ is a class of objects adapted for $F(-, N)$ and $\mathcal{Q} \subset \mathcal{B}$ is a class adapted for $F(M, -)$, there exist canonical isomorphisms in $D(\mathcal{C})$:

$$\mathrm{Tot}(F(P^\bullet, N^\bullet)) \cong \mathrm{Tot}(F(M^\bullet, Q^\bullet)) \cong \mathrm{Tot}(F(P^\bullet, Q^\bullet))$$

- Cite [SGA 4, XVII, 1.2.7] to justify that these isomorphisms are well-defined because complexes of adapted objects are "deployed".
- **Incorporate the “Everywhere Defined” Logic**
 - Include the “Finite Dimension” case from [SGA 4, XVII, Prop 1.2.10] to explain the conditions under which LF and RF can be extended to the unbounded derived category $D(\mathcal{A})$.
- **Technical Refinements**
 - Provide a precise proof or reference for the property: “If F is bi-right-exact, then projective objects are left-acyclic (adapted) for F .”
 - Define the term “Deployed in the first variable”: A complex $P^\bullet$ is deployed in the first variable if the family $(P^\bullet, N^\bullet)$ is deployed for the multifunctor F for every $N^\bullet \in K(\mathcal{A}_2)$.
 - Resolve the ♠ TODO regarding the existence of limits/colimits in **Sets** to ensure Ind/Pro categories are locally small.
- **Categorical Equivalence**
 - Add a Remark confirming the equivalence of categories $D(\mathcal{A}_1) \times D(\mathcal{A}_2) \simeq D(\mathcal{A}_1 \times \mathcal{A}_2)$, noting that quasi-isomorphisms are defined component-wise.
- **derived tensor product**
 - Do proposition 4.1.7. expose XVII

)

B.6. K -flat resolutions of chain complexes with respect to bi-additive functors.

Definition B.6.1. Let $\mathcal{A}$ be an abelian category (Definition .1.34).

1. ([?, After Theorem B]) Let $f : A^\bullet \rightarrow B^\bullet$ be a quasi-isomorphism (Definition .2.4) of chain complexes (Definition .2.1) of objects in $\mathcal{A}$. We say that $A^\bullet$ is a *left resolution of $B^\bullet$* and that $B^\bullet$ is a *right resolution of $A^\bullet$* .
2. Given a chain complex (Definition .2.1) $K^\bullet$ of objects in $\mathcal{A}$ and a class $\mathcal{Y}$ of chain complexes of objects in $\mathcal{A}$, an *$\mathcal{Y}$ -left resolution* may refer to a left resolution $Y^\bullet \rightarrow K^\bullet$ where $Y^\bullet \in \mathcal{Y}$ and an *$\mathcal{Y}$ -right resolution* may refer to a right resolution $K^\bullet \rightarrow Y^\bullet$ where $Y^\bullet \in \mathcal{Y}$.

Definition B.6.2. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be abelian categories (Definition .1.34), and let $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ be a bi-additive functor (Definition C.0.13).

Let us say that a complex (Definition .2.1) $K^\bullet$ of $\mathcal{A}$ (resp. of $\mathcal{B}$) is *K -flat with respect to F on the left (resp. right) for direct sum total complexes* if for every acyclic complex $M^\bullet$ of $\mathcal{B}$ (resp. of $\mathcal{A}$), the direct sum total complex $\mathrm{Tot}^\oplus(F(K^\bullet, M^\bullet))$ (resp. $\mathrm{Tot}^\oplus(F(M^\bullet, K^\bullet))$) is acyclic.

Similarly, let us say that $K^\bullet$ is K -flat with respect to F on the left (resp. right) for product total complexes if for every acyclic complex $M^\bullet$ of $\mathcal{B}$ (resp. of $\mathcal{A}$), the product total complex $\text{Tot}\Pi(F(K^\bullet, M^\bullet))$ (resp. $\text{Tot}\Pi(F(M^\bullet, K^\bullet))$) is acyclic.

Definition B.6.3. Let $\mathcal{A}$ be an additive category.

Given (cochain) complexes (Definition .2.1) $K^\bullet, L^\bullet$ of objects in $\mathcal{A}$, the *internal Hom complex* $\text{Hom}^\bullet(K^\bullet, L^\bullet)$, also called the *(product) total Hom complex* or the *complex of morphisms from $K^\bullet$ to $L^\bullet$* and also denoted by notations such as $\underline{\text{Hom}}(X, Y)$ depending on the context, is the product total complex $\text{Tot}\Pi(\text{Hom}(K^\bullet, L^\bullet))$ where $\text{Hom}(K^\bullet, L^\bullet)$ is the double complex associated to (??) the biadditive functor $\text{Hom} : \mathcal{A}^{\text{op}} \times \mathcal{A} \rightarrow \text{Ab}$. More explicitly, $\text{Hom}^\bullet(K^\bullet, L^\bullet)$ is the complex of objects in Ab given by

- Objects: In each degree $n \in \mathbb{Z}$, the object is the product

$$\text{Hom}^n(K^\bullet, L^\bullet) = \prod_{p \in \mathbb{Z}} \text{Hom}_{\mathcal{A}}(K^p, L^{p+n}).$$

An element $f \in \text{Hom}^n(K^\bullet, L^\bullet)$ is thus a family of morphisms $(f_p)_{p \in \mathbb{Z}}$ where $f_p : K^p \rightarrow L^{p+n}$.

- Differential: The differential $d^n : \text{Hom}^n(K^\bullet, L^\bullet) \rightarrow \text{Hom}^{n+1}(K^\bullet, L^\bullet)$ is defined for each component $f = (f_p)$ by

$$(d^n f)_p = d_L^{p+n} \circ f_p - (-1)^n f_{p+1} \circ d_K^p,$$

where d_K and d_L are the differentials of the complexes $K^\bullet$ and $L^\bullet$ respectively.

Under this construction, the 0-cocycles of $\text{Hom}^\bullet(K^\bullet, L^\bullet)$ correspond precisely to the chain maps from $K^\bullet$ to $L^\bullet$, and the 0-coboundaries correspond to those chain maps which are null-homotopic.

If $K_\bullet$ and $L_\bullet$ are chain complexes written with homological conventions instead, then the internal Hom complex should be denoted by $\underline{\text{Hom}}_\bullet(K_\bullet, L_\bullet)$ instead; explicitly, $\text{Hom}_\bullet(K_\bullet, L_\bullet)$ is the complex of objects in Ab given by

- Objects: In each degree $n \in \mathbb{Z}$, the object is the product

$$\text{Hom}_n(K_\bullet, L_\bullet) = \prod_{p \in \mathbb{Z}} \text{Hom}_{\mathcal{A}}(K_p, L_{p+n}).$$

An element $f \in \text{Hom}_n(K_\bullet, L_\bullet)$ is a family of morphisms $(f_p)_{p \in \mathbb{Z}}$ such that $f_p : K_p \rightarrow L_{p+n}$.

- Differential: The differential $d_n : \text{Hom}_n(K_\bullet, L_\bullet) \rightarrow \text{Hom}_{n-1}(K_\bullet, L_\bullet)$ is defined for a family $f = (f_p)$ by

$$(d_n f)_p = d_L^{p+n} \circ f_p - (-1)^n f_{p-1} \circ d_K^p,$$

where $d_K^p : K_p \rightarrow K_{p-1}$ and $d_L^q : L_q \rightarrow L_{q-1}$ are the homological differentials.

As in the cohomological case, the 0-cycles of $\text{Hom}_\bullet(K_\bullet, L_\bullet)$ are precisely the chain maps between the complexes, and the 0-boundaries are the null-homotopic chain maps.

Definition B.6.4. Let $\mathcal{A}$ be an abelian category. A chain complex $A^\bullet$ of objects in $\mathcal{A}$ is called *K-projective* (resp. *K-injective*) if for every acyclic complex $X^\bullet$ of objects in $\mathcal{A}$, the complex of abelian groups $\text{Hom}^\bullet(A^\bullet, X^\bullet)$ (Definition B.6.3) (resp. $\text{Hom}^\bullet(X^\bullet, A^\bullet)$) is acyclic.

Equivalently, $A^\bullet$ is *K-injective* if and only if it is *K-flat* with respect to (Definition B.6.2) Hom on the left for product total complexes and $A^\bullet$ is *K-projective* if and only if it is *K-flat* with respect to Hom on the right for product total complexes.

Proposition B.6.5 ([?, 1.2. Proposition]). Let $\mathcal{A}$ be an abelian category (Definition .1.34) and let $A^\bullet \in \mathbf{Ch}(\mathcal{A})$ be a complex (Definition .2.1) such that $A^i = 0$ for all $i \neq 0$. The complex (Definition .2.1) $A^\bullet$ is *K-projective* (resp. *K-injective*) if and only if A^0 is a projective (resp. injective) object of $\mathcal{A}$.

Proposition B.6.6 ([?, 1.3. Proposition]). Let $\mathcal{A}$ be an abelian category (Definition .1.34). If two of the vertices of a distinguished triangle (Definition B.1.2) of $K(\mathcal{A})$ (Definition B.1.1) are *K-projective* (Definition B.6.4) (resp. *K-injective* (Definition B.6.4)), then so is the third.

Proposition B.6.7 ([?, 1.3. Proposition]). Let $\mathcal{A}$ be an abelian category (Definition .1.34). A complex $A^\bullet$ of objects in $\mathcal{A}$ is *K-projective* (resp. *K-injective*) if and only if $A^\bullet[n]$ is as well, where n is any integer.

Proposition B.6.8 ([?, 1.4. Proposition, 1.5. Proposition]). Let $\mathcal{A}$ be an abelian category (Definition .1.34), and let $K(\mathcal{A})$ (Definition B.1.1) and $D(\mathcal{A})$ (Definition B.3.1) be its homotopy and derived categories, respectively.

1. For every complex (Definition .2.1) $A^\bullet$ in $\mathcal{A}$, the following conditions are equivalent:
 - (a) $A^\bullet$ is *K-projective* (Definition B.6.4).
 - (b) For every $S^\bullet \in K(\mathcal{A})$, the natural homomorphism

$$\text{Mor}_{K(\mathcal{A})}(A^\bullet, S^\bullet) \rightarrow \text{Mor}_{D(\mathcal{A})}(A^\bullet, S^\bullet)$$

is an isomorphism.

- (c) For every diagram in $K(\mathcal{A})$

$$\begin{array}{ccc} & & A^\bullet \\ & \swarrow g & \downarrow f \\ X^\bullet & \xrightarrow{s} & S^\bullet \end{array}$$

where s is a quasi-isomorphism (Definition .2.4), there exists a unique morphism $g : A^\bullet \rightarrow X^\bullet$ such that $s \circ g = f$ in $K(\mathcal{A})$.

- (d) For every quasi-isomorphism $u : S^\bullet \rightarrow A^\bullet$ in $K(\mathcal{A})$, there exists a morphism $v : A^\bullet \rightarrow S^\bullet$ such that $u \circ v = 1_{A^\bullet}$ in $K(\mathcal{A})$.
2. Similarly, for every $A^\bullet$ in $\mathcal{A}$, the following conditions are equivalent:
 - (a) $A^\bullet$ is *K-injective* (Definition B.6.4).
 - (b) For every $S^\bullet \in K(\mathcal{A})$, the natural homomorphism

$$\text{Mor}_{K(\mathcal{A})}(S^\bullet, A^\bullet) \rightarrow \text{Mor}_{D(\mathcal{A})}(S^\bullet, A^\bullet)$$

is an isomorphism.

(c) For every diagram in $K(\mathcal{A})$

$$\begin{array}{ccc} S^\bullet & \xrightarrow{s} & X^\bullet \\ f \downarrow & \swarrow g & \\ A^\bullet & & \end{array}$$

where s is a quasi-isomorphism, there exists a unique morphism $g : X^\bullet \rightarrow A^\bullet$ such that $g \circ s = f$ in $K(\mathcal{A})$.

(d) For every quasi-isomorphism $u : A^\bullet \rightarrow S^\bullet$ in $K(\mathcal{A})$, there exists a morphism $v : S^\bullet \rightarrow A^\bullet$ such that $v \circ u = 1_{A^\bullet}$ in $K(\mathcal{A})$.

Proposition B.6.9. Let $\mathcal{A}$ be an abelian category (Definition .1.34).

1. If $P_\bullet$ is a bounded above (Definition .2.2) homological chain complex (or equivalently a bounded below cohomological chain complex) of projective objects, then $P_\bullet$ is K -projective (Definition B.6.4).

$$\cdots \rightarrow P_{i+1} \rightarrow P_i \rightarrow 0 \rightarrow 0 \rightarrow \cdots$$

2. If $I^\bullet$ is a bounded below (Definition .2.2) cohomological chain complex (or equivalently a bounded above homological chain complex) of injective objects, then $I^\bullet$ is K -injective (Definition B.6.4).

$$\cdots \rightarrow 0 \rightarrow 0 \rightarrow I^i \rightarrow I^{i+1} \rightarrow \cdots$$

Proof. (♠ TODO: verify the following) We prove (1). By the characterization of K -projectivity, it suffices to show that for any acyclic complex $N^\bullet$, every chain map $f : P^\bullet \rightarrow N^\bullet$ is homotopic to zero ($f \sim 0$).

Assume without loss of generality that $P^n = 0$ for $n > 0$. We construct a homotopy $h^n : P^n \rightarrow N^{n-1}$ such that $f^n = d_N^{n-1}h^n + h^{n+1}d_P^n$ by descending induction on n .

Base Case: For $n = 0$, since $P^1 = 0$, the chain map condition $d_N^0 f^0 = f^1 d_P^0$ implies $d_N^0 f^0 = 0$. Thus $\text{im}(f^0) \subseteq \ker(d_N^0)$. By the acyclicity of $N^\bullet$, $\ker(d_N^0) = \text{im}(d_N^{-1})$. Because P^0 is projective, f^0 factors through the surjection $N^{-1} \rightarrow \text{im}(d_N^{-1})$, providing $h^0 : P^0 \rightarrow N^{-1}$ such that $d_N^{-1}h^0 = f^0$. Setting $h^1 = 0$, we have $f^0 = d_N^{-1}h^0 + h^1 d_P^0$.

Inductive Step: Suppose we have constructed $\{h^k\}_{k \geq n}$ satisfying the homotopy relation. Consider the map $g = f^n - h^{n+1}d_P^n : P^n \rightarrow N^n$. We check its image under d_N^n :

$$\begin{aligned} d_N^n g &= d_N^n f^n - d_N^n h^{n+1} d_P^n \\ &= f^{n+1} d_P^n - (f^{n+1} - h^{n+2} d_P^{n+1}) d_P^n \\ &= f^{n+1} d_P^n - f^{n+1} d_P^n + h^{n+2} d_P^{n+1} d_P^n = 0. \end{aligned}$$

Thus $\text{im}(g) \subseteq \ker(d_N^n)$. By acyclicity, $\ker(d_N^n) = \text{im}(d_N^{n-1})$. Since P^n is projective, g factors through the surjection d_N^{n-1} , yielding $h^n : P^n \rightarrow N^{n-1}$ such that $d_N^{n-1}h^n = g = f^n - h^{n+1}d_P^n$.

This construction provides a well-defined sequence of maps $(h^n)_{n \in \mathbb{Z}}$ defining a homotopy $h : f \sim 0$. Thus $\text{Mor}_{K(\mathcal{A})}(P^\bullet, N^\bullet) = 0$, proving $P^\bullet$ is K -projective. The proof for (2) is dual, using ascending induction starting from the lowest non-zero degree of $I^\bullet$. $\square$

Theorem B.6.10 ([?, 3.4. Theorem, 3.8. Proposition]). 1. Let $\mathcal{A}$ be an AB5 abelian category (Definition .1.34). Suppose that $\mathfrak{P}$ is a class of objects in $\mathbf{Ch}(\mathcal{A})$ satisfying the following:

Every (cohomologically) bounded above complex (Definition .2.2) $A^\bullet \in \mathbf{Ch}(\mathcal{A})$ has a left $\mathfrak{P}$ -resolution (Definition B.6.1).

Then every complex in $\mathbf{Ch}(\mathcal{A})$ has a left $\mathfrak{P}$ -resolution.

2. Dually, let $\mathcal{A}$ be an AB5* abelian category (Definition .1.34). Suppose that $\mathfrak{I}$ is a class of objects in $\mathbf{Ch}(\mathcal{A})$ satisfying the following:

Every (cohomologically) bounded below complex (Definition .2.2) $A^\bullet \in \mathbf{Ch}(\mathcal{A})$ has a left $\mathfrak{I}$ -resolution (Definition B.6.1).

Then every complex in $\mathbf{Ch}(\mathcal{A})$ has a right $\mathfrak{I}$ -resolution.

Corollary B.6.11 ([?, 3.5. Corollary, 3.9. Corollary]). 1. Let $\mathcal{A}$ be an AB5 abelian category (Definition .1.34). If $\mathcal{A}$ has enough projectives (Definition .1.37), then every complex in $\mathcal{C}$ has a left K -projective (Definition B.6.4) resolution (Definition B.6.1).

(♠ TODO: bounded complex of projectives is K -projective)

2. Dually, let $\mathcal{A}$ be an AB5* abelian category (Definition .1.34). If $\mathcal{A}$ has enough injectives (Definition .1.37), then every complex in $\mathcal{C}$ has a right K -injective (Definition B.6.4) resolution (Definition B.6.1).

Proof. 1. Since $\mathcal{A}$ has enough projectives, any bounded above homological complex in $\mathcal{A}$ has a left-resolution ?? by a bounded above homological complex of projective objects, which is K -projective by Proposition B.6.9. Therefore, any complex in $\mathcal{A}$ has a left-resolution by a K -projective object by Theorem B.6.10

2. This is dual to the previous part.

□

Theorem B.6.12 ([?, Theorem 3.13.]). Let $\mathcal{A}$ be a Grothendieck category.

For every chain complex (Definition .2.1) $A^\bullet$ in $\mathcal{A}$, there is a right resolution (Definition B.6.1) $A^\bullet \rightarrow I^\bullet$ such that $I^\bullet$ is K -injective (Definition B.6.4).

Lemma B.6.13. Let $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ be a bi-additive functor (Definition C.0.13) where $\mathcal{A}, \mathcal{B}, \mathcal{C}$ are additive categories. Let J be a category (Definition .1.1).

1. Assume that $F(X, -) : \mathcal{B} \rightarrow \mathcal{C}$ preserves colimits (Definition .1.13) of diagrams of shape J (Definition .1.33) for all objects $X \in \mathcal{A}$. Then the induced functor $F(X^\bullet, -) : \mathbf{Ch}(\mathcal{B}) \rightarrow \mathbf{DC}(\mathcal{C})$ (??) (Definition .2.1) also preserves colimits of diagrams of shape J for all objects $X^\bullet \in \mathbf{Ch}(\mathcal{A})$.
2. Dually, assume that $F(-, Y) : \mathcal{A} \rightarrow \mathcal{C}$ preserves colimits (Definition .1.13) of diagrams of shape J (Definition .1.33) for all objects $Y \in \mathcal{B}$. Then the induced functor $F(-, Y^\bullet) : \mathbf{Ch}(\mathcal{A}) \rightarrow \mathbf{DC}(\mathcal{C})$ (??) (Definition .2.1) also preserves colimits of diagrams of shape J for all objects $Y^\bullet \in \mathbf{Ch}(\mathcal{B})$.
3. Assume that $F(X, -) : \mathcal{B} \rightarrow \mathcal{C}$ preserves limits (Definition .1.13) of diagrams of shape J (Definition .1.33) for all objects $X \in \mathcal{A}$. Then the induced functor $F(X^\bullet, -) : \mathbf{Ch}(\mathcal{B}) \rightarrow \mathbf{DC}(\mathcal{C})$ (??) (Definition .2.1) also preserves limits of diagrams of shape J for all objects $X^\bullet \in \mathbf{Ch}(\mathcal{A})$.

4. Dually, assume that $F(-, Y) : \mathcal{A} \rightarrow \mathcal{C}$ preserves limits (Definition .1.13) of diagrams of shape J (Definition .1.33) for all objects $Y \in \mathcal{B}$. Then the induced functor $F(-, Y^\bullet) : \mathbf{Ch}(\mathcal{A}) \rightarrow \mathbf{DC}(\mathcal{C})$ (??) (Definition .2.1) also preserves limits of diagrams of shape J for all objects $Y^\bullet \in \mathbf{Ch}(\mathcal{B})$.

Proof. We prove that if $F(X, -) : \mathcal{B} \rightarrow \mathcal{C}$ preserves colimits of diagrams of shape J for all $X \in \mathcal{A}$, then so does $F(X^\bullet, -) : \mathbf{Ch}(\mathcal{B}) \rightarrow \mathbf{DC}(\mathcal{C})$ for all $X^\bullet \in \mathbf{Ch}(\mathcal{A})$. Given $X^\bullet \in \mathbf{Ch}(\mathcal{A})$ and $Y^\bullet \in \mathbf{Ch}(\mathcal{B})$, the term of $F(X^\bullet, Y^\bullet)$ at position (p, q) is $F(X^p, Y^q)$. Let $J \rightarrow \mathbf{Ch}(\mathcal{B}) : j \mapsto Y_{(j)}^\bullet$ be some diagram (Definition .1.33) whose colimit is $Y^\bullet$. In fact, since colimits of chain complexes are computed termwise (??), there are canonical isomorphisms

$$Y^q \cong \operatorname{colim}_{j \in J} Y_{(j)}^q$$

in $\mathcal{B}$.

We wish to show that $F(X^\bullet, Y^\bullet)$ is the colimit of the diagram $j \mapsto F(X^\bullet, Y_{(j)}^\bullet)$ in $\mathbf{DC}(\mathcal{C})$. The colimit in $\mathbf{DC}(\mathcal{C})$ is also computed termwise (note that $\mathbf{DC}(\mathcal{C})$ is identifiable as $\mathbf{Ch}(\mathbf{Ch}(\mathcal{C}))$), it suffices to verify the isomorphism at each position (p, q) and ensure it respects the differentials.

Consider the term at position (p, q) . The functor $F(X^\bullet, -)$ maps the colimit diagram in $\mathbf{Ch}(\mathcal{B})$ to a cocone (Definition .1.13) in $\mathbf{DC}(\mathcal{C})$. This induces a canonical comparison morphism:

$$\phi : \operatorname{colim}_{j \in J} F(X^\bullet, Y_{(j)}^\bullet) \longrightarrow F(X^\bullet, \operatorname{colim}_{j \in J} Y_{(j)}^\bullet) = F(X^\bullet, Y^\bullet).$$

At the position (p, q) , this morphism is the map:

$$\phi^{p,q} : \operatorname{colim}_{j \in J} F(X^p, Y_{(j)}^q) \longrightarrow F(X^p, Y^q).$$

By hypothesis, $F(X^p, -) : \mathcal{B} \rightarrow \mathcal{C}$ preserves colimits of shape J . Therefore, $\phi^{p,q}$ is an isomorphism for all $p, q \in \mathbb{Z}$.

Since a morphism of double complexes is an isomorphism if and only if it is an isomorphism at every term (p, q) , we conclude that ϕ is an isomorphism. Thus, $F(X^\bullet, -)$ preserves colimits of shape J .

□

Lemma B.6.14. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be abelian categories (Definition .1.34), and let $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ be a bi-additive functor (Definition C.0.13). Assume that $\mathcal{C}$ is Ab5.

1. Suppose that $F(X, -) : \mathcal{B} \rightarrow \mathcal{C}$ preserves filtered colimits (Definition .1.14) for all objects $X \in \mathcal{A}$. Let $P^\bullet$ be a (cohomologically) bounded above (Definition .2.2) complex of objects of $\mathcal{A}$ that are flat with respect to F on the left. The complex $P^\bullet$ is K -flat with respect to F on the left for direct sum total complexes (Definition B.6.2).
2. Suppose that $F(-, Y) : \mathcal{A} \rightarrow \mathcal{C}$ preserves filtered colimits (Definition .1.14) for all objects $Y \in \mathcal{B}$. Let $P^\bullet$ be a bounded above complex of objects in $\mathcal{B}$ that are flat with respect to F on the right. The complex $P^\bullet$ is K -flat with respect to F on the right for direct sum total complexes.

Proof. We show the claimed statement in the case that $P^\bullet$ is a complex of flat objects of $\mathcal{A}$ with respect to F on the left; the other case can be argued symmetrically. Let $M^\bullet$ be an acyclic complex of objects in $\mathcal{A}$. Writing $\tau_{\leq k} M^\bullet$ for the brutal truncation of $M^\bullet$, note that $M^\bullet$ is the filtered colimit of these truncations (??):

$$M^\bullet = \varinjlim_k \tau_{\leq k} M^\bullet.$$

Since $F(X, -) : \mathcal{B} \rightarrow \mathcal{C}$ preserves filtered colimits, so does $F(X^\bullet, -) : \mathbf{Ch}(\mathcal{B}) \rightarrow \mathbf{DC}(\mathcal{C})$ (Definition .2.1) by Lemma B.6.13. Therefore,

$$\mathrm{Tot}^\oplus F(P^\bullet, M^\bullet) \cong \mathrm{Tot}^\oplus F(P^\bullet, \varinjlim_k \tau_{\leq k} M^\bullet) \cong \varinjlim_k \mathrm{Tot}^\oplus F(P^\bullet, \tau_{\leq k} M^\bullet).$$

Since $\tau_{\leq k} M^\bullet$ is bounded above and is of flat objects, the double complex $F(P^\bullet, \tau_{\leq k} M^\bullet)$ is bounded above, and each row is acyclic. Since $\mathcal{C}$ satisfies Ab5, $\mathrm{Tot}^\oplus F(P^\bullet, \tau_{\leq k} M^\bullet)$ is acyclic by Lemma ???. That $\mathcal{C}$ satisfies AB5 further implies that

$$H^n(\mathrm{Tot}^\oplus F(P^\bullet, M^\bullet)) \cong \varinjlim_k H^n \mathrm{Tot}^\oplus F(P^\bullet, \tau_{\leq k} M^\bullet) = 0,$$

i.e. $\mathrm{Tot}^\oplus F(P^\bullet, M^\bullet)$ is acyclic. Therefore, $P^\bullet$ is K -flat as desired. $\square$

B.7. General conditions for defining derived functors.

Proposition B.7.1 (see [?, Tag 06XN, 05S8]). Suppose that $F : \mathcal{D} \rightarrow \mathcal{D}'$ is an exact functor (Definition B.1.3) of triangulated categories (Definition B.1.2) and that S is a saturated multiplicative system (Definition B.2.3) in $\mathcal{D}$ compatible with (Definition B.1.4) the structure of the triangulated category (Definition B.1.2) on $\mathcal{D}$, i.e. $s : X \rightarrow Y$ is in S if and only if $s[1] : X[1] \rightarrow Y[1]$ is in S and for every commutative diagram in $\mathcal{D}$ where the rows are distinguished triangles:

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \rightarrow & Z & \rightarrow & X[1] \\ \downarrow^{s_1} & & \downarrow^{s_2} & & \exists \downarrow^{s_3} & & \downarrow^{s_1[1]} \\ X' & \xrightarrow{f'} & Y' & \rightarrow & Z' & \rightarrow & X'[1] \end{array}$$

if $s_1, s_2 \in S$, there exists a morphism $s_3 \in S$ making the entire diagram commute.

If there exists a subcollection $\mathcal{I} \subset \mathrm{Ob}(\mathcal{D})$ such that

1. for all $X \in \mathrm{Ob}(\mathcal{D})$ there exists $s : X \rightarrow X'$ in S with $X' \in \mathcal{I}$, and
2. for every arrow $s : X \rightarrow X'$ in S with $X, X' \in \mathcal{I}$ the map $F(s) : F(X) \rightarrow F(X')$ is an isomorphism,

then RF (Definition B.5.4) is everywhere defined (Definition B.5.5) and every $X \in \mathcal{I}$ computes RF .

Dually, if there exists a subcollection $\mathcal{P} \subset \mathrm{Ob}(\mathcal{D})$ such that

1. for all $X \in \mathrm{Ob}(\mathcal{D})$ there exists $s : X' \rightarrow X$ in S with $X' \in \mathcal{P}$, and
2. for every arrow $s : X \rightarrow X'$ in S with $X, X' \in \mathcal{P}$ the map $F(s) : F(X) \rightarrow F(X')$ is an isomorphism,

then LF is everywhere defined and every $X \in \mathcal{P}$ computes LF . (♠ TODO: Proposition B.5.6 should be relevant)

Proposition B.7.2 (see [?, Tag 070K]). Let $\mathcal{A}$ be an abelian category (Definition .1.34). Assume that every complex (Definition .2.1) of objects in $\mathcal{A}$ has a right resolution (Definition B.6.1) by a K -injective (Definition B.6.4) complex. For any exact functor (Definition B.1.3) $F : K(\mathcal{A}) \rightarrow \mathcal{D}'$ (Definition B.1.1) of triangulated categories (Definition B.1.2), the right derived functor (Definition B.5.8)

$$RF : D(\mathcal{A}) \rightarrow \mathcal{D}'$$

is defined (Definition B.5.5) and $RF(I^\bullet) = F(I^\bullet)$ for K -injective complexes $I^\bullet$.

Proof. By Proposition B.7.1, it suffices to show that if $I^\bullet \rightarrow J^\bullet$ is a quasi-isomorphism of K -injective complexes, then $F(I^\bullet) \rightarrow F(J^\bullet)$ is an isomorphism in $\mathcal{D}'$ (♠ TODO: continue). $\square$

B.7.1. *Adjunction of derived functors of adjoint functors.*

Proposition B.7.3 ([?, Tag 0FND]). Let $F : \mathcal{D} \rightarrow \mathcal{D}'$ and $G : \mathcal{D}' \rightarrow \mathcal{D}$ be exact functors of triangulated categories (Definition B.1.3). Assume F is right adjoint (Definition .1.8) to G . Let S and S' respectively be multiplicative systems (Definition B.2.3) for $\mathcal{D}$ and $\mathcal{D}'$ compatible with the triangulated structures (Definition B.1.4). Let $K \in \text{Ob}(\mathcal{D})$ and $M \in \text{Ob}(\mathcal{D}')$. If RF (Definition B.5.4) is defined (Definition B.5.5) at K and LG is defined at M , then there is a canonical isomorphism

$$\text{Hom}_{\mathcal{D}'[(S')^{-1}]}(M, RF(K)) = \text{Hom}_{\mathcal{D}[S^{-1}]}(LG(M), K)$$

which is functorial in both variables on the triangulated subcategories (Definition B.1.2) of $\mathcal{D}[S^{-1}]$ and $\mathcal{D}'[(S')^{-1}]$ where RF and LG are defined.

Proposition B.7.4 ([?, Tag 0DVC]). Let $F : \mathcal{A} \rightarrow \mathcal{B}$ and $G : \mathcal{B} \rightarrow \mathcal{A}$ be functors of abelian categories (Definition .1.34) such that F is a right adjoint (Definition .1.8) to G ; in particular, F and G are additive (??). Let $K^\bullet$ be a complex (Definition .2.1) of $\mathcal{A}$ and let $M^\bullet$ be a complex of $\mathcal{B}$. If RF (Definition B.5.8) is defined (Definition B.5.5) at $K^\bullet$ and LG is defined at $M^\bullet$, then there is a canonical isomorphism

$$\text{Hom}_{D(\mathcal{B})}(M^\bullet, RF(K^\bullet)) = \text{Hom}_{D(\mathcal{A})}(LG(M^\bullet), K^\bullet)$$

This isomorphism is functorial in both variables on the triangulated subcategories (Definition B.1.2) of $D(\mathcal{A})$ (Definition B.3.1) and $D(\mathcal{B})$ where RF and LG are defined.

Proposition B.7.5 ([?, Tag 09T5]). Let $F : \mathcal{A} \rightarrow \mathcal{B}$ and $G : \mathcal{B} \rightarrow \mathcal{A}$ be functors of abelian categories (Definition .1.34) such that F is a right adjoint (Definition .1.8) to G . If the derived functors (Definition B.5.8) $RF : D(\mathcal{A}) \rightarrow D(\mathcal{B})$ and $LG : D(\mathcal{B}) \rightarrow D(\mathcal{A})$ exist, then RF is a right adjoint to LG .

B.8. Having enough K -injectives and flats for sheaves of modules.

Theorem B.8.1. Let $(\mathcal{C}, J, \mathcal{O})$ be a ringed site (Definition 6.1.5) where the underlying site $(\mathcal{C}, J)$ is essentially small. In particular, $\text{Mod}_{\mathcal{O}}$ is an abelian category (Definition .1.34) and in fact is a Grothendieck category (??).

For every chain complex (Definition .2.1) $A^\bullet$ in $\text{Mod}_{\mathcal{O}}$ (Definition 7.1.3) (resp. ${}_{\mathcal{O}}\text{Mod}$), there is a right resolution (Definition B.6.1) $A^\bullet \rightarrow I^\bullet$ such that $I^\bullet$ is K -injective (Definition B.6.4).

Proof. This follows from Theorem B.6.12. $\square$

Definition B.8.2. Let $(\mathcal{C}, J, \mathcal{O})$ be a ringed site (Definition 6.1.5) whose underlying site (Definition A.0.4) is essentially small.

A left (resp. right) $\mathcal{O}$ -module (Definition 7.1.3) $\mathcal{F}$ is called *flat* if the functor $- \otimes_{\mathcal{O}} \mathcal{F} : \text{Mod}_{\mathcal{O}} \rightarrow \text{Sh}(\mathcal{C}, J; \text{Ab})$ (resp. $\mathcal{F} \otimes_{\mathcal{O}} - : {}_{\mathcal{O}}\text{Mod} \rightarrow \text{Sh}(\mathcal{C}, J; \text{Ab})$) (??) (Definition 4.0.2) is exact (Definition .1.36).

Equivalently, $\mathcal{F}$ is flat if and only if it is flat with respect to $\otimes$.

Proposition B.8.3. Let $(\mathcal{C}, J, \mathcal{O})$ be a ringed site (Definition 6.1.5) whose underlying site (Definition A.0.4) is essentially small.

The categories $\text{Mod}_{\mathcal{O}}$ and ${}_{\mathcal{O}}\text{Mod}$ have enough flat $\mathcal{O}$ -modules (Definition B.8.2) on the left. (♠ TODO: prove)

Proposition B.8.4. Let $(\mathcal{C}, J, \mathcal{O})$ be a ringed site (Definition 6.1.5) whose underlying site (Definition A.0.4) is essentially small.

There are enough K -flat resolutions to the left of chain complexes (Definition .2.1) in of objects in $\text{Mod}_{\mathcal{O}}$ and ${}_{\mathcal{O}}\text{Mod}$. More precisely, for any chain complex (Definition .2.1) $A_\bullet \in \mathbf{Ch}(\text{Mod}_{\mathcal{O}})$ (resp. $\mathbf{Ch}({}_{\mathcal{O}}\text{Mod})$), there is a K -flat (Definition B.6.2) (with respect to $\otimes : \text{Mod}_{\mathcal{O}} \times {}_{\mathcal{O}}\text{Mod} \rightarrow \text{Sh}(\mathcal{C}, J; \text{Ab})$) chain complex $F_\bullet \in \text{Mod}_{\mathcal{O}}$ and a left resolution (Definition B.6.1) $F_\bullet \rightarrow A_\bullet$.

Proof. Since $\text{Mod}_{\mathcal{O}}$ and ${}_{\mathcal{O}}\text{Mod}$ have enough flats on the left by Proposition B.8.3, every (homologically) bounded below complex in $\text{Mod}_{\mathcal{O}}$ and ${}_{\mathcal{O}}\text{Mod}$ has a left resolution by a (homologically) bounded below complex of flat objects; a (homologically) bounded below complex of flats is K -flat with respect to $\otimes$ by Lemma B.6.14. To see that Lemma B.6.14 applies, note that $\otimes$ is a left adjoint by ?? and hence commutes with colimits by ??, and note that $\text{Sh}(\mathcal{C}, J; \text{Ab})$ is AB5. Theorem B.6.10 then applies to conclude that chain complex of objects in $\text{Mod}_{\mathcal{O}}$ and ${}_{\mathcal{O}}\text{Mod}$ has a left K -flat resolution. $\square$

B.9. Definitions of derived functors for modules of ringed sites.

Notation B.9.1. Let $(\mathcal{C}, J, \mathcal{O})$ be a ringed site (Definition 6.1.5) where the underlying site $(\mathcal{C}, J)$ is essentially small or let $(\mathcal{C}, \mathcal{O})$ be a ringed topos. In particular, $\text{Mod}_{\mathcal{O}}$ is an abelian category (Definition .1.34).

Denote by $D(\mathcal{O})$, $D^+(\mathcal{O})$, $D^-(\mathcal{O})$, $D^b(\mathcal{O})$ the derived categories $D(\mathcal{A})$, $D^+(\mathcal{A})$, $D^-(\mathcal{A})$, $D^b(\mathcal{A})$ (Definition B.3.1) where $\mathcal{A}$ is either the abelian category ${}_{\mathcal{O}}\text{Mod}$ or $\text{Mod}_{\mathcal{O}}$.

Definition B.9.2. Let $(\mathcal{C}, J, \mathcal{O})$ be a ringed site (Definition 6.1.5) whose underlying site (Definition A.0.4) is essentially small.

Given (cochain) complexes (Definition .2.1) $K^\bullet, L^\bullet$ of (say right) $\mathcal{O}$ -modules (Definition 7.1.3), the *internal sheaf/sheafy Hom complex* $\mathcal{H}om^\bullet(K^\bullet, L^\bullet)$, also called the *(product) total sheaf Hom complex*, but not to be confused with the internal hom complex (Definition B.6.3) $\text{Hom}^\bullet(K^\bullet, L^\bullet)$, is the product total complex $\text{Tot}^\Pi(\mathcal{H}om(K^\bullet, L^\bullet))$, where $\mathcal{H}om(K^\bullet, L^\bullet)$ is the double complex associated to (??) the sheaf hom functor

$$\mathcal{H}om : \text{Mod}_{\mathcal{O}}^{\text{op}} \times \text{Mod}_{\mathcal{O}} \rightarrow \text{Sh}(\mathcal{C}, J; \mathbb{Z})$$

(Definition 7.1.3) (Definition 4.0.2).

More explicitly, $\mathcal{H}om^\bullet(K^\bullet, L^\bullet)$ is the complex of objects in $\text{Sh}(\mathcal{C}, J; \mathbb{Z})$ given by

- **Objects:** In each degree $n \in \mathbb{Z}$, the sheaf of $\mathbb{Z}$ -modules is the product

$$\mathcal{H}om^n(K^\bullet, L^\bullet) = \prod_{p \in \mathbb{Z}} \mathcal{H}om_{\mathcal{O}}(K^p, L^{p+n}).$$

A local section $f \in \mathcal{H}om^n(K^\bullet, L^\bullet)(U)$ for an object $U \in \mathcal{C}$ is a family of $\mathcal{O}|_U$ -linear morphisms $f_p : K^p|_U \rightarrow L^{p+n}|_U$.

- **Differential:** The differential $d^n : \mathcal{H}om^n(K^\bullet, L^\bullet) \rightarrow \mathcal{H}om^{n+1}(K^\bullet, L^\bullet)$ is defined for a family of local morphisms $f = (f_p)$ by

$$(d^n f)_p = d_L^{p+n} \circ f_p - (-1)^n f_{p+1} \circ d_K^p,$$

where $d_K^p : K^p \rightarrow K^{p+1}$ and $d_L^q : L^q \rightarrow L^{q+1}$ are the cohomological differentials of the complexes.

Definition B.9.3 (cf. [?, Tag 08J7] for a discussion when $\mathcal{O}$ is assumed commutative). Let $(\mathcal{C}, J, \mathcal{O})$ be an essentially small ringed site (Definition 6.1.5) with a pretopology (Definition A.0.4). (♠ TODO: the “pretopology” part should be loosened)

The *derived (sheafy/sheaf) internal hom functor* is the functor $R\mathcal{H}om : D(\mathcal{O})^{\text{op}} \times D(\mathcal{O}) \rightarrow D(\text{Sh}(\mathcal{C}, J; \mathbb{Z}))$ (Notation B.9.1), also denoted by notations such as $R\mathcal{H}om(\mathcal{M}, \mathcal{N})$, $R\mathcal{H}om(\mathcal{M}, \mathcal{N})$, or $R\mathcal{H}om(\mathcal{M}, \mathcal{N})$, defined as the right derived functor (Definition B.5.8) of the sheaf hom functor

$$\mathcal{H}om : \text{Mod}_{\mathcal{O}}^{\text{op}} \times \text{Mod}_{\mathcal{O}} \rightarrow \text{Sh}(\mathcal{C}, J; \mathbb{Z}).$$

(♠ TODO: 07A5 for justifying why such a thing exists)

(♠ TODO: This is defined because...) Explicitly, given objects $L, M \in D(\mathcal{O})$, $R\mathcal{H}om(L, M)$ is given as follows: let $\mathcal{L}^\bullet$ be any complex of $\mathcal{O}$ -modules representing L and let $\mathcal{I}^\bullet$ be any (right) K -injective (Definition B.6.4) resolution (Definition B.6.1) of M , which exists by Theorem B.8.1. The object $R\mathcal{H}om(L, M)$ is then given as the object of $D(\text{Sh}(\mathcal{C}, J; \mathbb{Z}))$ represented by the internal sheaf hom object (Definition B.9.2) $\mathcal{H}om^\bullet(\mathcal{L}^\bullet, \mathcal{I}^\bullet)$.

Definition B.9.4 (cf. [?, Tag 06YU] for a simplified discussion when $\mathcal{O}$ is assumed commutative). Let $(\mathcal{C}, J, \mathcal{O})$ be an essentially small ringed site (Definition 6.1.5) with a pretopology (Definition A.0.4). (♠ TODO: the “pretopology” part should be loosened)

The *derived tensor product* is the functor

$$- \otimes_{\mathcal{O}}^L - : D(\text{Mod}_{\mathcal{O}}) \times D({}_{\mathcal{O}}\text{Mod}) \rightarrow D(\text{Sh}(\mathcal{C}, J; \text{Ab}))$$

(Definition 7.1.3) (Notation B.9.1) also denoted by notations such as $\overset{L}{\otimes}$ defined as the left derived functor (Definition B.5.8) of the tensor product functor

$$- \otimes_{\mathcal{O}} - : \text{Mod}_{\mathcal{O}} \times_{\mathcal{O}} \text{Mod} \rightarrow \text{Sh}(C, J; \text{Ab})$$

Explicitly, given objects $L, M \in D(\mathcal{O})$, $L \otimes_{\mathcal{O}} M$ is given as follows: let $\mathcal{L}^{\bullet}$ be any complex of right $\mathcal{O}$ -modules representing L and let $\mathcal{M}^{\bullet}$ be any complex of left $\mathcal{O}$ -modules representing M , such that $\mathcal{L}^{\bullet}$ is K -flat (Definition B.6.2) with respect to $\otimes_{\mathcal{O}}$ on the left for direct sum total complexes or $\mathcal{M}^{\bullet}$ is K -flat with respect to $\otimes_{\mathcal{O}}$ on the left for direct sum total complexes; note that such complexes exist by Proposition B.8.4. The object $L \otimes_{\mathcal{O}}^L M$ is then given as the object of $D(\text{Sh}(C, J; \text{Ab}))$ represented by the total complex $\text{Tot}^{\oplus}(\mathcal{L}^{\bullet} \otimes_{\mathcal{O}} \mathcal{M}^{\bullet})$. (♠ TODO: justify definedness, i.e. why does this actually define the derived functor)

Theorem B.9.5 (cf. [?, Tag 08J9]). Let $(\mathcal{C}, J, \mathcal{O})$ be an essentially small ringed site (Definition 6.1.5) with a pretopology (Definition A.0.4). (♠ TODO: this pretopology condition should be loosened)

Let K, L, M be objects of $D(\mathcal{O})$ (Notation B.9.1). There is a canonical isomorphism

$$R\mathcal{H}om(K, R\mathcal{H}om(L, M)) \cong R\mathcal{H}om(K \otimes_{\mathcal{O}}^L L, M)$$

(Definition B.9.3) (Definition B.9.4) in $D(\mathcal{O})$ functorial in K, L, M . Taking the zeroth sheaf cohomology (or equivalently global sections) of both sides, one obtains the adjunction

$$\text{Hom}_{D(\mathcal{O})}(K, R\mathcal{H}om(L, M)) \cong \text{Hom}_{D(\mathcal{O})}(K \otimes_{\mathcal{O}}^L L, M).$$

Definition B.9.6. Let $((\mathcal{C}, J), \mathcal{O}_{\mathcal{C}})$ and $((\mathcal{D}, K), \mathcal{O}_{\mathcal{D}})$ be ringed sites (Definition 6.1.5).

A *morphism of ringed sites* $f : ((\mathcal{C}, J), \mathcal{O}_{\mathcal{C}}) \rightarrow ((\mathcal{D}, K), \mathcal{O}_{\mathcal{D}})$ is a pair $(f, f^{\#})$ where:

1. $f : (\mathcal{C}, J) \rightarrow (\mathcal{D}, K)$ is a morphism of sites.
2. $f^{\#} : \mathcal{O}_{\mathcal{D}} \rightarrow f_* \mathcal{O}_{\mathcal{C}}$ (Definition 6.1.7) is a morphism of sheaves of rings (Definition 4.0.2) on the site $(\mathcal{D}, K)$.

Equivalently, the morphism $f^{\#}$ a morphism can be described by the adjoint morphism (??) $f^{\#} : f^{-1} \mathcal{O}_{\mathcal{D}} \rightarrow \mathcal{O}_{\mathcal{C}}$ of sheaves of rings on $(\mathcal{C}, J)$, where f^{-1} is the inverse image functor (Definition 6.1.10) of sheaves.

Definition B.9.7 (cf. [?, Tag 07A5]). Let $f : (\mathcal{C}, \mathcal{O}_{\mathcal{C}}) \rightarrow (\mathcal{D}, \mathcal{O}_{\mathcal{D}})$ be a morphism of ringed topoi.

The *(total) derived direct image* or *(total) derived pushforward functor* Rf_* is the functor

$$Rf_* : D(\mathcal{O}_{\mathcal{C}}) \rightarrow D(\mathcal{O}_{\mathcal{D}})$$

(Notation B.9.1) defined as the right derived functor (Definition B.5.8) of the direct image functor f_* . Explicitly, given an object $K \in D(\mathcal{O}_{\mathcal{C}})$, $Rf_*(K)$ is given as follows: Let $\mathcal{I}^{\bullet}$ be any (right) K -injective (Definition B.6.4) resolution (Definition B.6.1) of M , which exists by Theorem B.8.1. The object $Rf_*(K)$ is then given as the object of $D(\mathcal{O}_{\mathcal{D}})$ represented by the chain complex $f_*(\mathcal{I}^{\bullet})$ (Proposition B.4.2). This is well defined by Proposition B.7.2.

Definition B.9.8 (cf. [?, Tag 06YY]). Let $f : (\mathcal{C}, \mathcal{O}_{\mathcal{C}}) \rightarrow (\mathcal{D}, \mathcal{O}_{\mathcal{D}})$ be a morphism of ringed topoi.

The *(total) derived direct image* or *(total) derived pushforward functor* Lf^* is the functor

$$Lf^* : D(\mathcal{O}_{\mathcal{D}}) \rightarrow D(\mathcal{O}_{\mathcal{C}})$$

(Notation B.9.1) defined as the left derived functor (Definition B.5.8) of the inverse image functor f^* . Explicitly, given an object $K \in D(\mathcal{O}_{\mathcal{D}})$, $Lf^*(K)$ is given as follows: Let $\mathcal{F}^\bullet$ be any (left) K -flat (Definition B.6.2) (with respect to $\otimes_{\mathcal{O}_{\mathcal{D}}}$ either on the left or right, depending on whether we are considering right or left $\mathcal{O}_{\mathcal{D}}$ -modules) resolution (Definition B.6.1) of M , which exists by Proposition B.8.4 (♠ TODO: justify the well definedness)

Theorem B.9.9. Let $f : (\mathcal{C}, \mathcal{O}_{\mathcal{C}}) \rightarrow (\mathcal{D}, \mathcal{O}_{\mathcal{D}})$ be a morphism of ringed topoi.

The functors Rf_* (Definition B.9.7) and Lf^* (Definition B.9.8) are adjoint, i.e. for all $L \in D(\mathcal{O}_{\mathcal{C}})$ and $M \in D(\mathcal{O}_{\mathcal{D}})$, we have natural isomorphisms

$$\mathrm{Hom}_{D(\mathcal{O}_{\mathcal{C}})}(Lf^*M, L) \cong \mathrm{Hom}_{D(\mathcal{O}_{\mathcal{D}})}(M, Rf_*L)$$

(Definition B.9.7) (Definition B.9.8)

♠ TODO: total derived sheaf cohomology

Proposition B.9.10. The following are examples of abelian categories (Definition .1.34):

1. The category of R - S bimodules where R, S are (not necessarily commutative) rings (Definition C.0.8) (??).
2. The category **Ab** of abelian groups and group homomorphisms is abelian.
3. The category Vect_k of vector spaces over a field k and k -linear maps is abelian.
4. More generally, if R is a noetherian ring, then the category of finitely generated R -modules is abelian.
5. For a ringed space $(X, \mathcal{O}_X)$, the category of $\mathcal{O}_X$ -modules (Definition 7.1.3) is abelian. (♠ TODO: a quasi-coherent sheaf on a locally ringed space)
6. If X is a scheme (Definition .3.1) (or more generally a locally ringed space), the category of quasi-coherent sheaves on X is abelian.
7. For any essentially small category (Definition .1.26) $\mathcal{C}$ and any abelian category $\mathcal{A}$, the functor category $[\mathcal{C}, \mathcal{A}]$ (Definition .1.33) and the category $\mathrm{PreShv}(\mathcal{C}, \mathcal{A})$ of presheaves (Definition 4.0.1) are abelian. (♠ TODO: apparently, the essentially smallness condition is removable, provided that the sheafification functor exists. However, the essentially small assumption is needed to show that the category of sheaves of $\mathcal{O}$ -modules is a Grothendieck abelian category. Verify all this. Moreover, when working with a big site of a scheme, one typically fixes a universe or work relative to a cardinal cutoff to treat it as essentially small)
8. For any site (Definition A.0.4) $(\mathcal{C}, J)$ on an essentially small category (Definition .1.26) $\mathcal{C}$ and any abelian category $\mathcal{A}$, the category $\mathrm{Shv}(\mathcal{C}, \mathcal{A})$ of sheaves (Definition 4.0.2) is abelian.
9. For any site (Definition A.0.4) $(\mathcal{C}, J)$ on an essentially small category (Definition .1.26) $\mathcal{C}$ and a sheaf of rings (Definition 4.0.2) $\mathcal{O}$ on $\mathcal{C}$, the category **Mod**($\mathcal{O}$) of $\mathcal{O}$ -modules (Definition 7.1.3) is an abelian category.

Notation B.9.11. Let $((\mathcal{C}, J), \mathcal{O})$ be a ringed site (Definition 6.1.5) where $(\mathcal{C}, J)$ is essentially small. (♠ TODO: The essentially smallness hypothesis might not be strictly needed to have that the category of sheaves of $\mathcal{O}$ -modules is abelian.)

1. The homotopy category (Definition B.1.1) of the category of $\mathcal{O}$ -modules (Definition 7.1.3), which is (Proposition B.9.10) abelian, is often denoted by notations such as $K(\mathcal{C}, \mathcal{O})$ or $K(\mathcal{O})$. All of the usual superscripts apply — we may thus speak of $K^{?,??}(\mathcal{C}, \mathcal{O})$ or $K^{?,??}(\mathcal{O})$ for $? \in \{+, -, b\}$ and $?? \in \{+, -, b, (\text{blank})\}$.

If $(\mathcal{C}, J)$ is a site on some “space”, e.g. a scheme or topological space, X , then it is also usual to denote the homotopy category by $K(X, \mathcal{O})$. We may also speak of $K^{?,??}(X, \mathcal{O})$ for $? \in \{+, -, b\}$ and $?? \in \{+, -, b, (\text{blank})\}$.

2. Similarly, it is customary to denote the derived category (Definition B.3.1) of the category of $\mathcal{O}$ -modules (Definition 7.1.3) of the category of $\mathcal{O}$ -modules by notations such as $D(\mathcal{C}, \mathcal{O})$ or $D(\mathcal{O})$. All of the usual superscripts apply — we may thus speak of $D^?(\mathcal{C}, \mathcal{O})$ or $D^?(\mathcal{O})$ for $? \in \{+, -, b\}$.

If $(\mathcal{C}, J)$ is a site on some “space”, e.g. a scheme or topological space, X , then it is also usual to denote the homotopy category by $D(X, \mathcal{O})$. We may also speak of $D^?(X, \mathcal{O})$ for $? \in \{+, -, b\}$.

Lemma B.9.12. Let $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ be a biadditive functor (Definition C.0.13) of abelian categories (Definition .1.34). Assume that (small) filtered colimits which exist in $\mathcal{C}$ are exact (e.g. which holds if $\mathcal{C}$ satisfies Ab5).

Let $A \in \mathcal{A}$ and $B \in \mathcal{B}$ be objects.

1. Suppose that left resolutions (Definition .2.7) $P_{A,\bullet} \rightarrow A$ and $P_{B,\bullet} \rightarrow B$ exist such that $P_{A,i}$ and $P_{B,i}$ are flat with respect to F on the left and right respectively, i.e. $F(P_{A,i}, -) : \mathcal{B} \rightarrow \mathcal{C}$ and $F(-, P_{B,i}) : \mathcal{A} \rightarrow \mathcal{C}$ are exact for all i .

The complexes $F(P_{A,\bullet}, B)$ and $F(A, P_{B,\bullet})$ are quasi-isomorphic (Definition .2.4) to the complex $\text{Tot}(F(P_{A,\bullet}, P_{B,\bullet}))$.

2. Suppose that right resolutions (Definition .2.7) $A \rightarrow I^{A,\bullet}$ and $B \rightarrow I^{B,\bullet}$ exist such that $I^{A,i}$ and $I^{B,i}$ are flat with respect to F on the left and right respectively, $F(I^{A,i}, -) : \mathcal{B} \rightarrow \mathcal{C}$ and $F(-, I^{B,i}) : \mathcal{A} \rightarrow \mathcal{C}$ are exact for all i .

The complexes $F(I^{A,\bullet}, B)$ and $F(A, I^{B,\bullet})$ are quasi-isomorphic (Definition .2.4) to the complex $\text{Tot}(F(I^{A,\bullet}, I^{B,\bullet}))$.

Proof. We prove 1. The other part is the dual statement.

Choose resolutions $P_{A,\bullet} \xrightarrow{\varepsilon} A$ and $P_{B,\bullet} \xrightarrow{\eta} B$ such that $F(P_{A,i}, -) : \mathcal{B} \rightarrow \mathcal{C}$ and $F(-, P_{B,i}) : \mathcal{A} \rightarrow \mathcal{C}$ are exact for all i . Identifying A and B with complexes concentrated in degree 0, we can form the three double complexes $F(P_{A,\bullet}, P_{B,\bullet})$, $F(A, P_{B,\bullet})$, and $F(P_{A,\bullet}, B)$. Note that the augmentation morphisms ε and η induce morphisms $\text{Tot}(F(P_{A,\bullet}, P_{B,\bullet})) \rightarrow \text{Tot}(F(A \otimes P_{B,\bullet}), \text{Tot}(F(P_{A,\bullet}, B)))$.

Let C be the double complex of objects in $\mathcal{C}$ obtained from $F(P_{A,\bullet}, P_{B,\bullet})$ by adding $F(A, P_{B,\bullet}[-1])$ in the column $p = -1$. One can show that the translate $\text{Tot}(C)[1]$ is the mapping cone (Definition .2.5) of the map

$$\text{Tot}(F(P_{A,\bullet}, P_{B,\bullet})) \xrightarrow{\varepsilon \otimes \text{id}} \text{Tot}(F(A, P_{B,\bullet})) = F(A, P_{B,\bullet}).$$

Moreover, since each $F(-, P_{B,i})$ is an exact functor, every row of C is exact, so $\text{Tot}(C)$ is exact by ???. Therefore, $F(\varepsilon, \text{id})$ is a quasi-isomorphism and hence

$$H_*(\text{Tot}(F(P_{A,\bullet}, P_{B,\bullet}))) \xrightarrow{H_*(F(\varepsilon, P_{B,\bullet}))} H_*(F(A, P_{B,\bullet}))$$

is a natural isomorphism.

By symmetry, there is a natural isomorphism $H_*(\text{Tot}(F(P_{A,\bullet}, P_{B,\bullet}))) \rightarrow H_*(F(P_{A,\bullet}, B))$. $\square$

Proposition B.9.13. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be abelian categories (Definition .1.34), and let $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ be a biadditive functor (Definition C.0.13).

1. Suppose that $\mathcal{A}$ and $\mathcal{B}$ both have enough projectives (Definition .1.37). Given objects $M \in D^-(\mathcal{A})$ and $N \in D^-(\mathcal{A})$ (Definition B.3.1), the objects $LF(M, N)$ obtained as $(LF(M, -))(N)$ and $(LF(-, N))(M)$ are naturally isomorphic. Thus, the two definitions of $LF(M, N)$ in Definition B.9.14 are in agreement.
2. Dually, suppose that $\mathcal{A}$ and $\mathcal{B}$ both have enough injectives (Definition .1.37). Given objects $M \in D^+(\mathcal{A})$ and $N \in D^+(\mathcal{A})$ (Definition B.3.1), the objects $RF(M, N)$ obtained as $(RF(M, -))(N)$ and $(RF(-, N))(M)$ are naturally isomorphic. Thus, the two definitions of $RF(M, N)$ in Definition B.9.14 are in agreement.

Proof. We prove 1. The other part is dual. By Theorem B.3.4, note that $D^-(\mathcal{A})$ and $D^-(\mathcal{B})$ are respectively equivalent to the categories $K^-(\mathcal{P}_{\mathcal{A}})$ and $K^-(\mathcal{P}_{\mathcal{B}})$ (Definition B.3.3) of cohomologically bounded above complexes of projectives in $\mathcal{A}$ and $\mathcal{B}$ respectively. Let

$$P_{\bullet} \rightarrow M$$

and

$$Q_{\bullet} \rightarrow N$$

be projective resolutions. By Corollary B.4.4,

$$\begin{aligned} (LF(M, -))(N) &\cong q(K(F(M, -)))(N) \\ (LF(-, N))(M) &\cong q(K(F(-, N)))(M) \end{aligned}$$

The former is represented by the complex $F(M, Q_{\bullet})$ and the latter is represented by the complex $F(P_{\bullet}, N)$. These are quasi-isomorphic by Lemma B.9.12. $\square$

Definition B.9.14. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be abelian categories (Definition .1.34), and let $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ be a biadditive functor (Definition C.0.13).

There are notions of derived functors that we may consider depending on whether $\mathcal{A}$ or $\mathcal{B}$ has enough projectives or injectives or flats with respect to F . (♠ TODO: define for flats)

(♠ TODO: I really should be letting M and N be complexes, not just objects.)

1.

1. (a) Let $M \in \mathcal{A}$. Assume that $\mathcal{B}$ has enough projectives (Definition .1.37). By Corollary B.4.4, the functor $F(M, -) : \mathcal{B} \rightarrow \mathcal{C}$ has a left derived functor (Definition B.4.1) $L(F(M, -)) : D^-(\mathcal{B}) \rightarrow D(\mathcal{C})$ (Definition B.3.1), which we may write as $LF(M, -)$.

- (b) Symmetrically, let $N \in \mathcal{B}$. Assume that $\mathcal{A}$ has enough projectives (Definition .1.37). By Corollary B.4.4, the functor $F(-, N) : \mathcal{A} \rightarrow \mathcal{C}$ has a left derived functor (Definition B.4.1) $L(F(-, N)) : D^-(\mathcal{A}) \rightarrow D(\mathcal{C})$ (Definition B.3.1), which we may write as $LF(-, N)$.

Assuming that $\mathcal{A}$ and $\mathcal{B}$ have enough projectives, the notations $LF(M, N)$ above are in agreement (Proposition B.9.13)

2. (a) Let $M \in \mathcal{A}$. Assume that $\mathcal{B}$ has enough injectives (Definition .1.37). By Corollary B.4.4, the functor $F(M, -) : \mathcal{B} \rightarrow \mathcal{C}$ has a right derived functor (Definition B.4.1) $R(F(M, -)) : D^+(\mathcal{B}) \rightarrow D(\mathcal{C})$ (Definition B.3.1), which we may write as $RF(M, -)$.
- (b) Symmetrically, let $N \in \mathcal{B}$. Assume that $\mathcal{A}$ has enough injectives (Definition .1.37). By Corollary B.4.4, the functor $F(-, N) : \mathcal{A} \rightarrow \mathcal{C}$ has a right derived functor (Definition B.4.1) $R(F(-, N)) : D^+(\mathcal{A}) \rightarrow D(\mathcal{C})$ (Definition B.3.1), which we may write as $RF(-, N)$.

Assuming that $\mathcal{A}$ and $\mathcal{B}$ have enough injectives, the notations $RF(M, N)$ above are in agreement (Proposition B.9.13)

3. Let $M \in \mathcal{A}$. Assume that $\mathcal{B}$ has enough flats with respect to $\otimes$. Define $LF(M, -) : D^-(\mathcal{B}) \rightarrow D(\mathcal{C})$ as follows:

See Definition B.9.15 for notation used in the case that F is written as a tensor product.

Definition B.9.15. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be abelian categories (Definition .1.34), and let $\otimes : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ be a biadditive functor (Definition C.0.13) written as tensor product.

(♠ TODO: need a definition with enough flats)

1. Let $M \in \mathcal{A}$. Assume that $\mathcal{B}$ has enough projectives (Definition .1.37). We write $M \otimes^L -$ for $LF(M, -) : D^-(\mathcal{B}) \rightarrow D(\mathcal{C})$ (Definition B.9.14) in the case that $F = M \otimes - : \mathcal{B} \rightarrow \mathcal{C}$.
2. Symmetrically, let $N \in \mathcal{B}$. Assume that $\mathcal{A}$ has enough projectives (Definition .1.37). We write $- \otimes^L N$ for $LF(-, N) : D^+(\mathcal{A}) \rightarrow D(\mathcal{C})$ (Definition B.9.14) in the case that $F = - \otimes N : \mathcal{A} \rightarrow \mathcal{C}$.

(♠ TODO: show that projectives vs. flats yield the same thing) (♠ TODO: show that flats in each variable yield the same thing)

3. Let $M \in \mathcal{A}$. Assume that $\mathcal{B}$ has enough flats. We alternatively define $M \otimes^L - : D^-(\mathcal{B}) \rightarrow D(\mathcal{C})$ as follows — given an object $N \in D^-(\mathcal{B})$, say that $Q^\bullet$ is a complex of flat objects in $\mathcal{B}$ representing N (♠ TODO: show that such a thing exists), and let $M \otimes^L N$ be the object of $D(\mathcal{C})$ represented by the complex $M \otimes Q^\bullet$. (♠ TODO: show that this is well defined, i.e. does not depend on the choice of flat resolution)
4. Symmetrically, let $N \in \mathcal{B}$. Assume that $\mathcal{A}$ has enough flats. We alternatively define $- \otimes^L N : D^-(\mathcal{A}) \rightarrow D(\mathcal{C})$ as follows — given an object $M \in D^-(\mathcal{A})$, say that $P^\bullet$ is a complex of flat objects in $\mathcal{A}$ representing M (♠ TODO: show that such a thing exists), and let $M \otimes^L N$ be the object of $D(\mathcal{C})$ represented by the complex $P^\bullet \otimes N$. (♠ TODO: show that this is well defined, i.e. does not depend on the choice of flat resolution)

The first two notions agree assuming that $\mathcal{A}$ and $\mathcal{B}$ have enough projectives. (♠ TODO: comment on the next two notions agreeing)

APPENDIX C. MISCELLANEOUS DEFINITIONS

Definition C.0.1. A **commutative (unital) ring** is a ring (Definition C.0.8) $(R, +, \cdot)$ such that $\cdot$ is a commutative operation, i.e. $a \cdot b = b \cdot a$.

For many writers (e.g. “commutative” algebraists or number theorists), a **ring** refers to a commutative ring as above.

Definition C.0.2. Let R be a (not necessarily commutative, possibly nonunital) ring (Definition C.0.8). A **left ideal of R** is a subset $I \subseteq R$ such that

- $(I, +)$ is an additive subgroup of $(R, +)$,
- $RI \subseteq I$, i.e., for all $r \in R$ and $x \in I$, one has $rx \in I$.

Similarly, a **right ideal of R** is a subset $I \subseteq R$ such that

- $(I, +)$ is an additive subgroup of $(R, +)$,
- $IR \subseteq I$, i.e., for all $r \in R$ and $x \in I$, one has $xr \in I$.

A **two-sided ideal** (or simply an **ideal**) of R is a subset $I \subseteq R$ which is both a left ideal and a right ideal of R . We denote by $I \trianglelefteq R$ the relation expressing that I is a two-sided ideal of R .

Equivalently, an left/right/two-sided ideal of R is a submodule of R as an R -module (Definition C.0.11).

A left/right/two-sided ideal is said to be **proper** if it is strictly contained in R .

Note that every left or right ideal of a commutative ring is a two-sided ideal.

Definition C.0.3. Let R be a ring (Definition C.0.8) with unity, not necessarily commutative. The ring R is called a **local ring** if it has a unique maximal left ideal (Definition C.0.4). In this case, R also has a unique maximal right ideal, and these coincide with the Jacobson radical $J(R)$ of R .

The unique maximal left (and right) ideal of a local ring R may sometimes be denoted by $\mathfrak{m}_R$.

The following are each equivalent conditions for R to be a local ring:

1. $R/J(R)$ is a division ring
2. $R = J(R) \cup R^\times$

Definition C.0.4. Let R be a (not necessarily commutative) ring (Definition C.0.8). A proper two-sided ideal $P \trianglelefteq R$ (Definition C.0.2) is called a **prime ideal** if the following equivalent conditions holds:

1. If I, J are left ideals and $IJ \subseteq P$, then $I \subseteq P$ or $J \subseteq P$.
2. If I, J are right ideals and $IJ \subseteq P$, then $I \subseteq P$ or $J \subseteq P$.
3. If I, J are two-sided ideals and $IJ \subseteq P$, then $I \subseteq P$ or $J \subseteq P$.

4. If $x, y \in R$ with $xRy \subset \mathfrak{p}$, then $x \in \mathfrak{p}$ or $y \in \mathfrak{p}$.

A proper left/right/two-sided ideal $M \subsetneq R$ is called **maximal** if there exists no other left/right/two-sided ideal $J \leq R$ such that $M \subsetneq J \subsetneq R$. Equivalently,

- a left/right ideal M of R is maximal if and only if the quotient module R/M is a simple left/right R -module.
- a two-sided ideal M of R is maximal if and only if the quotient ring R/M is a simple ring.

Definition C.0.5. Let $(R, \mathfrak{m}_R)$ and $(S, \mathfrak{m}_S)$ be local rings (Definition C.0.21), not necessarily commutative. A ring homomorphism $\varphi : R \rightarrow S$ is called a **local morphism** (or **local homomorphism**) if $\varphi(\mathfrak{m}_R) \subseteq \mathfrak{m}_S$.

Definition C.0.6 (Topology). Let X be a set. A **topology on X** is a collection $\mathcal{T}$ of subsets of X such that:

1. $\emptyset \in \mathcal{T}$ and $X \in \mathcal{T}$,
2. For any collection $\{U_i\}_{i \in I} \subseteq \mathcal{T}$ (with I arbitrary), the union $\bigcup_{i \in I} U_i \in \mathcal{T}$,
3. For any finite collection $\{U_1, \dots, U_n\} \subseteq \mathcal{T}$, the intersection $U_1 \cap \dots \cap U_n \in \mathcal{T}$.

If $\mathcal{T}$ is a topology on X , the pair $(X, \mathcal{T})$ is called a **topological space**. Members of $\mathcal{T}$ are called **open sets**.

A subset $C \subseteq X$ is **closed** (also called a **closed subset**) if its complement $X \setminus C$ is an open set in $\mathcal{T}$.

One very often refers to X as a topological space, omitting the notation of the topology $\mathcal{T}$.

The collection of all topologies on a set X may be denoted by notations such as **Top(X)**, **Top(X)**, or **Top(X)**.

Definition C.0.7. Let X be a topological space (Definition C.0.6). A subset $Z \subseteq X$ is called a **locally closed subset** if Z can be written as the intersection $U \cap C$, where U is an open subset of X and C is a closed subset of X . Equivalently, Z is a locally closed subset if it is an open subset of its closure $\overline{Z}$ endowed with the subspace topology.

Definition C.0.8. A **ring** is a triple $(R, +, \cdot)$ where

1. $(R, +)$ is a commutative group, and
2. $(R, \cdot)$ is a monoid.
3. $\cdot$ is distributive over $+$, i.e. for all $a, b, c \in R$, we have

$$a \cdot (b + c) = a \cdot b + a \cdot c \quad \text{and} \quad (a + b) \cdot c = a \cdot c + b \cdot c.$$

Equivalently, a ring is a triple $(R, +, \cdot)$ where $+, \cdot : R \times R \rightarrow R$ are binary operations satisfying

1. $(a + b) + c = a + (b + c)$ and $(ab)c = a(bc)$ for all $a, b, c \in R$
2. There exists an element $0 \in R$ such that $a + 0 = a = 0 + a$ for all $a \in R$.

3. For every $a \in R$, there exists an element $-a \in R$ such that $a + (-a) = 0 = (-a) + a$ for all $a \in R$.
4. There exists an element $1 \in R$ such that $a \cdot 1 = a = 1 \cdot a$ for all $a \in R$.
5. For all $a, b, c \in R$, we have

$$a \cdot (b + c) = a \cdot b + a \cdot c \quad \text{and} \quad (a + b) \cdot c = a \cdot c + b \cdot c.$$

The operation $+$ is often called **addition** and the operation $\cdot$ is often called **multiplication**. Accordingly, the identity element 0 of $+$ is often called the **additive identity** and the identity element 1 of $\cdot$ is often called the **multiplicative identity**.

While other notations may be used to denote addition and multiplication, if no notation is specified explicitly, then $+$ and $\cdot$ are universally used as default notations for addition and multiplication in a ring.

Remark C.0.9. Some writers might not require a ring to have a multiplicative identity element, i.e. would define a ring so that $(R, +)$ is a commutative group, $(R, \cdot)$ is a semigroup, and $\cdot$ is distributive over $+$. Such writers would call the notion of ring in Definition C.0.8 a **unitary ring** to emphasize the existence of the multiplicative identity 1 .

Definition C.0.10 (Constant sheaf on a site). Let $\mathcal{C}$ be a category (Definition .1.1), let $\mathcal{A}$ be a category, and let A be an object of $\mathcal{A}$.

1. The **constant presheaf on $\mathcal{C}$ with value A** is the presheaf (Definition 4.0.1) P defined by

$$P(U) = A$$

for every object U of $\mathcal{C}$ such that every morphism $f : V \rightarrow U$ in $\mathcal{C}$ induces the identity map $A = P(U) \rightarrow P(V) = A$.

2. Let $\mathcal{C}$ be a site (Definition A.0.4) and assume that a sheafification functor (Definition 4.3.1)

$$a : \text{Shv}(\mathcal{C}, \mathcal{A}) \rightarrow \text{PreShv}(\mathcal{C}, \mathcal{A})$$

exists (e.g. see Theorem 4.3.2). The **constant sheaf on $\mathcal{C}$ with value A** , or the **constant sheaf on $\mathcal{C}$ associated to A** commonly denoted $\underline{A}$ or sometimes just A by abuse of notation, is the sheaf associated to (Theorem 4.3.2) the constant presheaf P with value A above.

3. Let $\mathcal{C}$ be a site. Let $\mathcal{O}$ be a sheaf of (not-necessarily commutative) rings on $\mathcal{C}$. Assume that the global sections ring $\Gamma(\mathcal{O})$ (Definition 7.1.1) exists. A **constant $\mathcal{O}$ -module** is an $\mathcal{O}$ -module (Definition 7.1.3) $\mathcal{F}$ which is isomorphic as a sheaf to the constant sheaf on $\mathcal{C}$ with value M where M is a module of the ring $\Gamma(\mathcal{O})$. Note that sheafification functors exist for presheaves/sheaves valued in Ab (Theorem 4.3.2).

In case that $\mathcal{O}$ is the constant sheaf associated to A for some (not-necessarily commutative) ring A , a constant $\mathcal{O}$ -module is simply called a **constant A -module**.

Definition C.0.11. Let R be a not-necessarily commutative ring (Definition C.0.8).

1. A **left R -module** is an abelian group $(M, +)$ together with an operation $R \times M \rightarrow M$, denoted $(r, m) \mapsto rm$, such that for all $r, s \in R$ and $m, n \in M$:
 - $r(m + n) = rm + rn$,

- $(r + s)m = rm + sm$,
 - $(rs)m = r(sm)$,
 - $1_R m = m$ where 1_R is the multiplicative identity of R .
2. A **right R -module** is defined similarly as an abelian group $(M, +)$ with an operation $M \times R \rightarrow M$, denoted $(m, r) \mapsto mr$, such that for all $r, s \in R$ and $m, n \in M$:
- $(m + n)r = mr + nr$,
 - $m(r + s) = mr + ms$,
 - $m(rs) = (mr)s$,
 - $m1_R = m$.
3. Let R and S be (not necessarily commutative) rings (Definition C.0.8).
 An **R - S -bimodule** (or an **R - S -module** or an (R, S) -module, etc.) is an abelian group $(M, +)$ equipped with
- (a) a left action of R :

$$R \times M \rightarrow M, \quad (r, m) \mapsto r \cdot m,$$

making M a left R -module (Definition C.0.11),

- (b) a right action of S :

$$M \times S \rightarrow M, \quad (m, s) \mapsto m \cdot s,$$

making M a right S -module,

such that the left and right actions commute; that is, for all $r \in R$, $s \in S$, and $m \in M$,

$$r \cdot (m \cdot s) = (r \cdot m) \cdot s.$$

4. A **two-sided R -module** (or **R -bimodule**) is an R - R -bimodule.

If R is a commutative ring (Definition C.0.1), then a left/right R -module can automatically be regarded as a two-sided R -module. As such, we simply talk about **R -modules** in this case.

Any abelian group is equivalent to a two-sided $\mathbb{Z}$ -module. Moreover, any left R -module is equivalent to an $R - \mathbb{Z}$ -bimodule (Definition C.0.11) and any right R -module is equivalent to an $\mathbb{Z} - R$ -bimodule (Definition C.0.11). Given a left/right/two-sided R -module, its **natural bimodule structure** will refer to its structure as a $R - \mathbb{Z} / \mathbb{Z} - R / R - R$ bimodule. In this way, many definitions associated with the notions of left/right/two-sided R -modules can be defined as special cases for definitions for R - S -bimodules.

Definition C.0.12 (Tensor product of bimodules). Let R, S, T be (not necessarily commutative) rings (Definition C.0.8), let M be an R - S bimodule (Definition C.0.11), and let N be an S - T bimodule. In the free abelian group $\mathbb{Z}[M \times N]$ generated by the Cartesian product $M \times N$, let U be the subgroup generated by elements of the form (♠ TODO: subgroup generated)

$$\begin{aligned} & (m + m', n) - (m, n) - (m', n), \\ & (m, n + n') - (m, n) - (m, n'), \\ & (m \cdot s, n) - (m, s \cdot n), \end{aligned}$$

for all $m, m' \in M$, $n, n' \in N$, and $s \in S$. The **tensor product of M and N over S** is the quotient abelian group

$$M \otimes_S N := \mathbb{Z}[M \times N] / U.$$

The image of an element of the form $(m, n) \in M \times N$ in $M \otimes_S N$ is denoted $m \otimes n$ and called a *pure tensor*. In general, the elements of $M \otimes_S N$ are finite sums

$$\sum_{i=1}^n m_i \otimes n_i \quad m_i \in M, n_i \in N$$

of pure tensors. Thus, the pure tensors satisfy the following relations:

$$\begin{aligned} (m + m') \otimes n &= m \otimes n + m' \otimes n \\ m \otimes (n + n') &= m \otimes n + m \otimes n' \\ (m \cdot s) \otimes n &= m \otimes (s \cdot n) \end{aligned}$$

This tensor product becomes naturally an R - T bimodule with left action and right action defined by

$$\begin{aligned} r \cdot (m \otimes n) &= (r \cdot m) \otimes n, \\ (m \otimes n) \cdot t &= m \otimes (n \cdot t), \end{aligned}$$

for all $r \in R$, $t \in T$, $m \in M$, and $n \in N$.

Inductively, given rings $R_0, \dots, R_k$ and $R_{i-1} - R_i$ -bimodules M_i for $i = 1, \dots, k$, we may speak of the tensor product

$$M_0 \otimes_{R_1} M_1 \otimes_{R_2} \cdots \otimes_{R_{k-1}} M_k;$$

tensor products are associative(♠ TODO:), so parentheses are not strictly needed to notate them. Its *pure tensors* are elements of the form $m_0 \otimes m_1 \otimes \cdots \otimes m_k$ for $m_i \in M_i$, and its general elements are finite sums

$$\sum_{j=1}^n m_{0j} \otimes m_{1j} \otimes \cdots \otimes m_{kj} \quad m_{ij} \in M_i.$$

of pure tensors. It also has a natural $R_0 - R_k$ -bimodule structure.

In general, $(M_0, \dots, M_k) \mapsto M_0 \otimes_{R_1} M_1 \otimes_{R_2} \cdots \otimes_{R_{k-1}} M_k$ defines a $(k+1)$ -ary additive functor (Definition C.0.13)

$${}_{R_0}\mathbf{Mod}_{R_1} \times \cdots \times {}_{R_{k-1}}\mathbf{Mod}_{R_k} \rightarrow {}_{R_0}\mathbf{Mod}_{R_k}$$

(??).

Given a ring R and a two-sided R -module M , we may also speak of the *n -fold tensor product* $M^{\otimes n} = M^{\otimes_R n}$

Definition C.0.13 (n-ary Additive Functor). Let I be a finite set with $|I| = n$. Let $\{\mathcal{A}_i\}_{i \in I}$ be additive categories and let $\mathcal{B}$ be an additive category. An *n -ary additive functor* (or *multilinear functor*)

$$F : \prod_{i \in I} \mathcal{A}_i \rightarrow \mathcal{B}$$

is a functor such that for each fixed collection of all but one variable, the resulting functor in the remaining variable is additive. Equivalently, for every $j \in I$ and objects $(A_i)_{i \in I}$ and

morphisms $f_1, f_2 : A_j \rightarrow A'_j$ in $\mathcal{A}_j$, we have

$$\begin{aligned} & F(A_1, \dots, A_{j-1}, f_1 + f_2, A_{j+1}, \dots, A_n) \\ &= F(A_1, \dots, A_{j-1}, f_1, A_{j+1}, \dots, A_n) \\ & \quad + F(A_1, \dots, A_{j-1}, f_2, A_{j+1}, \dots, A_n), \end{aligned}$$

and F preserves zero morphisms componentwise:

$$F(A_1, \dots, 0_{A_j, A'_j}, \dots, A_n) = 0_{F(A_1, \dots), F(A'_1, \dots)}.$$

A bifunctor that satisfies this property for $n = 2$ is simply called a **biadditive functor**.

Definition C.0.14. Let $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ be a morphism of schemes (Definition .3.2).

- The morphism f is called an **open immersion** or an **open embedding** if the underlying map of topological spaces induces a homeomorphism from X onto an open subset $V \subseteq Y$, and the induced map of sheaves (Definition 4.0.2) $f^\#|_V : \mathcal{O}_Y|_V \rightarrow f_*\mathcal{O}_X$ is an isomorphism of sheaves of rings on V .

(♠ TODO: surjective map of sheaves of sets)

- The morphism f is called a **closed immersion** or a **closed embedding** if the underlying map of topological spaces induces a homeomorphism from X onto a closed subset $Z \subseteq Y$, and the induced map of sheaves $f^\# : \mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$ is surjective.

Definition C.0.15. Let $\mathcal{C}$ be a category (Definition .1.1), let Z be an object, and let X, Y be objects of $\mathcal{C}$ over (Definition A.0.6) Z , i.e. morphisms $X \rightarrow Z$ and $Y \rightarrow Z$ are fixed. A **cartesian product of X and Y over Z in $\mathcal{C}$** (or **fiber product** or **pullback diagram**) is an object, often denoted by $X \times_Z Y$, with **projection morphisms** $X \times_Z Y \rightarrow X$ and $X \times_Z Y \rightarrow Y$ that are universal. More precisely, for any object T of $\mathcal{C}$ and morphisms $f_X : T \rightarrow X$, $f_Y : T \rightarrow Y$ such that the compositions $T \xrightarrow{f_X} X \rightarrow Z$ and $T \xrightarrow{f_Y} Y \rightarrow Z$ agree, there exists a unique morphism $u : T \rightarrow X \times_Z Y$ such that the following diagram commutes:

$$\begin{array}{ccccc} T & & & & \\ & \searrow^{f_X} & & \searrow & \\ & & X \times_Z Y & \longrightarrow & X \\ & \swarrow_{f_Y} & \downarrow & & \downarrow \\ & & Y & \longrightarrow & Z \end{array}$$

(Note: A dashed arrow labeled u points from T to $X \times_Z Y$.)

Equivalently, $X \times_Z Y$ is the limit (Definition .1.13) of the diagram (Definition .1.33)

$$\begin{array}{ccc} & X & \\ & \downarrow & \\ Y & \longrightarrow & Z \end{array}$$

in $\mathcal{C}$.

The commutative diagram

$$\begin{array}{ccc}
X \times_Z Y & \longrightarrow & X \\
\downarrow & & \downarrow \\
Y & \longrightarrow & Z
\end{array}$$

may be referred to as a **cartesian square**.

Definition C.0.16 (Locally Noetherian Scheme and Noetherian Scheme). Let X be a scheme (Definition .3.1).

- X is called **locally Noetherian** if it admits an open cover $\{U_i\}$ such that for each i , the ring $\mathcal{O}_X(U_i)$ of regular functions on U_i is a Noetherian ring. Equivalently, X is locally Noetherian if it is covered by open affine subschemes $\text{Spec } A_i$ with each A_i a Noetherian ring.
- X is called **Noetherian** if it is locally Noetherian and quasi-compact, i.e., X can be covered by finitely many affine opens $\text{Spec } A_i$ where each A_i is Noetherian.

Definition C.0.17 (Integral element over a ring). Let R be a commutative ring with unity (Definition C.0.1).

1. Let A be an R -algebra. An element $a \in A$ is called **integral over R** if there exists a monic polynomial

$$p(x) = x^n + r_{n-1}x^{n-1} + \cdots + r_1x + r_0$$

with coefficients $r_i \in R$ such that

$$p(a) = a^n + r_{n-1}a^{n-1} + \cdots + r_1a + r_0 = 0 \quad \text{in } A.$$

2. Let A be an extension ring of R . The ring extension A/R is called an **integral extension** if every element of A is integral over R .
3. Let A be an extension ring of R . The **integral closure of R in A** , sometimes denoted by $\tilde{A}$, is the subring

$$\tilde{A} = \{a \in A : a \text{ is integral over } R\}.$$

We say that R is integrally closed in A if $\tilde{A}$ coincides with A (considered as a subring of R).

4. Let R be an integral domain with field of fractions $K = \text{Frac}(R)$. We say that R is **integrally closed** if it is integrally closed as a subring of K .

Definition C.0.18. Let X be a locally Noetherian scheme (Definition C.0.16).

- The scheme X is **regular at a point $x \in X$** (or **nonsingular at a point $x \in X$**) if the local ring (Definition C.0.21) $\mathcal{O}_{X,x}$ (Definition C.0.23) is a regular local ring. Otherwise, X is said to be **singular at x** , or synonymously at x is a **singularity of X** .
- The scheme X is called a **regular scheme** (or **nonsingular scheme**) if it is regular at every point $x \in X$. Otherwise, X is said to be **singular**.

Definition C.0.19. Let $f : X \rightarrow Y$ be a morphism of schemes (Definition .3.2). (♠ **TODO: locally of finite type**) We say that f is a **finite type morphism** if the following equivalent conditions hold:

1. for every affine open $\text{Spec } B \subseteq Y$, the inverse image $f^{-1}(\text{Spec } B)$ is quasi-compact and has a covering $f^{-1}(\text{Spec } B) = \bigcup_j \text{Spec } A_j$ by affine opens such that each A_j is a **finitely generated B algebra**.
2. there exists an affine open covering $Y = \bigcup_i \text{Spec } B_i$ of Y such that the inverse image $f^{-1}(\text{Spec } B_i)$ has a finite cover $\{\text{Spec } A_{ij}\}_j$ by affine opens where each A_{ij} is a finitely generated B_i algebra.
3. f is quasi-compact and locally of finite type.

When X is equipped with a finite type morphism $f : X \rightarrow Y$, we say that X is a **finite type scheme over Y** or a **finite type Y -scheme** or a **Y -scheme of finite type** (Definition .3.3), etc.

Definition C.0.20 (Dimension of a Scheme). Let X be a scheme (Definition .3.1) with underlying topological space $|X|$.

- The **dimension at a point $x \in |X|$** , denoted $\dim_x(X)$, is the Krull dimension of the local ring (Definition C.0.21) $\mathcal{O}_{X,x}$ (Definition C.0.23), which is local by definition. This is the supremum of the lengths n of chains of prime ideals

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n \subseteq \mathcal{O}_{X,x}.$$

- The **dimension of the scheme X** is defined as

$$\dim(X) := \sup_{x \in |X|} \dim_x(X).$$

Equivalently, it is the supremum of the lengths of chains of distinct irreducible closed subsets of $|X|$ ordered by inclusion. It is also equivalent to the dimension of the underlying topological space $|X|$.

Definition C.0.21. Let R be a ring (Definition C.0.8) with unity, not necessarily commutative. The ring R is called a **local ring** if it has a unique maximal left ideal (Definition C.0.4). In this case, R also has a unique maximal right ideal, and these coincide with the Jacobson radical $J(R)$ of R .

The unique maximal left (and right) ideal of a local ring R may sometimes be denoted by $\mathfrak{m}_R$.

The following are each equivalent conditions for R to be a local ring:

1. $R/J(R)$ is a division ring
2. $R = J(R) \cup R^\times$

Definition C.0.22 (Presheaf on a topological space). Let X be a topological space (Definition C.0.6). Let $\mathcal{D}$ be a category.

A **presheaf (of objects of $\mathcal{D}$ /valued in $\mathcal{D}$) on X** is a rule $\mathcal{F}$ that assigns:

- to each open set $U \subseteq X$, an object $\mathcal{F}(U) \in \text{Ob } \mathcal{D}$, called the **sections of $\mathcal{F}$ over U** ,
- to each inclusion of open sets $V \subseteq U$, a morphism

$$\rho_V^U : \mathcal{F}(U) \rightarrow \mathcal{F}(V), \quad s \mapsto s|_V,$$

in the category $\mathcal{D}$ called the *restriction map* such that the following conditions hold:

- (Identity) For every open set $U \subseteq X$, the restriction map ρ_U^U is the identity on $\mathcal{F}(U)$.
- (Transitivity) For inclusions $W \subseteq V \subseteq U$ of open sets, one has

$$\rho_W^U = \rho_W^V \circ \rho_V^U.$$

For instance, we may speak of a *presheaf of sets/groups/rings/etc. on the topological space X* .

Equivalently, a presheaf on X (of objects in a category $\mathcal{D}$) is a functor

$$\mathbf{Open}(X)^{\mathrm{op}} \rightarrow \mathcal{D}$$

from the opposite of the category $\mathbf{Open}(X)$ of open subsets of X (see also Definition 4.0.1)

Equivalently, a presheaf on X is a presheaf on the category $\mathbf{Open}(X)$ in the sense of Definition 4.0.1 .

The sections object $\mathcal{F}(U)$ is also denoted by $\Gamma(U, \mathcal{F})$ (see Definition 7.1.1) . Moreover, the object $\mathcal{F}(X) = \Gamma(X, \mathcal{F})$ is called the *global sections object of $\mathcal{F}$* . This agrees with the notion of global sections as discussed in Definition 7.1.1.

Definition C.0.23 (Stalk of a sheaf). Let X be a topological space, and let $\mathcal{D}$ be a category (Definition .1.1) Let $\mathcal{F}$ be a presheaf on X valued in $\mathcal{D}$ (Definition C.0.22). For a point $x \in X$, the *stalk of $\mathcal{F}$ at x* , denoted $\mathcal{F}_x$, is defined as the filtered direct limit (Definition .1.14)

$$\mathcal{F}_x := \varinjlim_{x \in U} \mathcal{F}(U),$$

where the limit ranges over all open neighborhoods U of x in X ordered by inclusion, assuming that such a direct limit exists.

If $\mathcal{D}$ is some kind of category of sets (e.g. $\mathcal{D} = \mathbf{Sets}, \mathbf{Grps}, \mathbf{Rings}$), then an element of $\mathcal{F}_x$ is called a *germ of a section at x* . Concretely, a germ at x is given by a pair (U, s) with U an open neighborhood of x and $s \in \mathcal{F}(U)$, modulo the equivalence relation: $(U, s) \sim (V, t)$ if there exists an open neighborhood $W \subseteq U \cap V$ of x such that $s|_W = t|_W$, see also ??.

If $\mathcal{F}$ is a presheaf of groups, rings, or modules, then each stalk $\mathcal{F}_x$ inherits some corresponding algebraic structure. More generally, if $\mathcal{D}$ is so that $\mathcal{F}_x$ exists for every presheaf $\mathcal{F} \in \mathbf{PreShv}(X, \mathcal{D})$ (resp. every sheaf $\mathcal{F} \in \mathbf{Sh}(X, \mathcal{D})$), then $\mathcal{F} \mapsto \mathcal{F}_x$ is a functor $\mathbf{PreShv}(X, \mathcal{D}) \rightarrow \mathcal{D}$ (resp. $\mathbf{Sh}(X, \mathcal{D}) \rightarrow \mathcal{D}$).