

TOPOLOGY

March 27, 2026

CONTENTS

1. Basic topolgy	2
1.1. Topological spaces	2
1.1.1. Generating topologies from subsets	3
1.1.2. Basis of a topological space	4
1.1.3. Separation axioms	4
1.1.4. Subspaces of a topological space	5
1.2. Properties of topological spaces	6
1.2.1. Compact topological spaces	6
1.3. Continuous maps of topological spaces	7
1.4. Constructions of topological spaces from other topological spaces	8
1.5. Concrete examples of topological spaces	9
1.6. Compactly generated	11
1.7. Metrics, absolute values, and norms	11
1.7.1. Extended metric spaces	12
1.8. Convergence in extended metric spaces	13
1.8.1. Absolute values on fields	14
1.8.2. Norms on vector spaces over fields with absolute values	15
1.8.3. Inner product on a vector space over the reals or complex numbers	15
1.9. Euclidean space	16
1.10. Connectivity of topological spaces	16
1.11. Stone spaces, ultrafilters, and Boolean algebras	17
2. Covering spaces	19

3. Homotopy	21
4. Singular homology and cohomology	26
5. CW complexes	29
6. Fibrations	29
6.1. Hurewicz fibrations	29
6.2. Serre fibrations	30
6.3. Fiber bundles	30
Appendix A. Set theory	31
Appendix B. Category Theory	32
Appendix C. Group Theory	33
Appendix D. Ring Theory	34
Appendix E. Homological algebra	36
References	37

The purpose of this document is to compile common notions in basic topology.

1. BASIC TOPOLOGY

1.1. Topological spaces.

Definition 1.1.1 (Topology). Let X be a set. A **topology on X** is a collection $\mathcal{T}$ of subsets of X such that:

1. $\emptyset \in \mathcal{T}$ and $X \in \mathcal{T}$,
2. For any collection $\{U_i\}_{i \in I} \subseteq \mathcal{T}$ (with I arbitrary), the union $\bigcup_{i \in I} U_i \in \mathcal{T}$,
3. For any finite collection $\{U_1, \dots, U_n\} \subseteq \mathcal{T}$, the intersection $U_1 \cap \dots \cap U_n \in \mathcal{T}$.

If $\mathcal{T}$ is a topology on X , the pair $(X, \mathcal{T})$ is called a **topological space**. Members of $\mathcal{T}$ are called **open sets**.

A subset $C \subseteq X$ is **closed** (also called a **closed subset**) if its complement $X \setminus C$ is an open set in $\mathcal{T}$.

One very often refers to X as a topological space, omitting the notation of the topology $\mathcal{T}$.

The collection of all topologies on a set X may be denoted by notations such as **Top(X)**, **Top(X)**, or **Top(X)**.

Definition 1.1.2. Let X be a topological space. A subset $U \subseteq X$ is a **clopen subset** if U is both an open set and a closed set in the topology of X .

1.1.1. Generating topologies from subsets.

Definition 1.1.3. Let X be a set.

1. Let τ_1, τ_2 be topologies on (Definition 1.1.1) X . Say that τ_1 is **coarser than** (equivalently, **smaller than**) τ_2 if $\tau_1 \subseteq \tau_2$, and that τ_1 is **finer than** (equivalently, **larger than**) τ_2 if $\tau_2 \subseteq \tau_1$. This means that every subset of X that is open (Definition 1.1.1) with respect to τ_1 is also open with respect to τ_2 .

These relations are denoted by $\tau_1 \preceq \tau_2$ for “ τ_1 coarser than τ_2 ” and $\tau_1 \succeq \tau_2$ for “ τ_1 finer than τ_2 ”; their strict versions are $\tau_1 \prec \tau_2$ and $\tau_1 \succ \tau_2$, meaning proper inclusion.

2. Let $\mathcal{C}$ be some family of topologies on X . A topology $\tau \in \mathcal{C}$ is the **coarsest** (or **smallest**) element of $\mathcal{C}$ if for every $\sigma \in \mathcal{C}$ one has $\tau \subseteq \sigma$ (equivalently, $\tau \preceq \sigma$ for all $\sigma \in \mathcal{C}$). Dually, $\tau \in \mathcal{C}$ is the **finest** (or **largest**) element of $\mathcal{C}$ if for every $\sigma \in \mathcal{C}$ one has $\sigma \subseteq \tau$ (equivalently, $\sigma \preceq \tau$ for all $\sigma \in \mathcal{C}$).

Proposition 1.1.4. Let X be a set and let $f : (X, \tau_X) \rightarrow (Y, \tau_Y)$ be a function between topological spaces (Definition 1.1.1).

- If τ'_X is a topology (Definition 1.1.1) on X finer (Definition 1.1.3) than τ_X and f is continuous (Definition 1.3.1), then $f : (X, \tau'_X) \rightarrow (Y, \tau_Y)$ is also continuous.
- If τ'_Y is a topology on Y coarser than τ_Y and f is continuous, then $f : (X, \tau_X) \rightarrow (Y, \tau'_Y)$ is also continuous.

Proposition 1.1.5. Let X be a set and $\mathcal{S}$ a collection of subsets of X . The topology (Definition 1.1.1) $\tau(\mathcal{S})$ generated by (Definition 1.1.6) $\mathcal{S}$ is the coarsest topology (Definition 1.1.3) on X such that every element of $\mathcal{S}$ is an open (Definition 1.1.1) set.

Definition 1.1.6. Let X be a set and $\mathcal{S}$ a collection of subsets of X . The **topology on X generated by $\mathcal{S}$** , often denoted $\tau(\mathcal{S})$, is the intersection of all topologies on X that contain $\mathcal{S}$. It is the unique coarsest topology (Definition 1.1.3) on X in which every element of $\mathcal{S}$ is an open set.

The collection $\mathcal{S}$ is called a **subbasis for the topology $\tau(\mathcal{S})$** . Alternatively, one might generally call a family $\mathcal{S} \subseteq \mathcal{P}(X)$ a **subbasis on X** if $\bigcup_{S \in \mathcal{S}} S = X$ and we would consider $\mathcal{S}$ the subbasis for the topology $\tau(\mathcal{S})$.

Proposition 1.1.7. Let X be a set and $\mathcal{S}$ a subbasis (Definition 1.1.6) for a topology (Definition 1.1.1) τ . A subset $U \subseteq X$ belongs to τ if and only if U is the union of finite intersections of elements of $\mathcal{S}$. That is, the collection of all finite intersections of elements of $\mathcal{S}$ forms a basis (Definition 1.1.10) for τ .

Proposition 1.1.8. Let X and Y be topological spaces (Definition 1.1.1), and let $\mathcal{S}$ be a subbasis (Definition 1.1.6) for the topology (Definition 1.1.1) on Y . A function $f : X \rightarrow Y$ is continuous (Definition 1.3.1) if and only if $f^{-1}(V)$ is an open (Definition 1.1.1) set in X for every $V \in \mathcal{S}$.

1.1.2. Basis of a topological space.

Definition 1.1.9. Let $(X, \mathcal{T})$ be a topological space (Definition 1.1.1) and let $x \in X$.

- An *open neighborhood of x* is any open set $U \in \mathcal{T}$ such that $x \in U$.
- A *neighborhood of x* is a set $N \subseteq X$ for which there exists an open neighborhood $U \in \mathcal{T}$ of x such that $U \subseteq N$.
- A *neighborhood basis* (or *local base*) at x is a nonempty collection $\mathcal{B}_x$ of neighborhoods of x such that for every neighborhood N of x , there exists $B \in \mathcal{B}_x$ with $B \subseteq N$.
The elements of $\mathcal{B}_x$ are said to *form a base of neighborhoods* at x .

Definition 1.1.10. Let X be a set and let $\mathcal{B}$ be a collection of subsets of X . The collection $\mathcal{B}$ is called a *basis* (or *base*) for a topology (Definition 1.1.1) on X if the following two conditions hold:

1. For every $x \in X$, there exists at least one $B \in \mathcal{B}$ such that $x \in B$.
2. For any $B_1, B_2 \in \mathcal{B}$ and any $x \in B_1 \cap B_2$, there exists $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2$.

Given such a collection $\mathcal{B}$, the collection $\mathcal{T}$ of all unions of elements of $\mathcal{B}$ defines a topology on X , and it coincides with $\mathcal{T}_{\mathcal{B}}$, the topology generated by $\mathcal{B}$ (Definition 1.1.6). In other words,

$$\mathcal{T}_{\mathcal{B}} = \{U \subseteq X : \text{for every } x \in U, \text{ there exists } B \in \mathcal{B} \text{ with } x \in B \subseteq U\}.$$

Definition 1.1.11. Let $(X, \mathcal{T})$ be a topological space (Definition 1.1.1).

- The space $(X, \mathcal{T})$ is said to be *first countable* if every point $x \in X$ has a countable (Definition A.0.1) neighborhood basis (Definition 1.1.9), i.e., there exists a countable collection $\{U_n\}_{n \in \mathbb{N}}$ of open neighborhoods of x such that for any open neighborhood U of x , there exists $n \in \mathbb{N}$ with $U_n \subseteq U$.
- The space $(X, \mathcal{T})$ is said to be *second countable* if there exists a countable basis (Definition 1.1.10) $\mathcal{B} \subseteq \mathcal{T}$ for the topology $\mathcal{T}$, i.e., every open set $U \in \mathcal{T}$ can be written as a union of elements of $\mathcal{B}$.

1.1.3. Separation axioms.

Definition 1.1.12. Let $(X, \mathcal{T})$ be a topological space (Definition 1.1.1) and let $A \subseteq X$.

- The set A is called a *G_δ set* if there exists a countable collection of open sets $\{U_n\}_{n \in \mathbb{N}} \subseteq \mathcal{T}$ such that

$$A = \bigcap_{n \in \mathbb{N}} U_n.$$

- The set A is called an *F_σ set* if there exists a countable collection of closed sets $\{F_n\}_{n \in \mathbb{N}}$ such that

$$A = \bigcup_{n \in \mathbb{N}} F_n.$$

Definition 1.1.13 (Separation axioms). Let $(X, \mathcal{T})$ be a topological space (Definition 1.1.1).

- $(X, \mathcal{T})$ is T_0 (*Kolmogorov*) if for every pair of distinct points $x, y \in X$, there exists an open set $U \in \mathcal{T}$ such that, without loss of generality, $x \in U$ and $y \notin U$.
- $(X, \mathcal{T})$ is T_1 (*Fréchet*) if for every pair of distinct points $x, y \in X$, there exist open sets $U, V \in \mathcal{T}$ such that $x \in U$, $y \notin U$, and $y \in V$, $x \notin V$.
- $(X, \mathcal{T})$ is T_2 or *Hausdorff* if for every pair of distinct points $x, y \in X$, there exist disjoint open sets $U, V \in \mathcal{T}$ such that $x \in U$ and $y \in V$.
- $(X, \mathcal{T})$ is *regular* if it is T_1 and for each point $x \in X$ and closed set $F \subseteq X$ with $x \notin F$, there exist disjoint open sets $U, V \in \mathcal{T}$ such that $x \in U$ and $F \subseteq V$.
- $(X, \mathcal{T})$ is T_3 (regular Hausdorff) if it is T_1 and regular.
- $(X, \mathcal{T})$ is *completely regular* if for each closed set $F \subseteq X$ and $x \notin F$, there exists a continuous function $f : X \rightarrow [0, 1]$ such that $f(x) = 0$ and $f|_F = 1$.
- $(X, \mathcal{T})$ is $T_{3\frac{1}{2}}$ (completely regular Hausdorff) if it is T_1 and completely regular.
- $(X, \mathcal{T})$ is *normal* if it is T_1 and for each pair of disjoint closed sets $A, B \subseteq X$, there exist disjoint open sets $U, V \in \mathcal{T}$ such that $A \subseteq U$ and $B \subseteq V$.
- $(X, \mathcal{T})$ is T_4 (normal Hausdorff) if it is T_1 and normal.
- $(X, \mathcal{T})$ is T_5 (completely normal Hausdorff) if it is T_1 and completely normal.
- $(X, \mathcal{T})$ is *perfectly normal* if every closed set is a G_δ (Definition 1.1.12) (countable (Definition A.0.1) intersection of open sets) and the space is normal.
- $(X, \mathcal{T})$ is T_6 (perfectly normal Hausdorff) if it is T_1 and perfectly normal.

1.1.4. Subspaces of a topological space.

Definition 1.1.14 (Subspace topology and subspace). Let $(X, \mathcal{T})$ be a topological space and $Y \subseteq X$ a subset. The topology induced on Y by $\mathcal{T}$, called the *subspace topology*, is defined as

$$\mathcal{T}_Y := \{U \cap Y : U \in \mathcal{T}\}.$$

The pair $(Y, \mathcal{T}_Y)$ is called a *subspace of $(X, \mathcal{T})$* .

Convention 1.1.15. Given a topological space $(X, \mathcal{T})$ and a subset S of X , one almost always considers S as a subspace of X , i.e. considers S as a topological space equipped with the subspace topology.

Definition 1.1.16 (Discrete topological space). Let X be a set. The *discrete topology on X* is the topology $\mathcal{T}_{\text{disc}}$ defined by

$$\mathcal{T}_{\text{disc}} := \{U : U \subseteq X\},$$

i.e., every subset of X is open. The pair $(X, \mathcal{T}_{\text{disc}})$ is called a *discrete topological space*.

Definition 1.1.17 (Indiscrete (or trivial) topological space). Let X be a set. The *indiscrete topology* (also called the *trivial topology*) on X is the topology $\mathcal{T}_{\text{indisc}}$, also denoted by notations such as $\mathcal{T}_{\text{triv}}$, defined by

$$\mathcal{T}_{\text{indisc}} := \{\emptyset, X\},$$

i.e., only the empty set and the entire set are open. The pair $(X, \mathcal{T}_{\text{indisc}})$ is called an *indiscrete topological space*.

Definition 1.1.18 (Closure of a subset). Let $(X, \mathcal{T})$ be a topological space and $A \subseteq X$ a subset. The *closure of A in X* , denoted by $\overline{A}$, is defined as the intersection of all closed sets containing A , i.e.,

$$\overline{A} := \bigcap \{C \subseteq X : C \text{ is closed and } A \subseteq C\}.$$

Equivalently, $\overline{A}$ is the smallest closed set containing A .

1.2. Properties of topological spaces.

Definition 1.2.1 (Dense subset). Let $(X, \mathcal{T})$ be a topological space. A subset $D \subseteq X$ is called *dense in X* if the closure of D equals X , i.e.,

$$\overline{D} = X.$$

Equivalently, D is dense in X if every non-empty open set $U \in \mathcal{T}$ intersects D , that is,

$$\forall U \in \mathcal{T}, U \neq \emptyset \implies U \cap D \neq \emptyset.$$

Definition 1.2.2. A topological space X is *disconnected* if there exist two disjoint, non-empty open sets (Definition 1.1.1) U and V such that $X = U \cup V$. The space X is *connected* if it is not disconnected.

Definition 1.2.3. Let X be a topological space and x a point in X . The *connected component of x* is the union of all connected (Definition 1.2.2) subspaces (Definition 1.1.14) of X that contain x .

Definition 1.2.4. Let X be a topological space. X is *totally disconnected* if the connected component (Definition 1.2.3) of every point $x \in X$ consists only of the singleton set $\{x\}$.

Proposition 1.2.5. Let X be a compact (Definition 1.2.6) Hausdorff space (Definition 1.1.13). X is totally disconnected (Definition 1.2.4) if and only if the set of clopen subsets (Definition 1.1.2) of X forms a basis for the topology (Definition 1.1.10) of X .

1.2.1. Compact topological spaces.

Definition 1.2.6 (Compact topological space). A topological space $(X, \mathcal{T})$ is *compact* if every open cover of X admits a finite subcover; that is, for every collection $\{U_i\}_{i \in I}$ of open sets in $\mathcal{T}$ such that $X = \bigcup_{i \in I} U_i$, there exists a finite subcollection $\{U_{i_j}\}_{j=1}^n$ such that $X = \bigcup_{j=1}^n U_{i_j}$.

Some mathematicians, e.g. algebraic geometers, would refer to this property as *quasi-compactness*.

Here is an alternate, essentially equivalent definition, of compactness:

Definition 1.2.7 (Compactness). Let $(X, \mathcal{T})$ be a topological space (Definition 1.1.1). A subset $K \subseteq X$ is *compact* if for every collection $\{U_i\}_{i \in I}$ of open sets such that $K \subseteq \bigcup_{i \in I} U_i$, there exists a finite subcollection $\{U_{i_j}\}_{j=1}^n$ with $K \subseteq \bigcup_{j=1}^n U_{i_j}$.

Proposition 1.2.8. Let $(X, \mathcal{T})$ be a topological space (Definition 1.1.1). A subspace (Definition 1.1.14) $K \subseteq X$ is compact in the sense of Definition 1.2.7 if and only if the space K is compact in the sense of Definition 1.2.6.

Definition 1.2.9 (Locally compact). Let $(X, \mathcal{T})$ be a topological space (Definition 1.1.1). X is *locally compact* if for every $x \in X$, there exists an open set $U \in \mathcal{T}$ containing x and a compact set (Definition 1.2.6) $K \subseteq X$ such that $x \in U \subseteq K$.

Definition 1.2.10. Let $(X, \mathcal{T})$ be a topological space and let $A \subseteq X$.

- The *interior of A* , denoted by $\text{int}(A)$, is the union of all open sets contained in A :

$$\text{int}(A) = \bigcup \{U \in \mathcal{T} : U \subseteq A\}.$$

- The *boundary of A* , denoted by ∂A , is defined as

$$\partial A = \overline{A} \setminus \text{int}(A).$$

(Definition 1.1.18) Equivalently, a point $x \in X$ belongs to ∂A if every open neighborhood of x intersects both A and its complement $X \setminus A$.

1.3. Continuous maps of topological spaces.

Definition 1.3.1. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces (Definition 1.1.1). A map $f : X \rightarrow Y$ is called *continuous* if for every open set $V \in \mathcal{T}_Y$, the preimage $f^{-1}(V)$ is an open set in X , that is,

$$\forall V \in \mathcal{T}_Y, f^{-1}(V) \in \mathcal{T}_X.$$

Equivalently, f is continuous if and only if for every closed set $C \subseteq Y$, the preimage $f^{-1}(C)$ is closed in X .

A *map of topological spaces* usually refers to a continuous map between the topological spaces.

The set of continuous maps from X to Y is sometimes denoted by $\mathcal{C}(X, Y)$. Other standard notation include $\text{Hom}_{\text{Top}}(X, Y)$ or $\text{Top}(X, Y)$ coming from more general notation for morphisms between objects in a category (Definition B.0.1).

Definition 1.3.2. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces (Definition 1.1.1). A function $f : X \rightarrow Y$ is called a *homeomorphism* if it satisfies all of the following:

1. f is bijective (Definition A.0.2);
2. f is continuous (Definition 1.3.1) with respect to $\mathcal{T}_X$ and $\mathcal{T}_Y$; and
3. the inverse map $f^{-1} : Y \rightarrow X$ (Definition A.0.2) is also continuous.

(♠ **TODO: equivalence class**) If such a function exists, the spaces X and Y are said to be *homeomorphic*, or that X and Y have the *same topological type* — accordingly, the *topological type of a topological spaces* refers to its equivalence under the equivalence relation of homeomorphic spaces.

Definition 1.3.3 (Pointed topological space). Let X be a topological space (Definition 1.1.1) and let $x_0 \in X$ be a chosen element of X . A *pointed/based (topological) space* is a pair (X, x_0) consisting of the space X together with the distinguished point x_0 , called the *base point of X* . If the base point of a pointed space (X, x_0) is understood, then it may be suppressed from notation; in particular, X may be written as a pointed space as opposed to the full notation of (X, x_0) .

A **morphism of pointed spaces** (or **based map**) or **continuous map** between pointed spaces (X, x_0) and (Y, y_0) is a continuous map (Definition 1.3.1)

$$f : X \rightarrow Y$$

such that $f(x_0) = y_0$.

The collection of pointed spaces with their morphisms form a locally small (Definition B.0.2) category (Definition B.0.1), often called the **category of pointed spaces**. This category is often denoted by notations such as $\mathbf{Top}_*$, $\mathbf{Top}_\bullet$, $\mathbf{Top}_*$, $\mathbf{Top}_\bullet$, etc. The set of continuous maps from pointed spaces X to Y may be denoted by notations such as $C_*(X, Y)$, $C_\bullet(X, Y)$, $\mathbf{Top}_*(X, Y)$, $\mathbf{Top}_\bullet(X, Y)$, $\mathbf{Hom}_{\mathbf{Top}_\bullet}(X, Y)$, etc.

Definition 1.3.4 (Preimage of a subset under a map of sets). Let X and Y be sets and let $f : X \rightarrow Y$ be a function. Let $B \subseteq Y$ be a subset of the codomain Y . The **preimage of B under f** (also called the **inverse image of B under f**) is the subset of X defined by

$$f^{-1}(B) := \{x \in X : f(x) \in B\} \subseteq X.$$

If B is a singleton set $\{y\}$, then we often denote $f^{-1}(B)$ by $f^{-1}(y)$.

Definition 1.3.5 (Fiber of a map of topological spaces). Let X and Y be topological spaces (Definition 1.1.1) and let $f : X \rightarrow Y$ be a continuous map (Definition 1.3.1). For a point $y \in Y$, the **fiber of f over y** is the inverse image (Definition 1.3.4) $f^{-1}(y) = f^{-1}(\{y\})$ endowed with the subspace topology (Definition 1.1.14) induced from X . The fiber is also denoted by notations such as $\mathbf{Fib}_f(y)$ or X_y .

1.4. Constructions of topological spaces from other topological spaces.

Definition 1.4.1 (Product of Sets). Let I be a (possibly infinite (Definition A.0.1) but small) index set and let $\{A_i\}_{i \in I}$ be a family of sets indexed by I . The **Cartesian product of the family $\{A_i\}_{i \in I}$** , denoted by $\prod_{i \in I} A_i$, is defined as the set of all tuples/functions

$$\prod_{i \in I} A_i := \{(a_i)_{i \in I} \mid a_i \in A_i \text{ for all } i \in I\},$$

where $(a_i)_{i \in I}$ denotes a function from I to $\bigcup_{i \in I} A_i$ such that $(a_i)_{i \in I}(i) = a_i \in A_i$ for each $i \in I$.

The self product of a set A indexed by I is often denoted by A^I . Note that elements of A^I can be identified with functions $I \rightarrow A$. The finite self product of A taken n times is often denoted by A^n . For finitely many sets $A_1, \dots, A_n$, their Cartesian product is denoted by $A_1 \times \dots \times A_n$. Elements of such a finite product may be written as $(a_1, \dots, a_n)$.

Definition 1.4.2. Let I be an index set and $\{X_i\}_{i \in I}$ be a collection of topological spaces. Let $X = \prod_{i \in I} X_i$ be the set theoretic Cartesian product (Definition 1.4.1) of these spaces.

The **box topology on X** is the topology generated by (Definition 1.1.10) the base (Definition 1.1.10) consisting of all sets of the form $\prod_{i \in I} U_i$, where each U_i is an open subset (Definition 1.1.1) of X_i .

Definition 1.4.3. Let I be an index set and $\{X_i\}_{i \in I}$ be a collection of topological spaces. Let $X = \prod_{i \in I} X_i$ be the set theoretic Cartesian product (Definition 1.4.1) of these spaces.

The **product topology** (or **Tychonoff topology**) on X is the topology generated by (Definition 1.1.10) the base (Definition 1.1.10) consisting of all sets of the form $\prod_{i \in I} U_i$, where each U_i is an open subset (Definition 1.1.1) of X_i and $U_i = X_i$ for all but finitely many indices $i \in I$.

The **product** or the **product space of the X_i** refers to X equipped with the product topology.

For each $j \in I$, let $p_j : \prod_{i \in I} X_i \rightarrow X_j$ denote the canonical projection map. The product topology is the coarsest topology (Definition 1.1.3) on X such that all projection maps p_j are continuous (Definition 1.3.1).

Proposition 1.4.4. Let I be an index set and $\{X_i\}_{i \in I}$ be a collection of topological spaces. Let $X = \prod_{i \in I} X_i$ be the set theoretic Cartesian product (Definition 1.4.1) of these spaces.

If $X = \prod_{i \in I} X_i$ is endowed with the product topology (Definition 1.4.3), then a function $f : Z \rightarrow X$ from a topological space (Definition 1.1.1) Z is continuous (Definition 1.3.1) if and only if each component function $f_i = p_i \circ f : Z \rightarrow X_i$ is continuous.

This universal property generally fails for the box topology (Definition 1.4.2) when I is infinite (Definition A.0.1).

Theorem 1.4.5 (Tychonoff's theorem). Let I be an index set and let $\{X_i\}_{i \in I}$ be a collection of compact spaces (Definition 1.2.6).

The product space (Definition 1.4.2) $X = \prod_{i \in I} X_i$ is compact in the product topology (Definition 1.1.1).

1.5. Concrete examples of topological spaces.

Definition 1.5.1. For a positive integer n , let $\mathbb{R}^n$ denote the n -fold Cartesian product of the real line $\mathbb{R}$ with itself:

$$\mathbb{R}^n = \{(x_1, x_2, \dots, x_n) : x_i \in \mathbb{R} \text{ for all } i = 1, \dots, n\}.$$

The set $\mathbb{R}^n$ is called **Euclidean n-space**. A point $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ is associated with the **Euclidean norm**

$$\|x\| = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}.$$

The corresponding metric (Definition 1.7.1) $d : \mathbb{R}^n \times \mathbb{R}^n \rightarrow [0, \infty)$ is given by

$$d(x, y) = \|x - y\|.$$

This metric induces (Definition 1.7.4) the standard topology on $\mathbb{R}^n$, called the **Euclidean topology**.

The Euclidean topology on $\mathbb{R}^n$ coincides with the product topology (Definition 1.4.3) of $\prod_{i=1}^n \mathbb{R}$ where each copy of $\mathbb{R}$ is equipped with its Euclidean topology.

Definition 1.5.2 (Interval in the real line). Let $(\mathbb{R}, \leq)$ be the field of real numbers equipped with its usual total order. A subset $I \subseteq \mathbb{R}$ is called an *interval in the real line* if for all $x, y, z \in \mathbb{R}$ one has

$$x < z < y \text{ and } x \in I, y \in I \implies z \in I.$$

Equivalently, I is an interval if whenever $x, y \in I$ with $x < y$, then every $z \in \mathbb{R}$ satisfying $x < z < y$ also lies in I . By convention, the empty set $\emptyset$ and singletons $\{x\}$ for $x \in \mathbb{R}$ are also considered intervals, since they satisfy this property.

Let $a, b \in \mathbb{R}$ with $a \leq b$. We use the following standard notations for intervals in $\mathbb{R}$:

$$\begin{aligned} (a, b) &= \{x \in \mathbb{R} : a < x < b\}, \\ [a, b] &= \{x \in \mathbb{R} : a \leq x \leq b\}, \\ (a, b] &= \{x \in \mathbb{R} : a < x \leq b\}, \\ [a, b) &= \{x \in \mathbb{R} : a \leq x < b\}, \\ (a, \infty) &= \{x \in \mathbb{R} : a < x\}, \\ [a, \infty) &= \{x \in \mathbb{R} : a \leq x\}, \\ (-\infty, b) &= \{x \in \mathbb{R} : x < b\}, \\ (-\infty, b] &= \{x \in \mathbb{R} : x \leq b\}. \end{aligned}$$

We also regard $\mathbb{R}$ itself and the empty set $\emptyset$ as intervals when convenient.

An interval of the form (a, b) for $a \in \mathbb{R} \cup \{\text{infy}\}$ and $b \in \mathbb{R} \cup \{\infty\}$ is called an *open interval*. An interval of the form $[a, b]$ for $a \in \mathbb{R}$ and $b \in \mathbb{R}$ is called a *closed interval*. An interval of the form $(a, b]$ for $a \in \mathbb{R} \cup \{\text{infy}\}$ and $b \in \mathbb{R}$ or of the form $[a, b)$ for $a \in \mathbb{R}$ and $b \in \mathbb{R} \cup \{\infty\}$ is called a *half open interval*.

Definition 1.5.3. Let $n \geq 0$ be an integer. The *n-dimensional (unit) disk (or (unit) ball)*, often denoted by D^n (or B^n), is the subspace (Definition 1.1.14) of $\mathbb{R}^n$ (Definition 1.9.1) defined by:

$$D^n = \{x \in \mathbb{R}^n \mid \|x\| \leq 1\}$$

For $n = 1$, the disk D^1 is the closed interval (Definition 1.5.2) $I = [-1, 1]$, which is homeomorphically (Definition 1.3.2) to the standard unit interval $[0, 1]$.

One also often means by an *n-dimensional disk* any space homeomorphic to D^n .

Definition 1.5.4. Let $n \geq 0$ be an integer. The *n-dimensional (unit) sphere*, denoted by S^n , is the subspace of $\mathbb{R}^{n+1}$ defined by:

$$S^n = \{x \in \mathbb{R}^{n+1} \mid \|x\| = 1\}$$

The sphere S^n is the boundary (Definition 1.2.10) of the $(n + 1)$ -dimensional disk D^{n+1} (Definition 1.5.3). Specifically, S^0 consists of the two discrete points $\{-1, 1\}$.

S^1 is called the *unit circle*.

One also often means by an *n-dimensional sphere* any space homeomorphic to S^n .

Definition 1.5.5. The *standard unit interval*, denoted by I , is the subspace (Definition 1.1.14) of $\mathbb{R}$ (Definition 1.9.1) defined by:

$$I = [0, 1] = \{x \in \mathbb{R} \mid 0 \leq x \leq 1\}$$

(Definition 1.5.2)

Definition 1.5.6. The *n-dimensional unit cube* is the n -fold Cartesian product (Definition 1.4.1) $I^n = I \times \cdots \times I$, equipped with the product topology (Definition 1.4.3) where I is the standard unit interval (Definition 1.5.5) equipped with its usual topology. This topology on I^n coincides with the subspace topology (Definition 1.1.14) of I^n as a subset of $\mathbb{R}^n$ (Definition 1.9.1).

Definition 1.5.7. Let $n \geq 1$ be an integer. The *n-dimensional torus*, denoted by T^n , is the topological space (Definition 1.1.1) defined as the n -fold Cartesian product (Definition 1.4.1) of the circle (Definition 1.5.4):

$$T^n = \underbrace{S^1 \times S^1 \times \cdots \times S^1}_{n \text{ times}}$$

(♠ TODO: quotient space by a group action) Equivalently, T^n is the quotient space $\mathbb{R}^n / \mathbb{Z}^n$ under the action of translation by the integer lattice.

Definition 1.5.8. Let $n \geq 0$ be an integer and let $\mathbb{K}$ be the field of real numbers $\mathbb{R}$ or complex numbers $\mathbb{C}$. The *n-dimensional projective space over $\mathbb{K}$* , denoted by $\mathbb{K}P^n$, is the set of 1-dimensional vector subspaces of $\mathbb{K}^{n+1}$, equipped with the quotient topology induced by the map:

$$q : \mathbb{K}^{n+1} \setminus \{0\} \rightarrow \mathbb{K}P^n$$

which sends a non-zero vector to the line it spans. Explicitly, for $\mathbb{K} = \mathbb{R}$, this is the real projective space $\mathbb{R}P^n$; for $\mathbb{K} = \mathbb{C}$, this is the complex projective space $\mathbb{C}P^n$.

1.6. Compactly generated.

Definition 1.6.1. (♠ TODO: final topology) A topological space (Definition 1.1.1) X is said to be *compactly generated* (or a *k-space*) if a subset $U \subseteq X$ is open whenever for every compact (Definition 1.2.6) subset $K \subseteq X$, the intersection $U \cap K$ is open in the subspace K . Equivalently, X is compactly generated if and only if the topology of X is the final topology with respect to the collection of inclusions $K \hookrightarrow X$ for compact $K \subseteq X$.

Definition 1.6.2. A topological space (Definition 1.1.1) X is said to be *weakly Hausdorff* if for every continuous map (Definition 1.3.1) $f : K \rightarrow X$ from a compact (Definition 1.2.6) Hausdorff (Definition 1.2.7) space K , the image $\text{Im}(f)$ is closed in X . Equivalently, X is weakly Hausdorff if and only if for every compact Hausdorff K , the induced map $f : K \rightarrow X$ is a closed map.

1.7. Metrics, absolute values, and norms.

1.7.1. Extended metric spaces.

Definition 1.7.1 (Extended Metric). Let M be a set. An *extended metric* on M is a function

$$d : M \times M \rightarrow [0, \infty]$$

such that for all $x, y, z \in M$:

1. **Non-negativity:** $d(x, y) \geq 0$, and $d(x, y) = 0$ if and only if $x = y$.
2. **Symmetry:** $d(x, y) = d(y, x)$.
3. **Triangle inequality:** $d(x, z) \leq d(x, y) + d(y, z)$,

again adopting the convention that sums involving ∞ behave so that $a + \infty = \infty$. A set equipped with an extended metric is called an *extended metric space*.

If the range of the extended metric is contained in $[0, \infty)$, then the extended metric is a *metric* in the usual sense and V may be called a *metric space*.

Definition 1.7.2 (Isometry for extended metric spaces). Let (X, d_X) and (Y, d_Y) be extended metric spaces (Definition 1.7.1). A map $f : X \rightarrow Y$ is called an *isometry* if for all $x_1, x_2 \in X$,

$$d_Y(f(x_1), f(x_2)) = d_X(x_1, x_2).$$

An isometry is automatically injective. If, in addition, f is surjective, it is called an *isometric isomorphism*, and X and Y are said to be *isometric*.

Definition 1.7.3 (Extended Metric Induced by an Extended Norm). Let V be a vector space over a field F equipped with an absolute value (Definition 1.8.6)

$$|\cdot| : F \rightarrow [0, \infty),$$

and let $\|\cdot\| : V \rightarrow [0, \infty]$ be an extended norm on V (Definition 1.8.7). Then the *extended metric induced by the extended norm* is the function

$$d : V \times V \rightarrow [0, \infty]$$

defined by

$$d(x, y) := \|x - y\|.$$

It is indeed an extended metric. If $\|\cdot\|$ is a norm, then d is a metric.

Definition 1.7.4 (Topology induced by an extended metric). Let (X, d) be an extended metric space (Definition 1.7.1).

1. Given $\varepsilon \in (0, \infty)$ and $x \in X$, the set

$$B(x, \varepsilon) := \{y \in X : d(x, y) < \varepsilon\}$$

is called the *open ball of radius ε centered at x* .

2. The *topology induced by d on X* is the topology (Definition 1.1.1) defined by declaring a subset $U \subseteq X$ to be open if for every $x \in U$, there exists $\varepsilon \in (0, \infty)$ such that $B(x, \varepsilon)$ is contained in U .

1.8. Convergence in extended metric spaces.

Definition 1.8.1 (Convergence and Limits in an Extended Metric Space). Let (X, d) be an extended metric space (Definition 1.7.1). A sequence $(x_n)_{n=1}^{\infty}$ in X **converges to a point** $x \in X$ if for every $\varepsilon > 0$, there exists an integer $N \in \mathbb{N}$ such that for all $n \geq N$,

$$d(x_n, x) < \varepsilon.$$

In that case, x is called the **limit of the sequence** (x_n) , and we write

$$\lim_{n \rightarrow \infty} x_n = x \quad \text{or} \quad x_n \rightarrow x$$

Definition 1.8.2 (Cauchy sequence in an extended metric space). Let (M, d) be an extended metric space (Definition 1.7.1). A sequence $(x_n)_{n \in \mathbb{N}}$ in M is called a **Cauchy sequence** if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for all $m, n \geq N$, $d(x_m, x_n) < \epsilon$ (and thus, in particular, all pairwise distances $d(x_m, x_n)$ are finite for $m, n \geq N$).

Definition 1.8.3. Let (X, d) be an extended metric space (Definition 1.7.1). The space (X, d) is said to be **complete** if every Cauchy sequence in X (Definition 1.8.2) converges to a point in X (Definition 1.8.1); that is, for every sequence $(x_n)_{n \in \mathbb{N}}$ in X such that for all $\varepsilon > 0$ there exists $N \in \mathbb{N}$ with $d(x_m, x_n) < \varepsilon$ for all $m, n \geq N$, there exists $x \in X$ such that

$$\lim_{n \rightarrow \infty} d(x_n, x) = 0.$$

Definition 1.8.4. Let (X, d) be an extended metric space (Definition 1.7.1). A **completion** of (X, d) is a pair $(\widehat{X}, \widehat{d})$ together with an isometric (Definition 1.7.2) embedding $i : X \rightarrow \widehat{X}$, where:

- $(\widehat{X}, \widehat{d})$ is a complete extended metric space (Definition 1.8.1),
- $i(X)$ is dense in $\widehat{X}$ (Definition 1.2.1) (Definition 1.2.1) (with respect to the topologies induced by (Definition 1.7.4) the extended metrics),
- $d(x, y) = \widehat{d}(i(x), i(y))$ for all $x, y \in X$.

(♠ TODO: prove that completions of extended metric spaces exist) As such a space $\widehat{X}$ exist, and is unique up to unique isometry, we may refer to the space $\widehat{X}$ as called **the completion** of X (with respect to d).

Proposition 1.8.5. Let (X, d) be an extended metric space (Definition 1.7.1). A completion (Definition 1.8.4) $(\widehat{X}, \widehat{d})$ of (X, d) exists and is unique up to unique isometry. Moreover, if (X, d) is a metric space, then so is $(\widehat{X}, \widehat{d})$.

Proof. (♠ TODO: :) actually complete this proof Let $\mathcal{C}$ be the set of all Cauchy sequences in (X, d) :

$$\mathcal{C} = \{(x_n)_{n \in \mathbb{N}} : (x_n) \text{ is a Cauchy sequence in } (X, d)\},$$

and define a relation $\sim$ on $\mathcal{C}$ by

$$(x_n) \sim (y_n) \iff \lim_{n \rightarrow \infty} d(x_n, y_n) = 0.$$

This relation is well-defined and in fact an equivalence relation. (♠ TODO: Show that the relation is well-defined and an equivalence relation) Let $\widehat{X} = \mathcal{C} / \sim$ and let $\widehat{d} : \widehat{X} \times \widehat{X} \rightarrow [0, \infty]$ be defined by

$$\widehat{d}([(x_n)], [(y_n)]) := \lim_{n \rightarrow \infty} d(x_n, y_n).$$

This limit exists as a value in $[0, \infty]$ and is independent of representative sequences chosen. (♠ TODO: Demonstrate why the limit exists and is well defined)

It can be verified that $\widehat{d}$ is an extended metric on $\widehat{X}$ and that $(\widehat{X}, \widehat{d})$ is complete by construction.

Define an embedding

$$i : X \rightarrow \widehat{X}, \quad i(x) := [(x, x, x, \dots)],$$

which is an isometry:

$$\widehat{d}(i(x), i(y)) = \lim_{n \rightarrow \infty} d(x, y) = d(x, y).$$

Moreover, $i(X)$ is dense in $\widehat{X}$ because any equivalence class $[(x_n)]$ of a Cauchy sequence can be approximated arbitrarily closely by points $i(x_n)$ in the image of X .

Uniqueness:

Suppose $(\widehat{X}', \widehat{d}')$ is another completion of (X, d) with isometric embedding $i' : X \rightarrow \widehat{X}'$.

By the universal property of completions, there exists a unique isometry

$$\Phi : \widehat{X} \rightarrow \widehat{X}'$$

such that $\Phi \circ i = i'$.

The map Φ is onto since $i'(X)$ is dense in $\widehat{X}'$ and Φ preserves limits of Cauchy sequences.

Therefore, Φ is a unique isometry between the completions, and the completion is unique up to unique isometry. $\square$

1.8.1. Absolute values on fields.

Definition 1.8.6 (Absolute Value on a Field). Let F be a field. An *absolute value on F* is a function

$$|\cdot| : F \rightarrow \mathbb{R}_{\geq 0}$$

that satisfies the following properties for all $a, b \in F$:

1. Non-negativity: $|a| \geq 0$,
2. Positive-definiteness: $|a| = 0 \iff a = 0$,
3. Multiplicativity: $|ab| = |a| \cdot |b|$,
4. Triangle inequality: $|a + b| \leq |a| + |b|$.

Here, 0 denotes the additive identity of the field F , and the codomain $\mathbb{R}_{\geq 0}$ consists of non-negative real numbers.

1.8.2. Norms on vector spaces over fields with absolute values.

Definition 1.8.7 (Extended Norm). Let V be a vector space over a field F equipped with an absolute value (Definition 1.8.6)

$$|\cdot| : F \rightarrow [0, \infty).$$

An **extended norm on V** is a function

$$\|\cdot\| : V \rightarrow [0, \infty]$$

satisfying for all $x, y \in V$ and all scalars $\alpha \in F$:

1. **Positive definiteness:** $\|x\| = 0$ if and only if $x = 0$.
2. **Homogeneity:** $\|\alpha x\| = |\alpha| \cdot \|x\|$.
3. **Triangle inequality:** $\|x + y\| \leq \|x\| + \|y\|$,

where arithmetic is extended to allow sums involving ∞ with the convention that $a + \infty = \infty$ for any $a \in [0, \infty]$. A vector space with an extended norm over a field with an absolute value is called an **extended normed (vector) space**.

If the range of the extended norm is contained in $[0, \infty)$, then the extended norm is a **norm** in the usual sense and V may be called a **normed (vector) space**.

Definition 1.8.8 (Topology induced by a norm). Let V be a vector space over a field K with absolute value (Definition 1.8.6) $|\cdot|$, and let $\|\cdot\|$ be an extended norm on V (Definition 1.8.7). The **topology induced by the extended norm $\|\cdot\|$ on V** is defined by declaring a subset $U \subseteq V$ to be open if for every $x \in U$, there exists $\varepsilon > 0$ such that

$$B(x, \varepsilon) := \{y \in V : \|y - x\| < \varepsilon\}$$

is contained in U . The set $B(x, \varepsilon)$ is called the **open ball of radius ε around x** . The collection of all such open sets forms a topology on V .

Equivalently, the topology on V induced by the extended norm $\|\cdot\|$ is the topology on V induced by (Definition 1.7.4) the extended metric $d : V \times V \rightarrow [0, \infty]$ induced by $\|\cdot\|$ (Definition 1.7.3).

1.8.3. Inner product on a vector space over the reals or complex numbers.

Definition 1.8.9. An **inner product on a vector space V over a field $\mathbb{F}$** (either $\mathbb{R}$ or $\mathbb{C}$) is a function

$$\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{F}$$

satisfying, for all $u, v, w \in V$ and all scalars $\alpha \in \mathbb{F}$:

1. **Conjugate symmetry:** $\langle u, v \rangle = \overline{\langle v, u \rangle}$
2. **Linearity in the first argument:** $\langle \alpha u + v, w \rangle = \alpha \langle u, w \rangle + \langle v, w \rangle$
3. **Positive-definiteness:** $\langle v, v \rangle \geq 0$ and $\langle v, v \rangle = 0$ if and only if $v = 0$

A vector space over $\mathbb{F}$ equipped with such an inner product is called an **inner product space**.

Definition 1.8.10. Let V be a vector space over a field $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$ and let $\langle \cdot, \cdot \rangle$ be an inner product (Definition 1.8.9). The *norm induced by $\langle \cdot, \cdot \rangle$* is the norm (Definition 1.8.7)

$$\| \cdot \| : V \rightarrow \mathbb{R}_{\geq 0}$$

defined by

$$\|v\| = \sqrt{\langle v, v \rangle}$$

for any $v \in V$.

1.9. Euclidean space.

Definition 1.9.1. For a positive integer n , let $\mathbb{R}^n$ denote the n -fold Cartesian product of the real line $\mathbb{R}$ with itself:

$$\mathbb{R}^n = \{(x_1, x_2, \dots, x_n) : x_i \in \mathbb{R} \text{ for all } i = 1, \dots, n\}.$$

The set $\mathbb{R}^n$ is called *Euclidean n -space*. A point $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ is associated with the *Euclidean norm*

$$\|x\| = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}.$$

The corresponding metric (Definition 1.7.1) $d : \mathbb{R}^n \times \mathbb{R}^n \rightarrow [0, \infty)$ is given by

$$d(x, y) = \|x - y\|.$$

This metric induces (Definition 1.7.4) the standard topology on $\mathbb{R}^n$, called the *Euclidean topology*.

The Euclidean topology on $\mathbb{R}^n$ coincides with the product topology (Definition 1.4.3) of $\prod_{i=1}^n \mathbb{R}$ where each copy of $\mathbb{R}$ is equipped with its Euclidean topology.

1.10. Connectivity of topological spaces.

Definition 1.10.1. Let X be a topological space. It is said to be *path connected* if for any two points x_0, x_1 in X , there is some path (Definition 3.0.7) $\gamma : [0, 1] \rightarrow X$ in X such that $\gamma(0) = x_0$ and $\gamma(1) = x_1$.

Definition 1.10.2. Let X be a topological space. It is said to be *locally path connected* if for any $x \in X$, there is an open neighborhood (Definition 1.1.9) $x \in U \subseteq X$ that is path connected (Definition 1.10.1).

Definition 1.10.3. Let X be a topological space.

Let $x \in X$. The *path component of x in X* is the maximal path connected subspace of X containing x .

The set of path components in X is denoted $\pi_0(X)$. For any $x \in X$, $\pi_0(X)$ coincides, as a set, with the zeroth homotopy set (Definition 3.0.8) $\pi_0(X, x_0)$ of the pointed space (Definition 1.3.3) (X, x_0) .

Definition 1.10.4. A topological space X is *simply connected* if it is path-connected (Definition 1.10.1) and its fundamental group (Definition 3.0.8) $\pi_1(X, x_0)$ is trivial for some (and hence every) base (Definition 1.1.10) point $x_0 \in X$. Equivalently, X is simply connected if any two continuous maps (Definition 1.3.1) $f, g : [0, 1] \rightarrow X$ with $f(0) = g(0)$ and $f(1) = g(1)$ are homotopic relative to (Definition 3.0.1) $\{0, 1\}$.

Proposition 1.10.5. Let X be a topological space. X is simply connected (Definition 1.10.4) if and only if it is path-connected (Definition 1.10.1) and every continuous map (Definition 1.3.1) $S^1 \rightarrow X$ from the unit circle into X is null-homotopic.

Proposition 1.10.6. (♠ TODO:) Let X be a topological space. If X is contractible, then X is simply connected. The converse is not true in general; for example, the n -dimensional sphere S^n is simply connected for $n \geq 2$ but is not contractible.

Definition 1.10.7. A topological space X is *locally simply connected* if it admits a basis (Definition 1.1.10) of open sets (Definition 1.1.1) $\{U_\alpha\}$ such that each U_α is simply connected (Definition 1.2.2). That is, for every $x \in X$ and every open neighborhood (Definition 1.1.9) V of x , there exists a simply connected open neighborhood $U \subset V$ of x .

Proposition 1.10.8. Let X be a topological space. If X is locally simply connected (Definition 1.10.7), then X is both locally path-connected (Definition 1.10.2) and semi-locally simply connected (Definition 1.10.11).

Proposition 1.10.9. Every topological manifold, with or without boundary, is locally simply connected (Definition 1.10.7) because every point has a neighborhood homeomorphic to an open ball in $\mathbb{R}^n$ or $\mathbb{H}^n$, both of which are simply connected.

Definition 1.10.10. (♠ TODO: contractible) Let X be a topological space. X is *locally contractible* if every point $x \in X$ has a basis (Definition 1.1.10) of contractible open neighborhoods (Definition 1.1.9). Since every contractible space is simply connected, every locally contractible space is locally simply connected (Definition 1.10.7). (♠ TODO: simply connected space)

Definition 1.10.11. A topological space X is *semi-locally simply connected* if for every point $x \in X$, there exists an open neighborhood (Definition 1.1.9) U of x such that the inclusion-induced homomorphism (Definition C.0.2) on fundamental groups (Definition 3.0.8)

$$i_* : \pi_1(U, x) \rightarrow \pi_1(X, x)$$

is the trivial homomorphism.

Proposition 1.10.12. Let X be a topological space. The condition of semi-local simple connectivity (Definition 1.10.11) is equivalent to the requirement that every point $x \in X$ has an open neighborhood U such that every loop in U based at x is null-homotopic in X . Note that the loops are not required to be null-homotopic within U itself. (♠ TODO: null-homotopic)

1.11. Stone spaces, ultrafilters, and Boolean algebras.

Definition 1.11.1. A topological space X is a *Stone space* if it is compact (Definition 1.2.6), Hausdorff (Definition 1.1.13), and totally disconnected (Definition 1.2.4).

Definition 1.11.2. The *category of Stone spaces* is the category (Definition B.0.1) denoted by **Stone** whose objects are Stone spaces (Definition 1.11.1) (compact (Definition 1.2.6), Hausdorff (Definition 1.1.13), and totally disconnected (Definition 1.2.4) topological spaces (Definition 1.1.1)) and the morphisms are continuous maps (Definition 1.3.1) between them.

Definition 1.11.3. A *Boolean algebra* is a set B equipped with two binary operations $\vee$ (*join*) and $\wedge$ (*meet*), a unary operation $\neg$ (*complement*), and two distinct elements 0 (*bottom*) and 1 (*top*), such that for all $x, y, z \in B$:

- $x \vee (y \vee z) = (x \vee y) \vee z$ and $x \wedge (y \wedge z) = (x \wedge y) \wedge z$ (associativity).
- $x \vee y = y \vee x$ and $x \wedge y = y \wedge x$ (commutativity).
- $x \wedge (x \vee y) = x$ and $x \vee (x \wedge y) = x$ (absorption).
- $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$ and $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$ (distributivity).
- $x \vee (\neg x) = 1$ and $x \wedge (\neg x) = 0$ (complements).

Definition 1.11.4. Let A and B be Boolean algebras (Definition 1.11.3). A *morphism of Boolean algebras* is a function $f : A \rightarrow B$ such that for all $x, y \in A$:

- $f(x \vee y) = f(x) \vee f(y)$,
- $f(x \wedge y) = f(x) \wedge f(y)$,
- $f(\neg x) = \neg f(x)$,
- $f(0_A) = 0_B$ and $f(1_A) = 1_B$.

Definition 1.11.5. The *category of Boolean algebras* is the category (Definition B.0.1) whose objects are Boolean algebras (Definition 1.11.3) and the morphisms are Boolean algebra morphisms (Definition 1.11.4). It is often denoted by **Bool**.

Proposition 1.11.6. Let A and B be Boolean algebras (Definition 1.11.3). (♠ **TODO: corresponding boolean rings**) A function $f : A \rightarrow B$ is a morphism of Boolean algebras (Definition 1.11.4) if and only if it is a ring homomorphism between the corresponding Boolean rings.

Definition 1.11.7. Let B be a Boolean algebra (Definition 1.11.3). A *filter F on B* is a non-empty subset $F \subseteq B$ such that $0 \notin F$, if $x, y \in F$ then $x \wedge y \in F$, and if $x \in F$ and $x \leq y$ (where $x \leq y$ is defined as $x \wedge y = x$) then $y \in F$. An *ultrafilter* is a filter U that is maximal with respect to set inclusion.

Proposition 1.11.8. Let B be a Boolean algebra and $U \subseteq B$ a filter. U is an ultrafilter if and only if for every $x \in B$, either $x \in U$ or $\neg x \in U$.

Definition 1.11.9. Let B be a Boolean algebra (Definition 1.11.3). The *Stone space of B* , denoted by $\text{Spec}(B)$, is the set of all ultrafilters (Definition 1.11.7) on B equipped with the topology generated by (Definition 1.1.10) the basis (Definition 1.1.10) of sets

$$U_a = \{p \in \text{Spec}(B) : a \in p\}$$

for all $a \in B$.

Theorem 1.11.10 (Stone's representation theorem). Stone's representation theorem states that there is a contravariant equivalence of categories (a duality) between the category of Boolean algebras (Definition 1.11.5) **Bool** and the category of Stone spaces (Definition 1.11.2) **Stone**. This equivalence is realized by the following pair of contravariant functors:

- $\mathcal{S} : \mathbf{Bool} \rightarrow \mathbf{Stone}$, which maps a Boolean algebra (Definition 1.11.3) B to its Stone space $\text{Spec}(B)$ of ultrafilters (Definition 1.11.7).
- $\mathcal{C} : \mathbf{Stone} \rightarrow \mathbf{Bool}$, which maps a Stone space X to its Boolean algebra of clopen subsets (Definition 1.1.2) $\text{Clopen}(X)$.

For any Boolean algebra B , there is a natural isomorphism $B \cong \mathcal{C}(\mathcal{S}(B))$, and for any Stone space X , there is a natural homeomorphism (Definition 1.3.2) $X \cong \mathcal{S}(\mathcal{C}(X))$.

2. COVERING SPACES

Definition 2.0.1. 1. Let X be a topological space.

A *covering space of X* is a pair $(\tilde{X}, p)$ where $\tilde{X}$ is a topological space and $p : \tilde{X} \rightarrow X$ is a continuous (Definition 1.3.1) surjective (Definition A.0.2) map such that for every $x \in X$, there exists an open neighborhood (Definition 1.1.9) U of x (called an *evenly covered neighborhood*) for which the preimage $p^{-1}(U)$ is a disjoint union of open sets (Definition 1.1.1) $V_i \subset \tilde{X}$, each of which is mapped homeomorphically (Definition 1.3.2) onto U by the restriction $p|_{V_i}$.

2. Let (X, x_0) be a pointed topological space (Definition 1.3.3). A *(pointed) covering of (X, x_0)* is a pointed topological space (Definition 1.3.3) $(\tilde{X}, \tilde{x}_0)$ along with a covering $p : \tilde{X} \rightarrow X$ of X such that p is a map of pointed space (Definition 1.3.3).

Definition 2.0.2. 1. Let X be a topological space.

A *morphism of covering spaces of X from $p_1 : \tilde{X}_1 \rightarrow X$ to $p_2 : \tilde{X}_2 \rightarrow X$* is a continuous map (Definition 1.3.1) $f : \tilde{X}_1 \rightarrow \tilde{X}_2$ such that $p_2 \circ f = p_1$.

2. Let (X, x_0) be a pointed topological space (Definition 1.3.3). Let $p_1 : (\tilde{X}_1, \tilde{x}_1) \rightarrow (X, x_0)$ to $p_2 : (\tilde{X}_2, \tilde{x}_2) \rightarrow (X, x_0)$ be pointed coverings (Definition 2.0.1) of (X, x_0) . A *morphism $f : p_1 \rightarrow p_2$ of the pointed coverings* is a pointed map $f : (\tilde{X}_1, \tilde{x}_1) \rightarrow (\tilde{X}_2, \tilde{x}_2)$ such that $p_2 \circ f = p_1$ and $f(\tilde{x}_1) = \tilde{x}_2$.

Definition 2.0.3. Let $p : \tilde{X} \rightarrow X$ be a covering space (Definition 2.0.1) of a topological space X . A *deck transformation* (or *covering transformation*, or *automorphism*) of p is a homeomorphism (Definition 1.3.2) $f : \tilde{X} \rightarrow \tilde{X}$ such that $p \circ f = p$. The set of all deck transformations forms a group under composition, denoted $\text{Aut}(p)$ or $\text{Deck}(p)$.

Definition 2.0.4. Let X be a topological space. A *universal cover of X* is a pair $(\tilde{X}, p)$ where $p : \tilde{X} \rightarrow X$ is a covering space (Definition 2.0.1) such that $\tilde{X}$ is simply connected (Definition 1.10.4).

Theorem 2.0.5. Let $p : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$ be a covering space (Definition 2.0.1) of a path-connected (Definition 1.10.1) and locally path-connected space (Definition 1.10.2) X . The induced map on fundamental groups (Definition 3.0.8) $p_* : \pi_1(\tilde{X}, \tilde{x}_0) \rightarrow \pi_1(X, x_0)$ is an injective group homomorphism (Definition C.0.2).

Definition 2.0.6. Let $p : \tilde{X} \rightarrow X$ be a covering space (Definition 2.0.1) of a topological space X . Let $f : Y \rightarrow X$ be a continuous map (Definition 1.3.1) of topological spaces.

A *lift of f (via p)* is a continuous map $\tilde{f} : Y \rightarrow \tilde{X}$ satisfying $p \circ \tilde{f} = f$.

Similarly, for a pointed covering $p : (\tilde{X}, x_0) \rightarrow (X, x_0)$ and a pointed morphism (Definition 1.3.3) $f : (Y, y_0) \rightarrow (X, x_0)$, a *lift of f (via p)* is a pointed map $\tilde{f} : (Y, y_0) \rightarrow (\tilde{X}, x_0)$ satisfying $p \circ \tilde{f} = f$.

Proposition 2.0.7. Every simply connected (Definition 1.2.2) topological space is semi-locally simply connected (Definition 1.10.11). Furthermore, every locally simply connected space is semi-locally simply connected. However, there exist semi-locally simply connected spaces, such as the cone over the Hawaiian earring, that are not locally simply connected. *Remark:* (♠ TODO: define locally simply connected).

Definition 2.0.8. A covering space (Definition 2.0.1) $p : \tilde{X} \rightarrow X$ of a topological space X is *normal* (or *Galois*) if for every $\tilde{x}_0 \in \tilde{X}_0$, the image of the induced map $p_* : \pi_1(\tilde{X}, \tilde{x}_0) \rightarrow \pi_1(X, p(\tilde{x}_0))$ on fundamental groups (Definition 3.0.8) is the normal subgroup of $\pi_1(X, p(\tilde{x}_0))$.

Theorem 2.0.9. Let $p : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$ be a covering space (Definition 2.0.1) and let $f : (Y, y_0) \rightarrow (X, x_0)$ be a continuous map (Definition 1.3.1) where Y is path-connected (Definition 1.10.1) and locally path-connected (Definition 1.10.2). A lift $\tilde{f} : (Y, y_0) \rightarrow (\tilde{X}, \tilde{x}_0)$ exists if and only if the image of the fundamental group (Definition 3.0.8) of (Y, y_0) is contained in the image of the fundamental group of $(\tilde{X}, \tilde{x}_0)$, i.e.

$$f_*(\pi_1(Y, y_0)) \subseteq p_*(\pi_1(\tilde{X}, \tilde{x}_0))$$

If such a lift exists, it is unique.

See also Theorem 2.0.10.

Theorem 2.0.10. Let $p_X : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$ and $p_Y : (\tilde{Y}, \tilde{y}_0) \rightarrow (Y, y_0)$ be covering spaces of pointed topological spaces (Definition 1.3.3). Let $f : (X, x_0) \rightarrow (Y, y_0)$ be a continuous map (Definition 1.3.1), and assume $\tilde{X}$ is path-connected (Definition 1.10.1) and locally path-connected (Definition 1.10.2).

There exists a unique continuous lift $\tilde{f} : (\tilde{X}, \tilde{x}_0) \rightarrow (\tilde{Y}, \tilde{y}_0)$ such that $p_Y \circ \tilde{f} = f \circ p_X$ if and only if

$$f_*(p_{X*}(\pi_1(\tilde{X}, \tilde{x}_0))) \subseteq p_{Y*}(\pi_1(\tilde{Y}, \tilde{y}_0))$$

In the specific case where f is the identity on X , this provides the necessary and sufficient condition for the existence of a morphism of covering spaces (pointed or unpointed). Any such morphism between connected covering spaces is itself a covering map. See also Theorem 2.0.9.

Theorem 2.0.11. Let X be a connected (Definition 1.2.2), locally path-connected (Definition 1.10.2), and semi-locally simply connected (Definition 1.10.11) space with base (Definition 1.1.10) point x_0 . The universal cover (Definition 2.0.4) $\tilde{X}$ can be constructed as the space of homotopy classes (Definition 3.0.3) of paths (Definition 3.0.7) starting at x_0 :

$$\tilde{X} = \{[\gamma] \mid \gamma : [0, 1] \rightarrow X, \gamma(0) = x_0\}$$

with the covering map $p : \tilde{X} \rightarrow X$ given by $p([\gamma]) = \gamma(1)$. For any connected covering space (Definition 2.0.1) $q : Y \rightarrow X$, there exists a covering map $r : \tilde{X} \rightarrow Y$ such that $q \circ r = p$.

Proposition 2.0.12. Let X be a connected (Definition 1.2.2), locally path-connected (Definition 1.10.2), and semi-locally simply connected (Definition 1.10.11) space with base point (Definition 1.3.3) x_0 . There are bijective (Definition A.0.2) correspondences:

1. Between the set of isomorphism classes of connected (Definition 1.2.2) pointed covering spaces (Definition 2.0.1) $p : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$ and the set of subgroups of $\pi_1(X, x_0)$, via $((\tilde{X}, \tilde{x}_0), p) \mapsto p_*(\pi_1(\tilde{X}, \tilde{x}_0))$.
2. Between the set of isomorphism classes of connected covering spaces $p : \tilde{X} \rightarrow X$ and the set of conjugacy classes of subgroups of $\pi_1(X, x_0)$, via $p \mapsto [p_*(\pi_1(\tilde{X}, \tilde{x}_0))]$. (♠
TODO: conjugacy class)

Theorem 2.0.13. Let $p : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$ be a covering with $\tilde{X}$ (and hence X) connected (Definition 1.2.2). The group of deck transformations (Definition 2.0.3) $\text{Aut}(p)$ is isomorphic to the quotient $N(H)/H$, where $H = p_*(\pi_1(\tilde{X}, \tilde{x}_0))$ and $N(H)$ is its normalizer in $\pi_1(X, x_0)$. If the covering is normal (Definition 2.0.8) (i.e., H is a normal subgroup of $\pi_1(X, x_0)$), then:

$$\text{Aut}(p) \cong \pi_1(X, x_0)/p_*(\pi_1(\tilde{X}, \tilde{x}_0))$$

For the universal cover, $\text{Aut}(p) \cong \pi_1(X, x_0)$.

3. HOMOTOPY

The notions of homotopy have the following variants:

1. Absolute, on unpointed spaces
2. Relative, on unpointed spaces
3. Absolute, on pointed spaces
4. Relative, on pointed spaces.

Definition 3.0.1 (Homotopy of maps of topological spaces). Let X and Y be topological spaces and let $K \subseteq X$ be a subset. Let $C(X, Y)$ denote the set of all continuous maps $f : X \rightarrow Y$.

1. A *homotopy between two maps $f, g \in C(X, Y)$ relative to K* is a continuous map

$$H : X \times [0, 1] \rightarrow Y$$

such that for all $x \in X$,

$$H(x, 0) = f(x), \quad H(x, 1) = g(x),$$

and for all $x \in K$ and $t \in [0, 1]$,

$$H(x, t) = f(x) = g(x).$$

If such an H exists, we say f and g are *homotopic relative to K* , and we write $f \simeq g \text{ rel } K$ or $f \simeq_K g$; this is an equivalence relation.

In the case that $K = \emptyset$, a homotopy relative to K is simply called a *homotopy*. We write we write $f \simeq g$ if a homotopy between f and g exists.

- Let (X, x_0) and (Y, y_0) be pointed topological spaces (Definition 1.3.3) and let $K \subseteq X$ be a subset with $x_0 \in K$. Let $C_*(X, Y)$ denote the set of all continuous based maps $f : X \rightarrow Y$ satisfying $f(x_0) = y_0$.

A *homotopy of based maps* $f, g \in C_*(X, Y)$ *relative to* K is a continuous map

$$H : X \times [0, 1] \rightarrow Y$$

such that for all $x \in X$,

$$H(x, 0) = f(x), \quad H(x, 1) = g(x),$$

and for all $k \in K$ and $t \in [0, 1]$,

$$H(k, t) = f(k) = g(k),$$

in particular fixing the basepoint throughout,

$$H(x_0, t) = y_0 \quad \text{for all } t \in [0, 1].$$

If such an H exists, we say f and g are *based homotopic relative to* K , and we write $f \simeq g \text{ rel } K$ or $f \simeq_K g$. This is an equivalence relation.

When $K = \{x_0\}$, one calls a homotopy of based maps relative to K simply as a *homotopy of based maps* or *based homotopy*. We write $f \simeq g$ if such a homotopy exists.

Definition 3.0.2 (Homotopy equivalence). Let X and Y be topological spaces and let $K \subseteq X$.

- A continuous map (Definition 1.3.1) $f : X \rightarrow Y$ is a *homotopy equivalence relative to* K if there exists a continuous map $g : Y \rightarrow X$ such that

$$g \circ f \simeq \text{id}_X \text{ rel } K \quad \text{and} \quad f \circ g \simeq \text{id}_Y \text{ rel } f(K).$$

(Definition 3.0.1) That is, there exist homotopies (Definition 3.0.1) $H : X \times [0, 1] \rightarrow X$ and $G : Y \times [0, 1] \rightarrow Y$ satisfying

$$\begin{aligned} H(x, 0) &= (g \circ f)(x), & H(x, 1) &= x, & H(k, t) &= k \text{ for all } k \in K, t \in [0, 1], \\ G(y, 0) &= (f \circ g)(y), & G(y, 1) &= y, & G(f(k), t) &= f(k) \text{ for all } k \in K, t \in [0, 1]. \end{aligned}$$

In this case, the map g is called a *homotopy inverse relative to* K *of* f , and the pairs (X, K) and $(Y, f(K))$ are said to be *homotopy equivalent relative to* K , denoted $(X, K) \simeq (Y, f(K)) \text{ rel } K$. This is an equivalence relation.

A continuous map $f : X \rightarrow Y$ is a *homotopy equivalence* if it is a homotopy equivalence relative to $\emptyset$. A *homotopy inverse of* f is then simply a homotopy inverse of f relative to $\emptyset$. In this case, and the spaces X and Y are said to be *homotopy equivalent*, denoted $X \simeq Y$. This is an equivalence relation.

- Let (X, x_0) and (Y, y_0) be pointed topological spaces (Definition 1.3.3) and let $K \subseteq X$ be a subset with $x_0 \in K$.

A continuous map $f : (X, x_0) \rightarrow (Y, y_0)$ of pointed spaces is a *homotopy equivalence relative to* K if there exists a continuous map $g : (Y, y_0) \rightarrow (X, x_0)$ such that

$$g \circ f \simeq \text{id}_X \text{ rel } K \quad \text{and} \quad f \circ g \simeq \text{id}_Y \text{ rel } f(K),$$

where the homotopies (Definition 3.0.1) are homotopies of based maps relative to K and $f(K)$, respectively.

That is, there exist homotopies of based maps $H : X \times [0, 1] \rightarrow X$ and $G : Y \times [0, 1] \rightarrow Y$ satisfying

$$\begin{aligned} H(x, 0) &= (g \circ f)(x), & H(x, 1) &= x, & H(k, t) &= k \text{ for all } k \in K, t \in [0, 1], \\ G(y, 0) &= (f \circ g)(y), & G(y, 1) &= y, & G(f(k), t) &= f(k) \text{ for all } k \in K, t \in [0, 1], \end{aligned}$$

with $H(x_0, t) = x_0$ and $G(y_0, t) = y_0$ for all t .

The map g is called a *homotopy inverse relative to K of f* , and the pairs (X, K) and $(Y, f(K))$ are said to be *homotopy equivalent relative to K* , denoted $(X, K) \simeq (Y, f(K)) \text{ rel } K$. This is an equivalence relation.

A continuous map $f : (X, x_0) \rightarrow (Y, y_0)$ of pointed spaces is a *homotopy equivalence* if it is a homotopy equivalence relative to $\{x_0\}$. A *homotopy inverse of f* is then simply a homotopy inverse relative to $\{x_0\}$. In this case, the pointed spaces (X, x_0) and (Y, y_0) are said to be *homotopy equivalent*, denoted $(X, x_0) \simeq (Y, y_0)$.

In each case, topological spaces that are homotopy equivalent are said to be of the same *homotopy type*.

Definition 3.0.3 (Homotopy class of maps relative to a subset). Let X and Y be topological spaces (Definition 1.1.1) and let $K \subseteq X$. Let $C(X, Y)$ denote the set of all continuous maps (Definition 1.3.1) $f : X \rightarrow Y$.

1. Two maps $f, g \in C(X, Y)$ are said to be in the same *homotopy class relative to K* or that f and g are *homotopic relative to K* if there exists a homotopy relative to K (Definition 3.0.1)

$$H : X \times [0, 1] \rightarrow Y$$

such that

$$H(x, 0) = f(x), \quad H(x, 1) = g(x),$$

and

$$H(k, t) = f(k) = g(k) \quad \text{for all } k \in K, t \in [0, 1].$$

The *homotopy class of maps relative to K* containing a map $f : X \rightarrow Y$ is denoted by $[f]_K$.

Two maps $f, g \in C(X, Y)$ are said to be in the same *homotopy class* if they are in the same homotopy class relative to $\emptyset$.

The *homotopy class of maps* containing a map $f : X \rightarrow Y$ is denoted by $[f]$.

The set of homotopy classes of maps may often be denoted by $[X, Y]$.

2. Let (X, x_0) and (Y, y_0) be pointed topological spaces (Definition 1.3.3) and let $K \subseteq X$ be a subset containing x_0 . Let $C_*(X, Y)$ denote the set of all continuous based maps $f : X \rightarrow Y$ with $f(x_0) = y_0$.

Two based maps $f, g \in C_*(X, Y)$ are said to be in the same *homotopy class relative to K* or that f and g are *homotopic relative to K* if there exists a homotopy of based maps relative to K

$$H : X \times [0, 1] \rightarrow Y$$

such that for all $x \in X$,

$$H(x, 0) = f(x), \quad H(x, 1) = g(x),$$

and for all $k \in K$ and $t \in [0, 1]$,

$$H(k, t) = f(k) = g(k),$$

particularly ensuring the basepoint is fixed throughout,

$$H(x_0, t) = y_0 \quad \text{for all } t \in [0, 1].$$

In the case that $K = \{x_0\}$, one simply says that f and g are *homotopic (as based maps)*.

The *homotopy class relative to K* containing $f : (X, x_0) \rightarrow (Y, y_0)$ is denoted by $[f]_K$.

Two based maps $f, g \in C_*(X, Y)$ are said to be in the same *homotopy class* if they are in the same homotopy class relative to $\{x_0\}$.

The *homotopy class* containing a map $f : (X, x_0) \rightarrow (Y, y_0)$ is denoted by $[f]$.

The set of homotopy classes of pointed maps $(X, x_0) \rightarrow (Y, y_0)$ may often be denoted by $[(X, x_0), (Y, y_0)]$ or by notations such as $[X, Y]_*$, $[X, Y]_\bullet$, or $[X, Y]$ if the base points are clear.

Definition 3.0.4. The *homotopy category of topological spaces*, denoted $\mathbf{hTop}$, is the category whose objects are topological spaces (Definition 1.1.1) and whose morphisms are homotopy classes (Definition 3.0.3) of continuous maps (Definition 1.3.1). In other words, for objects X and Y , the set of morphisms is defined as $\text{Hom}_{\mathbf{hTop}}(X, Y) = [X, Y] = C(X, Y) / \simeq$.

Proposition 3.0.5. Composition in the homotopy category of topological spaces (Definition 3.0.4) is well-defined. If $f_1, f_2 : X \rightarrow Y$ are homotopic and $g_1, g_2 : Y \rightarrow Z$ are homotopic, then the compositions $g_1 \circ f_1$ and $g_2 \circ f_2$ are homotopic as maps from X to Z . That is, $[g] \circ [f] = [g \circ f]$ is independent of the choice of representatives.

Theorem 3.0.6. There exists a canonical functor $Q : \mathbf{Top} \rightarrow \mathbf{hTop}$ which is the identity on objects and maps each continuous map f to its homotopy class $[f]$. This functor is full and essentially surjective.

Definition 3.0.7 (Path in a topological space). Let X be a topological space (Definition 1.1.1), let $x_0, x_1 \in X$, and let $[0, 1] \subset \mathbb{R}$ be the closed unit interval equipped with the subspace topology from $\mathbb{R}$.

1. A *path in X from x_0 to x_1* is a continuous map $\gamma : [0, 1] \rightarrow X$ such that $\gamma(0) = x_0$ and $\gamma(1) = x_1$. More generally, a *path in X* is a path from some point $x_0 \in X$ to some point $x_1 \in X$ in this sense.
2. A *loop in X based at $x_0 \in X$* is a path $\gamma : [0, 1] \rightarrow X$ in X from x_0 to x_0 , that is, a continuous map γ such that $\gamma(0) = \gamma(1) = x_0$. When a basepoint x_0 is fixed and understood from the context, a *loop in X* means a loop in X based at this chosen basepoint.

Definition 3.0.8 (Homotopy groups of a pointed topological space). For any pointed topological space (Definition 1.3.3) (X, x_0) and integer $n \geq 0$, the *n -th homotopy set of X at x_0* , denoted $\pi_n(X, x_0)$ or simply $\pi_n(X)$ when the base point x_0 is understood or a choice of base point does not really matter, is defined as the set of all homotopy classes (rel. ∂I^n) (Definition 3.0.3) of paired maps

$$f : (I^n, \partial I^n) \rightarrow (X, \{x_0\}),$$

where $I^n = [0, 1]^n$ and ∂I^0 is taken to be the singleton I^0 instead of the literal boundary (Definition 1.2.10) thereof. Equivalently, one can also define it as the set of all homotopy classes of pointed maps (Definition 1.3.3)

$$f : (S^n, s_0) \rightarrow (X, x_0),$$

where S^n is the n -sphere (♠ TODO:) and $s_0 \in S^n$ is a choice of base point. In other words, $\pi_n(X, x_0)$ is by definition $[(S^n, s_0), (X, x_0)]_*$ (Definition 3.0.3).

For $n \geq 1$, $\pi_n(X, x_0)$ is a group under concatenation of pointed maps, and for $n \geq 2$, it is abelian. As such, for $n \geq 1$, it is appropriate to refer to $\pi_n(X, x_0)$ as the *n th homotopy group of (X, x_0)* .

The *fundamental group of (X, x_0)* refers to $\pi_1(X, x_0)$. Equivalently, it is the group of homotopy classes (rel. endpoints) of loops (Definition 3.0.7) $\gamma : [0, 1] \rightarrow X$ satisfying $\gamma(0) = \gamma(1) = x_0$.

For every $n \geq 0$, the assignment $(X, x_0) \mapsto \pi_n(X, x_0)$ is functorial — in fact

1. For $n = 0$, it is a functor $\text{Top}_* \rightarrow \mathbf{Sets}$,
2. For $n = 1$, it is a functor $\text{Top}_* \rightarrow \mathbf{Grp}$,
3. For $n \geq 2$, it is a functor $\text{Top}_* \rightarrow \mathbf{Ab}()$.

Given a map $f : (X, x_0) \rightarrow (Y, y_0)$ of pointed spaces, the induced map $\pi_n(X, x_0) \rightarrow \pi_n(Y, y_0)$ is often denoted by f_* .

The homotopy set $\pi_n(X, x_0)$ is naturally isomorphic to the relative homotopy set $\pi_n(X, \{x_0\}, x_0)$ (Definition 3.0.9).

We further note that the set $\pi_n(X, x_0) = [(S^n, s_0), (X, x_0)]_*$ is identifiable with $\pi_0(\text{Map}_*((S^n, s_0), (X, x_0)))$.

Definition 3.0.9 (Relative homotopy groups of pointed pairs). For any pointed pair of topological spaces (X, A, x_0) and an integer $n \geq 1$, the *n -th relative homotopy set of (X, A) at x_0* , denoted $\pi_n(X, A, x_0)$, is defined as the set of all homotopy classes (rel. J^{n-1}) (Definition 3.0.3) of maps of triples

$$f : (I^n, \partial I^n, J^{n-1}) \rightarrow (X, A, \{x_0\}),$$

where $I^n = [0, 1]^n$ is the n -cube, ∂I^n is its boundary (Definition 1.2.10), and

$$J^{n-1} := \overline{\partial I^n \setminus (I^{n-1} \times \{1\})} = (I^{n-1} \times \{0\}) \cup (\partial I^{n-1} \times I) \subseteq \partial I^n.$$

Explicitly, an element of $\pi_n(X, A, x_0)$ is represented by a continuous map (Definition 1.3.1) $f : I^n \rightarrow X$ such that:

1. $f(\partial I^n) \subseteq A$,
2. $f(J^{n-1}) = \{x_0\}$.

Equivalently, $\pi_n(X, A, x_0)$ can be defined as the set of all homotopy classes of maps of pairs

$$f : (D^n, S^{n-1}, s_0) \rightarrow (X, A, x_0),$$

where D^n is the n -disk, S^{n-1} is its boundary sphere, and $s_0 \in S^{n-1}$ is a base point.

For $n \geq 2$, $\pi_n(X, A, x_0)$ is a group under the concatenation of maps in the first coordinate, and for $n \geq 3$, it is abelian. As such:

1. For $n = 1$, the assignment $(X, A, x_0) \mapsto \pi_1(X, A, x_0)$ is a functor $\mathbf{Top}_{*,\text{pair}} \rightarrow \mathbf{Sets}$,
2. For $n = 2$, it is a functor $\mathbf{Top}_{*,\text{pair}} \rightarrow \mathbf{Grp}$,
3. For $n \geq 3$, it is a functor $\mathbf{Top}_{*,\text{pair}} \rightarrow \mathbf{Ab}$.

For $n \geq 2$, $\pi_n(X, A, x_0)$ is called the *n -th relative homotopy group of (X, A) at x_0* . Given a map $f : (X, A, x_0) \rightarrow (Y, B, y_0)$ of pointed pairs, the induced map $\pi_n(X, A, x_0) \rightarrow \pi_n(Y, B, y_0)$ is denoted by f_* .

Definition 3.0.10 (Weak homotopy equivalence). Let $f : X \rightarrow Y$ be a continuous map (Definition 1.3.1) between topological spaces. The map f is a *weak homotopy equivalence* if, for every choice of basepoint $x_0 \in X$ with $y_0 = f(x_0)$, the induced maps on homotopy groups (Definition 3.0.8)

$$f_* : \pi_n(X, x_0) \rightarrow \pi_n(Y, y_0)$$

are isomorphisms (Definition C.0.2) for all integers $n \geq 0$.

4. SINGULAR HOMOLOGY AND COHOMOLOGY

Definition 4.0.1. Let V be a real vector space of finite dimension. A *k -simplex in topology* (or a *geometric k -simplex*) is the convex hull of $k + 1$ affinely independent points $v_0, v_1, \dots, v_k \in V$, and is denoted by

$$[v_0, v_1, \dots, v_k] := \left\{ \sum_{i=0}^k t_i v_i \mid t_i \geq 0, \sum_{i=0}^k t_i = 1 \right\}.$$

It is also standard to talk of the *standard topological n -simplex* — the topological space, often denoted by notations such as Δ^n or $|\Delta^n|$ defined as the subset of Euclidean space $\mathbb{R}^{n+1}$ given by

$$|\Delta^n| = \left\{ (t_0, t_1, \dots, t_n) \in \mathbb{R}^{n+1} : \sum_{i=0}^n t_i = 1, \text{ and } t_i \geq 0 \text{ for all } i \right\}$$

equipped with the induced topology from the usual Euclidean topology on $\mathbb{R}^{n+1}$.

(♠ TODO: comment on how $|\Delta^n|$ makes sense via a geometric realization)

Definition 4.0.2 (Singular simplices). Let X be a topological space (Definition 1.1.1). For each integer $n \geq 0$, A *singular n -simplex in X* is a continuous map (Definition 1.3.1)

$$\sigma : \Delta^n \rightarrow X.$$

where Δ^n is the standard topological n -simplex (Definition 4.0.1). The set of all singular n -simplices in X is denoted by $S_n(X)$.

Definition 4.0.3 (Singular chain group with coefficients). Let X be a topological space (Definition 1.1.1), let $A \subseteq X$ be a subspace (Definition 1.1.14), and let $S_n(X)$ be the set of singular n -simplices in X (Definition 4.0.2).

1. Let R be a ring.

- (a) The *singular n -chain group of X with coefficients in R* is the free R -module $C_n(X; R)$ whose elements are finite formal linear combinations

$$\sum_i r_i \sigma_i, \quad \text{with } \sigma_i \in S_n(X), \quad r_i \in R.$$

Elements of $C_n(X; R)$ are called *singular n -chains in X with coefficients in R* .

- (b) If $A \subseteq X$ is a subspace, the quotient groups

$$C_n(X, A; R) = C_n(X; R) / C_n(A; R)$$

may be referred to as the *relative singular n -chain groups of X in A with coefficients in R* and elements of this group may be referred to as *relative singular n -chains in X relative to A with coefficients in R* .

In either case, when $R = \mathbb{Z}$, the ring R may be suppressed from notation, so we may write $C_n(X)$ and $C_n(X, A)$ for $C_n(X, \mathbb{Z})$ and $C_n(X, A, \mathbb{Z})$ respectively.

2. Let G be an abelian group (Definition C.0.1).

- (a) More generally, the *singular n -chain group of X with coefficients in G* is the abelian group

$$C_n(X; G) = C_n(X; \mathbb{Z}) \otimes_{\mathbb{Z}} G.$$

(??) In particular, an element of $C_n(X; G)$ is a finite formal sum

$$\sum_i g_i \sigma_i, \quad g_i \in G, \sigma_i \in S_n(X).$$

Elements of $C_n(X; G)$ are called *singular n -chains in X with coefficients in G* .

- (b) Similarly, the *relative singular n -chain group of X in A with coefficients in G* is the quotient group

$$C_n(X, A; G) = C_n(X; G) / C_n(A; G)$$

Elements of $C_n(X, A; G)$ are called *singular n -chains in X relative to A with coefficients in G* .

When $G = R$ is a ring, note that the notions of $C_n(X; R)$ where R is taken as a ring and where R is taken as an abelian group are equivalent; similarly for $C_n(X, A; R)$.

We note that, in general $C_n(X; G) \cong C_n(X, \emptyset; G)$.

Definition 4.0.4 (Singular chain complex with coefficients). Let X be a topological space (Definition 1.1.1), let $A \subseteq X$ be a subspace (Definition 1.1.14), and let G be an abelian group.

For each $n \geq 1$, define the *boundary operator*

$$\partial_n : C_n(X; G) \rightarrow C_{n-1}(X; G)$$

(Definition 4.0.3) by

$$\partial_n(\sigma) = \sum_{i=0}^n (-1)^i \sigma \circ \delta_i,$$

where $\delta_i : \Delta^{n-1} \rightarrow \Delta^n$ is the i -th face inclusion, and extend ∂_n to $C_n(X; G)$ linearly. For $n \leq 0$, we simply let $\partial_n = 0$

One can verify that $(C_n(X; G), \partial_n)$ forms a chain complex (Definition E.0.1) in the abelian category of abelian groups. This chain complex is called the *(absolute) singular chain complex of X with coefficients in G* and is often denoted by $C_\bullet(X; G)$.

Then the boundary maps above induce maps

$$C_n(X, A; G) \rightarrow C_{n-1}(X, A; G)$$

on the relative chain groups $C_n(X, A; G)$, yielding a chain complex of abelian groups whose boundary maps, often also denoted by ∂_n , are induced by the boundary operators ∂_n above; this chain complex may be called the *relative singular chain complex of the pair (X, A) with coefficients in G* and is often denoted by $C_\bullet(X, A; G)$.

In the case that $G = R$ is a (not necessarily commutative) ring, then the boundary maps of $C_\bullet(X; R)$ and $C_\bullet(X, A; R)$ are left R -module homomorphisms:

$$\partial \left(\sum_i r_i \sigma_i \right) = \sum_i r_i \partial(\sigma_i)$$

Definition 4.0.5 (Singular homology with coefficients). Let X be a topological space (Definition 1.1.1), let $A \subseteq X$ be a subspace (Definition 1.1.14), and let G be an abelian group.

The *(absolute) n -th singular homology group of X with coefficients in G* is the homology group (Definition E.0.2)

$$H_n(X; G) = H_n(C_*(X; G)) = \ker \partial_n / \operatorname{im} \partial_{n+1}$$

where $C_*(X; G)$ is the singular chain complex of X with coefficients in G and $\partial_n : C_n(X; G) \rightarrow C_{n-1}(X; G)$ are the boundary maps.

Given a subspace (Definition 1.1.14) $A \subseteq X$, the *n -th relative singular homology group of (X, A) with coefficients in G* is defined as

$$H_n(X, A; G) = H_n(C_*(X, A; G)) = \ker \partial_n / \operatorname{im} \partial_{n+1}$$

where $C_*(X, A; G)$ is the relative singular chain complex of (X, A) with coefficients in G and $\partial_n : C_n(X, A; G) \rightarrow C_{n-1}(X, A; G)$ are the boundary maps.

We may denote $H_n(X; \mathbb{Z})$ and $H_n(X, A; \mathbb{Z})$ by $H_n(X)$ and $H_n(X, A)$ respectively.

In the case that $G = R$ is a ring, $H_n(X; R)$ and $H_n(X, A; R)$ have the structure of left R -modules (Definition D.0.2) since the corresponding chain complexes are chain complexes of left R -modules..

Definition 4.0.6 (Singular cochain complex of a topological space). Let X be a topological space (Definition 1.1.1) and R a commutative ring (Definition D.0.1) with 1. The *singular n -cochain group of X with coefficients in R* is the R -module

$$C^n(X; R) = \operatorname{Hom}_R(C_n(X; R), R).$$

(Definition D.0.3) (Definition 4.0.3) We define *coboundary operators*

$$\delta^n : C^n(X; R) \rightarrow C^{n+1}(X; R), \quad \delta^n(\varphi) = \varphi \circ \partial_{n+1}.$$

The pair $(C^n(X; R), \delta^n)_n$ form a cochain complex (Definition E.0.1) called the *singular cochain complex of X with coefficients in R* . Given a subspace $A \subseteq X$, the *relative singular n -cochain group of (X, A) with coefficients in R* is the R -module

$$C^n(X, A; R) = \text{Hom}_R(C_n(X, A; R), R).$$

The coboundary operators δ^n induce maps $C^n(X, A; R) \rightarrow C^{n+1}(X, A; R)$ for which the $C^n(X, A; R)$ form a cochain complex, called the *relative singular cochain complex of (X, A) with coefficients in R*

Definition 4.0.7. Let X be a topological space (Definition 1.1.1) and R a commutative ring (Definition D.0.1) with 1. The *n -th singular cohomology group of X with coefficients in R* is

$$H^n(X; R) = \ker(\delta^n) / \text{im}(\delta^{n-1}).$$

For a subspace $A \subseteq X$ the *relative cohomology groups of (X, A)* are

$$H^n(X, A; R) = H^n(C^*(X, A; R)).$$

In other words, these singular cohomology groups are the cohomology R -modules (Definition E.0.2) of the singular cochain complex $C^*(X; R)$ and the relative singular cochain complex $C^*(X, A; R)$ (Definition 4.0.6) respectively.

5. CW COMPLEXES

Definition 5.0.1 (CW Complex). A **CW complex** is a cellular complex X that satisfies the following conditions:

- **Hausdorff:** X is a Hausdorff space (Definition 1.1.13).
- **C (Closure Finiteness):** The closure of each cell $\bar{e}_\alpha^n$ is contained in a finite union of cells of dimension less than or equal to n .
- **W (Weak Topology):** A subset $A \subseteq X$ is closed if and only if $A \cap \bar{e}_\alpha^n$ is closed for every cell e_α^n .

6. FIBRATIONS

6.1. Hurewicz fibrations.

Definition 6.1.1 (Homotopy Lifting Property). Let $p : E \rightarrow B$ be a continuous map. The map p is said to have the *homotopy lifting property (HLP) with respect to a topological space X* if, for every homotopy (Definition 3.0.1) $H : X \times I \rightarrow B$ and every map $f : X \rightarrow E$ such that $p \circ f = H(\cdot, 0)$, there exists a homotopy $\tilde{H} : X \times I \rightarrow E$ such that:

1. $\tilde{H}(\cdot, 0) = f$ (the lift starts at the given map);
2. $p \circ \tilde{H} = H$ (the diagram commutes).

In terms of diagrams, p satisfies the HLP with respect to X if every commutative solid square of the following form admits a dashed filler $\tilde{H}$:

$$\begin{array}{ccc} X \times \{0\} & \xrightarrow{f} & E \\ \downarrow & \nearrow \exists \tilde{H} & \downarrow p \\ X \times I & \xrightarrow{H} & B \end{array}$$

Definition 6.1.2. Let $p : E \rightarrow B$ be a continuous map of topological spaces (Definition 1.1.1). The map p is called a **Hurewicz fibration** (or simply a **fibration**) if it has the homotopy lifting property (Definition 6.1.1) with respect to all topological spaces X .

Equivalently, for every topological space X and every commutative diagram of the form:

$$\begin{array}{ccc} X \times \{0\} & \xrightarrow{f} & E \\ \downarrow i & \nearrow \exists \tilde{H} & \downarrow p \\ X \times I & \xrightarrow{H} & B \end{array}$$

where $I = [0, 1]$ is the unit interval and $i(x) = (x, 0)$, there exists a continuous map $\tilde{H} : X \times I \rightarrow E$ such that $p \circ \tilde{H} = H$ and $\tilde{H} \circ i = f$.

6.2. Serre fibrations.

Definition 6.2.1 (Serre Fibration). A continuous map (Definition 1.3.1) $p : E \rightarrow B$ of topological spaces (Definition 1.1.1) is called a **Serre fibration** (or a **weak fibration**) if it has the homotopy lifting property (Definition 6.1.1) with respect to all CW complexes (Definition 5.0.1).

Equivalently, for every $n \geq 0$, and every commutative diagram of the form:

$$\begin{array}{ccc} I^n \times \{0\} & \xrightarrow{f} & E \\ \downarrow i & \nearrow \exists \tilde{H} & \downarrow p \\ I^n \times I & \xrightarrow{H} & B \end{array}$$

where $I = [0, 1]$ is the unit interval and I^n is the n -dimensional cube, there exists a continuous map $\tilde{H} : I^n \times I \rightarrow E$ making the entire diagram commute.

Lemma 6.2.2. Any Hurewicz fibration (Definition 6.1.2) is a Serre fibration (Definition 6.2.1).

Proof. This holds trivially from the definitions — Hurewicz fibrations satisfy HLP (Definition 6.1.1) for all topological spaces whereas Serre fibrations satisfy HLP for all disks D^n . $\square$

6.3. Fiber bundles.

Definition 6.3.1. Let F be a topological space (Definition 1.1.1). A continuous map (Definition 1.3.1) $\pi : E \rightarrow B$ of topological spaces (Definition 1.1.1) is called a **fiber bundle with fiber F** if for every point $b \in B$, there exists an open neighborhood (Definition 1.1.9) $U \subseteq B$

containing b and a homeomorphism (Definition 1.3.2) $\phi : \pi^{-1}(U) \rightarrow U \times F$ such that the following diagram commutes:

$$\begin{array}{ccc} \pi^{-1}(U) & \xrightarrow{\phi} & U \times F \\ & \searrow \pi \quad \swarrow pr_1 & \\ & U & \end{array}$$

where $pr_1 : U \times F \rightarrow U$ is the projection onto the first factor. The space E is called the *total space*, B is the *base space*, and the homeomorphism ϕ is called a *local trivialization of the fiber bundle over U* .

It is also reasonable to write $F \hookrightarrow E \twoheadrightarrow B$ or simply $F \rightarrow E \rightarrow B$ to express the existence of a fiber bundle.

Definition 6.3.2. A topological space is *paracompact* if it is Hausdorff (Definition 1.1.13) and every open cover of X admits an open refinement that is locally finite.

See Theorem 6.3.3, which posits that nice enough topological spaces are paracompact.

Theorem 6.3.3. The following are paracompact (Definition 6.3.2) topological spaces:

1. Metric spaces (Definition 1.7.1) (Stone's theorem)
2. Topological manifolds
3. Compact (Definition 1.2.6) Hausdorff (Definition 1.1.13) spaces
4. CW-complexes (Definition 5.0.1)

Proposition 6.3.4. Let $\pi : E \rightarrow B$ be a fiber bundle (Definition 6.3.1) with fiber F . If B is a paracompact space (Definition 6.3.2), then π is a Hurewicz fibration (Definition 6.1.2).

Proposition 6.3.5. Let $\pi : E \rightarrow B$ be a fiber bundle (Definition 6.3.1). Then π is a Serre fibration (Definition 6.2.1).

APPENDIX A. SET THEORY

Definition A.0.1. Let A be a set.

- The set A is said to be *countably infinite* (or simply *countable*) if there exists a bijection $f : \mathbb{N} \rightarrow A$.
- The set A is said to be *finite* if there exists some $n \in \mathbb{N}$ and a bijection $g : \{1, 2, \dots, n\} \rightarrow A$.
- The set A is said to be *at most countable* if it is either finite or countably infinite.
- The set A is said to be *uncountable* if it is not at most countable.

Definition A.0.2. Let X and Y be sets and let $f : X \rightarrow Y$ be a function.

- The function f is said to be *injective* (or *one-to-one*) if for all $x_1, x_2 \in X$, $f(x_1) = f(x_2)$ implies $x_1 = x_2$.
- The function f is said to be *surjective* (or *onto*) if for every $y \in Y$ there exists $x \in X$ such that $f(x) = y$.

- The map f is *bijjective* if it is both injective and surjective. In this case, there exists a unique *inverse map* $f^{-1} : Y \rightarrow X$ such that for all $x \in X$ and $y \in Y$,

$$f^{-1}(f(x)) = x \text{ and } f(f^{-1}(y)) = y.$$

APPENDIX B. CATEGORY THEORY

Definition B.0.1 (Category). A *category* $\mathcal{C}$ consists of the following data:

- A class of *objects* denoted $\text{Ob}(\mathcal{C})$.
- For each pair of objects $X, Y \in \text{Ob}(\mathcal{C})$, a class

$$\text{Hom}_{\mathcal{C}}(X, Y)$$

of *morphisms* (also called *arrows* or *homs*). If the category $\mathcal{C}$ is clear, then this *hom-class* is also denoted by $\text{Hom}(X, Y)$. It may also be denoted by $\text{hom}_{\mathcal{C}}(X, Y)$ or $\text{hom}(X, Y)$, especially to distinguish from other types of hom's (e.g. internal hom's)

- For each triple of objects X, Y, Z , a composition law

$$\circ : \text{Hom}_{\mathcal{C}}(Y, Z) \times \text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{\mathcal{C}}(X, Z),$$

denoted $(g, f) \mapsto g \circ f$.

- For each object X , an *identity morphism*

$$\text{id}_X \in \text{Hom}_{\mathcal{C}}(X, X).$$

These data satisfy the following axioms:

- (Associativity) For all morphisms $f \in \text{Hom}_{\mathcal{C}}(X, Y)$, $g \in \text{Hom}_{\mathcal{C}}(Y, Z)$, and $h \in \text{Hom}_{\mathcal{C}}(Z, W)$,

$$h \circ (g \circ f) = (h \circ g) \circ f.$$

- (Identity) For all $f \in \text{Hom}_{\mathcal{C}}(X, Y)$,

$$\text{id}_Y \circ f = f = f \circ \text{id}_X.$$

One often writes $X \in \mathcal{C}$ synonymously with $X \in \text{Ob}(\mathcal{C})$, i.e. to denote that X is an object of $\mathcal{C}$.

We may call a category as above an *ordinary category* to distinguish this notion from the notions of *categories enriched in monoidal categories* or higher/ n -categories. (♠ **TODO: define n -categories**)

A category as defined above may be called called a *large category* or a *class category* to emphasize that the hom-classes may be proper classes rather than sets (note, however, that the possibility that hom-classes are sets is not excluded for large categories). Accordingly, a *category* may often refer to a locally small category (Definition B.0.2), which is a category whose hom-classes are all sets.

Definition B.0.2 (Locally small category). A (large) category (Definition B.0.1) $\mathcal{C}$ is called a *locally small category* if for every pair of objects $X, Y \in \text{Ob}(\mathcal{C})$, the collection $\text{Hom}_{\mathcal{C}}(X, Y)$ of morphisms between them is a (small) *set* (as opposed to a proper class). In other words, each hom-class is a set and may even be called a *hom-set*.

In some contexts, a locally small category may simply be called a *category*, especially when genuinely large categories are not considered.

A category $\mathcal{C}$ is called a *small category* if it is a locally small category and the class $\text{Ob}(\mathcal{C})$ of objects is a set.

Remark B.0.3. Many “concrete” categories considered in “classical mathematics” or outside of more “abstract” category theory tend to be locally small. For example, the categories of sets, groups, R -modules, vector spaces, topological spaces, schemes, manifolds, sheaves on “small enough” sites are all locally small.

APPENDIX C. GROUP THEORY

Definition C.0.1 (Groups). A *group* is a pair $(G, \cdot)$ where G is a set and $\cdot : G \times G \rightarrow G$ is a binary operation, subject to the following conditions:

1. (Associativity) For all $g, h, k \in G$ one has

$$(g \cdot h) \cdot k = g \cdot (h \cdot k).$$

2. (Identity element) There exists an element $e \in G$ such that for all $g \in G$,

$$e \cdot g = g \cdot e = g.$$

3. (Inverse element) For all $g \in G$ there exists an element $g^{-1} \in G$ such that

$$g \cdot g^{-1} = g^{-1} \cdot g = e.$$

The element e is called the *identity element of G* , and g^{-1} is called the *inverse of g* .

Equivalently, a group is a monoid with inverse elements.

A group $(G, \cdot)$ is often simply written as G , when the notation for the binary operation $\cdot$ is clear.

An *abelian group* or synonymously, a *commutative group*, is a group $(G, \cdot)$ whose binary operation $\cdot$ is *abelian* or *commutative*, i.e. satisfies

$$g \cdot h = h \cdot g$$

for all $g, h \in G$.

An abelian group is equivalent to a $\mathbb{Z}$ -module.

Definition C.0.2 (Group homomorphism). Let $(G, \cdot)$ and $(H, *)$ be groups (Definition C.0.1). A map $f : G \rightarrow H$ is called a **group homomorphism** if for all $g_1, g_2 \in G$ one has

$$f(g_1 \cdot g_2) = f(g_1) * f(g_2).$$

The collection of all groups with the group homomorphisms forms a locally small (Definition B.0.2) category (Definition B.0.1), called the **category of groups**.

If f is bijective (Definition A.0.2), then f is called a **group isomorphism**.

APPENDIX D. RING THEORY

Definition D.0.1. A **commutative (unital) ring** is a ring $(R, +, \cdot)$ such that $\cdot$ is a commutative operation, i.e. $a \cdot b = b \cdot a$.

For many writers (e.g. “commutative” algebraists or number theorists), a **ring** refers to a commutative ring as above.

Definition D.0.2. Let R be a not-necessarily commutative ring.

1. A **left R -module** is an abelian group $(M, +)$ together with an operation $R \times M \rightarrow M$, denoted $(r, m) \mapsto rm$, such that for all $r, s \in R$ and $m, n \in M$:
 - $r(m + n) = rm + rn$,
 - $(r + s)m = rm + sm$,
 - $(rs)m = r(sm)$,
 - $1_R m = m$ where 1_R is the multiplicative identity of R .
2. A **right R -module** is defined similarly as an abelian group $(M, +)$ with an operation $M \times R \rightarrow M$, denoted $(m, r) \mapsto mr$, such that for all $r, s \in R$ and $m, n \in M$:
 - $(m + n)r = mr + nr$,
 - $m(r + s) = mr + ms$,
 - $m(rs) = (mr)s$,
 - $m1_R = m$.
3. Let R and S be (not necessarily commutative) rings.

An **R - S -bimodule** (or an **R - S -module** or an (R, S) -module, etc.) is an abelian group (Definition C.0.1) $(M, +)$ equipped with

- (a) a left action of R :

$$R \times M \rightarrow M, \quad (r, m) \mapsto r \cdot m,$$

making M a left R -module (Definition D.0.2),

- (b) a right action of S :

$$M \times S \rightarrow M, \quad (m, s) \mapsto m \cdot s,$$

making M a right S -module,

such that the left and right actions commute; that is, for all $r \in R$, $s \in S$, and $m \in M$,

$$r \cdot (m \cdot s) = (r \cdot m) \cdot s.$$

4. A **two-sided R -module** (or **R -bimodule**) is an R - R -bimodule.

If R is a commutative ring (Definition D.0.1), then a left/right R -module can automatically be regarded as a two-sided R -module. As such, we simply talk about **R -modules** in this case.

Any abelian group is equivalent to a two-sided $\mathbb{Z}$ -module. Moreover, any left R -module is equivalent to an $R - \mathbb{Z}$ -bimodule (Definition D.0.2) and any right R -module is equivalent to an $\mathbb{Z} - R$ -bimodule (Definition D.0.2). Given a left/right/two-sided R -module, its ***natural bimodule structure*** will refer to its structure as a $R - \mathbb{Z} / \mathbb{Z} - R / R - R$ bimodule. In this way, many definitions associated with the notions of left/right/two-sided R -modules can be defined as special cases for definitions for R - S -bimodules.

Definition D.0.3 (Hom of left/right/bi-modules). Let R, S, T be (not necessarily commutative) rings.

1. Let M and N be left R -modules (Definition D.0.2). The ***homomorphism group of left R -modules from M to N*** is the abelian group

$$\text{Hom}(M, N) = \text{Hom}_R(M, N) = \text{Hom}_{R-\mathbb{Z}}(M, N) := \{f : M \rightarrow N \mid f \text{ is a left } R\text{-module homomorphism}\}.$$

2. Let M and N be right R -modules (Definition D.0.2). The ***homomorphism group of right R -modules from M to N*** is the abelian group

$$\text{Hom}(M, N) = \text{Hom}_R(M, N) = \text{Hom}_{\mathbb{Z}-R}(M, N) := \{f : M \rightarrow N \mid f \text{ is a right } R\text{-module homomorphism}\}.$$

3. Let S be a (not necessarily commutative ring) and let M and N be $R - S$ -bimodules (Definition D.0.2). The ***homomorphism group of R - S -bimodules from M to N*** is the abelian group

$$\text{Hom}(M, N) = \text{Hom}_{R-S}(M, N) := \{f : M \rightarrow N \mid f \text{ is a } R - S\text{-bimodule homomorphism}\}$$

In each case, $\text{Hom}(M, N)$ has a natural structure of an ***abelian group*** given by ***pointwise addition***: for $f, g \in \text{Hom}(M, N)$,

$$(f + g)(m) := f(m) + g(m),$$

and the zero morphism **0** given by $0(m) := 0_N$ acts as the identity element. The additive inverse **$-f$** is defined by $(-f)(m) := -f(m)$. Moreover, depending on bi-module structures that M and N may be carrying, $\text{Hom}(M, N)$ may itself carry additional module structures:

- In case that M is a $R - S$ -bimodule and N is a $R - T$ -bimodule, $\text{Hom}_R(M, N)$, the group of left R -module homomorphisms, is an $S - T$ -bimodule as follows:

$$(s \cdot f \cdot t)(m) = f(m \cdot s) \cdot t \quad f \in \text{Hom}_R(M, N), s \in S, t \in T.$$

- Dually, in case that M is a $S - R$ -bimodule and N is a $T - R$ -bimodule, $\text{Hom}_R(M, N)$, the group of right R -module homomorphisms, is an $S - T$ -bimodule as follows:

$$(s \cdot f \cdot t)(m) = f(s \cdot m) \cdot t \quad f \in \text{Hom}_R(M, N), s \in S, t \in T.$$

Some cases of interest may be when R, S , or T is in fact $\mathbb{Z}$ — these allow us to see module structures on $\text{Hom}(M, N)$ even when M and N are one-sided modules.

(♠ TODO: state this as a theorem) We furthermore note that $\text{Hom}_R(-, -)$ yields biadditive functors

$$\begin{aligned}\text{Hom}_R(-, -) &: {}_R\mathbf{Mod}_S^{\text{op}} \times {}_R\mathbf{Mod}_T \rightarrow {}_S\mathbf{Mod}_T \\ \text{Hom}_R(-, -) &: {}_S\mathbf{Mod}_R^{\text{op}} \times {}_T\mathbf{Mod}_R \rightarrow {}_S\mathbf{Mod}_T.\end{aligned}$$

APPENDIX E. HOMOLOGICAL ALGEBRA

Definition E.0.1 (Chain complex in a preadditive category). Let $\mathcal{A}$ be a preadditive category and let I be a totally ordered set (typically $\mathbb{Z}$, but $I \subseteq \mathbb{Z}$ is also allowed).

1. A *chain complex* (or *homological (chain) complex*) $(K_\bullet, d_\bullet)$ in $\mathcal{A}$ indexed by I is the homological convention for sequences with decreasing degrees. It consists of:
 - Objects $\{K_i\}_{i \in I}$ in $\mathcal{A}$, called the *terms in degree i* ,
 - Morphisms $d_i : K_i \rightarrow K_{i-1}$ in $\mathcal{A}$, called the *boundary maps* or *differentials in degree i* ,
 such that for every $i \in I$, $d_{i-1} \circ d_i = 0$. That is,

$$K_\bullet : \cdots \xrightarrow{d_{i+1}} K_i \xrightarrow{d_i} K_{i-1} \xrightarrow{d_{i-1}} K_{i-2} \rightarrow \cdots$$

with $d_{i-1}d_i = 0$ for all i . We typically use the notation $K_\bullet = (K_i, d_i)_{i \in I}$.

2. Dually, a *cochain complex* (or *cohomological complex*) $(K^\bullet, d^\bullet)$ in $\mathcal{A}$ follows the cohomological convention with increasing degrees. It consists of objects $\{K^i\}_{i \in I}$ and *coboundary maps* $d^i : K^i \rightarrow K^{i+1}$ such that $d^{i+1} \circ d^i = 0$:

$$K^\bullet : \cdots \xrightarrow{d^{i-1}} K^i \xrightarrow{d^i} K^{i+1} \xrightarrow{d^{i+1}} K^{i+2} \rightarrow \cdots$$

We typically use the notation $K^\bullet = (K^i, d^i)_{i \in I}$.

3. Let $K_\bullet = (K_i, d_i^K)$ and $L_\bullet = (L_i, d_i^L)$ be chain complexes (Definition E.0.1) in $\mathcal{A}$ indexed by the same set I . A *morphism of chain complexes* (or *chain map*)

$$f_\bullet : K_\bullet \rightarrow L_\bullet$$

consists of morphisms $f_i : K_i \rightarrow L_i$ for all $i \in I$, such that for every $i \in I$, the following diagram commutes:

$$\begin{array}{ccc} K_i & \xrightarrow{d_i^K} & K_{i-1} \\ \downarrow f_i & & \downarrow f_{i-1} \\ L_i & \xrightarrow{d_i^L} & L_{i-1} \end{array}$$

i.e., $d_i^L \circ f_i = f_{i-1} \circ d_i^K$.

A *morphism of cochain complexes* $f^\bullet : K^\bullet \rightarrow L^\bullet$ is defined similarly, satisfying the commutativity condition $d_L^i \circ f^i = f^{i+1} \circ d_K^i$.

The collection of these objects and morphisms forms a category. Notation for these categories is as follows:

- $\mathbf{Ch}(\mathcal{A})$ or $\mathbf{Ch}(\mathcal{A})$ is often used as a general term.
- To be explicit about the indexing convention, one uses $\mathbf{Ch}_\bullet(\mathcal{A})$ for chain complexes and $\mathbf{Ch}^\bullet(\mathcal{A})$ (or sometimes $\mathbf{CoCh}(\mathcal{A})$) for cochain complexes.

- The set of chain maps between two complexes is denoted by $\mathrm{Hom}_{\mathrm{Ch}(\mathcal{A})}(K_\bullet, L_\bullet)$; it is an abelian group under pointwise addition $(f + g)_i = f_i + g_i$.

Notations used to write chain complexes are *homological conventions* whereas notations used to write cochain complexes are *cohomological conventions*. In homological conventions, indices are written as subscripts and decrease when “following the differential maps”. In cohomological conventions, indices are written as superscripts and increase when “following the differential maps”. The theories of chain complexes and of cochain complexes are not genuinely different, however, as the conventions are essentially notational — starting from a chain complex $(K_\bullet, d_\bullet)$, one can obtain a cochain complex $(K^\bullet, d^\bullet)$ simply by defining $K^i = K_{-i}$ and d^i as d_{-i} and vice versa. As such, cochain complexes may be simply called *chain complexes* when careful distinctions between the conventions are unnecessary. Homological and cohomological notational conventions persist throughout homological algebra, see e.g. spectral sequences, double complexes, etc, but again the distinction between such conventions are merely notational.

Definition E.0.2 (Chain complexes and their (co)homology objects). Let $\mathcal{A}$ be an abelian category.

- For a cochain complex $K^\bullet$, its *cohomology object in degree i* is defined as the quotient of the object of i -cocycles by the object of i -coboundaries:

$$H^i(K^\bullet) := Z^i(K)/B^i(K) = \ker(d^i)/\mathrm{im}(d^{i-1}).$$

In fact, each H^i is a functor $H^i : \mathbf{Ch}^\bullet(\mathcal{A}) \rightarrow \mathcal{A}$ (Definition E.0.1); given a morphism $f^\bullet : K^\bullet \rightarrow L^\bullet$ of cochain complexes, it is thus customary to write $H^i(f^\bullet) : H^i(K^\bullet) \rightarrow H^i(L^\bullet)$ for the induced morphism on the i th cohomology objects.

- For a chain complex (Definition E.0.1) $K_\bullet$, its *homology object in degree i* is defined as the quotient of the object of i -cycles by the object of i -boundaries:

$$H_i(K_\bullet) := Z_i(K)/B_i(K) = \ker(d_i)/\mathrm{im}(d_{i+1}).$$

In fact, each H_i is a functor $H_i : \mathbf{Ch}_\bullet(\mathcal{A}) \rightarrow \mathcal{A}$ (Definition E.0.1); given a morphism $f_\bullet : K_\bullet \rightarrow L_\bullet$ of chain complexes, it is thus customary to write $H_i(f_\bullet) : H_i(K_\bullet) \rightarrow H_i(L_\bullet)$ for the induced morphism on the i th homology objects.

REFERENCES

- [AP07] Jeffrey D. Achter and Rachel Pries. The integral monodromy of hyperelliptic and trielliptic curves. *Mathematische Annalen*, 338(1):187–206, 2007.
- [Arm92] Brumer Armand. The average rank of elliptic curves i. *Inventiones mathematicae*, 109(1):445–472, 1992.
- [Ayo23] Joseph Ayoub. Counterexamples to F. Morel’s conjecture on $\pi_0^{\geq 1}$. *Comptes Rendus. Mathématique*, 361:1087–1090, 2023.
- [BBD82] Alexander A. Beilinson, Joseph Bernstein, and Pierre Deligne. Analyse et topologie sur les espaces singuliers (i). *Astérisque*, 100, 1982.
- [BBDG18] Alexander A. Beilinson, Joseph Bernstein, Pierre Deligne, and Ofer Gabber. *Faisceaux pervers*, volume 4. Société mathématique de France Paris, 2018.

- [BBK⁺23] Barinder S. Banwait, Armand Brumer, Hyun Jong Kim, Zev Klagsbrun, Jacob Mayle, Padmavathi Srinivasan, and Isabel Vogt. Computing nonsurjective primes associated to galois representations of genus 2 curves. *LuCaNT: LMFDB, Computation, and Number Theory*, 796:129, 2023.
- [BC19] Tilman Bauer and Magnus Carlson. Tensor products of affine and formal abelian groups. *Documenta Mathematica*, 24:2525–2582, 2019.
- [BFK⁺17] Valentin Blomer, Étienne Fouvry, Emmanuel Kowalski, Philippe Michel, and Djordje Milićević. Some applications of smooth bilinear forms with kloosterman sums. *Proceedings of the Steklov Institute of Mathematics*, 296:18–29, 2017.
- [BGI71] Pierre Berthelot, Alexander Grothendieck, and Luc Illusie. *Théorie des Intersections et Théorème de Riemann-Roch (SGA6)*, volume 225 of *Lecture Notes in Mathematics*. Springer-Verlag, 1971.
- [BH12] Salman Baig and Chris Hall. Experimental data for goldfeld’s conjecture over function fields. *Experimental Mathematics*, 21(4):362–374, 2012.
- [BLGHT11] Tom Barnet-Lamb, David Geraghty, Michael Harris, and Richard Taylor. A family of Calabi–Yau varieties and potential automorphy ii. *Publications of the Research Institute for Mathematical Sciences*, 47(1):29–98, 2011.
- [Bor12] Armand Borel. *Linear algebraic groups*, volume 126 of *Graduate Texts in Mathematics*. Springer New York, 2012.
- [BS15a] Manjul Bhargava and Arul Shankar. Binary quartic forms having bounded invariants, and the boundedness of the average rank of elliptic curves. *Annals of Mathematics*, pages 191–242, 2015.
- [BS15b] Manjul Bhargava and Arul Shankar. Ternary cubic forms having bounded invariants, and the existence of a positive proportion of elliptic curves having rank 0. *Annals of Mathematics*, pages 587–621, 2015.
- [BS15c] Bhargav Bhatt and Peter Scholze. The pro-étale topology for schemes. *Astérisque*, 360:99–201, 2015.
- [BSD65] Bryan John Birch and Peter Francis Swinnerton-Dyer. Notes on elliptic curves. ii. *Journal für die reine und angewandte Mathematik*, 218:79–108, 1965.
- [BSS18] Bhargav Bhatt, Christian Schnell, and Peter Scholze. Vanishing theorems for perverse sheaves on abelian varieties, revisited. *Selecta Mathematica*, 24:63–84, 2018.
- [Cho08] Utsav Choudhury. Homotopy theory of schemes and a^1 -fundamental groups. Master’s thesis, Università degli Studi di Padova, 2008.
- [CHT08] Laurent Clozel, Michael Harris, and Richard Taylor. Automorphy for some l -adic lifts of automorphic mod l galois representations. *Publications mathématiques*, 108:1–181, 2008.
- [DA73] Pierre Deligne and Michael Artin. *Théorie des Topos et Cohomologies Étale des Schémas. Séminaire de Géométrie Algébrique due Bois-Marie 1963-1964 (SGA 4)*. Lecture Notes in Mathematics. Springer Berlin, 1973.
- [DBG⁺77] Pierre Deligne, Jean-François Boutot, Alexander Grothendieck, Luc Illusie, and Jean-Louis Verdier. *Étale Cohomology. Séminaire de Géométrie Algébrique due Bois-Marie 1963-1964 (SGA 4 1/2)*. Lecture Notes in Mathematics. Springer-Verlag, 1977.
- [Del80] Pierre Deligne. La conjecture de Weil : II. *Publications Mathématiques de l’IHÉS*, 52:137–252, 1980.
- [Del89] Pierre Deligne. Le groupe fondamental de la droite projective moins trois points. In *Galois Groups over $\mathbb{Q}$: Proceedings of a Workshop Held March 23–27, 1987*, pages 79–297. Springer, 1989.
- [Die02] Luis V. Dieulefait. Explicit determination of the images of the Galois representations attached to abelian surfaces with $\text{End}(A) = \mathbb{Z}$. *Experiment. Math.*, 11(4):503–512 (2003), 2002.
- [DR04] Luis V. Dieulefait and Victor Rotger. The arithmetic of qm -abelian surfaces through their galois representations. *Journal of Algebra*, 281:124–143, 2004.
- [Dri89] Vladimir Gershonovich Drinfeld. Cohomology of compactified manifolds of modules of f -sheaves. *Journal of Soviet Mathematics*, 46(2):1789–1821, 1989.
- [DZ19] Alexander Dunn and Alexandru Zaharescu. Sums of Kloosterman sums over primes in an arithmetic progression. *The Quarterly Journal of Mathematics*, 70(1):319–342, 2019.

- [Eke07] Torsten Ekedahl. *On The Adic Formalism*, pages 197–218. Birkhäuser Boston, Boston, MA, 2007.
- [ELS20] Jordan S. Ellenberg, Wanlin Li, and Mark Shusterman. Nonvanishing of hyperelliptic zeta functions over finite fields. *Algebra & Number Theory*, 14(7):1895–1909, 8 2020.
- [EVW16] Jordan S. Ellenberg, Akshay Venkatesh, and Craig Westerland. Homology stability for Hurwitz spaces and the Cohen-Lenstra conjecture over function fields. *Annals of Mathematics*, 183:729–786, 2016.
- [FFK25] Arthur Forey, Javier Fresán, and Emmanuel Kowalski. Arithmetic fourier transforms over finite fields: generic vanishing, convolution, and equidistribution, 2025.
- [FK12] Sergey Finashin and Viatcheslav Kharlamov. Abundance of Real Lines on Real Projective Hypersurfaces. *International Mathematics Research Notices*, 2013(16):3639–3646, 06 2012.
- [FLR23] Tony Feng, Aaron Landesman, and Eric M. Rains. The geometric distribution of Selmer groups of elliptic curves over function fields. *Mathematische Annalen*, 387:615–687, 2023.
- [Fu15] Lei Fu. *Etale Cohomology Theory*, volume 14 of *Nankai Tracts in Mathematics*. World Scientific, 2015.
- [FvdG04] Carel Faber and Gerard van der Geer. Complete subvarieties of moduli spaces and the Prym map. *Journal für die reine und angewandte Mathematik*, 2004(573):117–137, 2004.
- [GL96] Ofer Gabber and François Loeser. Faisceaux pervers ℓ -adiques sur un tore. *Duke Math J.*, 83(3):501–606, 1996.
- [Gol06] Dorian Goldfeld. Conjectures on elliptic curves over quadratic fields. In *Number Theory Carbon-dale 1979: Proceedings of the Southern Illinois Number Theory Conference Carbondale, March 30 and 31, 1979*, pages 108–118. Springer, 2006.
- [GR04] Alexander Grothendieck and Michèle Raynaud. Revêtements étales et groupe fondamental (SGA 1). eprint arXiv matyh/0206203, 2004. Updated edition of the book of the same title published by Springer-Verlag in 1971 as volume 224 of the series Lecture Notes in Mathematics.
- [Gro77] Alexander Grothendieck. *Cohomologie l -adique et fonctions L* Séminaire de Géométrie Algébrique due Bois-Marie 1965-1966 (SGA 5), volume 589 of *Springer Lecture Notes*. Springer-Verlag, 1977. Avec la collaboration de I. Bucur, C. Houzel, L. Illusie, J.-P. Jouanolou, et J.-P. Serre.
- [GV72] Alexander Grothendieck and Jean-Louis Verdier. *Theorie des Topos et Cohomologie Etale des Schemas. Seminaire de Geometrie Algebrique du Bois-Marie 1963-1964 (SGA 4)*. Lecture Notes in mathematics. Springer-Verlag Berlin Heidelberg, 1 edition, 1972.
- [HB04] David Heath-Brown. The average analytic rank of elliptic curves. *Duke Math. J.*, 122:591–623, 2004.
- [HK25] Chris Hall and Hyun Jong Kim. Independence of ℓ (title to be determined). In progress, 2025.
- [HM73] Dale Husemoller and John Milnor. *Symmetric Bilinear Forms*, volume 73 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 2. Folge*. Springer Berlin Heidelberg, 1973.
- [HSBT05] Michael Harris, Nick Shepherd-Barron, and Richard Taylor. Ihara’s lemma and potential automorphy, 2005.
- [Hub97] Annette Huber. Mixed perverse sheaves for schemes over number fields. *Compositio Mathematica*, 108:107–121, 1997.
- [Ibu22] Tomoyoshi Ibukiyama. Supersingular loci of low dimensions and parahoric subgroups. *Osaka Journal of Mathematics*, 59:703–726, 2022.
- [Jor05] Andrei Jorza. The birch and swinnerton-dyer conjecture for abelian varieties over number fields, 2005.
- [Kat90] Nicholas M. Katz. *Exponential sums and differential equations*, volume 124 of *Annals of Mathematics Studies*. Princeton University Press, 1990.
- [Kat96] Nicholas M. Katz. *Rigid Local Systems*, volume 139 of *annals of Mathematics Studies*. Princeton University Press, 1996.
- [Kat98] Nicholas M. Katz. *Gauss Sums, Kloosterman Sums, and Monodromy Groups*, volume 116 of *Annals of Mathematics Studies*. Princeton University Press, 1998.
- [Kat12] Nicholas M. Katz. *Convolution and Equidistribution Sato-Tate Theorems for Finite-Field Mellin Transforms*, volume 180 of *Annals of Mathematics Studies*. Princeton University Press, 2012.

- [Kim23] Hyun Jong Kim. `trouver`, 2023. GitHub repository: <https://github.com/hyunjongkimmath/trouver>.
- [Kim24] Hyun Jong Kim. *Cohen-Lenstra heuristics and vanishing of zeta functions for superelliptic curves over finite fields*. PhD thesis, University of Wisconsin-Madison, 2024.
- [KL85] Nicholas M. Katz and Gérard Laumon. Transformation de fourier et majoration de sommes exponentielles. *Publications Mathématiques de l’IHÉS*, 62:145–202, 1985.
- [KLSW23] Jesse Leo Kass, Marc Levine, Jake P. Solomon, and Kirsten Wickelgren. A quadratically enriched count of rational curves. arXiv 2307.01936, 2023.
- [KM23] Seoyoung Kim and M. Ram Murty. From the Birch and Swinnerton-Dyer conjecture to Nagao’s conjecture. *Mathematics of Computation*, 92(339):385–408, 2023.
- [KMS17] Emmanuel Kowalski, Philippe Michel, and Will Sawin. Bilinear forms with Kloosterman sums and applications. *Annals of Mathematics*, 186:413–500, 2017.
- [KP24] Hyun Jong Kim and Sun Woo Park. Global A^1 degrees of covering maps between modular curves, 2024.
- [Krä14] Thomas Krämer. Perverse sheaves on semiabelian varieties. *Rendiconti del Seminario Matematico della Università di Padova*, 132:83–102, 2014.
- [KS99] Nicholas M. Katz and Peter Sarnak. *Random matrices, Frobenius eigenvalues, and monodromy*, volume 45. American Mathematical Society, 1999.
- [KS22] Timo Keller and Michael Stoll. Exact verification of the strong bsd conjecture for some absolutely simple abelian surfaces. *Comptes Rendus Mathématique*, 360:483–489, 2022.
- [KW13] Reinhardt Kiehl and Rainer Weissauer. *Weil Conjectures, Perverse Sheaves and ℓ -adic Fourier Transform*, volume 42 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics*. Springer Berlin, Heidelberg, 2013.
- [KW15] Thomas Krämer and Rainer Weissauer. Vanishing theorems for constructible sheaves on abelian varieties. *J. Algebraic Geometry*, 24:531–568, 2015.
- [KW19] Jesse Leo Kass and Kirsten Wickelgren. The class of Eisenbud-Khimshiashvili-Levine is the local A^1 – Brouwer degree. *Duke Mathematical Journal*, 168(3):429–469, 2019.
- [KW21] Jesse Leo Kass and Kirsten Wickelgren. An arithmetic count of the lines on a smooth cubic surface. *Compositio Mathematica*, 157(4):677–709, 2021.
- [Laf02] Laurent Lafforgue. Chtoucas de Drinfeld et correspondance de Langlands. *Inventiones mathematicae*, 147:1–241, 2002.
- [Lom17] Davide Lombardo. Galois representations attached to abelian varieties of cm type. *Bulletin de la Société mathématique de France*, 145(3):469–501, 2017.
- [May99] Jon Peter May. *A Concise Course in Algebraic Topology*. Chicago Lectures in Mathematics. University of Chicago Press, 1999.
- [Mil80] James S. Milne. *Etale cohomology*. Number 33 in Princeton Mathematical Series. Princeton university press, 1980.
- [Mil07] James S. Milne. Quotients of Tannakian categories. *Theory and Applications of Categories*, 18(21):654–664, 2007.
- [Mil13] James S. Milne. Lie algebras, algebraic groups, and lie groups, 2013. Available at www.jmilne.org/math/.
- [Mil17] James S. Milne. *Algebraic Groups*, volume 170 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 2017.
- [Mit14] Howard H. Mitchell. The subgroups of the quaternary abelian linear group. *Trans. Amer. Math. Soc.*, 15(4):379–396, 1914.
- [Mor06] Fabien Morel. A^1 -algebraic topology. In *International Congress of Mathematicians*, volume 2, pages 1035–1059, 2006.
- [Mor12] Fabien Morel. *A^1 -Algebraic Topology over a field*. Lecture Notes in Mathematics. Springer Berlin, Heidelberg, 2012.
- [MV99] Fabien Morel and Vladimir Voevodsky. A^1 -homotopy theory of schemes. *Publications Mathématiques de l’IHÉS*, 90:45–143, 1999.
- [Nag97] Koh-ichi Nagao. $Q(t)$ -rank of elliptic curves and certain limit coming from the local points. *Manuscripta mathematica*, 92(1):13–32, 1997.

- [nLa25a] nLab authors. geometric morphism. <https://ncatlab.org/nlab/show/geometric+morphism>, July 2025. Revision 61.
- [nLa25b] nLab authors. homotopy group of a spectrum. <https://ncatlab.org/nlab/show/homotopy+group+of+a+spectrum>, June 2025. Revision 7.
- [nLa25c] nLab authors. Introduction to Stable homotopy theory – 1-1. <https://ncatlab.org/nlab/show/Introduction+to+Stable+homotopy+theory+--+1-1>, June 2025. Revision 43.
- [nLa25d] nLab authors. model structure on topological sequential spectra. <https://ncatlab.org/nlab/show/model+structure+on+topological+sequential+spectra>, June 2025. Revision 61.
- [nLa25e] nLab authors. point of a topos. <https://ncatlab.org/nlab/show/point+of+a+topos>, July 2025. Revision 53.
- [nLa25f] nLab authors. sheafification. <https://ncatlab.org/nlab/show/sheafification>, September 2025. Revision 40.
- [nLa25g] nLab authors. stable homotopy category. <https://ncatlab.org/nlab/show/stable+homotopy+category>, June 2025. Revision 31.
- [OT14] Christian Okonek and Andrei Teleman. Intrinsic signs and lower bounds in real algebraic geometry. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2014(688):219–241, 2014.
- [Poo18] Bjorn Poonen. Heuristics for the arithmetic of elliptic curves. In *Proceedings of the International Congress of Mathematicians: Rio de Janeiro 2018*, pages 399–414. World Scientific, 2018.
- [PR12] BJORN POONEN and ERIC RAINS. Random maximal isotropic subspaces and selmer groups. *Journal of the American Mathematical Society*, 25(1):245–269, 2012.
- [Pri24] Rachel Pries. The Torelli locus and Newton polygons, 2 2024. Lecture Notes for the 2024 Arizona Winter School.
- [Pri25] Rachel Pries. Some cases of Oort’s conjecture about Newton polygons of curves. *Nagoya Mathematical Journal*, 257:93–103, 2025.
- [PW21] Sabrina Pauli and Kirsten Wickelgren. Applications to $\mathbb{A}^1$ -enumerative geometry of the $\mathbb{A}^1$ -degree. *Research in the Mathematical Sciences*, 8(24):24–29, 2021.
- [Ros02] Michael Rosen. *Number Theory in Function Fields*, volume 210 of *Graduate Texts in Mathematics*. Springer-Verlag New York, 2002.
- [RS98] Michael Rosen and Joseph H. Silverman. On the rank of an elliptic surface. *Inventiones mathematicae*, 133:43–67, 1998.
- [Rud87] Walter Rudin. *Real and Complex Analysis*. Mathematics Series. McGraw-Hill Book Company, 3 edition, 1987.
- [Saw24] Will Sawin. General multiple Dirichlet series from perverse sheaves. *Journal of Number Theory*, 262:408–453, 2024.
- [Ser72] Jean-Pierre Serre. Propriétés galoisiennes des points d’ordre fini des courbes elliptiques. *Invent. math.*, 15:259–331, 1972.
- [Ser00] Jean-Pierre Serre. Lettre à Marie-France Vignéras du 10/2/1986. *Oeuvres–Collected Papers*, 4:38–55, 2000.
- [SFFK23] Will Sawin, Arthur Forey, Javier Fresán, and Emmanuel Kowalski. Quantitative sheaf theory. *Journal of the American Mathematical Society*, 36(3):653–726, 2023.
- [Sil89] Joseph H. Silverman. Elliptic curves of bounded degree and height. *Proceedings of the American Mathematical Society*, 105(3):540–545, 1989.
- [Sil09] Joseph H. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Graduate Texts in Mathematics*. Springer-Verlag, 2 edition, 2009.
- [ST21] Will Sawin and Jacob Tsimerman. Bounds for the stalks of perverse sheaves in characteristic p and a conjecture of Shende and Tsimerman. *Inventiones mathematicae*, 224(1):1–32, 2021.
- [Sta25] The Stacks project authors. The stacks project. <https://stacks.math.columbia.edu>, 2025.
- [Tat65] John T. Tate. Algebraic cycles and poles of zeta functions. In *Arithmetical Algebraic Geometry (Proc. Conf. Purdue Univ., 1963)*, pages 93–110. Harper & Row, 1965. Also in *Collected works of John Tate* (2 vols.), Amer. Math. Soc. (2016), vol. 2.

- [Tat66] John Tate. On the conjectures of Birch and Swinnerton-Dyer and a geometric analog. In *Séminaire Bourbaki : années 1964/65 1965/66, exposés 277-312*, number 9 in Astérisque, pages 415–440. Société mathématique de France, 1966. talk:306.
- [Tay08] Richard Taylor. Automorphy for some l -adic lifts of automorphic mod l galois representations. ii. *Publications mathématiques*, 108:183–239, 2008.
- [Voe98] Vladimir Voevodsky. A1-homotopy theory. In *Proceedings of the international congress of mathematicians*, volume 1, pages 579–604. Berlin, 1998.
- [Wei94] Charles A. Weibel. *An Introduction to Homological Algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 1994. First paperback edition 1995 Reprinted 1997.
- [Wik25] Wikipedia contributors. Frobenius endomorphisms#frobenius for schemes — Wikipedia, the free encyclopedia, 2025. [Online; accessed 08-July-2025].
- [You06] Matthew Young. Low-lying zeros of families of elliptic curves. *Journal of the American Mathematical Society*, 19(1):205–250, 2006.
- [Yu97] Jiu-Kang Yu. Toward a proof of the Cohen-Lenstra conjecture in the function field case. preprint, 1997.