

# Incremental learning, game theory, and applications

## Lecture 1: Finite zero-sum games

Rida Laraki and Guillaume Vigeral

Master IASD, 2024

- 1 Introduction ; value in pure strategies
- 2 Value in Mixed Strategies (in the finite case)
- 3 Learning to Play Optimal

- 1 Introduction ; value in pure strategies
- 2 Value in Mixed Strategies (in the finite case)
- 3 Learning to Play Optimal

Zero-sum games are two-person games where the players have opposite evaluations of outcomes, hence the sum of the payoff functions is 0.

## Definition

Zero-sum games are two-person games where the players have opposite evaluations of outcomes, hence the sum of the payoff functions is 0.

### Definition

A *zero-sum game*  $G$  in *strategic form* is defined by a triple  $(I, J, g)$ , where  $I$  (resp.  $J$ ) is the non-empty set of strategies of player 1 (resp. player 2) and  $g : I \times J \rightarrow \mathbb{R}$  is the payoff function of player 1.

Zero-sum games are two-person games where the players have opposite evaluations of outcomes, hence the sum of the payoff functions is 0.

### Definition

A *zero-sum game*  $G$  in *strategic form* is defined by a triple  $(I, J, g)$ , where  $I$  (resp.  $J$ ) is the non-empty set of strategies of player 1 (resp. player 2) and  $g : I \times J \rightarrow \mathbb{R}$  is the payoff function of player 1.

The interpretation is as follows :

Zero-sum games are two-person games where the players have opposite evaluations of outcomes, hence the sum of the payoff functions is 0.

### Definition

A *zero-sum game*  $G$  in *strategic form* is defined by a triple  $(I, J, g)$ , where  $I$  (resp.  $J$ ) is the non-empty set of strategies of player 1 (resp. player 2) and  $g : I \times J \rightarrow \mathbb{R}$  is the payoff function of player 1.

The interpretation is as follows :

- Player 1 chooses  $i$  in  $I$  and player 2 chooses  $j$  in  $J$ , in an independent way.

Zero-sum games are two-person games where the players have opposite evaluations of outcomes, hence the sum of the payoff functions is 0.

### Definition

A *zero-sum game*  $G$  in *strategic form* is defined by a triple  $(I, J, g)$ , where  $I$  (resp.  $J$ ) is the non-empty set of strategies of player 1 (resp. player 2) and  $g : I \times J \rightarrow \mathbb{R}$  is the payoff function of player 1.

The interpretation is as follows :

- Player 1 chooses  $i$  in  $I$  and player 2 chooses  $j$  in  $J$ , in an independent way.
- The payoff of player 1 is then  $g(i, j)$  and that of player 2 is  $-g(i, j)$ .



Zero-sum games are two-person games where the players have opposite evaluations of outcomes, hence the sum of the payoff functions is 0.

### Definition

A zero-sum game  $G$  in strategic form is defined by a triple  $(I, J, g)$ , where  $I$  (resp.  $J$ ) is the non-empty set of strategies of player 1 (resp. player 2) and  $g : I \times J \rightarrow \mathbb{R}$  is the payoff function of player 1.

The interpretation is as follows :

- Player 1 chooses  $i$  in  $I$  and player 2 chooses  $j$  in  $J$ , in an independent way.
- The payoff of player 1 is then  $g(i, j)$  and that of player 2 is  $-g(i, j)$ .
- Player 1 wants to maximize  $g$  and is called the maximizing player. Player 2 is the minimizing player.

Zero-sum games are two-person games where the players have opposite evaluations of outcomes, hence the sum of the payoff functions is 0.

## Definition

A *zero-sum game*  $G$  in *strategic form* is defined by a triple  $(I, J, g)$ , where  $I$  (resp.  $J$ ) is the non-empty set of strategies of player 1 (resp. player 2) and  $g : I \times J \rightarrow \mathbb{R}$  is the payoff function of player 1.

The interpretation is as follows :

- Player 1 chooses  $i$  in  $I$  and player 2 chooses  $j$  in  $J$ , in an independent way.
- The payoff of player 1 is then  $g(i, j)$  and that of player 2 is  $-g(i, j)$ .
- Player 1 wants to maximize  $g$  and is called the maximizing player. Player 2 is the minimizing player.
- With the notations of the introduction, the strategy sets are  $S^1 = I$  and  $S^2 = J$  and the payoff functions are  $g^1 = g = -g^2$ .

- $G = (I, J, g)$  is a *finite* zero-sum game when  $I$  and  $J$  are finite.
- The game is then represented by an  $I \times J$  matrix  $A$ , where player 1 chooses the row  $i \in I$ , player 2 chooses the column  $j \in J$  and the entry  $A_{ij}$  of the matrix corresponds to the payoff  $g(i, j)$ .

- $G = (I, J, g)$  is a *finite* zero-sum game when  $I$  and  $J$  are finite.
- The game is then represented by an  $I \times J$  matrix  $A$ , where player 1 chooses the row  $i \in I$ , player 2 chooses the column  $j \in J$  and the entry  $A_{ij}$  of the matrix corresponds to the payoff  $g(i, j)$ .

## 'Matching Pennies'

	$L$	$R$
$T$	1	-1
$B$	-1	1

- Formulation 1 : A zero-sum game

	$L$	$R$
$T$	1	-1
$B$	-1	1

- Formulation 1 : A zero-sum game

	$L$	$R$
$T$	1	-1
$B$	-1	1

- Formulation 2 : bi-matrix game

	$L$	$R$
$T$	1, -1	-1, 1
$B$	-1, 1	1, -1

- Formulation 1 : A zero-sum game

	$L$	$R$
$T$	1	-1
$B$	-1	1

- Formulation 2 : bi-matrix game

	$L$	$R$
$T$	1, -1	-1, 1
$B$	-1, 1	1, -1

Saying that the game is zero-sum avoid to specify the payoffs of players 2.

Let  $w \in \bar{\mathbb{R}} = \mathbb{R} \cup \{-\infty, +\infty\}$ .

## Definition

P1 guarantees  $w$  if :  $\exists i \in I, \quad \forall j \in J, \quad g(i, j) \geq w$ .



Let  $w \in \bar{\mathbb{R}} = \mathbb{R} \cup \{-\infty, +\infty\}$ .

## Definition

P1 guarantees  $w$  if :  $\exists i \in I, \quad \forall j \in J, \quad g(i, j) \geq w$ .

## Definition

P2 can guarantee  $w$  if :  $\exists j \in J, \quad \forall i \in I, \quad g(i, j) \leq w$ .

Because player 2 is minimizing, the inequalities are reversed.

Let  $G = (I, J, g)$  be a zero-sum game.

## Definition

The *maxmin* of  $G$  is the supremum of quantities that P1 can guarantee. We denote it  $\max \min(G)$ , or  $\underline{v}$ . We have :  $\underline{v} = \sup_{i \in I} \inf_{j \in J} g(i, j)$ .

Let  $G = (I, J, g)$  be a zero-sum game.

## Definition

The *maxmin* of  $G$  is the supremum of quantities that P1 can guarantee. We denote it  $\max\min(G)$ , or  $\underline{v}$ . We have :  $\underline{v} = \sup_{i \in I} \inf_{j \in J} g(i, j)$ .

Interpretation : if P1 plays before P2 and if P2 observes what P1 did before choosing his action, then the rational outcome of the game is  $\underline{v}$  for player 1 and  $-\underline{v}$  for player 2.

Let  $G = (I, J, g)$  be a zero-sum game.

## Definition

The *maxmin* of  $G$  is the supremum of quantities that P1 can guarantee. We denote it  $\max \min(G)$ , or  $\underline{v}$ . We have :  $\underline{v} = \sup_{i \in I} \inf_{j \in J} g(i, j)$ .

Interpretation : if P1 plays before P2 and if P2 observes what P1 did before choosing his action, then the rational outcome of the game is  $\underline{v}$  for player 1 and  $-\underline{v}$  for player 2.

## Definition

The *minmax* of  $G$  is the infimum of the quantities that P2 can guarantee. It is denoted  $\min \max(G)$ , or  $\bar{v}$ . we have  $\bar{v} = \inf_{i \in I} \sup_{j \in J} g(i, j)$ .

Interpretation : if P2 plays before P1, the rational outcome must be  $\bar{v}$  for p1 and  $-\bar{v}$  for P2.

## Proposition

$$\underline{v} \leq \bar{v}.$$

The jump  $\bar{v} - \underline{v}$  is called the *duality gap*.

## Proposition

$$\underline{v} \leq \bar{v}.$$

The jump  $\bar{v} - \underline{v}$  is called the *duality gap*.

## Definition

The game  $G$  has a *value* if  $\underline{v} = \bar{v}$ . In this case the value of  $G$  is denoted  $v$ .

## Proposition

$$\underline{v} \leq \bar{v}.$$

The jump  $\bar{v} - \underline{v}$  is called the *duality gap*.

## Definition

The game  $G$  has a *value* if  $\underline{v} = \bar{v}$ . In this case the value of  $G$  is denoted  $v$ .

## Proposition

If  $w$  can be guaranteed by both players, then  $w$  is unique and it is the value.

	$L$	$R$
$T$	1	-1
$B$	-1	1

$\underline{v} = -1 < 1 = \bar{v}$  : No value.



	$L$	$R$
$T$	1	-1
$B$	-1	1

$\underline{v} = -1 < 1 = \bar{v}$  : No value.

	$L$	$R$
$T$	1	2
$B$	-1	1

$\underline{v} = 1 = \bar{v}$  : the value exists and is  $v = 1$ .

Suppose that the game has a value  $v$ .

## Definition

- A strategy of player 1 is  $\varepsilon$ -optimal if it guarantees  $v - \varepsilon$ .  
A strategy of player 2 is  $\varepsilon$ -optimal if it guarantees  $v + \varepsilon$ .
- The 0-optimal strategies are called optimal. Let  $I^*$  be the optimal strategies for player 1,  $J^*$  be the optimal for player 2.

Suppose that the game has a value  $v$ .

## Definition

- A strategy of player 1 is  $\varepsilon$ -optimal if it guarantees  $v - \varepsilon$ .  
A strategy of player 2 is  $\varepsilon$ -optimal if it guarantees  $v + \varepsilon$ .
- The 0-optimal strategies are called optimal. Let  $I^*$  be the optimal strategies for player 1,  $J^*$  be the optimal for player 2.

- If  $(i^*, j^*) \in I^* \times J^*$ , then  $v = g(i^*, j^*)$ .
- If a finite game has a value, each player has an optimal strategy.
- Not true when the game is infinite : If  $G = (\mathbf{N}, \mathbf{N}, g)$ , where  $g(i, j) = 1/(i + j + 1)$ , what are the  $\varepsilon$ -optimal strategies of player 1? player 2?

Let  $G = (g, I, J)$  be a zero-sum game which has a value. Then  $(i^*, j^*)$  belongs to  $I^* \times J^*$  if and only if it is a saddle point of  $g$ , that is :

$$\forall (i, j) \in I \times J, \quad g(i, j^*) \leq g(i^*, j^*) \leq g(i^*, j),$$

Meaning that  $i^*$  is the best response against  $j^*$  et  $j^*$  is the best response against  $i^*$ , e.g.  $(i^*, j^*)$  is a Nash equilibrium.

Let  $G = (g, I, J)$  be a zero-sum game which has a value. Then  $(i^*, j^*)$  belongs to  $I^* \times J^*$  if and only if it is a saddle point of  $g$ , that is :

$$\forall (i, j) \in I \times J, \quad g(i, j^*) \leq g(i^*, j^*) \leq g(i^*, j),$$

Meaning that  $i^*$  is the best response against  $j^*$  et  $j^*$  is the best response against  $i^*$ , e.g.  $(i^*, j^*)$  is a Nash equilibrium.

## Careful !

An optimal strategy is not necessarily a best reply against any strategy of the opponent !... An example ?

## Definition

- A strategy  $a$  of Player 1 is strongly dominated by another strategy  $b$  if  $g(a, j) < g(b, j)$  for every action  $j$  of Player 2.
- A strategy  $a$  of Player 1 is weakly dominated by another strategy  $b$  if  $g(a, j) \leq g(b, j)$  for every action  $j$  of Player 2, and the inequality is strict for at least one  $j$ .

This is basically a way to express the fact that a strategy is worse than another one. It should not be a big surprise that it is not worthwhile to play dominated strategies.

## Proposition

- The value,  $\text{infsup}$ ,  $\text{supinf}$  of a game don't change when one removes weakly dominated strategies.
- If the  $\text{supinf}$  and  $\text{infsup}$  are  $\text{maxmin}$  and  $\text{minmax}$  (for example if the game is finite) then a strongly dominated strategy cannot be optimal.

## Careful !

A weakly dominated strategy can be optimal, even in a finite game.

- 1 Introduction ; value in pure strategies
- 2 Value in Mixed Strategies (in the finite case)
- 3 Learning to Play Optimal

- In many simultaneous games, knowing the strategy of the opponent before his play can be an advantage.



- In many simultaneous games, knowing the strategy of the opponent before his play can be an advantage.
- A good player must thus hide his strategy. He must be unpredictable.

- In many simultaneous games, knowing the strategy of the opponent before his play can be an advantage.
- A good player must thus hide his strategy. He must be unpredictable.
- To be unpredictable, players must play at random : instead of choosing deterministically an element in  $I$  or  $J$ , they choose a probability distribution on  $I$  or  $J$ , that we call a *mixed strategy*.

- In many simultaneous games, knowing the strategy of the opponent before his play can be an advantage.
- A good player must thus hide his strategy. He must be unpredictable.
- To be unpredictable, players must play at random : instead of choosing deterministically an element in  $I$  or  $J$ , they choose a probability distribution on  $I$  or  $J$ , that we call a *mixed strategy*.
- For example, if we are playing Matching Pennies, or describing an algorithm that will play it “online”, it is clearly interesting to select each strategy with probability  $1/2$ .

If  $E$  is finite with cardinality  $n$ , let's denote :

$$\Delta(E) = \{x \in \mathbb{R}_+^n; \sum_{e \in E} x^e = 1\}.$$

If  $E$  is finite with cardinality  $n$ , let's denote :

$$\Delta(E) = \{x \in \mathbb{R}_+^n; \sum_{e \in E} x^e = 1\}.$$

## Definition

The mixed extension of a **finite game**  $G = (I, J, g)$  is the game

$$\Gamma = (\Delta(I), \Delta(J), g),$$

Where

$$g(x, y) = \mathbb{E}_{x \otimes y} g = \sum_{i,j} x^i y^j g(i, j).$$

- Elements of  $I$  and  $J$  are called *pure strategies* of  $G$ ,

- Elements of  $I$  and  $J$  are called *pure strategies* of  $G$ ,
- Elements of  $\Delta(I)$  and  $\Delta(J)$  are *mixed strategies* of  $G$ ,

- Elements of  $I$  and  $J$  are called *pure strategies* of  $G$ ,
- Elements of  $\Delta(I)$  and  $\Delta(J)$  are *mixed strategies* of  $G$ ,
- The pure strategy  $i \in I$  is identified with the Dirac mass at  $i$  (denoted  $\delta_i$ )  
With this identification,  $I \subset \Delta(I)$  and  $J \subset \Delta(J)$ ,



- Elements of  $I$  and  $J$  are called *pure strategies* of  $G$ ,
- Elements of  $\Delta(I)$  and  $\Delta(J)$  are *mixed strategies* of  $G$ ,
- The pure strategy  $i \in I$  is identified with the Dirac mass at  $i$  (denoted  $\delta_i$ )  
With this identification,  $I \subset \Delta(I)$  and  $J \subset \Delta(J)$ ,
- The *support* of the mixed strategy  $x$  is :  $\text{supp}(x) = \{i \in I : x^i > 0\}$ ,

- Elements of  $I$  and  $J$  are called *pure strategies* of  $G$ ,
- Elements of  $\Delta(I)$  and  $\Delta(J)$  are *mixed strategies* of  $G$ ,
- The pure strategy  $i \in I$  is identified with the Dirac mass at  $i$  (denoted  $\delta_i$ )  
With this identification,  $I \subset \Delta(I)$  and  $J \subset \Delta(J)$ ,
- The *support* of the mixed strategy  $x$  is :  $\text{supp}(x) = \{i \in I : x^i > 0\}$ ,
- Let  $A = (g(i, j))_{(i, j) \in I \times J}$  and  $(x, y) \in \Delta(I) \times \Delta(J)$ . Then we have

$$g(x, y) = xAy := x^t Ay$$

- For any  $y \in \Delta(J)$ , one has :  $\max_{x \in \Delta(I)} g(x, y) = \max_{i \in I} g(i, y)$

- For any  $y \in \Delta(J)$ , one has :  $\max_{x \in \Delta(I)} g(x, y) = \max_{i \in I} g(i, y)$
- For any  $x \in \Delta(I)$ , one has :  $\min_{y \in \Delta(J)} g(x, y) = \min_{j \in J} g(x, j)$

- For any  $y \in \Delta(J)$ , one has :  $\max_{x \in \Delta(I)} g(x, y) = \max_{i \in I} g(i, y)$
- For any  $x \in \Delta(I)$ , one has :  $\min_{y \in \Delta(J)} g(x, y) = \min_{j \in J} g(x, j)$

### Proposition

The duality gap is smaller in  $\Gamma$ . If a player can guarantee  $w \in \mathbb{R}$  in  $G$ , he can guarantee  $w$  in  $\Gamma$ , using the same strategy. In particular, if  $G$  has a value, then  $\Gamma$  has the same value.

The converse is false !

## Theorem

Let  $A$  be a real valued matrix indexed by  $I \times J$ . There exists  $(x^*, y^*, v)$  in  $\Delta(I) \times \Delta(J) \times \mathbb{R}$  such that :

$$\forall y \in \Delta(J), x^* A y \geq v \text{ et } \forall x \in \Delta(I), x A y^* \leq v.$$

Said differently, the mixed extension of a game has a value, and players have optimal strategies.

## Theorem

Let  $A$  be a real valued matrix indexed by  $I \times J$ . There exists  $(x^*, y^*, v)$  in  $\Delta(I) \times \Delta(J) \times \mathbb{R}$  such that :

$$\forall y \in \Delta(J), x^* A y \geq v \text{ et } \forall x \in \Delta(I), x A y^* \leq v.$$

Said differently, the mixed extension of a game has a value, and players have optimal strategies.

Moreover we have :

$$\begin{aligned} v &= \max_{x \in \Delta(I)} \min_{y \in \Delta(J)} x A y = \min_{y \in \Delta(J)} \max_{x \in \Delta(I)} x A y \\ &= \max_{x \in \Delta(I)} \min_{j \in J} x A j = \min_{y \in \Delta(J)} \max_{i \in I} i A y. \end{aligned}$$

Let  $A \in \mathbb{R}^{I \times J}$  be a matrix game. Let  $X(A) \subset \Delta(I)$  and  $Y(A) \subset \Delta(J)$  be optimal for players 1 and 2 respectively.

## Proposition

- $X(A)$  and  $Y(A)$  are non empty polytopes,



Let  $A \in \mathbb{R}^{I \times J}$  be a matrix game. Let  $X(A) \subset \Delta(I)$  and  $Y(A) \subset \Delta(J)$  be optimal for players 1 and 2 respectively.

## Proposition

- $X(A)$  and  $Y(A)$  are non empty polytopes,
- $X(A) \times Y(A)$  is the set of saddle-points of  $A$ , e.g. elements of  $(x^*, y^*)$  on  $\Delta(I) \times \Delta(J)$  such that :

$$x A y^* \leq x^* A y^* \leq x^* A y \quad \forall (x, y) \in \Delta(I) \times \Delta(J).$$

Let  $A \in \mathbb{R}^{I \times J}$  be a matrix game. Let  $X(A) \subset \Delta(I)$  and  $Y(A) \subset \Delta(J)$  be optimal for players 1 and 2 respectively.

## Proposition

- $X(A)$  and  $Y(A)$  are non empty polytopes,
- $X(A) \times Y(A)$  is the set of saddle-points of  $A$ , e.g. elements of  $(x^*, y^*)$  on  $\Delta(I) \times \Delta(J)$  such that :

$$x A y^* \leq x^* A y^* \leq x^* A y \quad \forall (x, y) \in \Delta(I) \times \Delta(J).$$

- Let  $(x^*, y^*) \in X(A) \times Y(A)$ . Then for all  $i \in \text{supp}(x^*)$  and  $j \in \text{supp}(y^*)$ ,

$$i A y^* = v = x^* A j = x^* A y^* \quad (\text{complementarity}).$$

1	-2
-1	3

Here  $v = 1/7$ . Player 1 optimal strategy : play Top with probab  $(4/7, 3/7)$  on  $(T, B)$ . Player 2 optimal strategy :  $(5/7, 2/7)$  on  $(L, R)$ .

1	-2
-1	3

Here  $v = 1/7$ . Player 1 optimal strategy : play Top with probab  $(4/7, 3/7)$  on  $(T, B)$ . Player 2 optimal strategy :  $(5/7, 2/7)$  on  $(L, R)$ .

1	2
0	$t$

For all  $t \in \mathbb{R}$ , the game has value  $v = 1$ , and each player has a unique optimal strategy, which is pure : Top for player 1, Left for player 2.

1	-2
-1	3

Here  $v = 1/7$ . Player 1 optimal strategy : play Top with probab  $(4/7, 3/7)$  on  $(T, B)$ . Player 2 optimal strategy :  $(5/7, 2/7)$  on  $(L, R)$ .

1	2
0	$t$

For all  $t \in \mathbb{R}$ , the game has value  $v = 1$ , and each player has a unique optimal strategy, which is pure : Top for player 1, Left for player 2.

$a$	$b$
$c$	$d$

In the case where each player has two actions, either there exists a pair of pure optimal strategies (and then the value is one of the numbers  $a, b, c, d$ ) or the optimal strategies are completely mixed and the value is given by

$$v = \frac{ad - bc}{a + d - b - c}.$$

## Theorem

Let  $A$  be an  $n \times m$  matrix,  $b$  an  $1 \times m$  vector and  $c$  a  $n \times 1$  vector with real coefficients. The two dual linear programs

$$\begin{array}{ll} \min \langle c, x \rangle & \max \langle y, b \rangle \\ (\mathcal{P}_1) \quad xA \geq b & (\mathcal{P}_2) \quad Ay \leq c \\ \quad x \geq 0 & \quad y \geq 0 \end{array}$$

have the same value as soon as they are feasible, i.e. when the sets  $\{xA \geq b; x \geq 0\}$  and  $\{Ay \leq c; y \geq 0\}$  are non-empty.

## Proof of minmax theorem from duality

By considering  $A + tE$  with  $t \geq 0$  and  $E$  being the matrix with  $E_{ij} = 1, \forall (i, j) \in I \times J$ , one can assume  $A \gg 0$ , of dimension  $m \times n$ .

## Proof of minmax theorem from duality

By considering  $A + tE$  with  $t \geq 0$  and  $E$  being the matrix with  $E_{ij} = 1, \forall (i, j) \in I \times J$ , one can assume  $A \gg 0$ , of dimension  $m \times n$ .  
Let us consider the dual programs

$$\begin{array}{ll} \min \langle X, c \rangle & \max \langle b, Y \rangle \\ (\mathcal{P}_1) \quad XA \geq b & (\mathcal{P}_2) \quad AY \leq c \\ \quad X \geq 0 & \quad Y \geq 0 \end{array}$$



## Proof of minmax theorem from duality

By considering  $A + tE$  with  $t \geq 0$  and  $E$  being the matrix with  $E_{ij} = 1, \forall (i, j) \in I \times J$ , one can assume  $A \gg 0$ , of dimension  $m \times n$ .  
Let us consider the dual programs

$$\begin{array}{ll} \min \langle X, c \rangle & \max \langle b, Y \rangle \\ (\mathcal{P}_1) \quad XA \geq b & (\mathcal{P}_2) \quad AY \leq c \\ \quad X \geq 0 & \quad Y \geq 0 \end{array}$$

where the variables satisfy  $X \in \mathbb{R}^m, Y \in \mathbb{R}^n$  and the parameters are given by  $c \in \mathbb{R}^m, c_i = 1, \forall i$  and  $b \in \mathbb{R}^n, b_j = 1, \forall j$ .

## Proof of minmax theorem from duality

By considering  $A + tE$  with  $t \geq 0$  and  $E$  being the matrix with  $E_{ij} = 1, \forall (i, j) \in I \times J$ , one can assume  $A \gg 0$ , of dimension  $m \times n$ .  
Let us consider the dual programs

$$\begin{array}{ll} \min \langle X, c \rangle & \max \langle b, Y \rangle \\ (\mathcal{P}_1) \quad XA \geq b & (\mathcal{P}_2) \quad AY \leq c \\ \quad X \geq 0 & \quad Y \geq 0 \end{array}$$

where the variables satisfy  $X \in \mathbb{R}^m, Y \in \mathbb{R}^n$  and the parameters are given by  $c \in \mathbb{R}^m, c_i = 1, \forall i$  and  $b \in \mathbb{R}^n, b_j = 1, \forall j$ .

## Proof of minmax theorem from duality

By considering  $A + tE$  with  $t \geq 0$  and  $E$  being the matrix with  $E_{ij} = 1, \forall (i, j) \in I \times J$ , one can assume  $A \gg 0$ , of dimension  $m \times n$ .  
Let us consider the dual programs

$$\begin{array}{ll} \min \langle X, c \rangle & \max \langle b, Y \rangle \\ (\mathcal{P}_1) \quad XA \geq b & (\mathcal{P}_2) \quad AY \leq c \\ \quad X \geq 0 & \quad Y \geq 0 \end{array}$$

where the variables satisfy  $X \in \mathbb{R}^m, Y \in \mathbb{R}^n$  and the parameters are given by  $c \in \mathbb{R}^m, c_i = 1, \forall i$  and  $b \in \mathbb{R}^n, b_j = 1, \forall j$ .

$(\mathcal{P}_2)$  is feasible with  $Y = 0$ , as is  $(\mathcal{P}_1)$  by taking  $X$  large enough.

# Proof of minmax theorem from duality

By considering  $A + tE$  with  $t \geq 0$  and  $E$  being the matrix with  $E_{ij} = 1, \forall (i, j) \in I \times J$ , one can assume  $A \gg 0$ , of dimension  $m \times n$ .  
Let us consider the dual programs

$$\begin{array}{ll} \min \langle X, c \rangle & \max \langle b, Y \rangle \\ (\mathcal{P}_1) \quad XA \geq b & (\mathcal{P}_2) \quad AY \leq c \\ \quad X \geq 0 & \quad Y \geq 0 \end{array}$$

where the variables satisfy  $X \in \mathbb{R}^m, Y \in \mathbb{R}^n$  and the parameters are given by  $c \in \mathbb{R}^m, c_i = 1, \forall i$  and  $b \in \mathbb{R}^n, b_j = 1, \forall j$ .

$(\mathcal{P}_2)$  is feasible with  $Y = 0$ , as is  $(\mathcal{P}_1)$  by taking  $X$  large enough.

Thus by the duality theorem there exists a triple  $(X^*, Y^*, w)$  with :

$$X^* \geq 0, Y^* \geq 0, X^*A \geq b, AY^* \leq c, \quad \sum_i X_i^* = \sum_j Y_j^* = w.$$

# Proof of minmax theorem from duality

By considering  $A + tE$  with  $t \geq 0$  and  $E$  being the matrix with  $E_{ij} = 1, \forall (i, j) \in I \times J$ , one can assume  $A \gg 0$ , of dimension  $m \times n$ .  
Let us consider the dual programs

$$\begin{array}{ll} \min \langle X, c \rangle & \max \langle b, Y \rangle \\ (\mathcal{P}_1) \quad XA \geq b & (\mathcal{P}_2) \quad AY \leq c \\ \quad X \geq 0 & \quad Y \geq 0 \end{array}$$

where the variables satisfy  $X \in \mathbb{R}^m, Y \in \mathbb{R}^n$  and the parameters are given by  $c \in \mathbb{R}^m, c_i = 1, \forall i$  and  $b \in \mathbb{R}^n, b_j = 1, \forall j$ .

$(\mathcal{P}_2)$  is feasible with  $Y = 0$ , as is  $(\mathcal{P}_1)$  by taking  $X$  large enough.

Thus by the duality theorem there exists a triple  $(X^*, Y^*, w)$  with :

$$X^* \geq 0, Y^* \geq 0, X^*A \geq b, AY^* \leq c, \quad \sum_i X_i^* = \sum_j Y_j^* = w.$$

$X^* \neq 0$  implies  $w > 0$ , hence dividing  $X^*$  and  $Y^*$  by  $w$ , one obtains the existence of  $(x^*, y^*) \in \Delta(I) \times \Delta(J)$  with

$$x^*Ae^j \geq 1/w, \forall j, \quad e^iAy^* \leq 1/w, \forall i.$$

# Proof of minmax theorem from duality

By considering  $A + tE$  with  $t \geq 0$  and  $E$  being the matrix with  $E_{ij} = 1, \forall (i, j) \in I \times J$ , one can assume  $A \gg 0$ , of dimension  $m \times n$ .  
Let us consider the dual programs

$$\begin{array}{ll} \min \langle X, c \rangle & \max \langle b, Y \rangle \\ (\mathcal{P}_1) \quad XA \geq b & (\mathcal{P}_2) \quad AY \leq c \\ & X \geq 0 \quad Y \geq 0 \end{array}$$

where the variables satisfy  $X \in \mathbb{R}^m, Y \in \mathbb{R}^n$  and the parameters are given by  $c \in \mathbb{R}^m, c_i = 1, \forall i$  and  $b \in \mathbb{R}^n, b_j = 1, \forall j$ .

$(\mathcal{P}_2)$  is feasible with  $Y = 0$ , as is  $(\mathcal{P}_1)$  by taking  $X$  large enough.

Thus by the duality theorem there exists a triple  $(X^*, Y^*, w)$  with :

$$X^* \geq 0, Y^* \geq 0, X^*A \geq b, AY^* \leq c, \quad \sum_i X_i^* = \sum_j Y_j^* = w.$$

$X^* \neq 0$  implies  $w > 0$ , hence dividing  $X^*$  and  $Y^*$  by  $w$ , one obtains the existence of  $(x^*, y^*) \in \Delta(I) \times \Delta(J)$  with

$$x^* A e^j \geq 1/w, \forall j, \quad e^i A y^* \leq 1/w, \forall i.$$

Hence there is a value, namely  $1/w$ , and  $x^*$  and  $y^*$  are optimal strategies.

- 1 Introduction ; value in pure strategies
- 2 Value in Mixed Strategies (in the finite case)
- 3 Learning to Play Optimal

Fictitious play (FP) : is a learning process (an online algorithm) which allows to prove the minmax theorem and compute optimal strategies.



Fictitious play (FP) : is a learning process (an online algorithm) which allows to prove the minmax theorem and compute optimal strategies.

### Principle

- We start from any  $(i_1, j_1)$  in  $I \times J$ ,
- At each stage  $n \geq 2$ , each player will play a best response to the average past behavior of the opponent up to stage  $n - 1$ .

Fictitious play (FP) : is a learning process (an online algorithm) which allows to prove the minmax theorem and compute optimal strategies.

## Principle

- We start from any  $(i_1, j_1)$  in  $I \times J$ ,
- At each stage  $n \geq 2$ , each player will play a best response to the average past behavior of the opponent up to stage  $n - 1$ .

## Definition

A sequence  $(i_n, j_n)_{n \geq 1}$  is a realisation of FP process if for each  $n \geq 1$  :

- $i_{n+1}$  is a best response of player 1 against  $y_n := \frac{1}{n} \sum_{t=1}^n j_t \in \Delta(J)$ ,
- $j_{n+1}$  is a best response of player 2 against  $x_n := \frac{1}{n} \sum_{t=1}^n i_t \in \Delta(I)$ .

## Theorem (Robinson, 1951)

Let  $(i_n, j_n)_{n \geq 1}$  be the realization of a fictitious play process for the matrix  $A$ .  
Then :

- 1) The distance from  $(x_n, y_n)$  to the set of optimal strategies  $X(A) \times Y(A)$  goes to 0 as  $n \rightarrow \infty$ . Explicitly :

$$\forall \varepsilon > 0, \exists N, \forall n \geq N, \forall x \in \Delta(I), \forall y \in \Delta(J)$$

$$x_n A y \geq \text{val}(A) - \varepsilon \quad \text{and} \quad x A y_n \leq \text{val}(A) + \varepsilon.$$

- 2) The average payoff on the trajectory, namely  $\frac{1}{n} \sum_{t=1}^n A_{i_t, j_t}$ , converges to  $\text{val}(A)$ .

## Theorem (Robinson, 1951)

Let  $(i_n, j_n)_{n \geq 1}$  be the realization of a fictitious play process for the matrix  $A$ .  
Then :

- 1) The distance from  $(x_n, y_n)$  to the set of optimal strategies  $X(A) \times Y(A)$  goes to 0 as  $n \rightarrow \infty$ . Explicitly :

$$\forall \varepsilon > 0, \exists N, \forall n \geq N, \forall x \in \Delta(I), \forall y \in \Delta(J)$$

$$x_n A y \geq \text{val}(A) - \varepsilon \quad \text{and} \quad x A y_n \leq \text{val}(A) + \varepsilon.$$

- 2) The average payoff on the trajectory, namely  $\frac{1}{n} \sum_{t=1}^n A_{i_t, j_t}$ , converges to  $\text{val}(A)$ .

This is an alternative (and constructive) proof of the minmax theorem.

- Initial proof : by induction

- Initial proof : by induction
- Modern proof : go to continuous time (this lecture).

## Continuous Fictitious Play

Take as variables the empirical frequencies  $x_n$  and  $y_n$ , so that the discrete dynamics for player 1 reads as

$$x_{n+1} = \frac{1}{n+1} [i_{n+1} + nx_n] \quad \text{with} \quad i_{n+1} \in \text{BR}^1(y_n)$$

Take as variables the empirical frequencies  $x_n$  and  $y_n$ , so that the discrete dynamics for player 1 reads as

$$x_{n+1} = \frac{1}{n+1} [i_{n+1} + nx_n] \quad \text{with} \quad i_{n+1} \in \text{BR}^1(y_n)$$

and hence satisfies

$$x_{n+1} - x_n \in \frac{1}{n+1} [\text{BR}^1(y_n) - x_n].$$



Take as variables the empirical frequencies  $x_n$  and  $y_n$ , so that the discrete dynamics for player 1 reads as

$$x_{n+1} = \frac{1}{n+1} [i_{n+1} + nx_n] \quad \text{with} \quad i_{n+1} \in \text{BR}^1(y_n)$$

and hence satisfies

$$x_{n+1} - x_n \in \frac{1}{n+1} [\text{BR}^1(y_n) - x_n].$$

The corresponding system in continuous time is

$$\dot{x}(t) \in \frac{1}{t} [\text{BR}^1(y(t)) - x(t)].$$

Take as variables the empirical frequencies  $x_n$  and  $y_n$ , so that the discrete dynamics for player 1 reads as

$$x_{n+1} = \frac{1}{n+1} [i_{n+1} + nx_n] \quad \text{with} \quad i_{n+1} \in \text{BR}^1(y_n)$$

and hence satisfies

$$x_{n+1} - x_n \in \frac{1}{n+1} [\text{BR}^1(y_n) - x_n].$$

The corresponding system in continuous time is

$$\dot{x}(t) \in \frac{1}{t} [\text{BR}^1(y(t)) - x(t)].$$

This is a differential inclusion which defines, with a similar condition for player 2, the process CFP. The theory of differential inclusions implies existence of a CFP process.

Take as variables the empirical frequencies  $x_n$  and  $y_n$ , so that the discrete dynamics for player 1 reads as

$$x_{n+1} = \frac{1}{n+1} [i_{n+1} + nx_n] \quad \text{with} \quad i_{n+1} \in \text{BR}^1(y_n)$$

and hence satisfies

$$x_{n+1} - x_n \in \frac{1}{n+1} [\text{BR}^1(y_n) - x_n].$$

The corresponding system in continuous time is

$$\dot{x}(t) \in \frac{1}{t} [\text{BR}^1(y(t)) - x(t)].$$

This is a differential inclusion which defines, with a similar condition for player 2, the process CFP. The theory of differential inclusions implies existence of a CFP process.

**Theorem (Harris (1998); Hofbauer and Sorin (2006))**

*For the CFP process, the duality gap converges to 0 at a speed  $O(1/t)$  and  $(x(t), y(t))$  to the set of optimal strategies  $X(A) \times Y(A)$*

Make the time change  $z(t) = x(\exp(t))$ , which leads to the autonomous differential inclusion

$$\dot{x}(t) \in [\text{BR}^1(y(t)) - x(t)] , \quad \dot{y}(t) \in [\text{BR}^2(x(t)) - y(t)]$$

Make the time change  $z(t) = x(\exp(t))$ , which leads to the autonomous differential inclusion

$$\dot{x}(t) \in [\text{BR}^1(y(t)) - x(t)] , \quad \dot{y}(t) \in [\text{BR}^2(x(t)) - y(t)]$$

known as the *best response dynamics* (Gilboa and Matsui (1991)).

Make the time change  $z(t) = x(\exp(t))$ , which leads to the autonomous differential inclusion

$$\dot{x}(t) \in [\text{BR}^1(y(t)) - x(t)], \quad \dot{y}(t) \in [\text{BR}^2(x(t)) - y(t)]$$

known as the *best response dynamics* (Gilboa and Matsui (1991)).

Write the payoff as  $g(x, y) = xAy$  and for  $(x, y) \in \Delta(I) \times \Delta(J)$ , let

$$L(y) = \max_{x' \in \Delta(I)} g(x', y) \quad M(x) = \min_{y' \in \Delta(J)} g(x, y').$$

Make the time change  $z(t) = x(\exp(t))$ , which leads to the autonomous differential inclusion

$$\dot{x}(t) \in [\text{BR}^1(y(t)) - x(t)], \quad \dot{y}(t) \in [\text{BR}^2(x(t)) - y(t)]$$

known as the *best response dynamics* (Gilboa and Matsui (1991)).

Write the payoff as  $g(x, y) = xAy$  and for  $(x, y) \in \Delta(I) \times \Delta(J)$ , let

$$L(y) = \max_{x' \in \Delta(I)} g(x', y) \quad M(x) = \min_{y' \in \Delta(J)} g(x, y').$$

Thus the duality gap at  $(x, y)$  is defined as  $W(x, y) = L(y) - M(x) \geq 0$  and the pair  $(x, y)$  defines optimal strategies in  $A$  if and only if  $W(x, y) = 0$ .

Make the time change  $z(t) = x(\exp(t))$ , which leads to the autonomous differential inclusion

$$\dot{x}(t) \in [\text{BR}^1(y(t)) - x(t)], \quad \dot{y}(t) \in [\text{BR}^2(x(t)) - y(t)]$$

known as the *best response dynamics* (Gilboa and Matsui (1991)).

Write the payoff as  $g(x, y) = xAy$  and for  $(x, y) \in \Delta(I) \times \Delta(J)$ , let

$$L(y) = \max_{x' \in \Delta(I)} g(x', y) \quad M(x) = \min_{y' \in \Delta(J)} g(x, y').$$

Thus the duality gap at  $(x, y)$  is defined as  $W(x, y) = L(y) - M(x) \geq 0$  and the pair  $(x, y)$  defines optimal strategies in  $A$  if and only if  $W(x, y) = 0$ .

Let now  $(x(t), y(t))_{t \geq 0}$  be a solution of CFP.



Make the time change  $z(t) = x(\exp(t))$ , which leads to the autonomous differential inclusion

$$\dot{x}(t) \in [\text{BR}^1(y(t)) - x(t)], \quad \dot{y}(t) \in [\text{BR}^2(x(t)) - y(t)]$$

known as the *best response dynamics* (Gilboa and Matsui (1991)).

Write the payoff as  $g(x, y) = xAy$  and for  $(x, y) \in \Delta(I) \times \Delta(J)$ , let

$$L(y) = \max_{x' \in \Delta(I)} g(x', y) \quad M(x) = \min_{y' \in \Delta(J)} g(x, y').$$

Thus the duality gap at  $(x, y)$  is defined as  $W(x, y) = L(y) - M(x) \geq 0$  and the pair  $(x, y)$  defines optimal strategies in  $A$  if and only if  $W(x, y) = 0$ .

Let now  $(x(t), y(t))_{t \geq 0}$  be a solution of CFP.

Denote by

$$w(t) = W(x(t), y(t))$$

the evaluation of the duality gap on the trajectory,

Make the time change  $z(t) = x(\exp(t))$ , which leads to the autonomous differential inclusion

$$\dot{x}(t) \in [\text{BR}^1(y(t)) - x(t)], \quad \dot{y}(t) \in [\text{BR}^2(x(t)) - y(t)]$$

known as the *best response dynamics* (Gilboa and Matsui (1991)).

Write the payoff as  $g(x, y) = xAy$  and for  $(x, y) \in \Delta(I) \times \Delta(J)$ , let

$$L(y) = \max_{x' \in \Delta(I)} g(x', y) \quad M(x) = \min_{y' \in \Delta(J)} g(x, y').$$

Thus the duality gap at  $(x, y)$  is defined as  $W(x, y) = L(y) - M(x) \geq 0$  and the pair  $(x, y)$  defines optimal strategies in  $A$  if and only if  $W(x, y) = 0$ .

Let now  $(x(t), y(t))_{t \geq 0}$  be a solution of CFP.

Denote by

$$w(t) = W(x(t), y(t))$$

the evaluation of the duality gap on the trajectory,  
and write

$$\alpha(t) = x(t) + \dot{x}(t) \in \text{BR}^1(y(t)) \quad \text{and} \quad \beta(t) = y(t) + \dot{y}(t) \in \text{BR}^2(x(t)).$$

$$\alpha(t) = x(t) + \dot{x}(t) \in \text{BR}^1(y(t)) \quad \text{and} \quad \beta(t) = y(t) + \dot{y}(t) \in \text{BR}^2(x(t)).$$

$$\alpha(t) = x(t) + \dot{x}(t) \in \text{BR}^1(y(t)) \quad \text{and} \quad \beta(t) = y(t) + \dot{y}(t) \in \text{BR}^2(x(t)).$$

We have  $L(y(t)) = g(\alpha(t), y(t))$ , thus

$$\frac{d}{dt}L(y(t)) = \dot{\alpha}(t)Ay(t) + \alpha(t)A\dot{y}(t).$$

$$\alpha(t) = x(t) + \dot{x}(t) \in \text{BR}^1(y(t)) \quad \text{and} \quad \beta(t) = y(t) + \dot{y}(t) \in \text{BR}^2(x(t)).$$

We have  $L(y(t)) = g(\alpha(t), y(t))$ , thus

$$\frac{d}{dt}L(y(t)) = \dot{\alpha}(t)Ay(t) + \alpha(t)A\dot{y}(t).$$

The envelope theorem shows that the first term collapses hence we obtain

$$\alpha(t) = x(t) + \dot{x}(t) \in \text{BR}^1(y(t)) \quad \text{and} \quad \beta(t) = y(t) + \dot{y}(t) \in \text{BR}^2(x(t)).$$

We have  $L(y(t)) = g(\alpha(t), y(t))$ , thus

$$\frac{d}{dt}L(y(t)) = \dot{\alpha}(t)Ay(t) + \alpha(t)A\dot{y}(t).$$

The envelope theorem shows that the first term collapses hence we obtain

$$\dot{w}(t) = \frac{d}{dt}L(y(t)) - \frac{d}{dt}M(x(t))$$

$$\alpha(t) = x(t) + \dot{x}(t) \in \text{BR}^1(y(t)) \quad \text{and} \quad \beta(t) = y(t) + \dot{y}(t) \in \text{BR}^2(x(t)).$$

We have  $L(y(t)) = g(\alpha(t), y(t))$ , thus

$$\frac{d}{dt}L(y(t)) = \dot{\alpha}(t)Ay(t) + \alpha(t)A\dot{y}(t).$$

The envelope theorem shows that the first term collapses hence we obtain

$$\begin{aligned} \dot{w}(t) &= \frac{d}{dt}L(y(t)) - \frac{d}{dt}M(x(t)) \\ &= \alpha(t)A\dot{y}(t) - \dot{x}(t)A\beta(t) \end{aligned}$$

$$\alpha(t) = x(t) + \dot{x}(t) \in \text{BR}^1(y(t)) \quad \text{and} \quad \beta(t) = y(t) + \dot{y}(t) \in \text{BR}^2(x(t)).$$

We have  $L(y(t)) = g(\alpha(t), y(t))$ , thus

$$\frac{d}{dt}L(y(t)) = \dot{\alpha}(t)Ay(t) + \alpha(t)A\dot{y}(t).$$

The envelope theorem shows that the first term collapses hence we obtain

$$\begin{aligned} \dot{w}(t) &= \frac{d}{dt}L(y(t)) - \frac{d}{dt}M(x(t)) \\ &= \alpha(t)A\dot{y}(t) - \dot{x}(t)A\beta(t) \\ &= x(t)A\dot{y}(t) - \dot{x}(t)Ay \end{aligned}$$



$$\alpha(t) = x(t) + \dot{x}(t) \in \text{BR}^1(y(t)) \quad \text{and} \quad \beta(t) = y(t) + \dot{y}(t) \in \text{BR}^2(x(t)).$$

We have  $L(y(t)) = g(\alpha(t), y(t))$ , thus

$$\frac{d}{dt}L(y(t)) = \dot{\alpha}(t)Ay(t) + \alpha(t)A\dot{y}(t).$$

The envelope theorem shows that the first term collapses hence we obtain

$$\begin{aligned} \dot{w}(t) &= \frac{d}{dt}L(y(t)) - \frac{d}{dt}M(x(t)) \\ &= \alpha(t)A\dot{y}(t) - \dot{x}(t)A\beta(t) \\ &= x(t)A\dot{y}(t) - \dot{x}(t)Ay \\ &= x(t)A\beta(t) - \alpha(t)Ay \end{aligned}$$

$$\alpha(t) = x(t) + \dot{x}(t) \in \text{BR}^1(y(t)) \quad \text{and} \quad \beta(t) = y(t) + \dot{y}(t) \in \text{BR}^2(x(t)).$$

We have  $L(y(t)) = g(\alpha(t), y(t))$ , thus

$$\frac{d}{dt}L(y(t)) = \dot{\alpha}(t)Ay(t) + \alpha(t)A\dot{y}(t).$$

The envelope theorem shows that the first term collapses hence we obtain

$$\begin{aligned} \dot{w}(t) &= \frac{d}{dt}L(y(t)) - \frac{d}{dt}M(x(t)) \\ &= \alpha(t)A\dot{y}(t) - \dot{x}(t)A\beta(t) \\ &= x(t)A\dot{y}(t) - \dot{x}(t)Ay \\ &= x(t)A\beta(t) - \alpha(t)Ay \\ &= M(x(t)) - L(y(t)) \end{aligned}$$

$$\alpha(t) = x(t) + \dot{x}(t) \in \text{BR}^1(y(t)) \quad \text{and} \quad \beta(t) = y(t) + \dot{y}(t) \in \text{BR}^2(x(t)).$$

We have  $L(y(t)) = g(\alpha(t), y(t))$ , thus

$$\frac{d}{dt}L(y(t)) = \dot{\alpha}(t)Ay(t) + \alpha(t)A\dot{y}(t).$$

The envelope theorem shows that the first term collapses hence we obtain

$$\begin{aligned} \dot{w}(t) &= \frac{d}{dt}L(y(t)) - \frac{d}{dt}M(x(t)) \\ &= \alpha(t)A\dot{y}(t) - \dot{x}(t)A\beta(t) \\ &= x(t)A\dot{y}(t) - \dot{x}(t)Ay \\ &= x(t)A\beta(t) - \alpha(t)Ay \\ &= M(x(t)) - L(y(t)) \\ &= -w(t). \end{aligned}$$

$$\alpha(t) = x(t) + \dot{x}(t) \in \text{BR}^1(y(t)) \quad \text{and} \quad \beta(t) = y(t) + \dot{y}(t) \in \text{BR}^2(x(t)).$$

We have  $L(y(t)) = g(\alpha(t), y(t))$ , thus

$$\frac{d}{dt}L(y(t)) = \dot{\alpha}(t)Ay(t) + \alpha(t)A\dot{y}(t).$$

The envelope theorem shows that the first term collapses hence we obtain

$$\begin{aligned} \dot{w}(t) &= \frac{d}{dt}L(y(t)) - \frac{d}{dt}M(x(t)) \\ &= \alpha(t)A\dot{y}(t) - \dot{x}(t)A\beta(t) \\ &= x(t)A\dot{y}(t) - \dot{x}(t)Ay \\ &= x(t)A\beta(t) - \alpha(t)Ay \\ &= M(x(t)) - L(y(t)) \\ &= -w(t). \end{aligned}$$

Thus  $w(t) = w(0) e^{-t}$ . There is convergence of  $w(t)$  to 0 at exponential speed, hence convergence to 0 at a speed  $O(1/t)$  in the original problem.

Let  $C$  be a non-empty closed convex subset of  $\mathbb{R}^k$  (endowed with the Euclidean norm) and  $\{x_n\}$  a bounded sequence in  $\mathbb{R}^k$ .

Let  $C$  be a non-empty closed convex subset of  $\mathbb{R}^k$  (endowed with the Euclidean norm) and  $\{x_n\}$  a bounded sequence in  $\mathbb{R}^k$ .

For  $x \in \mathbb{R}^k$ ,  $\Pi_C(x)$  stands for the projection of  $x$  on  $C$  and  $\bar{x}_n$  is the Cesàro mean up to stage  $n$  of the sequence  $\{x_i\}$  :

$$\bar{x}_n = \frac{1}{n} \sum_{i=1}^n x_i.$$

Let  $C$  be a non-empty closed convex subset of  $\mathbb{R}^k$  (endowed with the Euclidean norm) and  $\{x_n\}$  a bounded sequence in  $\mathbb{R}^k$ .

For  $x \in \mathbb{R}^k$ ,  $\Pi_C(x)$  stands for the projection of  $x$  on  $C$  and  $\bar{x}_n$  is the Cesàro mean up to stage  $n$  of the sequence  $\{x_i\}$  :

$$\bar{x}_n = \frac{1}{n} \sum_{i=1}^n x_i.$$

Say that  $\{x_n\}$  is a Blackwell  $C$ -sequence if it satisfies :

$$\langle x_{n+1} - \Pi_C(\bar{x}_n), \bar{x}_n - \Pi_C(\bar{x}_n) \rangle \leq 0, \quad \forall n.$$

Let  $C$  be a non-empty closed convex subset of  $\mathbb{R}^k$  (endowed with the Euclidean norm) and  $\{x_n\}$  a bounded sequence in  $\mathbb{R}^k$ .

For  $x \in \mathbb{R}^k$ ,  $\Pi_C(x)$  stands for the projection of  $x$  on  $C$  and  $\bar{x}_n$  is the Cesàro mean up to stage  $n$  of the sequence  $\{x_i\}$  :

$$\bar{x}_n = \frac{1}{n} \sum_{i=1}^n x_i.$$

Say that  $\{x_n\}$  is a Blackwell  $C$ -sequence if it satisfies :

$$\langle x_{n+1} - \Pi_C(\bar{x}_n), \bar{x}_n - \Pi_C(\bar{x}_n) \rangle \leq 0, \quad \forall n.$$

## Theorem

*If  $\{x_n\}$  is a Blackwell  $C$ -sequence then  $d_n = d(\bar{x}_n, C)$  converges to 0.*



Let  $y_n = \Pi_C(\bar{x}_n)$ . Then

Let  $y_n = \Pi_C(\bar{x}_n)$ . Then

$$d_{n+1}^2 \leq \|\bar{x}_{n+1} - y_n\|^2 = \|\bar{x}_n - y_n\|^2 + \|\bar{x}_{n+1} - \bar{x}_n\|^2 + 2\langle \bar{x}_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle.$$

Let  $y_n = \Pi_C(\bar{x}_n)$ . Then

$$d_{n+1}^2 \leq \|\bar{x}_{n+1} - y_n\|^2 = \|\bar{x}_n - y_n\|^2 + \|\bar{x}_{n+1} - \bar{x}_n\|^2 + 2\langle \bar{x}_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle.$$

Decompose

$$\langle \bar{x}_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle = \left( \frac{1}{n+1} \right) \langle x_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle$$

Let  $y_n = \Pi_C(\bar{x}_n)$ . Then

$$d_{n+1}^2 \leq \|\bar{x}_{n+1} - y_n\|^2 = \|\bar{x}_n - y_n\|^2 + \|\bar{x}_{n+1} - \bar{x}_n\|^2 + 2\langle \bar{x}_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle.$$

Decompose

$$\begin{aligned}\langle \bar{x}_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle &= \left( \frac{1}{n+1} \right) \langle x_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle \\ &= \left( \frac{1}{n+1} \right) (\langle x_{n+1} - y_n, \bar{x}_n - y_n \rangle - \|\bar{x}_n - y_n\|^2).\end{aligned}$$

Let  $y_n = \Pi_C(\bar{x}_n)$ . Then

$$d_{n+1}^2 \leq \|\bar{x}_{n+1} - y_n\|^2 = \|\bar{x}_n - y_n\|^2 + \|\bar{x}_{n+1} - \bar{x}_n\|^2 + 2\langle \bar{x}_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle.$$

Decompose

$$\begin{aligned} \langle \bar{x}_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle &= \left( \frac{1}{n+1} \right) \langle x_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle \\ &= \left( \frac{1}{n+1} \right) (\langle x_{n+1} - y_n, \bar{x}_n - y_n \rangle - \|\bar{x}_n - y_n\|^2). \end{aligned}$$

Using the hypothesis we obtain

$$d_{n+1}^2 \leq \left( 1 - \frac{2}{n+1} \right) d_n^2 + \left( \frac{1}{n+1} \right)^2 \|x_{n+1} - \bar{x}_n\|^2.$$

Let  $y_n = \Pi_C(\bar{x}_n)$ . Then

$$d_{n+1}^2 \leq \|\bar{x}_{n+1} - y_n\|^2 = \|\bar{x}_n - y_n\|^2 + \|\bar{x}_{n+1} - \bar{x}_n\|^2 + 2\langle \bar{x}_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle.$$

Decompose

$$\begin{aligned} \langle \bar{x}_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle &= \left( \frac{1}{n+1} \right) \langle x_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle \\ &= \left( \frac{1}{n+1} \right) (\langle x_{n+1} - y_n, \bar{x}_n - y_n \rangle - \|\bar{x}_n - y_n\|^2). \end{aligned}$$

Using the hypothesis we obtain

$$d_{n+1}^2 \leq \left( 1 - \frac{2}{n+1} \right) d_n^2 + \left( \frac{1}{n+1} \right)^2 \|x_{n+1} - \bar{x}_n\|^2.$$

From

$$\|x_{n+1} - \bar{x}_n\|^2 \leq 2\|x_{n+1}\|^2 + 2\|\bar{x}_n\|^2 \leq 4M^2,$$

Let  $y_n = \Pi_C(\bar{x}_n)$ . Then

$$d_{n+1}^2 \leq \|\bar{x}_{n+1} - y_n\|^2 = \|\bar{x}_n - y_n\|^2 + \|\bar{x}_{n+1} - \bar{x}_n\|^2 + 2\langle \bar{x}_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle.$$

Decompose

$$\begin{aligned} \langle \bar{x}_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle &= \left( \frac{1}{n+1} \right) \langle x_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle \\ &= \left( \frac{1}{n+1} \right) (\langle x_{n+1} - y_n, \bar{x}_n - y_n \rangle - \|\bar{x}_n - y_n\|^2). \end{aligned}$$

Using the hypothesis we obtain

$$d_{n+1}^2 \leq \left( 1 - \frac{2}{n+1} \right) d_n^2 + \left( \frac{1}{n+1} \right)^2 \|x_{n+1} - \bar{x}_n\|^2.$$

From

$$\|x_{n+1} - \bar{x}_n\|^2 \leq 2\|x_{n+1}\|^2 + 2\|\bar{x}_n\|^2 \leq 4M^2,$$

we deduce  $d_{n+1}^2 \leq \left( \frac{n-1}{n+1} \right) d_n^2 + \left( \frac{1}{n+1} \right)^2 4M^2$ .

Let  $y_n = \Pi_C(\bar{x}_n)$ . Then

$$d_{n+1}^2 \leq \|\bar{x}_{n+1} - y_n\|^2 = \|\bar{x}_n - y_n\|^2 + \|\bar{x}_{n+1} - \bar{x}_n\|^2 + 2\langle \bar{x}_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle.$$

Decompose

$$\begin{aligned} \langle \bar{x}_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle &= \left( \frac{1}{n+1} \right) \langle x_{n+1} - \bar{x}_n, \bar{x}_n - y_n \rangle \\ &= \left( \frac{1}{n+1} \right) (\langle x_{n+1} - y_n, \bar{x}_n - y_n \rangle - \|\bar{x}_n - y_n\|^2). \end{aligned}$$

Using the hypothesis we obtain

$$d_{n+1}^2 \leq \left( 1 - \frac{2}{n+1} \right) d_n^2 + \left( \frac{1}{n+1} \right)^2 \|x_{n+1} - \bar{x}_n\|^2.$$

From

$$\|x_{n+1} - \bar{x}_n\|^2 \leq 2\|x_{n+1}\|^2 + 2\|\bar{x}_n\|^2 \leq 4M^2,$$

we deduce  $d_{n+1}^2 \leq \left( \frac{n-1}{n+1} \right) d_n^2 + \left( \frac{1}{n+1} \right)^2 4M^2$ .

Thus by induction  $d_n \leq \frac{2M}{\sqrt{n}}$ .



## Learning to play optimally using Blackwell

Let  $A$  be an  $I \times J$  matrix and assume that the minmax is 0 :

$$\bar{v} = \min_{s \in \Delta(I)} \max_{i \in I} e^i A t = 0.$$

## Learning to play optimally using Blackwell

Let  $A$  be an  $I \times J$  matrix and assume that the minmax is 0 :

$$\bar{v} = \min_{s \in \Delta(I)} \max_{i \in I} e^i A t = 0.$$

Let us define by induction a sequence  $x_n \in \mathbb{R}^k$  for player 1 as follows, where  $k = |J|$ .

## Learning to play optimally using Blackwell

Let  $A$  be an  $I \times J$  matrix and assume that the minmax is 0 :

$$\bar{v} = \min_{s \in \Delta(I)} \max_{i \in I} e^i A t = 0.$$

Let us define by induction a sequence  $x_n \in \mathbb{R}^k$  for player 1 as follows, where  $k = |J|$ .

The first term  $x_1$  is any row of the matrix  $A$ . Given  $x_1, x_2, \dots, x_n$ , define  $x_{n+1}$  as follows :

## Learning to play optimally using Blackwell

Let  $A$  be an  $I \times J$  matrix and assume that the minmax is 0 :

$$\bar{v} = \min_{s \in \Delta(I)} \max_{i \in I} e^i A t = 0.$$

Let us define by induction a sequence  $x_n \in \mathbb{R}^k$  for player 1 as follows, where  $k = |J|$ .

The first term  $x_1$  is any row of the matrix  $A$ . Given  $x_1, x_2, \dots, x_n$ , define  $x_{n+1}$  as follows :

- Let  $\bar{x}_n^+$  be the vector with  $j^{th}$  coordinate equal to  $\max(\bar{x}_n^j, 0)$ .

Let  $A$  be an  $I \times J$  matrix and assume that the minmax is 0 :

$$\bar{v} = \min_{s \in \Delta(I)} \max_{i \in I} e^i A s = 0.$$

Let us define by induction a sequence  $x_n \in \mathbb{R}^k$  for player 1 as follows, where  $k = |J|$ .

The first term  $x_1$  is any row of the matrix  $A$ . Given  $x_1, x_2, \dots, x_n$ , define  $x_{n+1}$  as follows :

- Let  $\bar{x}_n^+$  be the vector with  $j^{th}$  coordinate equal to  $\max(\bar{x}_n^j, 0)$ .
- If  $\bar{x}_n = \bar{x}_n^+$  then choose  $x_{n+1}$  as any row of  $A$ .

Let  $A$  be an  $I \times J$  matrix and assume that the minmax is 0 :

$$\bar{v} = \min_{s \in \Delta(I)} \max_{i \in I} e^i A t = 0.$$

Let us define by induction a sequence  $x_n \in \mathbb{R}^k$  for player 1 as follows, where  $k = |J|$ .

The first term  $x_1$  is any row of the matrix  $A$ . Given  $x_1, x_2, \dots, x_n$ , define  $x_{n+1}$  as follows :

- Let  $\bar{x}_n^+$  be the vector with  $j^{th}$  coordinate equal to  $\max(\bar{x}_n^j, 0)$ .
- If  $\bar{x}_n = \bar{x}_n^+$  then choose  $x_{n+1}$  as any row of  $A$ .
- Otherwise let  $a > 0$  such that

$$t_{n+1} = \frac{\bar{x}_n^+ - \bar{x}_n}{a} \in \Delta(J).$$

Let  $A$  be an  $I \times J$  matrix and assume that the minmax is 0 :

$$\bar{v} = \min_{s \in \Delta(I)} \max_{i \in I} e^i A s = 0.$$

Let us define by induction a sequence  $x_n \in \mathbb{R}^k$  for player 1 as follows, where  $k = |J|$ .

The first term  $x_1$  is any row of the matrix  $A$ . Given  $x_1, x_2, \dots, x_n$ , define  $x_{n+1}$  as follows :

- Let  $\bar{x}_n^+$  be the vector with  $j^{th}$  coordinate equal to  $\max(\bar{x}_n^j, 0)$ .
- If  $\bar{x}_n = \bar{x}_n^+$  then choose  $x_{n+1}$  as any row of  $A$ .
- Otherwise let  $a > 0$  such that

$$t_{n+1} = \frac{\bar{x}_n^+ - \bar{x}_n}{a} \in \Delta(J).$$

- Since  $\bar{v} = 0$ , there exists an  $i_{n+1} \in I$  such that  $i_{n+1} A t_{n+1} \geq 0$ .

Let  $A$  be an  $I \times J$  matrix and assume that the minmax is 0 :

$$\bar{v} = \min_{s \in \Delta(I)} \max_{i \in I} e^i A s = 0.$$

Let us define by induction a sequence  $x_n \in \mathbb{R}^k$  for player 1 as follows, where  $k = |J|$ .

The first term  $x_1$  is any row of the matrix  $A$ . Given  $x_1, x_2, \dots, x_n$ , define  $x_{n+1}$  as follows :

- Let  $\bar{x}_n^+$  be the vector with  $j^{th}$  coordinate equal to  $\max(\bar{x}_n^j, 0)$ .
- If  $\bar{x}_n = \bar{x}_n^+$  then choose  $x_{n+1}$  as any row of  $A$ .
- Otherwise let  $a > 0$  such that

$$t_{n+1} = \frac{\bar{x}_n^+ - \bar{x}_n}{a} \in \Delta(J).$$

- Since  $\bar{v} = 0$ , there exists an  $i_{n+1} \in I$  such that  $i_{n+1} A t_{n+1} \geq 0$ .
- $x_{n+1}$  is then the  $i_{n+1}$ th row of the matrix  $A$ .



Let  $A$  be an  $I \times J$  matrix and assume that the minmax is 0 :

$$\bar{v} = \min_{s \in \Delta(I)} \max_{i \in I} e^i A t = 0.$$

Let us define by induction a sequence  $x_n \in \mathbb{R}^k$  for player 1 as follows, where  $k = |J|$ .

The first term  $x_1$  is any row of the matrix  $A$ . Given  $x_1, x_2, \dots, x_n$ , define  $x_{n+1}$  as follows :

- Let  $\bar{x}_n^+$  be the vector with  $j^{th}$  coordinate equal to  $\max(\bar{x}_n^j, 0)$ .
- If  $\bar{x}_n = \bar{x}_n^+$  then choose  $x_{n+1}$  as any row of  $A$ .
- Otherwise let  $a > 0$  such that

$$t_{n+1} = \frac{\bar{x}_n^+ - \bar{x}_n}{a} \in \Delta(J).$$

- Since  $\bar{v} = 0$ , there exists an  $i_{n+1} \in I$  such that  $i_{n+1} A t_{n+1} \geq 0$ .
- $x_{n+1}$  is then the  $i_{n+1}$ th row of the matrix  $A$ .

## Theorem

$\{x_n\}$  is a Blackwell  $C$ -sequence with  $C = \{x \in \mathbb{R}^k; x \geq 0\}$ . Consequently, there is  $s \in \Delta(I)$  with  $s A t \geq 0$ , for all  $t \in \Delta(J)$ .

By construction :  $\langle i_{n+1}A, t_{n+1} \rangle \geq 0$ ,

By construction :  $\langle i_{n+1}A, t_{n+1} \rangle \geq 0$ ,

thus

$$\langle x_{n+1}, \bar{x}_n^+ - \bar{x}_n \rangle \geq 0.$$

By construction :  $\langle i_{n+1}A, t_{n+1} \rangle \geq 0$ ,

thus

$$\langle x_{n+1}, \bar{x}_n^+ - \bar{x}_n \rangle \geq 0.$$

Since  $\langle \bar{x}_n^+, \bar{x}_n - \bar{x}_n^+ \rangle = 0$

By construction :  $\langle i_{n+1}A, t_{n+1} \rangle \geq 0$ ,

thus

$$\langle x_{n+1}, \bar{x}_n^+ - \bar{x}_n \rangle \geq 0.$$

Since  $\langle \bar{x}_n^+, \bar{x}_n - \bar{x}_n^+ \rangle = 0$

we get

$$\langle x_{n+1} - \bar{x}_n^+, \bar{x}_n - \bar{x}_n^+ \rangle \leq 0$$

By construction :  $\langle i_{n+1}A, t_{n+1} \rangle \geq 0$ ,

thus

$$\langle x_{n+1}, \bar{x}_n^+ - \bar{x}_n \rangle \geq 0.$$

Since  $\langle \bar{x}_n^+, \bar{x}_n - \bar{x}_n^+ \rangle = 0$

we get

$$\langle x_{n+1} - \bar{x}_n^+, \bar{x}_n - \bar{x}_n^+ \rangle \leq 0$$

which is what we want since  $\bar{x}_n^+ = \Pi_C(\bar{x}_n)$ .

By construction :  $\langle i_{n+1}A, t_{n+1} \rangle \geq 0$ ,

thus

$$\langle x_{n+1}, \bar{x}_n^+ - \bar{x}_n \rangle \geq 0.$$

Since  $\langle \bar{x}_n^+, \bar{x}_n - \bar{x}_n^+ \rangle = 0$

we get

$$\langle x_{n+1} - \bar{x}_n^+, \bar{x}_n - \bar{x}_n^+ \rangle \leq 0$$

which is what we want since  $\bar{x}_n^+ = \Pi_C(\bar{x}_n)$ .

To conclude. Consider the empirical frequencies arising in  $\bar{x}_n$  as a mixed strategy of player 1 and use compactness of  $\Delta(I)$  to deduce that its limit provides a strategy  $s \in \Delta(I)$  which satisfies  $sAt \geq 0$ , for all  $t \in \Delta(J)$ .