

Online Learning in Games

Lecture 2: Zero-sum games with infinitely many actions

Rida Laraki and Guillaume Vigeral

Master IASD, 2024

From previous lecture

- A **zero-sum game** is a two player game $G = (I, J, g)$ where g is a bounded function from $I \times J$ to \mathbb{R} that player 1 maximizes and player 2 minimizes.

From previous lecture

- A **zero-sum game** is a two player game $G = (I, J, g)$ where g is a bounded function from $I \times J$ to \mathbb{R} that player 1 maximizes and player 2 minimizes.
- In this lecture, we are interested by the infinite case (I and/or J is infinite). **Examples : duels, biggest integer, electoral competition, GANs.**

From previous lecture

- A **zero-sum game** is a two player game $G = (I, J, g)$ where g is a bounded function from $I \times J$ to \mathbb{R} that player 1 maximizes and player 2 minimizes.
- In this lecture, we are interested by the infinite case (I and/or J is infinite). **Examples : duels, biggest integer, electoral competition, GANs.**
- The **maxmin** \underline{v} is the highest quantity player 1 can guarantee

$$\underline{v} = \sup_{i \in I} \inf_{j \in J} g(i, j)$$

The **minmax** \bar{v} is :

$$\bar{v} = \inf_{j \in J} \sup_{i \in I} g(i, j)$$

From previous lecture

- A **zero-sum game** is a two player game $G = (I, J, g)$ where g is a bounded function from $I \times J$ to \mathbb{R} that player 1 maximizes and player 2 minimizes.
- In this lecture, we are interested by the infinite case (I and/or J is infinite). **Examples : duels, biggest integer, electoral competition, GANs.**
- The **maxmin** \underline{v} is the highest quantity player 1 can guarantee

$$\underline{v} = \sup_{i \in I} \inf_{j \in J} g(i, j)$$

The **minmax** \bar{v} is :

$$\bar{v} = \inf_{j \in J} \sup_{i \in I} g(i, j)$$

- The **game has a value** v if $\underline{v} = \bar{v} = v$.

From previous lecture

- A **zero-sum game** is a two player game $G = (I, J, g)$ where g is a bounded function from $I \times J$ to \mathbb{R} that player 1 maximizes and player 2 minimizes.
- In this lecture, we are interested by the infinite case (I and/or J is infinite). **Examples : duels, biggest integer, electoral competition, GANs.**
- The **maxmin** \underline{v} is the highest quantity player 1 can guarantee

$$\underline{v} = \sup_{i \in I} \inf_{j \in J} g(i, j)$$

The **minmax** \bar{v} is :

$$\bar{v} = \inf_{j \in J} \sup_{i \in I} g(i, j)$$

- The **game has a value** v if $\underline{v} = \bar{v} = v$.
- When a value exists, a strategy that achieves the argmax in the maxmin (resp. the argmin in the minmax) is called **optimal** for P1 (resp. P2).

From previous lecture

- A **zero-sum game** is a two player game $G = (I, J, g)$ where g is a bounded function from $I \times J$ to \mathbb{R} that player 1 maximizes and player 2 minimizes.
- In this lecture, we are interested by the infinite case (I and/or J is infinite). **Examples : duels, biggest integer, electoral competition, GANs.**
- The **maxmin** \underline{v} is the highest quantity player 1 can guarantee

$$\underline{v} = \sup_{i \in I} \inf_{j \in J} g(i, j)$$

The **minmax** \bar{v} is :

$$\bar{v} = \inf_{j \in J} \sup_{i \in I} g(i, j)$$

- The **game has a value** v if $\underline{v} = \bar{v} = v$.
- When a value exists, a strategy that achieves the argmax in the maxmin (resp. the argmin in the minmax) is called **optimal** for P1 (resp. P2).
- Question : **general sufficient conditions for existence of the value** (either in pure or mixed strategies).

Contents

- 1 Pure Strategies
- 2 Mixed Strategies
- 3 Fictitious Play
- 4 Application to GANs

Berge lemma

Lemma (Berge, 1965)

Let C_1, \dots, C_n be non-empty convex compact subsets of a Euclidean space. Assume that the union $\bigcup_{i=1}^n C_i$ is convex and that for each $j = 1, \dots, n$, $\bigcap_{i \neq j} C_i$ is non-empty.

Berge lemma

Lemma (Berge, 1965)

Let C_1, \dots, C_n be non-empty convex compact subsets of a Euclidean space. Assume that the union $\bigcup_{i=1}^n C_i$ is convex and that for each $j = 1, \dots, n$, $\bigcap_{i \neq j} C_i$ is non-empty. *Then $\bigcap_{i=1}^n C_i$ is non-empty.*

Proof of Berge lemma 1

- The lemma is clear if $n = 1$. Let $n \geq 2$.

Proof of Berge lemma 1

- The lemma is clear if $n = 1$. Let $n \geq 2$.
- By contradiction, suppose the lemma holds for $n - 1$ but not for n .

Proof of Berge lemma 1

- The lemma is clear if $n = 1$. Let $n \geq 2$.
- By contradiction, suppose the lemma holds for $n - 1$ but not for n .
- Let C_1, \dots, C_n satisfy the hypotheses but have empty intersection.

Proof of Berge lemma 1

- The lemma is clear if $n = 1$. Let $n \geq 2$.
- By contradiction, suppose the lemma holds for $n - 1$ but not for n .
- Let C_1, \dots, C_n satisfy the hypotheses but have empty intersection.
- Then C_n and $\bigcap_{i=1}^{n-1} C_i := D_n$ are $\neq \emptyset$, disjoint, convex and compact.

Proof of Berge lemma 1

- The lemma is clear if $n = 1$. Let $n \geq 2$.
- By contradiction, suppose the lemma holds for $n - 1$ but not for n .
- Let C_1, \dots, C_n satisfy the hypotheses but have empty intersection.
- Then C_n and $\bigcap_{i=1}^{n-1} C_i := D_n$ are $\neq \emptyset$, disjoint, convex and compact.
- By Hahn–Banach, C_n and D_n can be strictly separated by a hyperplane H .

Proof of Berge lemma 2

Proof of Berge lemma 2

Proof of Berge lemma 2

Proof of Berge lemma 2

Proof of Berge lemma 2

- By Hahn–Banach, C_n and D_n can be strictly separated by a hyperplane H .

Proof of Berge lemma 2

- Define $\tilde{C}_i = C_i \cap H$ for $i = 1, \dots, n-1$, and $\tilde{C} = (\bigcup_{i=1}^n C_i) \cap H$.

Proof of Berge lemma 2

- Since $C_n \cap H = \emptyset = D_n \cap H$, we have $\bigcup_{i=1}^{n-1} \tilde{C}_i = \tilde{C}$ and $\bigcap_{i=1}^{n-1} \tilde{C}_i = \emptyset$.

Proof of Berge lemma 2

- By the induction hypothesis, $\exists j \in \{1, \dots, n-1\}$ such that $\bigcap_{i \neq j, n} \tilde{C}_i = \emptyset$.

Proof of Berge lemma 2

- Let $K = \bigcap_{i \neq j, n} C_i$. Then $D_n \subset K$ and $C_n \cap K \neq \emptyset$.

Proof of Berge lemma 2

- As K is convex, we must have $K \cap H \neq \emptyset$.

Proof of Berge lemma 2

- But $K \cap H = \bigcap_{i \neq j, n} \tilde{C}_i = \emptyset$, a contradiction.

Definitions

- Let E be a convex subset of a topological vector space.

Definitions

- Let E be a convex subset of a topological vector space.
- A real valued function $f : E \rightarrow \mathbb{R}$, is called **quasi-convex** if for all $\lambda \in \mathbb{R}$, $\{x \in E, f(x) \leq \lambda\}$ is convex.

Definitions

- Let E be a convex subset of a topological vector space.
- A real valued function $f : E \rightarrow \mathbb{R}$, is called **quasi-convex** if for all $\lambda \in \mathbb{R}$, $\{x \in E, f(x) \leq \lambda\}$ is convex.
- f is **quasi-concave** if for all $\lambda \in \mathbb{R}$, $\{x \in E, f(x) \geq \lambda\}$ is convex.

Definitions

- Let E be a convex subset of a topological vector space.
- A real valued function $f : E \rightarrow \mathbb{R}$, is called **quasi-convex** if for all $\lambda \in \mathbb{R}$, $\{x \in E, f(x) \leq \lambda\}$ is convex.
- f is **quasi-concave** if for all $\lambda \in \mathbb{R}$, $\{x \in E, f(x) \geq \lambda\}$ is convex.
- f is **lower-semicontinuous** (l.s.c) if for all λ , $\{x \in E, f(x) \leq \lambda\}$ is closed.

Definitions

- Let E be a convex subset of a topological vector space.
- A real valued function $f : E \rightarrow \mathbb{R}$, is called **quasi-convex** if for all $\lambda \in \mathbb{R}$, $\{x \in E, f(x) \leq \lambda\}$ is convex.
- f is **quasi-concave** if for all $\lambda \in \mathbb{R}$, $\{x \in E, f(x) \geq \lambda\}$ is convex.
- f is **lower-semicontinuous** (l.s.c) if for all λ , $\{x \in E, f(x) \leq \lambda\}$ is closed.
- f is **upper-semicontinuous** (u.s.c.) if for all λ , $\{x \in E, f(x) \geq \lambda\}$ is closed.

Definitions

- Let E be a convex subset of a topological vector space.
- A real valued function $f : E \rightarrow \mathbb{R}$, is called **quasi-convex** if for all $\lambda \in \mathbb{R}$, $\{x \in E, f(x) \leq \lambda\}$ is convex.
- f is **quasi-concave** if for all $\lambda \in \mathbb{R}$, $\{x \in E, f(x) \geq \lambda\}$ is convex.
- f is **lower-semicontinuous** (l.s.c) if for all λ , $\{x \in E, f(x) \leq \lambda\}$ is closed.
- f is **upper-semicontinuous** (u.s.c.) if for all λ , $\{x \in E, f(x) \geq \lambda\}$ is closed.
- **Remark** : if E is compact and f u.s.c (resp l.s.c), then f achieves its maximum on E (resp. minimum).

Sion minmax theorem

Theorem (Sion, 1958)

Let $G = (I, J, g)$ be a zero-sum game satisfying :

- (i) I and J are convex;

Sion minmax theorem

Theorem (Sion, 1958)

Let $G = (I, J, g)$ be a zero-sum game satisfying :

- (i) I and J are convex;
- (ii) I or J is compact;

Sion minmax theorem

Theorem (Sion, 1958)

Let $G = (I, J, g)$ be a zero-sum game satisfying :

- (i) I and J are convex;
- (ii) I or J is compact;
- (iii) for each j in J , $g(\cdot, j)$ is quasi-concave u.s.c. in i ,

Sion minmax theorem

Theorem (Sion, 1958)

Let $G = (I, J, g)$ be a zero-sum game satisfying :

- (i) I and J are convex;
- (ii) I or J is compact;
- (iii) for each j in J , $g(\cdot, j)$ is quasi-concave u.s.c. in i ,
- (iv) for each i in I , $g(i, \cdot)$ is quasi-convex l.s.c. in j .

Sion minmax theorem

Theorem (Sion, 1958)

Let $G = (I, J, g)$ be a zero-sum game satisfying :

- (i) I and J are convex;
- (ii) I or J is compact;
- (iii) for each j in J , $g(\cdot, j)$ is quasi-concave u.s.c. in i ,
- (iv) for each i in I , $g(i, \cdot)$ is quasi-convex l.s.c. in j .

Then G has a value :

$$\sup_{i \in I} \inf_{j \in J} g(i, j) = \inf_{j \in J} \sup_{i \in I} g(i, j).$$

If I or J is compact, the corresponding player has an optimal strategy.

Sion minmax theorem

Theorem (Sion, 1958)

Let $G = (I, J, g)$ be a zero-sum game satisfying :

- (i) I and J are convex;
- (ii) I or J is compact;
- (iii) for each j in J , $g(\cdot, j)$ is quasi-concave u.s.c. in i ,
- (iv) for each i in I , $g(i, \cdot)$ is quasi-convex l.s.c. in j .

Then G has a value :

$$\sup_{i \in I} \inf_{j \in J} g(i, j) = \inf_{j \in J} \sup_{i \in I} g(i, j).$$

If I or J is compact, the corresponding player has an optimal strategy.

Proof Sion 1

- Assume I compact. By contradiction suppose there is v such that

$$\sup_{i \in I} \inf_{j \in J} g(i, j) < v < \inf_{j \in J} \sup_{i \in I} g(i, j).$$

Remark : if g is bilinear this is over because this contradicts von Neumann result in the finite case.

Proof Sion 1

- Assume I compact. By contradiction suppose there is v such that

$$\sup_{i \in I} \inf_{j \in J} g(i, j) < v < \inf_{j \in J} \sup_{i \in I} g(i, j).$$

- Define for each j in J the set $I_j = \{i \in I, g(i, j) < v\}$.

Remark : if g is bilinear this is over because this contradicts von Neumann result in the finite case.

Proof Sion 1

- Assume I compact. By contradiction suppose there is v such that

$$\sup_{i \in I} \inf_{j \in J} g(i, j) < v < \inf_{j \in J} \sup_{i \in I} g(i, j).$$

- Define for each j in J the set $I_j = \{i \in I, g(i, j) < v\}$.
- $(I_j)_{j \in J}$ is an open covering of I , hence \exists a finite subset J_0 of J s.t.

$$I = \bigcup_{j \in J_0} I_j$$

Remark : if g is bilinear this is over because this contradicts von Neumann result in the finite case.

Proof Sion 1

- Assume I compact. **By contradiction** suppose there is v such that

$$\sup_{i \in I} \inf_{j \in J} g(i, j) < v < \inf_{j \in J} \sup_{i \in I} g(i, j).$$

- Define for each j in J the set $I_j = \{i \in I, g(i, j) < v\}$.
- $(I_j)_{j \in J}$ is **an open covering of I** , hence \exists a finite subset J_0 of J s.t.

$$I = \bigcup_{j \in J_0} I_j$$

- Define $J' = \text{co}(J_0)$, then J' is compact and $\max_{i \in I} \inf_{j \in J'} g(i, j) < v < \inf_{j \in J'} \sup_{i \in I} g(i, j)$.

Remark : if g is bilinear this is over because this contradicts von Neumann result in the finite case.

Proof Sion 1

- Assume I compact. **By contradiction** suppose there is v such that

$$\sup_{i \in I} \inf_{j \in J} g(i, j) < v < \inf_{j \in J} \sup_{i \in I} g(i, j).$$

- Define for each j in J the set $I_j = \{i \in I, g(i, j) < v\}$.
- $(I_j)_{j \in J}$ is **an open covering of I** , hence \exists a finite subset J_0 of J s.t.

$$I = \bigcup_{j \in J_0} I_j$$

- Define $J' = \text{co}(J_0)$, then J' is compact and $\max_{i \in I} \inf_{j \in J'} g(i, j) < v < \inf_{j \in J'} \sup_{i \in I} g(i, j)$.
- Similarly, \exists a finite subset I_0 of I s.t.**

$$\forall i \in \text{co}(I_0), \exists j \in J_0, \quad g(i, j) < v,$$

$$\forall j \in \text{co}(J_0), \exists i \in I_0, \quad g(i, j) > v.$$

Remark : if g is bilinear this is over because this contradicts von Neumann result in the finite case.

Proof Sion 2

- We proved that, \exists a finite subset S_0 of S s.t.

$$\forall i \in \text{co}(I_0), \exists j \in J_0, \quad g(i, j) < v,$$

$$\forall j \in \text{co}(J_0), \exists i \in I_0, \quad g(i, j) > v.$$

Proof Sion 2

- We proved that, \exists a finite subset S_0 of S s.t.

$$\forall i \in \text{co}(I_0), \exists j \in J_0, \quad g(i, j) < v,$$

$$\forall j \in \text{co}(J_0), \exists i \in I_0, \quad g(i, j) > v.$$

- Let (I_0, J_0) be a minimal pair (for inclusion) satisfying the property.

Proof Sion 2

- We proved that, \exists a finite subset S_0 of S s.t.

$$\forall i \in \text{co}(I_0), \exists j \in J_0, \quad g(i, j) < v,$$

$$\forall j \in \text{co}(J_0), \exists i \in I_0, \quad g(i, j) > v.$$

- Let (I_0, J_0) be a minimal pair (for inclusion) satisfying the property.
- For each $i \in I_0$, define now $A_i = \{j \in \text{co}(J_0), g(i, j) \leq v\}$.

Proof Sion 2

- We proved that, \exists a finite subset S_0 of S s.t.

$$\forall i \in \text{co}(I_0), \exists j \in J_0, \quad g(i, j) < v,$$

$$\forall j \in \text{co}(J_0), \exists i \in I_0, \quad g(i, j) > v.$$

- Let (I_0, J_0) be a minimal pair (for inclusion) satisfying the property.
- For each $i \in I_0$, define now $A_i = \{j \in \text{co}(J_0), g(i, j) \leq v\}$.
- $\bigcap_{i \in I_0} A_i = \emptyset$

Proof Sion 2

- We proved that, \exists a finite subset S_0 of S s.t.

$$\forall i \in \text{co}(I_0), \exists j \in J_0, \quad g(i, j) < v,$$

$$\forall j \in \text{co}(J_0), \exists i \in I_0, \quad g(i, j) > v.$$

- Let (I_0, J_0) be a minimal pair (for inclusion) satisfying the property.
- For each $i \in I_0$, define now $A_i = \{j \in \text{co}(J_0), g(i, j) \leq v\}$.
- $\bigcap_{i \in I_0} A_i = \emptyset$
- By minimality of I_0 , $\bigcap_{i \in I_0 \setminus \{i_0\}} A_i \neq \emptyset$ for each i_0 in I_0 .

Proof Sion 2

- We proved that, \exists a finite subset S_0 of S s.t.

$$\forall i \in \text{co}(I_0), \exists j \in J_0, \quad g(i, j) < v,$$

$$\forall j \in \text{co}(J_0), \exists i \in I_0, \quad g(i, j) > v.$$

- Let (I_0, J_0) be a minimal pair (for inclusion) satisfying the property.
- For each $i \in I_0$, define now $A_i = \{j \in \text{co}(J_0), g(i, j) \leq v\}$.
- $\bigcap_{i \in I_0} A_i = \emptyset$
- By minimality of I_0 , $\bigcap_{i \in I_0 \setminus \{i_0\}} A_i \neq \emptyset$ for each i_0 in I_0 .
- By Berge lemma, the union $\bigcup_{i \in S_0} A_i$ is not convex.

Proof Sion 2

- We proved that, \exists a finite subset S_0 of S s.t.

$$\forall i \in \text{co}(I_0), \exists j \in J_0, \quad g(i, j) < v,$$

$$\forall j \in \text{co}(J_0), \exists i \in I_0, \quad g(i, j) > v.$$

- Let (I_0, J_0) be a minimal pair (for inclusion) satisfying the property.
- For each $i \in I_0$, define now $A_i = \{j \in \text{co}(J_0), g(i, j) \leq v\}$.
- $\bigcap_{i \in I_0} A_i = \emptyset$
- By minimality of I_0 , $\bigcap_{i \in I_0 \setminus \{i_0\}} A_i \neq \emptyset$ for each i_0 in I_0 .
- By Berge lemma, the union $\bigcup_{i \in S_0} A_i$ is not convex.
- Consequently $\exists j_0$ in $\text{co}(J_0) \setminus \bigcup_{i \in I_0} A_i$.

Proof Sion 2

- We proved that, \exists a finite subset S_0 of S s.t.

$$\forall i \in \text{co}(I_0), \exists j \in J_0, \quad g(i, j) < v,$$

$$\forall j \in \text{co}(J_0), \exists i \in I_0, \quad g(i, j) > v.$$

- Let (I_0, J_0) be a minimal pair (for inclusion) satisfying the property.
- For each $i \in I_0$, define now $A_i = \{j \in \text{co}(J_0), g(i, j) \leq v\}$.
- $\bigcap_{i \in I_0} A_i = \emptyset$
- By minimality of I_0 , $\bigcap_{i \in I_0 \setminus \{i_0\}} A_i \neq \emptyset$ for each i_0 in I_0 .
- By Berge lemma, the union $\bigcup_{i \in S_0} A_i$ is not convex.
- Consequently $\exists j_0$ in $\text{co}(J_0) \setminus \bigcup_{i \in I_0} A_i$.
- Thus, $\forall i \in I_0$, $g(i, j_0) > v$ and by quasi-concavity, $g(i, j_0) > v \quad \forall i \in \text{co}(I_0)$.

Proof Sion 2

- We proved that, \exists a finite subset S_0 of S s.t.

$$\forall i \in \text{co}(I_0), \exists j \in J_0, \quad g(i, j) < v,$$

$$\forall j \in \text{co}(J_0), \exists i \in I_0, \quad g(i, j) > v.$$

- Let (I_0, J_0) be a minimal pair (for inclusion) satisfying the property.
- For each $i \in I_0$, define now $A_i = \{j \in \text{co}(J_0), g(i, j) \leq v\}$.
- $\bigcap_{i \in I_0} A_i = \emptyset$
- By minimality of I_0 , $\bigcap_{i \in I_0 \setminus \{i_0\}} A_i \neq \emptyset$ for each i_0 in I_0 .
- By Berge lemma, the union $\bigcup_{i \in S_0} A_i$ is not convex.
- Consequently $\exists j_0$ in $\text{co}(J_0) \setminus \bigcup_{i \in I_0} A_i$.
- Thus, $\forall i \in I_0, g(i, j_0) > v$ and by quasi-concavity, $g(i, j_0) > v \quad \forall i \in \text{co}(I_0)$.
- Similarly, $\exists s_0 \in \text{co}(I_0)$ s.t. $g(s_0, j) < v$ for each j in $\text{co}(J_0)$.
A contradiction.

Weakening the assumptions

We can weaken the topological conditions by strengthening the convexity hypothesis on $g(i, \cdot)$.

Proposition

Let $G = (I, J, g)$ be a zero-sum game such that :

- (i) I is convex and compact ;*

Weakening the assumptions

We can weaken the topological conditions by strengthening the convexity hypothesis on $g(i, \cdot)$.

Proposition

Let $G = (I, J, g)$ be a zero-sum game such that :

- (i) I is convex and compact ;
- (ii) J is convex ;

Weakening the assumptions

We can weaken the topological conditions by strengthening the convexity hypothesis on $g(i, \cdot)$.

Proposition

Let $G = (I, J, g)$ be a zero-sum game such that :

- (i) I is convex and compact ;
- (ii) J is convex ;
- (iii) for each j in J , $g(\cdot, j)$ is quasi-concave u.s.c., and for each i in I , $g(i, \cdot)$ is convex.

Weakening the assumptions

We can weaken the topological conditions by strengthening the convexity hypothesis on $g(i, \cdot)$.

Proposition

Let $G = (I, J, g)$ be a zero-sum game such that :

- (i) I is convex and compact ;
- (ii) J is convex ;
- (iii) for each j in J , $g(\cdot, j)$ is quasi-concave u.s.c., and for each i in I , $g(i, \cdot)$ is convex.

Then G has a value : $\sup_{i \in I} \inf_{j \in J} g(i, j) = \inf_{j \in J} \sup_{i \in I} g(i, j)$, and player 1 has an optimal strategy.

Contents

- 1 Pure Strategies
- 2 Mixed Strategies
- 3 Fictitious Play
- 4 Application to GANs

Mixed Extension

- We can convexify a set X either by considering the set $\Delta_f(X)$ of probabilities with finite support over X or, if X has a topological structure, by considering the set $\Delta(X)$ of regular probabilities over X .

Mixed Extension

- We can convexify a set X either by considering the set $\Delta_f(X)$ of probabilities with finite support over X or, if X has a topological structure, by considering the set $\Delta(X)$ of regular probabilities over X .
- Given a game (I, J, g) , the payoff function g can be multi-linearly extended to (finite support or regular) probability distributions :

$$g(\sigma, \tau) = \mathbb{E}_{\sigma \otimes \tau}(g) = \int_I \int_J g(i, j) d\sigma(i) d\tau(j)$$

Mixed Extension

- We can convexify a set X either by considering the set $\Delta_f(X)$ of probabilities with finite support over X or, if X has a topological structure, by considering the set $\Delta(X)$ of regular probabilities over X .
- Given a game (I, J, g) , the payoff function g can be multi-linearly extended to (finite support or regular) probability distributions :

$$g(\sigma, \tau) = \mathbb{E}_{\sigma \otimes \tau}(g) = \int_I \int_J g(i, j) d\sigma(i) d\tau(j)$$

- σ and τ are called mixed strategies.

Mixed Extension

- We can convexify a set X either by considering the set $\Delta_f(X)$ of probabilities with finite support over X or, if X has a topological structure, by considering the set $\Delta(X)$ of regular probabilities over X .
- Given a game (I, J, g) , the payoff function g can be multi-linearly extended to (finite support or regular) probability distributions :

$$g(\sigma, \tau) = \mathbb{E}_{\sigma \otimes \tau}(g) = \int_I \int_J g(i, j) d\sigma(i) d\tau(j)$$

- σ and τ are called mixed strategies.
- Elements of I and J are pure strategies.

Mixed Extension

- We can convexify a set X either by considering the set $\Delta_f(X)$ of probabilities with finite support over X or, if X has a topological structure, by considering the set $\Delta(X)$ of regular probabilities over X .
- Given a game (I, J, g) , the payoff function g can be multi-linearly extended to (finite support or regular) probability distributions :

$$g(\sigma, \tau) = \mathbb{E}_{\sigma \otimes \tau}(g) = \int_I \int_J g(i, j) d\sigma(i) d\tau(j)$$

- σ and τ are called mixed strategies.
- Elements of I and J are pure strategies.
- If X has a topological structure, $\Delta(X)$ is usually endowed with the weak* topology (the weakest topology such that $\hat{\phi} : \mu \mapsto \int_X \phi d\mu$ is continuous for each real continuous function ϕ on X).

Mixed Minmax (Sion)

Theorem

Let $G = (I, J, g)$ be a zero-sum game such that :

- (i) I and J are compact Hausdorff topological spaces ;

Mixed Minmax (Sion)

Theorem

Let $G = (I, J, g)$ be a zero-sum game such that :

- (i) I and J are compact Hausdorff topological spaces ;
- (ii) for each j in J , $g(\cdot, j)$ is u.s.c., and for each i in I , $g(i, \cdot)$ is l.s.c. ;

Mixed Minmax (Sion)

Theorem

Let $G = (I, J, g)$ be a zero-sum game such that :

- (i) I and J are compact Hausdorff topological spaces ;
- (ii) for each j in J , $g(\cdot, j)$ is u.s.c., and for each i in I , $g(i, \cdot)$ is l.s.c. ;
- (iii) g is bounded and measurable with respect to the product Borel σ -algebra $\mathcal{B}_I \otimes \mathcal{B}_J$.

Mixed Minmax (Sion)

Theorem

Let $G = (I, J, g)$ be a zero-sum game such that :

- (i) I and J are compact Hausdorff topological spaces ;
- (ii) for each j in J , $g(\cdot, j)$ is u.s.c., and for each i in I , $g(i, \cdot)$ is l.s.c. ;
- (iii) g is bounded and measurable with respect to the product Borel σ -algebra $\mathcal{B}_I \otimes \mathcal{B}_J$.

Then the mixed extension $(\Delta(I), \Delta(J), g)$ of G has a value. Each player has a mixed optimal strategy, and for each $\varepsilon > 0$ each player has an ε -optimal strategy with finite support.

Mixed Minmax (Sion)

Theorem

Let $G = (I, J, g)$ be a zero-sum game such that :

- (i) I and J are compact Hausdorff topological spaces ;
- (ii) for each j in J , $g(\cdot, j)$ is u.s.c., and for each i in I , $g(i, \cdot)$ is l.s.c. ;
- (iii) g is bounded and measurable with respect to the product Borel σ -algebra $\mathcal{B}_I \otimes \mathcal{B}_J$.

Then the mixed extension $(\Delta(I), \Delta(J), g)$ of G has a value. Each player has a mixed optimal strategy, and for each $\varepsilon > 0$ each player has an ε -optimal strategy with finite support.

Proof : a direct consequence of Sion minmax theorem in pure strategies.

Mixed minmax theorem (second version)

Proposition

Consider a zero-sum game (I, J, g) satisfying :

- (i) I is a compact Hausdorff topological space;

Mixed minmax theorem (second version)

Proposition

Consider a zero-sum game (I, J, g) satisfying :

- (i) I is a compact Hausdorff topological space ;
- (ii) for each j in J , $g(\cdot, j)$ is u.s.c.

Mixed minmax theorem (second version)

Proposition

Consider a zero-sum game (I, J, g) satisfying :

- (i) I is a compact Hausdorff topological space ;
- (ii) for each j in J , $g(\cdot, j)$ is u.s.c.

Then the game $(\Delta(I), \Delta_f(J), g)$ has a value and player 1 has an optimal strategy.

Mixed minmax theorem (second version)

Proposition

Consider a zero-sum game (I, J, g) satisfying :

- (i) I is a compact Hausdorff topological space ;
- (ii) for each j in J , $g(\cdot, j)$ is u.s.c.

Then the game $(\Delta(I), \Delta_f(J), g)$ has a value and player 1 has an optimal strategy.

Proof :

- When I is compact and $g(\cdot, t)$ u.s.c., then $\Delta(I)$ (endowed with the weak* topology) is compact and $(\sigma \mapsto g(\sigma, j) = \int_S g(i, j)\sigma(di))$ is u.s.c.

Mixed minmax theorem (second version)

Proposition

Consider a zero-sum game (I, J, g) satisfying :

- (i) I is a compact Hausdorff topological space ;
- (ii) for each j in J , $g(\cdot, j)$ is u.s.c.

Then the game $(\Delta(I), \Delta_f(J), g)$ has a value and player 1 has an optimal strategy.

Proof :

- When I is compact and $g(\cdot, t)$ u.s.c., then $\Delta(I)$ (endowed with the weak* topology) is compact and $(\sigma \mapsto g(\sigma, j) = \int_S g(i, j) \sigma(di))$ is u.s.c.
- Moreover, $g(\sigma, \tau)$ is well defined on $\Delta(I) \times \Delta_f(J)$ and is bilinear.

Mixed minmax theorem (second version)

Proposition

Consider a zero-sum game (I, J, g) satisfying :

- (i) I is a compact Hausdorff topological space ;
- (ii) for each j in J , $g(\cdot, j)$ is u.s.c.

Then the game $(\Delta(I), \Delta_f(J), g)$ has a value and player 1 has an optimal strategy.

Proof :

- When I is compact and $g(\cdot, t)$ u.s.c., then $\Delta(I)$ (endowed with the weak* topology) is compact and $(\sigma \mapsto g(\sigma, j) = \int_S g(i, j)\sigma(di))$ is u.s.c.
- Moreover, $g(\sigma, \tau)$ is well defined on $\Delta(I) \times \Delta_f(J)$ and is bilinear.
- Assumptions of last proposition are satisfied.

Contents

- 1 Pure Strategies
- 2 Mixed Strategies
- 3 Fictitious Play
- 4 Application to GANs

Fictitious play : Danskin (1954-1981)

Fictitious play can be defined for any zero sum game $G = (I, J, g)$, not necessarily finite.

Fictitious play : Danskin (1954-1981)

Fictitious play can be defined for any zero sum game $G = (I, J, g)$, not necessarily finite.

Principle

- We start from any (i_1, j_1) in $I \times J$.
- At each stage $n \geq 2$, each player will play an approximate best response to the sum of past behavior of the opponent up to stage $n - 1$.

Fictitious play : Danskin (1954-1981)

Fictitious play can be defined for any zero sum game $G = (I, J, g)$, not necessarily finite.

Principle

- We start from any (i_1, j_1) in $I \times J$.
- At each stage $n \geq 2$, each player will play an approximate best response to the sum of past behavior of the opponent up to stage $n - 1$.

Definition

A sequence $(i_n, j_n)_{n \geq 1}$ is an η -FP process for some fixed $\eta \geq 0$ if for each $n \geq 1$:

- i_{n+1} is an η -best response of player 1 against $y_n := \frac{1}{n} \sum_{t=1}^n \delta_{j_t} \in \Delta(J)$,
- j_{n+1} is an η -best response of player 2 against $x_n := \frac{1}{n} \sum_{t=1}^n \delta_{i_t} \in \Delta(I)$.

Fictitious play : Theorem

Theorem (Danskin, 1954-1981)

Let $(i_n, j_n)_{n \geq 1}$ be the realization of a η -fictitious play process. If the game is compact and continuous and if $val(g)$ denotes its value in mixed strategies, then $\forall \varepsilon > 0, \exists N, \forall n \geq N, \forall x \in \Delta(I), \forall y \in \Delta(J)$

$$g(x_n, y) \geq val(g) - \varepsilon - \eta \quad \text{and} \quad g(x, y_n) \leq val(g) + \varepsilon + \eta.$$

Contents

- 1 Pure Strategies
- 2 Mixed Strategies
- 3 Fictitious Play
- 4 Application to GANs

Generative adversarial networks

Generative adversarial networks

Generative adversarial networks

- The **generator** (G) tries to trick the **classifier** (C) into classifying its generated fake data as true samples from a target distribution.

Generative adversarial networks

- The **generator** (G) tries to trick the **classifier** (C) into classifying its generated fake data as true samples from a target distribution.
- **Mathematically** : let (X, P) be a probability space representing the target data distribution, and let (Z, Q) denote the generator's distribution (for example, noise).

Generative adversarial networks

- The **generator** (G) tries to trick the **classifier** (C) into classifying its generated fake data as true samples from a target distribution.
- **Mathematically** : let (X, P) be a probability space representing the target data distribution, and let (Z, Q) denote the generator's distribution (for example, noise).
- A **generator strategy** may then be viewed as a function $g \in \mathcal{G}$, where \mathcal{G} is a fixed subset of the set of measurable functions from Z to X .

Generative adversarial networks

- The **generator** (G) tries to trick the **classifier** (C) into classifying its generated fake data as true samples from a target distribution.
- **Mathematically** : let (X, P) be a probability space representing the target data distribution, and let (Z, Q) denote the generator's distribution (for example, noise).
- A **generator strategy** may then be viewed as a function $g \in \mathcal{G}$, where \mathcal{G} is a fixed subset of the set of measurable functions from Z to X .
- A **classifier strategy** is a function $c \in \mathcal{C}$, where \mathcal{C} is a fixed subset of measurable functions from X to $[0, 1]$.

Generative adversarial networks

- The **generator** (G) tries to trick the **classifier** (C) into classifying its generated fake data as true samples from a target distribution.
- **Mathematically** : let (X, P) be a probability space representing the target data distribution, and let (Z, Q) denote the generator's distribution (for example, noise).
- A **generator strategy** may then be viewed as a function $g \in \mathcal{G}$, where \mathcal{G} is a fixed subset of the set of measurable functions from Z to X .
- A **classifier strategy** is a function $c \in \mathcal{C}$, where \mathcal{C} is a fixed subset of measurable functions from X to $[0, 1]$.
- The **payoff function for this zero-sum game** is given by is :

$$\Phi(g, c) = \int_X \log(c(x)) dP(x) + \int_Z \log(1 - c(g(z))) dQ(z) \quad (1)$$

$$= \int_X \log(c(x)) dP(x) + \int_X \log(1 - c(x)) dQ(g^{-1}(x)) \quad (2)$$

Generative adversarial networks

- The **generator** (G) tries to trick the **classifier** (C) into classifying its generated fake data as true samples from a target distribution.
- **Mathematically** : let (X, P) be a probability space representing the target data distribution, and let (Z, Q) denote the generator's distribution (for example, noise).
- A **generator strategy** may then be viewed as a function $g \in \mathcal{G}$, where \mathcal{G} is a fixed subset of the set of measurable functions from Z to X .
- A **classifier strategy** is a function $c \in \mathcal{C}$, where \mathcal{C} is a fixed subset of measurable functions from X to $[0, 1]$.
- The **payoff function for this zero-sum game** is given by is :

$$\Phi(g, c) = \int_X \log(c(x)) dP(x) + \int_Z \log(1 - c(g(z))) dQ(z) \quad (1)$$

$$= \int_X \log(c(x)) dP(x) + \int_X \log(1 - c(x)) dQ(g^{-1}(x)) \quad (2)$$

- The G-player minimizes with respect to g , the C-player maximizes in c .

Resolution

Proposition

If \mathcal{G} is the whole set of measurable functions from Z to X , then the game has a value in pure strategies and optimal strategies are

- $c^*(x) = 1/2 \forall x$ for player 1
- g^* such that $P = Q \circ (g^*)^{-1}$

Resolution

Proposition

If \mathcal{G} is the whole set of measurable functions from Z to X , then the game has a value in pure strategies and optimal strategies are

- $c^*(x) = 1/2 \forall x$ for player 1
- g^* such that $P = Q \circ (g^*)^{-1}$

Proof : (c^*, g^*) is a saddle point.

Resolution

Proposition

If \mathcal{G} is the whole set of measurable functions from Z to X , then the game has a value in pure strategies and optimal strategies are

- $c^*(x) = 1/2 \forall x$ for player 1
- g^* such that $P = Q \circ (g^*)^{-1}$

Proof : (c^*, g^*) is a saddle point.

Proposition

Assume that \mathcal{G} is any set, and that \mathcal{C} is convex compact. Then the game has a value in mixed strategy and player 1 has a pure optimal strategy.

Resolution

Proposition

If \mathcal{G} is the whole set of measurable functions from Z to X , then the game has a value in pure strategies and optimal strategies are

- $c^*(x) = 1/2 \forall x$ for player 1
- g^* such that $P = Q \circ (g^*)^{-1}$

Proof : (c^*, g^*) is a saddle point.

Proposition

Assume that \mathcal{G} is any set, and that \mathcal{C} is convex compact. Then the game has a value in mixed strategy and player 1 has a pure optimal strategy.

Proof :

Apply Sion's theorem to $I = \mathcal{C}$ and $J = \Delta_f(\mathcal{G})$. Observe that Φ is continuous, and concave in c .