

Serre vanishing theorem in the twisted primitive cohomology

NCG Summer School at DUT

Hao Zhuang

BICMR, PKU

July 16, 2026

Motivation in complex geometry

Suppose that

- ◇ X is a compact complex manifold,
- ◇ L is a positive holomorphic line bundle over X ,
- ◇ E is a holomorphic vector bundle over X .

Theorem (Serre, 1955)

There is a $p_0 \in \mathbb{N}$ such that the cohomology (of the sheaf of sections)

$$H^k(X, L^{\otimes p} \otimes E) = 0$$

for all $p \geq p_0$ and $k \geq 1$.

Motivation in complex geometry

Symplectic:

- ◇ primitive cohomology (Tseng-Yau, 2012),
- ◇ filtered cohomology (Tsai-Tseng-Yau, 2016),

Almost complex:

- ◇ generalized Dolbeault cohomology (Cirici-Wilson, 2021).

...

Question

Does Serre's vanishing theorem generalize to these “complex-like” cases?

Theorem (Z, 2026)

YES for the twisted Tseng-Yau primitive cohomology of a compact symplectic manifold.

Twisted primitive cohomology

We let

- ◇ (M, ω) : compact symplectic manifold,
- ◇ d : de Rham exterior derivative.

We obtain a cochain complex $(0 \leq k \leq 1 + \dim M)$:

$$\begin{aligned} d^\omega : \Omega^k(M) \oplus \Omega^{k-1}(M) &\rightarrow \Omega^{k+1}(M) \oplus \Omega^k(M) \\ (\alpha, \beta) &\mapsto (d\alpha + \omega \wedge \beta, -d\beta). \end{aligned}$$

Definition & Theorem (Tseng-Yau, 2012; Tanaka-Tseng, 2018)

The cohomology defined by $\Omega^\bullet(M) \oplus \Omega^{\bullet-1}(M)$ and d^ω is called the primitive cohomology of (M, ω) .

Twisted primitive cohomology

Originally by Tseng-Yau: Via the Lefschetz decomposition.

$\Leftrightarrow$

Current mapping cone model by Tanaka-Tseng.

How about the “twist”?

Definition (Tseng-Zhou, 2022)

For a connection ∇ on a smooth vector bundle $E \rightarrow M$, if there is a $\Phi \in \Gamma(\text{End}(E))$ such that

$$\begin{aligned}\nabla^2 + \omega\Phi &= 0, \\ \Phi\nabla &= \nabla\Phi,\end{aligned}$$

then we say (E, ∇, Φ) is symplectically flat.

Twisted primitive cohomology

For symplectically flat (E, ∇, Φ) , we have the cochain complex $(0 \leq k \leq 1 + \dim M)$:

$$\begin{aligned} \mathbb{A} : \Omega^k(M, E) \oplus \Omega^{k-1}(M, E) &\rightarrow \Omega^{k+1}(M, E) \oplus \Omega^k(M, E) \\ (\alpha, \beta) &\mapsto (\nabla\alpha + \omega \wedge \beta, \Phi\alpha - d\beta). \end{aligned}$$

Definition & Theorem (Tseng-Zhou, 2022)

The cohomology defined by $\Omega^\bullet(M, E) \oplus \Omega^{\bullet-1}(M, E)$ and $\mathbb{A}$ is called the primitive cohomology of (M, ω) twisted by (E, ∇, Φ) .

This twisted cohomology also admits a Lefschetz decomposition model.

Vanishing theorems

Given a symplectically flat (E, ∇, Φ) over (M, ω) :

Question

What if Φ is an automorphism?

The answer is a vanishing theorem.

Theorem (Tseng-Zhou, 2022)

When Φ is invertible, the k -th primitive cohomology group of (M, ω) twisted by (E, ∇, Φ) vanishes for all $0 \leq k \leq 1 + \dim M$.

This cohomology vanishes from $k = 0$, not just from $k = 1$.

Question

What if Φ is just an endomorphism?

Vanishing theorems

(L, ∇', Φ') :

- ◇ Symplectically flat (real) line bundle over (M, ω) .
- ◇ $\Phi' \neq 0$ everywhere.

Naturally induced on $E \otimes L^{\otimes p}$:

- ◇ a connection ∇'' ,
- ◇ an endomorphism $\Phi'' \in \Gamma(\text{End}(E \otimes L^{\otimes p}))$.

The answer is a Serre type vanishing theorem:

Theorem (Z, 2026)

There is a $p_0 \in \mathbb{N}$ such that when $p \geq p_0$, the k -th primitive cohomology group of (M, ω) twisted by $(E \otimes L^{\otimes p}, \nabla'', \Phi'')$ vanishes for all k .

You can always replace ω by any closed 2-form.

Replacement of the holomorphicity

Why do we care about the symplectic flatness?

Remark

The symplectic flatness is an analogue of the holomorphicity.

Complex case:

Theorem (Atiyah, 1957)

Holomorphic connection exists $\Leftrightarrow$ Atiyah class vanishes.

Symplectic case:

Question

Does the symplectic flatness admit an associated characteristic class? For example, an analogue of the Atiyah class?

Primitive characteristic classes

Recall the map

$$\begin{aligned}\mathbb{A} : \Omega^k(M, E) \oplus \Omega^{k-1}(M, E) &\rightarrow \Omega^{k+1}(M, E) \oplus \Omega^k(M, E) \\ (\alpha, \beta) &\mapsto (\nabla\alpha + \omega \wedge \beta, \Phi\alpha - d\beta).\end{aligned}$$

Symplectic flatness $\Leftrightarrow \mathbb{A}^2 = 0$.

This $\mathbb{A}^2$ should be viewed as a “curvature”.

Theorem (Z, 2025)

With or without symplectic flatness, $\text{tr}(e^{-\mathbb{A}^2})$ satisfies transgression.

Primitive characteristic classes

We call $\text{tr}(e^{-\mathbb{A}^2})$ the primitive Chern character.

Corollary

Symplectically flat connection $\Rightarrow$ 1st primitive Chern class vanishes.

It may not be the final counterpart of the Atiyah class.

Yang-Mills theory perspective:

Theorem (Tseng-Zhou, 2025)

Symplectically flat connections on principal bundles $\Leftrightarrow$ conjugacy classes of a certain type of group homomorphisms.

“holomorphic-like” perspective?

Thank the organizers for the invitation.

Thank you for your attention.

Slides are available at:

https://hzhmp.github.io/events_en.html/ncg_dut.pdf