

Note on Utility Functions of Lexicographic Preferences

Wenjun Zheng

Abstract

In this note, we provide the general condition for the existence of lexicographic preferences.

Keywords: lexicographic preferences; utility representation

1 Introduction

Lexicographic preferences derive from reality. A typical example is Chinese College Application. Student should first choose the order of universities and then their desiring majors. We may know that lexicographic preferences don't have utility representation, however this argument de facto suffers fallacies. Consider the following two examples.

Example 1 (Lexicographic preference on $\mathbb{N} \times \mathbb{R}$). Consider lexicographic preference on $\mathbb{N} \times \mathbb{R}$, then it has utility function

$$u(x_1, y_1) = x_1 + \frac{\arctan(y_1)}{\pi}.$$

Proof. Consider pairs (x_1, y_1) and (x_2, y_2) with $x_2 > x_1$. Then,

$$\begin{aligned} u(x_2, y_2) - u(x_1, y_1) &= x_2 + \arctan(y_2) - x_1 - \arctan(y_1) \\ &\geq x_2 - \frac{\pi}{2} - x_1 - \frac{\pi}{2} \\ &= x_2 - x_1 - 1 \geq 0. \end{aligned}$$

□

Example 2 (Lexicographic preference on more fancy setting). Consider lexicographic preference on $([0, 1] \cap \mathbb{Q}) \times [0, 1]$, then for all $(x, y) \in ([0, 1] \cap \mathbb{Q}) \times [0, 1]$ it has utility function

$$u(x, y) = \sum_{k=1}^{l-1} 2^{-k} + \frac{2^{-l}}{2} y, \text{ where } x^l = x.$$

where $[0, 1] \times \mathbb{Q} = \{x^1, x^2, \dots\}$.

Proof. We discuss it into three cases.

- (i) if $(x_1, y_1) = (x_2, y_2)$, then $u(x_1, y_1) = u(x_2, y_2)$;
- (ii) if $x_1 > x_2$: let $x^{l_1} = x_1$ and $x^{l_2} = x_2$.

$$\begin{aligned} u(x_1, y_1) &\geq u(x_1, 0) \geq \sum_{k=1}^{l_2} 2^{-k} \\ u(x_2, y_2) &\leq u(x_2, 1) = \sum_{k=1}^{l_2-1} 2^{-k} + 2^{-l_2-1} \end{aligned}$$

clearly, $u(x_1, y_1) \geq u(x_2, y_2)$.

- (iii) if $x_1 = x_2$ and $y_1 > y_2$, the conclusion is obvious.

□

These two examples potentially demonstrate that for $A \subset \mathbb{R}$ and $B \subset \mathbb{R}$, lexicographic preferences on $A \times B$ might admit a utility representation as long as A is countable. The next theorem pins down our conjecture.

Theorem 1. Consider $X_1, \dots, X_n \subset \mathbb{R}$ and every X_i is infinite. The lexicographic preferences on $X_1 \times \dots \times X_n$ admit utility representation if and only if X_1, \dots, X_{n-1} are countable.

Proof. We first prove the “if” part, then “only if” part.

- (i) “ \leftarrow ”:
- (ii) “ \rightarrow ”: assume \succsim admits a utility representation, we would like to prove by contradiction. Without loss of generality, assume $X_i \in \{X_1, \dots, X_{n-1}\}$ is uncountable. Then, consider

$$(\bar{x}_1, \dots, \bar{x}_i, a, \dots, \bar{x}_n), (\bar{x}_1, \dots, \bar{x}_i, b, \dots, \bar{x}_n), \quad a, b \in X_{i+1} \text{ and } a > b.$$

Hence,

$$u(\bar{x}_1, \dots, \bar{x}_i, a, \dots, \bar{x}_n) > u(\bar{x}_1, \dots, \bar{x}_i, b, \dots, \bar{x}_n).$$

Pick one rational number from interval

$$q(\bar{x}_i) \in [u(\bar{x}_1, \dots, \bar{x}_i, b, \dots, \bar{x}_n), u(\bar{x}_1, \dots, \bar{x}_i, a, \dots, \bar{x}_n)]$$

Further notice that for $\bar{x}'_i > \bar{x}_i$, we have

$$\begin{aligned} q(\bar{x}'_i) &\geq u(\bar{x}_1, \dots, \bar{x}'_i, b, \dots, \bar{x}_n) \\ &> u(\bar{x}_1, \dots, \bar{x}_i, a, \dots, \bar{x}_n) \\ &\geq q(\bar{x}_i). \end{aligned}$$

Consider a mapping from X_i to the set of all $q(x_i)$, where $x_i \in X_i$. It is injection by above argument and surjection by our construction of codomain. Hence, we form a bijection from uncountable set to a countable set¹, contradiction yields.

□

¹Infinite subset of a countable set is still countable.