

# Research Article

Malays. j. med. biol. res.



## Antimicrobial Resistance and Sensitivity among Isolates of *Escherichia coli* from Urine Samples in Denizli, Turkey

Selma Kirac<sup>1\*</sup>, Dilek Keskin<sup>2</sup>, F. Banu Karahasanoğlu<sup>3</sup><sup>1</sup>Denizli Health Services Vocational High School, Pamukkale University, Denizli, TURKEY<sup>2</sup>Cine Vocational High School, Adnan Menderes University, 09500- Cine-Aydin, TURKEY<sup>3</sup>ER-PA Special Health Hospital, Denizli, TURKEY\*Email for Correspondence: [skirac@pau.edu.tr](mailto:skirac@pau.edu.tr)

### ABSTRACT

**Objective:** The research was carried out with isolate and determines the antimicrobial sensitivity in *E. coli* from urinary tract infections in special hospital in Denizli and recorded at specimens.

**Methods:** Urine samples (n=21) were collected from patients with signs and symptoms of Urinary tract infections. Bacteria were isolated and identified by conventional biochemical profile. Antibiotic resistance pattern of *E. coli* against different antibiotic was determined by Kirby-Baur method.

**Results:** The results revealed that sensitivity rate of antimicrobial agents were in the range of meropenem (100%), norfloxacin and ciprofloxacin (86%), cefotaxime (80%), aztreonam (76%). None of the samples showed no resistance to amikacin, ceftazidime, aztreonam, amoxicillin/clavulanic acid, and meropenem. Out of 21 isolates, 3(14%) isolates showed Multiple Antibiotic Resistance ten to thirteen antibiotics.

**Conclusion:** It is concluded that most of the urinary tract infections in human are caused by *E.coli* exhibited highest resistance to meropenem (100%), followed by norfloxacin and ciprofloxacin (86%).

**Key words:** *Escherichia coli*, Antibiotic, Resistance, Prevalence, MDR

1/20/2016

Source of Support: None, No Conflict of Interest: Declared

This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

Attribution-NonCommercial (CC BY-NC) license lets others remix, tweak, and build upon work non-commercially, and although the new works must also acknowledge &amp; be non-commercial.



### INTRODUCTION

Antibiotic resistance is recognized worldwide as a major problem in the management of patients hospitalized with serious infections (Swartz, 1994). Turkish hospitals also face increasing numbers of antibiotic-resistant organisms including *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, Methicillin resistance in *Staphylococcus aureus*, *Escherichia coli* (Toroglu and Keskin, 2011, Toroglu *et al.*, 2013, Yakupoglu *et al.*, 2006).

*E. coli* are gram negative, facultative bacteria that ferment glucose and are members of the family Enterobacteriaceae (Feng and Weagant, 2009). They are mainly allocated in the intestine of animals and forms part of the normal intestinal flora that maintains the physiology of a healthy animal (Conway and Macfarlane, 1995). Thus, most *E.coli* strains are nonpathogenic but pathogenic strains that cause gastrointestinal illness in humans and opportunistic ones that normally affect immune compromised patients exists (Nataro and Kaper, 1998). For example, more than 80% of urinary tract infections occur in outpatients and *E.coli* accounts for more than 50% of the infections in these patients (Blomgran *et al.*, 2004, Jha and Bapat, 2005). In rare cases, virulent strains are also responsible for Haemolytic Uremic Syndrome (HUS), peritonitis, mastitis septicemia, and gram-negative pneumonia Olowe *et al.*, 2003). It is one of the organisms most frequently isolated from different clinical cases of diarrhea and others (Okeke *et al.*, 1999, Tobih *et al.*, 2006).

We aimed in these research to determine the status of antibiotic resistance, underlying conditions, and isolation of *E.coli* isolates with from a special hospital in Denizli, Turkey.

## SUBJECTS AND METHODS

**Isolation of bacterial strains and identification:** 21 isolates were determined from special hospital patients in Denizli July and August in 2013 and recorded at specimens. Mac Conkey agar and EMB agar (Eosin Methylene Blue) agar used for *E.coli* isolation. Isolates were considered to be presumptive *Escherichia* spp. Gram-Negative bacilli, mucoid colonies and lactose positive. Confirmation of isolates was performed by using classic chemical tests (motility test, ure hydrolysis, acid production from mannitol, production of H<sub>2</sub>S, IMVIC (Indol, Metil Red, Voges-Proskauer and Citrate) (Prakash *et al.*, 2011, Cowan and Steel, 1970).

### Antibiotic resistance activity

Antibiotic resistance was determined by an agar disc diffusion test (Bauer *et al.*, 1996) using Mueller-Hinton agar (Difco) according to Clinical and Laboratory Standards Institute (CLSI., 2005) recommendations. Twenty different antibiotics were used. For antibiotic resistance determination, the isolates were grown in Luria- Bertani (LB) broth until the turbidity equal to the 0.5 Mc Farland standart. Cultures were swabbed on to the Mueller-Hinton agar and all isolates were tested against Meropenem (MEM, 10µg/ml), Piperacillin/ tazobactam (TZP, 110µg/ml), Ampicillin/Sulbactam (SAM, 20µg/ml), Amikacin (AK, 30µg/ml), Ceftazidime (CAZ, 30µg/ml), Tobramycin (TOB,10µg/ml), Amoxycillin /clavulanic acid (AMC, 30µg/ml), Gentamycin (CN, 10µg/ml), Aztreonam (ATM., 30µg/ml), Cefepime (FEP PM, 30µg/ml), Cefotaxime (CTX, 30µg/ml), Cefuroxime (CXM, 30µg/ml), Ceftriaxone (CRO, 30µg/ml), Sulphamethazol/Trimetoprim (SXT,25µg/ml), Ciprofloxacin (CIP,5µg/ml), Cefoperozone (CFP, 75µg/ml), Cephazolin (CZ, 30µg/ml), Norfloxacin (NOR, 10µg/ml), Ampicillin (AM, 10µg/ml), Cefixime (CFM, 5µg/ml).

The isolates those grown in inoculation were evaluated as resistant, and the others were evaluated as susceptible (Bauer *et al.*, 1996). The antibiotic discs were dispensed sufficiently separated from each other so as to avoid overlapping of inhibition zones. The plates were incubated at 37°C, and the diameters of the inhibition zones were measured after 18 h. All susceptibility tests were carried out in duplicate and were repeated twice if discordant results had been obtained.

### Multiple Antibiotic Resistance Index:

For all isolates, we calculated the MAR index values (a/b, where a represents the number of antibiotics the isolate was resistant to, b represents the total number of antibiotics the isolate tested against). A MAR index value  $\geq 0.2$  is observed when isolates are exposed to high-risk sources of human or animal contamination, where antibiotics use is common; in contrast a MAR index value  $<0.2$  observed when antibiotics are seldom or never used (Krumperman, 1985, Matyar *et al.*, 2008).

**Results** The sensitivity of *E.coli* isolates to antimicrobial agents (n=21) gave high sensitive rates found that *E.coli* isolates diffusion tests for meropenem (100%), norfloxacin and ciprofloxacin (86%), cefotaxime (80%), aztreonam (76%). None of the samples showed no resistance to amikacin, ceftazidime, aztreonam, amoxicillin/clavulanic acid, and meropenem. The results were given table 1.

## DISCUSSION

Carbapenems, mainly meropenem, sensitivity rate of meropenem was showed in 100 %. Some researchers have reported meropenem sensitivity rate to *E.coli* from 100% to 90.9% (İnan and Gurler, 2004, Yılmaz *et al.*, 2010, Barisic *et al.*, 2003). Our results were similar to Yılmaz *et al.*,(2010) who also reported that percentage of meropenem sensitivity was 100% in Turkey. It can be suggested that meropenem can be used for infections based on *E.coli*.

Table 1: Antibiotic resistance pattern of *Escherichia coli* isolated from urine samples

Antibiotics	Sensitive	Intermediate	Resistance
MEM	21(%100)	(%0)	(%0)
NOR	18(%86)	(%0)	3(%14)
CIP	18(%86)	(%0)	3(%14)
CTX	17(%80)	3(%14)	1(%5)
ATM	16(%76)	5(%24)	(%0)
CAZ	15(%71)	5(%24)	(%0)
FEP PM	15(%71)	5(%24)	1 (%5)
CFM	13(%62)	5(%24)	3(%14)
CRO	12(%57)	6(%29)	3(%14)
TZP	12(%57)	9(%42)	(%0)
SXT	11(%57)	4( %19)	6(%29)
CFP	10(%48)	8(%4)	3(%14)
CXM	9(%42)	8(%38)	4(%19)
AMC	8(%38)	13(%62)	(%0)
AK	7(%33)	14(%67)	(%0)
CN	6(%29)	12(%57)	5(%21)
TOB	4(%19)	16(%76)	1(%5)
CZ	4(%19)	12(%57)	5(%21)
SAM	3(%14)	17(%80)	1(%5)
AM	1(%5)	10(%21)	10(%48)

Meropenem =MEM, Piperacillin/ tazobactam =TZP, Ampicillin/Sulbactam= SAM, Amikacin =AK, Ceftazidime =CAZ, Tobramycin= TOB, Amoxycillin /clavulanic acid =AMC, Gentamycin =CN, Aztreonam =ATM, Cefepime= FEP PM, Cefotaxime =CTX, Cefuroxime=CXM, Ceftriaxone= CRO, Sulphamethazol/Trimethoprim =SXT, Ciprofloxacin=CIP, Cefoperazone=CFP, Cephazolin= CZ, Norfloxacin=NOR, Ampicillin=AM, Cefixime= CFM

When it comes to norfloxacin, sensitivity rate, it was %86. Many researchers tested sensitivity of norfloxacin to *E.coli* (Karki *et al.*, 2001, Ay *et al.*, 2003).

Sensitivity rate of ciprofloxacin was showed in 86%. Many researchers have tested the sensitivity of ciprofloxacin to *E.coli* (Turnidge *et al.*, 2002, Aiyegoro *et al.*, 2007, Mansouri *et al.*, 2002). Our results were similar to Mansouri *et al.*, (2002) who also reported that percentage of ciprofloxacin sensitivity was 84.62% in Iran. It can be suggested that ciprofloxacin can be used for infections based on *E.coli*.

As for the sensitivity rate of cefotaxime, it was 80%. Many researchers have tested resistance of cefotaxime to *E.coli* (Jones *et al.*, 2004, Gönüllü *et al.*, 2008. Our results were similar to Gonullu *et al* (2008) who also reported that sensitivity rate of cefotaxime was 84%.

Sensitivity rate of aztreonam was showed in 76 %. Many researchers have tested the resistance of aztreonam to *E.coli* (Cho *et al.*, 2011, Gonlugur *et al.*, 2004, Iqbal *et al.*, 2002) Our results were similar to Iqbal *et al* (2002) who also reported that aztreonam sensitivity rate was 74% .

When it comes to ceftazidime sensitivity rate, it was 71%. Some researchers have reported that ceftazidime sensitivity rate from 99.6% to 4.5% (Rhonberg and Jones, 2007, Ozsahin *et al.*, 2005). Our results were in compliance with previous researchers

As for the sensitivity rate of cefepime, it was 71%. Some researchers have reported sensitivity rate cefepime to *E.coli* in clinical samples (Iqbal *et al* 2002, Ashgar, 2006, Albayrak and Kaya, 2009, Kumarasinghe, 2001). Our results were similar to Iqbal *et al* (2002) who also reported that cefepime sensitivity rate was 80% .

As for sensitivity rate of cefixime was 62%. Cefixime has a broad antibacterial spectrum and it is active against a wide variety at gram-negative organisms, including *E.coli*. Our results were similar to Iqbal *et al.* (2002) who also reported that cefixime sensitivity rate was 61% .

As for sensitivity rate of ceftriaxone, it was 57%. Many researchers have tested the resistance of ceftriaxone to *E.coli*. According to previous studies resistance of *E.coli* was from 0% to 100% (Koken *et al.*, 2008, Matute *et al.*, 2004, Ateş, 2007, Yuluğkiral, 2007, Kalantar *et al.*, 2008, Uzun *et al.*, 2006).

As for the sensitivity rate of piperacillin/ tazobactam, it was 57%. Tazobactam seems to be the most promising beta-lactamase inhibitor, which has unlike clavulanic acid and sulbactam, its own antibiotic activity (Blahova *et al.*, 1995).

Sensitivity rate of sulfamethoxazole/trimethoprim was showed in 52 %. Some researchers have reported sulfamethoxazole/trimethoprim sensitivity rate to *E.coli* in clinical samples (Gonlugur *et al.*, 2004, Eryılmaz *et al.*, 2010, Rawat *et al.*, 2010) Our results were similar to Gonlugur *et al.* (2004)<sup>33</sup> who also reported that sulfamethoxazole /trimethoprim sensitivity rate was 55%.

As for the sensitivity rate of cefoperazone, it was 48 %. Some researchers have reported sensitivity rate cefoperazone to *E.coli* in clinical samples (Ozsahin *et al.*, 2005, Dean *et al.*, 2008)

Sensitivity rate of cefuroxime was showed in 42 %. Some researchers have reported sensitivity rate cefuroxime to *E.coli* in clinical samples (İnan and Gurler 2004, Duman *et al.*, 2010). In our study resistance of cefuroxime was detected 19%. Our results were similar to Inan and Gurler (2004) who also reported that resistance of cefuroxime showed 14 % in *E.coli* strains isolated from children with urinary tract infections.

As for the sensitivity rate of amoxicillin-clavunat, it was %38. Some researchers have reported amoxicillin-clavunat sensitivity rate to *E.coli* ( Uzun *et al.*, 2006, Kader and Kumar, 2004).

As for the sensitivity rate of amikacin, it was 33%. Some researchers have reported amikacin sensitivity rate to *E.coli* (Karki *et al.*, 2001, Ekim *et al.*, 1998, Çetin *et al.*, 2006, Giray *et al.*, 2012). In our study resistance of amikacin was detected 0%. Our results were similar to Giray *et al.*, (2012) who also reported that resistance of amikacin showed 0 % in *E.coli* strains obtained from urology outpatient clinic of Ege Medical Faculty in Izmir.

Among the aminoglycosides group, gentamycin sensitivity rate was 29%. Some researchers have reported gentamycin sensitivity rate to *E.coli* from 1.5% to 54% (Koksaldi-Motor *et al.*, 2010, Kutlu, 2007, Pieboji *et al.*, 2004, Kalem *et al.*, 2008). Our results were compliance with previous researchers. Koksaldi-Motor *et al.*, (2010) reported that when it compared to previous year's data susceptibility of *E.coli* isolated from urine to gentamycin were decreased and also different resistance rate occurred different locations.

As for the sensitivity rate of tobramycin, it was 19%. Some researchers have reported tobramycin sensitivity rate to *E.coli* (Gonlugur *et al.*, 2004, Sucu *et al.*, 2004). Our results were similar to Gonlugur *et al.* (2004) who also reported that tobramycin sensitivity rate was 16.6%.

As for the sensitivity rate of cefazolin, it was 19%. Some researchers have reported that cefazolin sensitivity (Vlieghe, 2009, Arıkan *et al.*, 1995, Frederick, 2011). Our results were in compliance with previous researchers.

As for sensitivity rate of ampicillin/sulbactam was 14%. Many researchers have tested sensitivity ampicillin/sulbactam to *E.coli* (Toroglu *et al.*, 2013, Khan and Zaman, 2006).

As for sensitivity rate of ampicillin was 5%. Many researchers have tested the resistance of ampicillin to *E.coli* (Leblebicioğlu *et al.*, 1994, Rawat *et al.*, 2010, Ahmed *et al.*, 2000). Our results were similar to Rawat *et al.* (2004) who also reported that aztreonam sensitivity rate was 5.71% from Kumaun region.

In the present study, the lowest MAR index was 0 obtained from urine samples from female and male. In contrast to the highest MAR index was 0.65 obtained from a female. Out of 21 isolates, 3 (14%) isolates showed Multiple Antibiotic Resistance ten to thirteen antibiotics. The MAR index were determined 0.25 and above (Table 2). Some researchers have reported Multi-Drug Resistance rate to *E.coli* from 2% to 97% (Al-Mardeni *et al.*, 2009, Mathai *et al.*, 2008, Al-Tawfiq, 2006). Our results were in compliance with previous researchers. Study shows that multiple resistance is a common hospital pathogen.

Table 2: Multiple antibiotic resistance (MAR) index among 21 *Escherichia coli* urine samples

Source of isolates	Total isolates	MAR index
Male	1	0.5
Male	2	0.15
Male	1	0.45
Female	2	0.05
Female	1	0.65
Female	1	0.05
Female	3	0.1
Female	9	0
Male	1	0
Total	21	

In conclusion, it is suggested that meropenem, norfloxacin and ciprofloxacin, cefotaxime, aztreonam could be better for the treatment of infections based on *E.coli* according to the present study. Ampicillin/sulbactam and ampicillin were not-advisable antibiotics for *E.coli* infections according to results of the present study.

## REFERENCES

- Ashgar, AH. 2006. Frequency and antimicrobial susceptibility patterns of bacterial pathogens isolated from septicemic patients in Makkah hospitals. Saudi Med J, 27(3):361-367.
- Ahmed AA, Osman H, Mansour AM, Musa HA, Ahmed AB, Karrar Z and Hassan HS.. Antimicrobial agent resistance in bacterial isolates from patients with diarrhea and urinary tract infection in Sudan. The American Soc Trop Med Hygiene, 63(5, 6), pp. 259-263, 2000.
- Aiyegoro, O.A., O.O. Igbinosa, I.N. Ogunmwoyoni, E.E. Odjadre, O.E. Igbinosa and A.I. Okoh, 2007. Incidence of urinary tract infections (UTI) among children and adolescents in Ile-Ife, Nigeria. African J. Microbiol. Res., 013-019
- Albayrak, N. and S. Kaya.. Extended spectrum beta lactamases production and antimicrobial resistance ratio of the *Escherichia coli* and *Klebsiella pneumoniae* strains isolated from various clinical specimens. Türk Microbiol Cem. Derg., 39: 16-21, 2009.
- Al-Mardeni RI, Batarseh A, Omaish L, Shraideh M, Batarseh B, Unis N. Empirical treatment for pediatric urinary tract infection and resistance patterns of uropathogens, in Queen Alias Hospital and Prince A'isha Military Center. Jordan. Saudi J Kidney Dis Transplant; 20: 135-9, 2009.
- Al-Tawfiq JA. Increasing Antibiotic Resistance Among Isolates of *Escherichia coli* Recovered From Inpatients and Outpatients in a Saudi Arabian Hospital. Infect Control Hosp Epidemiol; 27:748-753, 2006.
- Arıkan S, Gür D, Hayran M, ve ark Hastane dışı enfeksiyonlara yol açan gram negatif bakterilere karşı cefamet'in in vitro etkinliği. Mikrobiol Bül 1995; 29:14-19.
- Ateş F. The analyse of urine culture results in patients with lower urinary tract infection, Türk Üroloji Derg; 33(2):223-7, 2007
- Ay S, İşeri LA, Duman B. Antibiotic Susceptibilities of Gram Negative Microorganisms Isolated from Urine Samples. İnönü Üniv Tıp Fak Derg., 10(2) 59-62, 2003
- Barišić Z, Babić-Erceg A, Borzić E, Zoranić V, Klaiterna V, Carev M. Urinary tract infections in South Croatia: aetiology and antimicrobial resistance. Int J Antimicrob Agents, 22, 61-64, 2003.
- Bauer, A.W, Kirby, WM.M, Sherris, JC., Turck, M. Antibiotic susceptibility testing by a standart.. CLSI., Performance standards for antimicrobial susceptibility testing; 15th informational supplement. 1966.



- Blahova J, Hupkova M, Kremery V Sr. The effectiveness of so called potentiated penicillins (augmentin and tazobactam) in vitro. *Cas Lek cesk* 1995; 134: 558-61
- Blomgran, R., Zheng L, Stendhal O. 2004. Uropathogenic *Escherichia coli* trigger oxygen-dependent apoptosis in human neutrophils through the cooperative effect of type 1 fimbriae and lipopolysaccharide. *Infect. Immun.*, 72:4570-4578.
- Cetin ES, Demirci M, Kaya S, Aridogan BC, Adiloglu AK, Goksu Y, Gonulates N. Antibiotic susceptibilities of *Escherichia coli*, *Klebsiella pneumoniae* and *Enterobacter* strains isolated from blood specimens. *Turk Mikrobiyol Cem Derg* 36 (1) : 20 - 24, 2006
- Cho SH, Lim YS, Park MS, Kim SH a, Yeon-Ho Kang YH. 2011. Prevalence of Antibiotic Resistance in *Escherichia coli* Fecal Isolates From Healthy Persons and Patients With Diarrhea. *Public Health Res Perspect* 2(1), 41-45.
- CLSI., 2005. Performance standards for antimicrobial susceptibility testing; 15th informational supplement. CLSI / NCCLS M 100-S 15. Clinical and Laboratory Standards Institute, Wayne, PA().
- Conway P and G. Macfarlane, 1995. Microbial ecology of the Human large intestine. In: human Colonic Bacteria: Role in nutrition, Physiology and pathology, Gibson G and G. Macfarlane (Eds.) CRC Press, London
- Cowan SF, Steel KJ (1970). Manual for the Identification of the Medical Bacteria Cambridge: Cambridge University Press. pp. 7-122
- Dean DC., Krahe D., Wareham D. 2008. Antimicrobial resistance in community and nosocomial *Escherichia coli* urinary tract isolates, London 2005-2006. *Annals of Clinical Microbiology and Antimicrobials* 7:13,1-4.
- Duman Y., Güllüer N., Serindag A., Tekerekoglu MS. Antimicrobial susceptibility of *E.coli* strains and presence of extended-spectrum beta lactamase. *Firat Tıp Derg.*, 15(4) 197-200, 2010.
- Ekim M., Kuluglu Z., Aysev D., Can S. 1998. Changes in antibiotic sensitivity in urinary tract infection caused by *Escherichia coli* *Journal of the Turkish Nephrology*, 3 141-144.
- Eryilmaz M., Eylül -Bozkurt M., Yildiz MM., Akin A. Antimicrobial resistance of urinary *Escherichia coli* isolates. *Trop J of Pharm Res*, 9(2) 205-209, 2010.
- Feng P and S.D. Weagant., 2009. Bacteriological analytical manual 8th. Edn., Chapter 4 <http://www.fda.gov/Food/scienceResearch/laboratoryMethods/BacteriologicalAnalyticalManualBAM/UCM064948>.
- Frederick A., 2011. *Escherichia coli*, its prevalence and antibiotic resistance in Malaysia: A mini review. *Microbiology Journal* 1(2) 47-53.
- Giray B., Ucar FB., Aydemir SS. 2012. Characterization of uropathogenic *Escherichia coli* strains obtained from urology outpatient clinic of Ege Medical Faculty in Izmir. *Turk J Med Sci.* 42(1) 1328-1337.
- Gonlugur, U., M.Z. Bakici, I. Akkurt and T. Efeoglu, 2004. Antibiotic susceptibility patterns among respiratory isolates of Gram negative bacilli in a Turkey university hospital. *BMC Microbiol.*, 4: 1-5
- Gönüllü N., Canberk MB., Filiz O., Altinkum S., Kucükbasmacı O., Aygun G., Altas K.. Antimicrobial susceptibilities and beta lactam resistance phenotypes of *Escherichia coli* strains isolated from various clinical samples. *Ankem Derg* 22(2) 64-68, 2008.
- Iqbal M, Patel IK, Shah SH, Ain Q, Barney N, Kiani Q, Rabbani KZ, Zaidi G, Mehdi B. Susceptibility patterns of *Escherichia coli*: prevalence of multidrug-resistant isolates and extended spectrum beta-lactamase phenotype *J Pak Med Assoc.* 2002 Sep;52(9):407-11.
- İnan N U, Gurler N. Investigation of Antibiotic Resistance and Some Virulence Factors of *Escherichia coli* Strains Isolated from Children with Urinary Tract Infections. *Ankem Derg*; 18(2):89-96, 2004.
- Jha, N and Bapat SK. 2005. A study of sensitivity and resistance of pathogenic microorganisms causing UTI in Kathmandau valley. *Kathmandu Univ. Med J.*, 3:123-129.
- Jones ME, DC, Draghi, C, Thornsberry, JA, Karlowsky, DF, Sahm, RP, Wenzel. 2004. Emerging resistance among bacterial pathogens in the intensive care unit a European and North American Surveillance study. *Ann Clin Microbiol*, 3(14) 1-11.
- Kader, A.A. and A. Kumar. Prevalence and antimicrobial susceptibility of extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* in a general hospital in Saudi Arabia. *Saudi Med J.*, 25: 570-574, 2004.
- Kalantar E, Motlagh ME, Lornejad H, Reshadmanesh N. Prevalence of urinary tract pathogens and antimicrobial susceptibility patterns in children at hospitals in Iran. *Iranian Journal of Clinical Infectious Diseases*, 2008;3(3):149-153.
- Kalem F., Gunden NS., Arslan U., Tuncer I. Antimicrobial susceptibility of *Escherichia coli* strains isolated in urine samples. *ANKEM Derg* 22: 193-197, 2008.
- Karki T., Truuaslu K., Vainumae, Mikellsaar M. Antibiotic Susceptibility Patterns of Community- and Hospital-acquired *Staphylococcus aureus* and *Escherichia coli* in Estonia. *Scand J Infect Dis* 33: 333-338, 2001.
- Khan AU and Zaman MS. Multiple drug resistance pattern in Urinary Tract Infection patients in Aligarh. *Biomedical Research* 2006; 17 (3): 179-181.
- Koken G, Asik G, Ciftci H, Cetinkaya Z, Aktepe O.C, Yilmazer M. Efficiency of Fosfomycin Trometamol on *Escherichia coli* Strains from Community Acquired Urinary Tract Infections. *Ankem Derg*, 22(1):23-27, 2008.
- Koksaldi-Motor V., Tutanc M., Arica V., Arica S., Ay B. Susceptibility of Uropathogen *Escherichia coli* strains to commonly used antibacterial agents in urinary tract infections. *Ankem Derg*, 2010, 24(4) 198-201, 2010.
- Krumperman, PH. 1985. Multiple antibiotic resistance indexing of *Escherichia coli* to identify high-risk sources of fecal contamination of foods. *App. Environ. Microbiol.*, 46:165-170.
- Kumarasinghe G, Chow C, Tambyah PA. Widespread resistance to new antimicrobials in a university hospital before clinical use. *Int J Antimicrob Agents*, Oct;18(4):391-393. 2001.
- Kutlu S, Kutlu M. Antibiotic susceptibility of *Escherichia coli* strains isolated in urinary tract infections in Didim, Turkey. *Turk J Infect*; 21(2): 81-83, 2007.
- Leblebicioglu H., Gunaydin H., Sanic A., Buyukalpelli R. 1994. Comparative in-vitro susceptibility of betalactam and betalactam/betalactamase inhibitor antibiotics against gram negative rods isolated from urine cultures. *Mikrobiyol Bült.*, 28 218-222.
- Lim KT., Yasin R., Yeo CC., Puthucheary S., Thong K.L. Characterization of multidrug-resistant ESBL producing *Escherichia coli* isolates from hospitals in Malaysia. *J. Biomed Biotechnol.*, 1-10, 2009

- Mansouri S, Shareif S, Ahmad K (2002). Antimicrobial resistance pattern of *E. coli* causing urinary tract infections and that of human faecal flora in southeast of Iran. *Iran Med. Journal* 8: 123-128.
- Mathai E, Chandy S, Thomas K, Antoniswamy B, Joseph I, Mathai M, Sorensen TL, Holloway K.. Antimicrobial resistance surveillance among commensal *Escherichia coli* in rural and urban areas in Southern India. *Trop Med Int Health*. Jan;13(1):41-5, 2008.
- Matute AJ, Hak E, Schurink CA, McArthur A, Alonso E, Paniagua M, Van Asbeck E, Roskott AM, Froeling F, Rozenberg-Arska M, Hoepelman IM. Resistance of uropathogens in symptomatic urinary tract infections in León, Nicaragua. *Int J Antimicrob Agents*.,23(5):506-9,2004.
- Matyar, F., Kaya, A and Dinçer, S. 2008. Antibacterial agents and heavy metal resistance in Gram-negative bacteria isolated from seawater, shrimp and sediment in Iskenderun Bay, Turkey. *Sci. Total Environ.*, 407:279-285.
- Nataro J.P and J.B Kaper,1998. Diarrheagenic *Escherichia coli* .In: *Foodborne Disease Handbook*, Hui,Y.H., JR. Gorham, K.D.Murrell and D.O Cliver ( Eds). Marcel Decker, Inc., New York, pp:169-213.
- Okeke IN, Lamikanra A, Edelman R. 1999. Socioeconomic and behavioral factors leading to acquired bacterial resistance to antibiotics in developing countries. *Emerg.Infect. Dis*. 5:18-27.
- Olowe OA, Olayemi AB, Eniola KIT and Adeyeba AO.2003.Aetiological agents of diarrhoea in children under 5 years of age in Osogbo. *Afr.J. Clin and Exp. Microbiol* 4(3):62-66.
- Ozsahin D., Digrak M., Kiran OE.2005. The Investigation on resistance Properties of *Escherichia coli* against Beta Lactam Groups Antibiotics. *KSU, Journal Of science and Engineering* 8(2)8-12.
- Piéboji JG, Koulla-Shiro S, Ngassam P, Adiogo D, Njine T, Ndumbe P.Antimicrobial resistance of Gram-negative bacilli isolates from inpatients and outpatients at Yaounde Central Hospital, Cameroon. *Int J Infect Dis*, 8(3):147-54, 2004.
- Prakash S, Dayalan JA and Edison N 2011. Prevalence of Bacteriuria In Jeyaseharan Hospital of South India and their antibiogram. *Asian Pacific Journal of Tropical Biomedicine*, 105-108.Cambridge University Press. pp. 7-122 .
- Rawat DV, Umesh D, Paul MP. 2010. Antibiotic Resistance Pattern of Urinary Tract Isolates of *Escherichia coli* from Kumaun Region. *National Journal of Integrated Research in Medicine* 1(4): 43-46.
- Rhonberg PR and Jones RN.2007. Contemporary activity of meropenem and comparator broad-spectrum agents: MYSTIC program report from the United States component (2005). *Diagn Microbiol Infect Dis*. 2007 Feb;57(2):207-15
- Sucu N, Boz GA, Bayraktar O, Caylan R, Aydın K, Koksall İ. Th e change of antibiotic susceptibilities of uropathogen *Escherichia coli* strains in years. *Klinik Dergisi* 2004; 17: 128-131.
- Swartz MN,1994.Hospital-acquired diseases with increasingly limited therapies. *Proc Natl Acad Sci USA*, 91:2420-2427.
- Tobih JE, Taiwo SS, Olowe OA, Olaosun OA, Adejumo SO.2006. Microbiological profiles of discharging ears in Osogbo, Nigeria. *Trop. Doc*. 36(3):165-166.
- Toroglu S and Keskin D. Antimicrobial Resistance and Sensitivity among Isolates of *Klebsiella pneumoniae* from Hospital Patients in Turkey. *Int J Agri Biol*, 13(6) 941-946,2011
- Toroglu, H. Avan, D. Keskin. Beta- Lactamases production and Antimicrobial resistance ratio of *Pseudomonas aeruginosa* from hospitalized patients in Kahramanmaraş, Turkey, *Journal of Environmental Biology*, 34 (4), 695-700, 2013.
- Turnidge J., Bell J., Biedenbach J and Jones R.N, 2002Pathogen occurrence and antimicrobial resistance trends among urinary tract infection isolates in the Asia-Western Pacific Region: report from the SENTRY Antimicrobial Surveillance Program, 1998–1999. *International Journal of Antimicrobial Agents* 2002, 20, 10-17.
- Uzun K., Teke T., Yavuz Z. ., G..Surveillance of antimicrobial resistance and susceptibility in bacterai isolated from pulmonart critical care. *Tip Araşt Derg.*, 4(3) 8-13, 2006.
- Vlieghe E., Phoba MF., Muyembe Tamfun JJ., Jacobs J.. Antibiotic resistance among bacterial pathogens in Central africa: a review of the published literatue between 1995 and 2008. *Int J Antimicrob Agents*, 34, 295-303, 2009.
- Yakupogulari, Y., A, Gunduz., M, Ozcan., M, Dogukan., A, Seyrek., M, Yilmaz. 2006. Susceptibility of *Staphylococcus aureus* Strains to Ciprofloxacin, Ofloxacin, Levofloxacin and Moxifloxacin. *Fırat Tıp Derg*; 11(1): 45-47.
- Yılmaz N., Köse Ş., Ağuş N., Ece G., Akkoçlu G., Kıraklı C. Microorganisms Isolated from Blood Cultures of Intensive Care Unit Patients, their Antimicrobial Susceptibility and Etiological Agents in Nosocomial Bacteremia. *Ankem Derg* 24(1):12-19,2010.
- Yuluğkural Z, Mutlu B. Susceptibility of *Escherichia coli* strains isolated from urine cultures to some commonly used antibacterial agents *Medical Journal of Trakya University*; 24: 6-11, 2007

#### Authors Contributions

All the authors have contributed significantly in study design, experimentation, data analysis and manuscript drafting.