

Vol 1, No. 1, 2014

Fish Culture Practice and Socio Economic Status in Northern Middle Part of Bangladesh

Samir Kumer Sarker¹, Md. Tawheed Hasan², Simon Kumar Das³, & Sabuj Kanti Mazumder⁴

ABSTRACT

The present study was carried out in Bhaluka upazilla, the northern middle part of Bangladesh to observe the culture technique of fish culture using of supplementary feeds and socio-economic condition of the fish farm owners. Data were collected from 39 fish farms in Bhaluka upazilla during the period from September 2010 to August 2011. Most of the studied farm owners carried out polyculture system and culture, mostly carps (Indian major carps and exotic carps), pangus and tilapia using supplementary feeds. The feeding rate varied from 1-7% of body weight of fish with frequency 1-3 times/day. The highest production was recorded as 56,810 kg/ha/yr using company feed and the lowest production was recorded as 14,210 kg/ha/yr with an average production of 34,350 kg/ha/yr. The average production of 23,520 kg/ha/yr was recorded (farm made feed). Most of the farmers did not use fertilizer and manure in their ponds, but in winter season sometimes they used poultry litter to avoid high priced feed. The overall management practice was not smoothly developed in the monoculture and polyculture farms. It may be concluded that we have to reduce all of the constraints of management practices of aquaculture to achieve sustainable growth in future aquaculture of Bangladesh.

Key words: Beel, Fish farm, Owner, Stocking density and fry

crossref DOI: 10.15590/mjmbr/2014/v1i1/53719 License: CC-BY-NC

INTRODUCTION

Bangladesh is an agro-based riverine country enriched with vast fisheries resources. It lies in the north-eastern part of the South Asia, between 20°34' and 26°38' north latitude and 88°01' and 92°41' east longitude. The country is surrounded by India on the west, the north and the northwest and Myanmar on the southeast and the Bay of Bengal on the south. Its total area is covering 147,570 sq. km. Fish and Fisheries play a significant role in the economy of Bangladesh in terms of nutrition, income, employment and earning foreign exchange. Bangladesh is a highly populated country. The total population of this country is about 142,319 thousand (BBS, 2011). So the nutrition requirement is very high in our country. Fish is the principal source of animal protein for the people of Bangladesh.

¹Quality Assurance Officer, BEST Project, Upazilla Fisheries Office, Satkhira, **Bangladesh**

²Lecturer, Departemt of Aquaculture, Sylhet Agricultural University, **Bangladesh**

³Senior Lecturer, School of Environmental and Natural Resource Sciences, UKM, Malaysia

⁴Assistant Professor, Dept. of Aquatic Resource Management, Sylhet Agricultural University, **Bangladesh**

Fish provides about 60% animal protein for the people of Bangladesh. Fisheries sector is contributing about 3.74% to GDP and 3.00% to nation's foreign exchange (DoF, 2010). There are about 12.5 million people's livelihood depends directly and indirectly on the fisheries sector.

Bangladesh is very rich in inland capture fish production. But now-a-days capture fish production is decreasing day by day due to different causes. Such as agricultural pollution, industrial pollution, natural calamity etc. Fish of the beels along with other aquatic organisms are silent victims of chlorine sub-lethal toxicity resulting from different types of pollutants matters (Bernet *et.al.*, 1999). Fish production is also decreasing due to natural causes like flood, drought etc. (Chakraborty 2009). So due to high demand of fish its needs to increase culture based fishery in the country. There are about 265 indigenous species and 12 exotic species of fish, 24 species of prawn and 36 species of shrimp (DoF, 2010). Above these species some are used as culture species. The commercially important culture species are Rui, Catla, Mrigal, Bighead carp, Mirror carp, Minor carp, Silver carp, Sharpunti, Tilapia, Pangas, Thai koi, Shing, Magur etc.

In Bangladesh fish culture is very popular mostly in Bhaluka upazila, the northern middle part of Bangladesh. About 80% people of Bhaluka upazila engage with agricultural activities. Due to the environmental suitability and the economic profit fish culture is increasing very rapidly. The soil, water and climate are very much suitable for aquaculture in Bhaluka upazila. The temperature ranges from 7 to 36°C (BBS, 2011). That is very essential for the fish culture. But the management practice is not maintained properly. They practiced fish culture with very high stocking density and used different types of supplementary feeds to get the more production. Most of the farmers of this area thought about the production, but never think about the management practice. In the Bhaluka upazilla there are many large water bodies known as beel. The management technique of these water bodies is very simple. Different farmers used poultry manure in their water bodies, but they have no idea about the effects of poultry manure in water bodies. Taking all the issues in concern the present study is conducted with a view to serve the following objectives- 1. To know the socioeconomic status of the fish farm owners; 2. To observe the management technique of fish culture; 3. To know the water quality parameters of the existing culture system and 4. To find out a suitable methods of feeding rate and use of quality feed for more production of fish.

METHODOLOGY

Profile of the study area

The study was conducted in Bhaluka Upazilla lies between 24°38′N and 91°33′E situated in Mymensing District in Dhaka Division Bangladesh (Figure 1). The total area of Bhaluka Upazilla is 444.05 sq km. and water area is 9026ha. Most of the area was related to fish culture. The topography is irregular. The soil type of these water bodies is loams to clay loams. The color of the soil is grey to brownish. The northern side of the upazilla is bounded by Fulbaria and Trisal upazilla and eastern side is Gaffargoan upazilla under the Mymensingh district. The southern side attached with Sreepur Upazilla under Gazipur district and the western side is covered by Sakhipur and Ghatail upazilla under the Tangail district. Main rivers of the upazilla are Khiru and Sutia (Fig. 1).




Figure 1: map of Bhaluka Upazilla

Questionnaire preparation and data collection

On the basis of objectives the questionnaire was made to consist of all necessary questions that were essential for detail information about the farm. The questionnaire has the two parts, one is personal information and another is farm information. Then the schedule was pre-tested to verify the relevance of the questions and the nature of the sample producers. After pre-testing and necessary adjustment, a final survey schedule was developed. For the interest of study 3 categories (small, medium and large farms) of farm was selected in the different area of Bhaluka upazilla.

The data were collected for the period of 12 months between September 2010 to August 2011 in Bhaluka upazilla, Bangladesh. The selected number of fish farmer was 39. A random sampling technique was followed to select the location of the different farm. The direct monitoring was conducted frequently for the accurate information about the fish culture. The essential information was noted during interview. The questions were asked to the farmers very systematically. The explanations of the farmers are recorded in the questionnaire. For the each respondent a set of questionnaire was prepared.

Processing and analysis of data

The collected data were summarized, tabulated and analyzed according to the objectives of the study. The data were analyzed through one way ANOVA using SPSS and Excel followed to find out whether any significant difference was existed among different studies (Duncan, 1955; Zar, 1984).

RESULTS

Socio-economic characteristics of the fish farm owners

Out of 39 farm owners the highest level of the total farm owners were belonged to the age group range of 20-30 years (35%) and the lowest level farm owners were belongs to age group range 51-60 years (15%) and other 30% and 20% were age group ranges between 31-40 and 41-50 years respectively. In the case of educational status, 12% of the total man's power are illiterate but had the ability of signature. About 70% of the man's power had primary and high school education. About 14% manpower of the farm had SSC to HSC level and other 4% had a Graduate level education. The occupation of the fish farm owners is divided into four groups. The highest percentage was found in group fish farm and agriculture (35.90%) whereas the lowest was in group fish farm and job (Table 1).

Table I: Occupation status of the fish farm owners of the studied area.

Type of occupation	Monoculture (N=08)	Polyculture (N=31)	, ,		Cumulative (%)
Only Fish farm business	2	10	12	30.76	_
Fish farm + Job	3	6	9	23.08	53.84
Fish farm+ Agriculture	2	12	14	35.90	89.74
Fish farm + Others	1	3	4	10. <mark>2</mark> 6	100

About 8% Owners mentioned that the transportation facilities of farm was not good due to narrow road, 62% mentioned as good, while 30% owners mentioned the facility as excellent. Farmers were trained from the different organization. Out of 39 fish farm owners 20.51% had no training while 30.77% received training from BFRI, Department of Fisheries (DoF) and NGO's. About 48.72% owners gained their knowledge on fish farm operation through personal contact. The farm owners receive technical assistance from DoF and other organizations. Local people were benefited from fish farm through various ways. It was found that the benefits were employment opportunity, poverty alleviation, increased consumption of fish and socio-economic status (Table 2).

Table II: Percentage distribution fish farms given benefit to people

Benefit	Number of cases (N=39)	Percentage
Employment opportunity	14	35.89
Poverty alleviation	11	28.22
Increased consumption of fish	08	20.51
Obtain their money through leasing	06	15.38

Feature of the water body

Both monoculture and polyculture were practiced in Bhaluka upazilla but polyculture practice is very advanced. During the study period it seems that about 79.5% was found in polyculture and 20.5% was in monoculture. The area of fish farms of the study area was classified into three categories that are Small (<1.00), Medium (1.1-3.00) and Large (3.1>) ha. Out of 39 farms 38.5% were in small categories, 33.3% were in medium categories and rest of the farms 28.2% were in large categories (Table 3).

Table III: Farm size and their frequencies in the study area.

Area	Number of diffe	erent fish farms	Total fish farms	Percentage	Cummulative
(ha)	Monoculture (N=08)	Polyculture (N=31)	(N=39)	(%)	(%)
Small (>1.00)	03	12	15	38.5	00
Medium (1.1-3.00)	03	10	13	33.3	71.8
Large (3.1<)	02	9	11	28.2	100

In the studied area, 30% of farm owners having single ownership, 15% having multiple ownership and 55% having leased water body (Figure 2).




Figure 2: Showing ownership of the water body in the study area.

In the study area about 80% of the water bodies were irregular shaped due to tectonic origin and 20% water bodies were regular shaped. The minimum size of the pond is 0.16 ha and the maximum size of the pond is 37.24 ha where as average pond size was 5.78 ha. The depth of the water body varies from summer to winter. The average depth of the water body in winter season was 1.61±0.15 meter and in summer season was 2.32±0.15 meter.

The farmers were commonly used shallow and deep tube wells. About 74.36% farm owners had Shallow tube well, 20.52% farm owners had Deep tube well and only 5.12% farm owners usually depended on the rainfall (Table 4).

Table IV: Source of water in the respective fish farm

Sources of water	Number of Different farms Monoculture N=08	Total N=39 Polyculture N=31	Percentage (%)	_'
Shallow tube well	7	22	29	74.36
Deep tube well	1	7	8	20.52
Rain fall	0	2	2	5.12

Water quality parameter

The water quality parameters like water temperature, transparency, pH, dissolved oxygen were measured to know whether it is suitable or not (Table 5).

Table V. Water quality parameters of the farms in the study areas.

Parameters	Small farm	Medium farm	Large farm
Tomoroustumo (oC)	25.98±4.26	25.94±5.06	25.81±7.14
Temperature (°C)	(14.01-32.40)	(14.15-32.40)	(14.20-31.40)
Transparency (cm)	28.48±2.79	29.30±3.64	36.24±4.61
Transparency (cm)	(23.80-37.50)	(25.40-41.40)	(24.10-45.20)
pН	8.45±0.30	8.08±0.33	8.17±0.35
_	(7.70-8.40)	(7.80-8.60)	(7.70-8.55)
Dissolved oxygen (mg.L-1)	4.72±0.66	4.47±0.70	4.05±0.88
Dissolved oxygen (mg.L-1)	(3.80-6.12)	(3.50-6.10)	(3.20-5.43)

Management of fish culture

Pre-stocking management

About 90% fish farmers dried their ponds 2-4 year's interval. But rest of others dried their ponds 4-5 year's interval because the very large water body. 84% farmer used lime in their pond. The dose of lime varies from 0.25 kg/dec. to 1.50 kg/dec. About 55% farmers don't use fertilizer and manure in their water bodies. They used compost as a fertilizer and in some cases used poultry litter in their pond. About 80% farm owners knew it was harmful for aquatic environment but they used it for the save of feed cost. The dose of the poultry litter was indiscriminate.

Stocking management

The maximum fish farm owner used fry in their fish culture farm. At first the fry are stocked in the nursing pond. After nursing, the fingerlings are stocked in the culture pond. The highest size of the pangus and bangla fish were 3-4 inch and the tilapia was 1.5 inch. It was found that higher density of pangus, tilapia and indian major carps were maintained in the farms of Bhaluka Upazilla (Table 6).

Table VI: Stocking density of fry in monoculture and polyculture systems in the study area.

Stocking density (No/ha)	Monoculture (N=08)	Percentage (%)	Polyculture (N=31)	Percentage (%)
50,000-75,000	6	75	2	6.45
75,001-1,00,000	1	12.5	1	3.22
1,00,001-1,25,000	1	12.5		-
1,25,001-1,50,000	-	-	4	12.91
1,50,001-175,000	-	-	6	19.35
1,75,001-2,00,000	-	-	1	3.22
2,00,001-2,25,000	-			-
2,25,001-2,50,000	-		2	6.45
2,50,001-2,75,000	-		4	12.91
2,75,001-3,00,000	-		5	16.13
3,00,001-3,25,000	-	-	-	-
3,25,001-3,50,000	-	- 1	-	-
3,50,001-3,75,000		_	4	12.91
3,75,001-4,00,000	- 🔠	-	2	6.45

Post-stocking management

In the studied area, about 5.12% monoculture and 17.96% polyculture farm owners sampled weekly, 7.69% and 20.51% farm owners sampled fortnightly, 5.12% and 25.65% farm owner sampled monthly and 2.56% and 15.39% farm owner sampled irregularly. 90% farm used supplementary ready made feed and rest of 10% farm owner used home made feed. The different types of company supplied feeds in the farms. They don't maintain actual feed ingredients. Farm owner used Saudi Bangla fish feed, Mega feed, ACI fish feed, CP, Biswas fish feed, Aftab fish feed etc. Farmers used two types of fish feed; sinking feed and floating feed. The cost of supplementary man made feed were range to be 22.0 to 23.0 BDT./kg and the price of company ready made floating and sinking feeds were found to be 25.0 to 33.0 BTK/kg (Table 7).

Table VII: Commercial supplementary feed used in different fish farms of the study area

Company Name	User percentage (%)	Price/Bag (BDT)
ACI fish feed	10.28	680
Saudi Bangla fish feed	12.85	800
Mega fish feed	17.95	820
Aftab fish feed	7.70	740
CP fish feed	5.12	750
National fish feed	7.70	720
Fresh fish feed	10.28	660
Quality fish feed	7.70	650
Biswas fish feed	10.28	650

In monoculture practice feeding rates were found to be 5-6% and 6-7% of fish body weight for small fish, 1-2% and 2-3% for large fish. In polyculture feeding rates were found to be 5-6% and 6-7% of fish body weight for small fish, 1-2% and 2-3% for large fish. 35.89% farmer used feeding frequency 1 time/day, 53.85% used 2 times/day and 10.26% used 3 times/day, respectively (Table 8).

Table VIII: Feeding frequency of the farms

Feeding frequency	Monoculture	Monoculture Polyculture		Percentage (%)
	N=08	N=31		
1 time/day	3	11	14	35.89
2 times/day	4	17	21	53.85
3 times/day	1	3	4	10.26

In monoculture practice, farm owners harvest the fish yearly in a culture period and in poly culture system harvesting was practiced two to four times yearly. Among 31 polyculture farms 67.75% had done partial harvesting.

Fish production

In case of man made feed, the highest production was recorded at 31,820 kg/ha/yr where lowest production was found to be 10,157 kg/ha/yr with an average production of 23,520 kg/ha/yr. On the other hand, in case of company feed the highest production was recorded at 56,810 kg/ha/yr and the lowest production was 14,210 kg/ha/yr with an average production of 34,350 kg/ha/yr (Table 9).

Table IX: Fish production in the study area.

Feed	Highest production (kg/ha/yr)	Lowest Production (kg/ha/yr)	Average
Farmmade feed	31,820	10,157	23,520
Company feed	56,810	14,210	34,350

Feed Ingredients analysis

The different types of feed used in the Bhaluka upazilla were analyzed. Proximate composition of the feeds was analyzed according to AOAC (1995) method, nitrogen free extract (NFE) by subtraction. Daily ration was adjusted by estimating of standing crop once weekly, fortnightly and monthly (Table 10).

Table X: Analyzed proximate composition (± SE) of different types feed.

Type of the Feed	Moisture		Lipid		Protein		Ash	
Type of the reed	Range	Mean	Range	Mean	Range	Mean	Range	Mean
Abdul Mannan farm	16.33-17.27	16.83	9.26-14.06	11.79	21.13-25.63	23.70	11.40-13.40	12.67
made feed	10.55-17.27	±0.27	7.20-14.00	±1.39	21.13-23.03	±1.33	11.40-15.40	±0.63
Md. Suruj Mia farm	11.11-13.48	12.51	8.94-12.94	10.47	21.12-27.05	24.57	13.42-17.20	15.44
made feed	11.11-15.40	±0.71	0.74-12.74	±1.24	21.12-27.03	±1.78	13.42-17.20	±1.10
Md.Azad farm made	15.68-25.75	19.36	13.21-15.60	14.32	20.56-23.92	22.20	8.40-15.78	12.51
feed	13.06-23.73	±3.21	13.21-13.00	±0.69	20.30-23.92	±0.96		±2.17
Mega feed	10.29-14.83	12.085	8.50-13.30	11.303	26.92-30.60	28.714	12.69-18.81	15.597
wiega ieeu		±1.392		±1.442		±2.217		±1.773
Soudi Bangla feed	13.58-14.80	14.34	12.28-16.25	14.47	24.88-29.46	26.54	11.76-14.18	13.35
Joudi Dangia leed	13.36-14.60	±0.38	12.20-10.25	±1.96		±1.04		±0.79
ACI feed	11.36-12.12	11.812	8.35-10.12	9.392	26.75-29.83	27.546	19.29-23.59	21.609
ACTIEEU	11.30-12.12	±0.233	6.55-10.12	±0.532		±1.201		±1.252
Fresh feed	19.76-20.53	20.053	15.83-18.60	16.940	20.14-25.60	23.103	10.13-12.66	11.492
rresn ieeu	19.76-20.55	±0.239	15.83-18.60	±0.845		±1.026		±0.734
Biggreen food	11.59-12.28	11.931	8.93-9.28	9.123	27.07-30.76	28.707	14 05-16 97	15.439
Biswas feed		±0.199	8.93-9.28	±0.101	27.07-30.76	±1.085		± 0.845

DISCUSSION

In the study area it was found that most of the farm owners are young belongs to the age group range of 20-30 years (35%) where 15% were related to fish culture belongs to the age group 50-60 years which is very much similar to Hossain (2007). Educational status of the fish farmer plays a vital role for the proper management and utilization of the new technology. About 12% of the total men power are illiterate but had the ability of signature. About 70% of the men power had primary and high school education respectively. About 14% and 4% manpower of the farm had SSC to HSC and Graduate level education, respectively. Quddus *et. al.* (2000) stated that educational levels of the pond farmers in Demra, Dhaka was below 43% in SSC, below 38% in Bachelor and above were 19% respectively. About 30% of farm owners having single ownership, 15% having multiple ownership and 65% having leased water body. It is evident from many studies that multiple ownership is one of the main problems to improve the management of culture system as well as efficient use of resource for fish cultivation (Ali and Rahman, 1986; Mollah *et al.* 1990).

In the study area about 80% of the water bodies were irregular shaped due to tectonic origin and 20% water bodies were regular. The minimum size of the pond was 0.162 ha and maximum size of the pond was 37.247 ha where as the average pond size was 5.78 ha. Bangla pedia (2006) stated that the topography of the area was rough and irregular. The minimum size of the water bodies were 0.06 ha. and maximum size of the water body was 7ha. Ali *et al.* (2008) stated that an average pond size was 0.13 ha with single (64%) and multiple ownerships (36%). According to Islam (2005) average pond size was 0.16 ha (40 decimals) with a range from 0.04 ha (11decimals) to 0.81 ha (200 decimals).

During the study period water quality of different ponds was measured. It was found that the temperature ranges from 14 to 32°C, pH ranges from 7.7.-8.5, transparency ranges from 23-45 cm and the dissolve oxygen ranges from 3.8-6.12 mg.L⁻¹. Similar works have been done by Rahman *et al.* (1982) measured 25.05-32.89°C and 26.5-32.22 °C for surface and bottom temperature respectively. They recorded DO with a range of 0.40-8.60 mg/l. Wahab *et al.* (1995) recorded the water temperature varied between 27.2 to 32.4°C, secchi depth between 26 and 50 cm, pH was always around 6.0 and dissolved oxygen was always low varied between 2.2 to 7.5 mg/l.

It was found that the stocking density of pond was very high. The highest stocking density was 3, 90,000/ha and the lowest was 54,340/ha. The average stocking density was found between 1, 40,000 and 1, 60,000/ ha. Islam (2005) found that average stocking density to be 17, 370/ha/year and Saha (2003) found average stocking density to be 16,561/ha. NFEP-II (1998) suggested the stocking density about 14,820/ha is suitable.

In the study area feeding rate was different for growing fish and aged fish. In monoculture practice feeding rate were found to be 5-6% to 6-7% of fish body weight for small fish, 1-2% and 2-3% for large fish. In polyculture feeding rate was found 5-6% and 6-7% of fish body weight for small fish, 1-2% and 2-3% for large fish. 35.89% farmer used feeding frequency 1 time/day, 53.85% used 2 times/day and 10.26% used 3 times/day. The Department of Fisheries (DoF) 2009 recommended feeding rate is 3-6% of the total body weight and feeding frequency will be two or three times per day. Hossain *et al.* (1997) reported the rate of 5% of total body weight, daily in two to three times.

The ingredients are analyzed and found moisture, lipid, protein, and ash. The highest protein (28.714%) was found in Mega feed, lipid (16.94%) found in Fresh feed, moisture (20.053%) found in Fresh feed and ash (21.61%) found in ACI feed.

In case of man made feed the highest production was 31,820 kg/ha/yr where lowest production was 10,157 kg/ha/yr with an average production of 23,520 kg/ha/yr. The farm where company feed was used the highest production was recorded to be 56,810 kg/ha/yr and the lowest production was 14,210 kg/ha/yr with an average production 34,350 kg/ha/yr. The farmer who used the company feed found the best result. According to Islam (2005) yield of fish production was found to be 2,609 kg/ha/year. Rahman (2003) stated that the average production of carp was estimated about 2,925 kg/ha/yr. Islam and Dewan (1986) observed in North-West Bangladesh that the average annual yield was 1,025 kg/ha.

All the previous results are lower than the present study. So, It is indicated that production of the studied area is higher than any other part of the country. Apparently in the present study, overall production was higher to lower on remarkable company feed. Therefore, the result of the study suggested that a good quality of feed with a recommended stocking density is resulted a good production.

CONCLUSION

The fish culture system is developed in Bhaluka upazilla, Bangladesh which changes the socioeconomic status of the farmers and contributes a remarkable part of inland fish production in Bangladesh. As the fisheries sector plays a vital role in the socio-economic development, opportunity for employment, poverty alleviation of a large number of population and earning foreign currency of Bangladesh the following recommendations should be made to develop fish culture practice. 1. The Government should need to take steps to train up interested farm owners in the area of water quality management; quality feed production; production of good quality seed; fish health management, disease control and management and other farm management techniques; 2. More research needed to find out a quality feed and feeding frequency with a recommended stocking density. 3. Control measure should be taken against the indiscriminate use of fertilizers, chemicals, disinfectants, antibiotics, hormones and other toxic substances in the ponds. and 4. Effective linkage should be developed more between Upazilla fisheries office, NGO and fish pond owners.

REFERENCES

- Ali, M.H. and Rahman, M.I. (1986). An investigation on some socio-economic and technical problems in fish pond culture in two Districts of Bangladesh. Bangladesh. *J. Aquacult.*, 8(1).
- Ali, M.H., Hossain, M.D., Hasan, A.N.G.M. and Bashar, M.A. (2008). Assessment of the livelihood status of the fish farmers in some selected areas of Bagmara upazilla under Rajshahi district. *J. Bangladesh Agril. Univ.* 6 (2): 367-374pp.
- AOAC International (1995). In: Howritz, W. (Ed.), Official Methods of Analysis, 16th edn. AOAC, Washington, DC. 1018 pp.
- Bangla pedia (2006). Asiatic society of Bangladesh, Agroecological zone of Bangladesh. www.baglapedia.org.
- BBS (2011). Statical yearbook of Bangladesh, Bagladesh Bureau of Statistics, Statistical division, Government of the peoples of Republic of Bangladesh, Dhaka. 580p.
- Bernet, Schmidt D.H., Meier, W., Burkhardt, P., Holm and wahli, T. (1999). Histopathology in fish, Proposal for a protocol to asses aquatic pollution, *J. of Fish Dis.* **22**(1): 25-34.
- Chakraborty, B. K. (2009). Aquatic biodiversity of Someswari and Nethai River and Gharia and Nidaya beel of Northern Bangladesh. In: Takumi K. Nakamura (Ed.). Aquaculture Research Progress, Nova Science, Publishers, New York, USA.: 3231-268.

- DoF. (2009). Department of Fisheries, Ministry of Fisheries and Livestock, Matshya Pakkhya Saranika 13p.
- DoF. (2010). Department of Fisheries, Ministry of Fisheries and Livestock, Matshya Pakkhya Saranika pp122-124.
- Duncan, D.B. 1955. Multiple range and multiple *F*-testes. Biometrics 11: 1-42pp.
- Hossain, M.A., Ahmed, M. and Islam, M.N. (1997). Mixed culture of fishes in seasonal ponds through fertilizer and feeding. *Bangladesh J. Fish. Res.*, 1 (2):9-18pp.
- Hossain, M.M. (2007). Utilization of Mokash beel for Livelihood Management of Local Fishermen and their Socio-economic Conditions. M.S. Thesis, Department of Aquaculture, BAU, Mymensingh. 85p.
- Islam, M.S. (2005). Socio-economic status of pond fish farming in some selected areas of Dinajpur district. M.S. Thesis, Department of Fisheries Management, Bangladesh Agricultural University, Mymensingh, 65p.
- Islam, M.S. and Dewan, S. (1986). Economics of fpond fish culture in some selected areas of Bangladesh. *Bangladesh J. Aquaculture.*, 8 (1): 57-61pp.
- Mollah, A.R., Chowdhury S.N.I. and Habib M.A.B. (1990). Input output relations in fish production under various pond sizes, ownership pattern and constraints. *Bangladesh J. Train. Dev.*, 3(2): 87-101pp.
- NFEP-II (1998). Aquaculture manual for literate and semi-literate fish farmers. Northwest Fisheries Extension Project-2, Parbatipur, Dinajpur. 81p.
- Quddus M.A., Rahman M.S. and Moniruzzaman, M. (2000). Socio-economic eonditions of the pond owners of Demra, Dhaka. *Bangladesh J. Fish. Res.*, 4(2):203-207pp.
- Rahman M.S., Chowdhury M.Y., Haque A.K.M. and Haque M.S. (1982). Limnological studies of four ponds. *Bangladesh J. Fish.*, 2-5(1-2):25-35pp.
- Rahman, M.M. (2003). Socio-economic aspects of carp culture development of Gazipur, Bangladesh. M.S. thesis, Department of Fisheries Management, Bangladesh Agricultural University, Mymensingh, 72p.
- Saha, M.K. (2003). A study on fish production technology in North-West Bangladesh. M.S. thesis. Department of Aquaculture, Bangladesh Agricultural University, Mymensingh, 71p.
- Wahab, M.A., Ahmed Z.F., Islam A. and Rahmatullah S.M. (1995). Effect of introduction of common carp *Cyprinus carpio* (L) on the pond ecology and growth of fish in polyculture. *Aquacult. Res.*, 26:619 628pp.
- Zar, J.H. (1984). Biostatistics: Prentice-Hall, Inc. Englewood Cliffs, New Jersey, USA. 718p.

Asian Business Consortium realizes the meaning of fast publication to researchers, particularly to those working in competitive and dynamic fields. Hence, we offer an exceptionally fast publication schedule including prompt peer-review by the experts in the field and immediate publication upon acceptance.

Submit your next manuscript at- www.abcjournals.net

ABC Journals is a unique forum to offer open access to all of its articles.

Now ABC Journals's portfolio is over ten journals, which publish both online and in print.



- Pantidalam, Kuala Lampur, Malaysia
- Rd 4, Shyamoli, Dhaka-1207, Bangladesh
- 3900 Woodhue Place, Alexandria, VA 22309, USA

www.abcreorg.weebly.com / www.abcjournals.net

Asian Business Consortium (ABC) is a multi-disciplinary research, training, publishing, digital library supporting and service house. Though founded in 2010 as the Business and Computing organization of Asia, it was reconstituted as the ABC in 2011. It has been working for creating and nurturing talents in USA, Malaysia and Bangladesh since its inception. As ABC is going global, it intends to open chapters in Australia, Germany, Japan, Pakistan, and other Asian countries in near future. The objectives of consortium are solely centered round the welfare and humane attitude of the founders who enthusiastically took up this noble cause and materialized it with a view to promote research and educational activities for the encouragement of scholars to develop their knowledge, to publish their analysis oriented scientific researches in international Journals, books, the task of organizing workshops, seminars, conferences, training, personality development programs and allied services.

In addition to research activities, ABC provides a good number of scholarships to the poor and meritorious students at various levels of education throughout the world. It plays an important role in the field of research by funding research projects and publishing the research papers. This consortium will unquestionably become the mouth-piece of the dark horses and unacknowledged scholar whose endowed and commendable contributions shall be provided an outlet keeping in mind the greater good of the larger society of the world.

ABC runs the following international referred journals for creating a platform to share the thoughts of professionals, scholars and academicians throughout the world.

ABC Publications (ABC Journals)

- Asian Accounting and Auditing Advancement (4A Journal)
- Asian Business Review (ABR)
- Asian Journal of Applied Sciences and Engineering (AJASE)
- Global Disclosure of Economics and Business (GDEB)
- ABC Journal of Advanced Research (ABC-JAR)
- International Journal of Reciprocal Symmetry and Theoretical Physics (IJRSTP)
- American Journal of Trade and Policy (AJTP)
- Asian Journal of Humanity, Art and Literature (AJHAL)
- Malaysian Journal of Medical and Biological Research (MJMBR)
- Asia Pacific Journal of Energy and Environment (APJEE)
- Engineering International (EI)
- ABC Research Alert (Online)

Each journal home page provides specific information for potential authors and subscribers. Open access policy, the quick review process, rich editorial boards and quality publications have already made ABC Journals unique. **ABC Journals** are published under the direct supervisions of renowned academicians of the world.

Collaboration in Conference: ABC considers high-quality conference papers for publication. Please contact us for detailed information.

Collaboration in Publishing: If you like to start writing a book, propose a new journal or advertise in ABC journals, please feel free to contact us.