

Optimal bound on the quantum Fisher information

Based on few initial expectation values

Iagoba Apellaniz¹, Matthias Kleinmann¹, Otfried Gühne²,
& Géza Tóth^{1,3,4}

iagoba.apellaniz@gmail.com

¹Department of Theoretical Physics, University of the Basque Country, Spain

²Naturwissenschaftlich-Technische Fakultät, Universität Siegen, Germany

³IKERBASQUE, Basque Foundation for Science, Spain

⁴Wigner Research Centre for Physics, Hungarian Academy of Sciences, Hungary

Recent Advances in Quantum Metrology; Warsaw - 2016

Outline

- 1 Introduction and Motivation
- 2 QFI based on expectation values: Are they optimal?
 - Optimisation problem
- 3 Case study
 - Fidelities
 - Spin-squeezed states
 - Unpolarised Dicke states
- 4 Conclusion and outlook

- **Many inequalities** have been proposed to lower bound the quantum Fisher Information.

Bounds for qFI

$$F_Q[\varrho, J_z] \geq \frac{\langle J_x \rangle^2}{(\Delta J_y)^2}, \quad F_Q[\varrho, J_y] \geq \beta^{-2} \frac{\langle J_x^2 + J_z^2 \rangle}{(\Delta J_z)^2 + \frac{1}{4}},$$


$$F_Q[\varrho, J_z] \geq \frac{4(\langle J_x^2 + J_y^2 \rangle)^2}{2\sqrt{(\Delta J_x^2)^2 (\Delta J_y^2)^2 + \langle J_x^2 \rangle - 2\langle J_y^2 \rangle(1 + \langle J_x^2 \rangle) + 6\langle J_y J_x^2 J_y \rangle}}$$

[L. Pezzé & A. Smerzi, PRL **102**, 100401 (2009)]

[Z. Zhang & L.-M. Duan, NJP **16**, 103037 (2014)]

[I.A., B. Lücke, J. Peise, C. Klempt & G. Toth, NJP **17**, 083027 (2015)]

- *Many inequalities* have been proposed to lower bound the quantum Fisher Information.
- For large systems, we only have *a couple of expectation values* to characterise the state.



- *Many inequalities* have been proposed to lower bound the quantum Fisher Information.
- For large systems, we only have *a couple of expectation values* to characterise the state.



- *Many inequalities* have been proposed to lower bound the quantum Fisher Information.
- For large systems, we only have *a couple of expectation values* to characterise the state.
- The archetypical criteria that shows *metrologically useful entanglement*.

$$F_Q[\varrho, J_z] \geq \frac{\langle J_x \rangle}{(\Delta J_z)^2}$$

[L. Pezzé & A. Smerzi, PRL **102**, 100401 (2009)]

- *Many inequalities* have been proposed to lower bound the quantum Fisher Information.
- For large systems, we only have *a couple of expectation values* to characterise the state.
- The archetypical criteria that shows *metrologically useful entanglement*.
- It is essential either to *verify them or find new ones* for different set of expectation values.

- 1 Introduction and Motivation
- 2 QFI based on expectation values: Are they optimal?
 - Optimisation problem
- 3 Case study
 - Fidelities
 - Spin-squeezed states
 - Unpolarised Dicke states
- 4 Conclusion and outlook

The non-trivial exercise of computing the qFI

- Different forms of the qFI

$$F_Q[\varrho, J_z] = 2 \sum_{\lambda, \gamma} \frac{(p_\lambda - p_\gamma)^2}{p_\lambda + p_\gamma} |\langle \lambda | J_z | \gamma \rangle|^2$$

Alternatively, as convex roof

$$F_Q[\varrho, J_z] = \min_{\{p_k, |\Psi_k\rangle\}} 4 \sum_k p_k (\Delta J_z)_{|\Psi_k\rangle}^2$$

[M.G.A. Paris, Int. J. Quant. Inf. **7**, 125 (2009)]

[G. Tóth & D. Petz, PRA **87**, 032324 (2013)]

[S. Yu, arXiv:1302.5311]

The non-trivial exercise of computing the qFI

- Different forms of the qFI

$$F_Q[\varrho, J_z] = 2 \sum_{\lambda, \gamma} \frac{(p_\lambda - p_\gamma)^2}{p_\lambda + p_\gamma} |\langle \lambda | J_z | \gamma \rangle|^2$$

Alternatively, as convex roof

$$F_Q[\varrho, J_z] = \min_{\{p_k, |\Psi_k\rangle\}} 4 \sum_k p_k (\Delta J_z)_{|\Psi_k\rangle}^2$$

[M.G.A. Paris, Int. J. Quant. Inf. **7**, 125 (2009)]

[G. Tóth & D. Petz, PRA **87**, 032324 (2013)]

[S. Yu, arXiv:1302.5311]

- For pure states it's extremely simple

$$F_Q[\varrho, J_z] = 4 (\Delta J_z)^2$$

Optimisation based on the Legendre Transform

- When $g(\varrho)$ is a *convex roof*

$$g(\varrho) \geq \mathcal{B}(w := \text{Tr}[\varrho W]) = \sup_r (rw - \sup_{|\psi\rangle} [r\langle W \rangle - g(|\psi\rangle)]).$$

[O. Gühne, M. Reimpell & R.F. Werner, PRL **98**, 110502 (2007)]

[J. Eisert, F.G.S.L. Brandão & K.M.R. Audenaert, NJP **9**, 46 (2007)]

Optimisation for the qFI

The *simplicity* of qFI for pure states leads to

$$\mathcal{F}(w) = \sup_r (rw - \sup_{\mu} [\lambda_{\max}(rW - 4(J_z - \mu)^2)]).$$

For more parameters



$$\mathcal{F}(\mathbf{w}) = \sup_{\mathbf{r}} (\mathbf{r} \cdot \mathbf{w} - \sup_{\mu} [\lambda_{\max}(\mathbf{r} \cdot \mathbf{W} - 4(J_z - \mu)^2)]).$$

[I.A., M. Kleinmann, O. Gühne & G. Tóth, arXiv:1511.05203]



- 1 Introduction and Motivation
- 2 QFI based on expectation values: Are they optimal?
 - Optimisation problem
- 3 Case study
 - Fidelities
 - Spin-squeezed states
 - Unpolarised Dicke states
- 4 Conclusion and outlook

- Measuring F_{GHZ} and F_{Dicke}

- Measuring F_{GHZ} and F_{Dicke}



- Measuring F_{GHZ} and F_{Dicke}



- For fidelity of GHZ \Rightarrow *analytic solution*

$$\mathcal{F} = \Theta(F_{\text{GHZ}} - 0.5)(2F_{\text{GHZ}} - 1)^2 N^2$$

Measuring $\langle J_z \rangle$ and $(\Delta J_x)^2$ for Spin Squeezed States

- 3 operators $\{J_z, J_x, J_x^2\}$

Measuring $\langle J_z \rangle$ and $(\Delta J_x)^2$ for Spin Squeezed States

- 3 operators $\{J_z, J_x, J_x^2\}$
- Reducing one dimension of \mathcal{F} on the $\langle J_x \rangle$ direction

$$\mathcal{F} \geq \mathcal{F}(\langle J_x \rangle = 0)$$

\Downarrow

$$\mathcal{F}(\langle J_z \rangle, (\Delta J_x)^2) := \mathcal{F}(\langle J_z \rangle, \langle J_x^2 \rangle)$$

Measuring $\langle J_z \rangle$ and $(\Delta J_x)^2$ for Spin Squeezed States

- 3 operators $\{J_z, J_x, J_x^2\}$
- Reducing one dimension of \mathcal{F} on the $\langle J_x \rangle$ direction



$$\mathcal{F} \geq \mathcal{F}(\langle J_x \rangle = 0)$$

$$\Downarrow$$

$$\mathcal{F}(\langle J_z \rangle, (\Delta J_x)^2) := \mathcal{F}(\langle J_z \rangle, \langle J_x^2 \rangle)$$

- Pezze-Smerzi bound, $F_Q \geq \langle J_z \rangle^2 / (\Delta J_x)^2$, can be verified.

• 4-particle system



Left: For $(\Delta J_x)^2 < 1.5$ it almost coincides with the P-S bound $F_Q \geq \langle J_z \rangle^2 / (\Delta J_x)^2$. **Right:** The measurement of $\langle J_x^4 \rangle$ improve the bound.

[I.A., M. Kleinmann, O. Güne & G. Tóth, arXiv:1511.05203]

Scaling the result for large systems

Experimental setup \rightarrow [C. Gross *et al.*, Nature **464**, 1165 (2010)]

$$N = 2300 \quad \xi_s^2 = -8.2\text{dB} = 0.1514$$

Scaling the result for large systems

Experimental setup \rightarrow [C. Gross *et al.*, Nature **464**, 1165 (2010)]

$$N = 2300 \quad \xi_s^2 = -8.2\text{dB} = 0.1514$$

- We choose

$$\langle J_z \rangle = 0.85 \frac{N}{2}$$

Scaling the result for large systems

Experimental setup \rightarrow [C. Gross *et al.*, Nature **464**, 1165 (2010)]

$$N = 2300 \quad \xi_s^2 = -8.2\text{dB} = 0.1514$$


- We choose

$$\langle J_z \rangle = 0.85 \frac{N}{2}$$

- P-S bound results is

$$\frac{F_Q}{N} \geq \frac{1}{\xi_s^2} = 6.605$$

- Starting from small systems, and assuming bosonic symmetry.
- The results with our method converge to P-S bound!




- Starting from small systems.

Metrology with unpolarised Dicke states

- 3 operators $\{J_x^2, J_y^2, J_z^2\}$; Constraint: $\langle J_x^2 \rangle = \langle J_y^2 \rangle$.

Metrology with unpolarised Dicke states

- 3 operators $\{J_x^2, J_y^2, J_z^2\}$; Constraint: $\langle J_x^2 \rangle = \langle J_y^2 \rangle$.
- For $\sum_I \langle J_I^2 \rangle = \frac{N}{2}(\frac{N}{2} + 1)$ and 6 particles:



Realistic characterisation of Dicke state

Experiment \rightarrow [B. Lücke *et al.*, PRL **112**, 155304 (2014)]


$$N = 7900$$

$$\langle J_z^2 \rangle = 112 \pm 31$$

$$\langle J_x^2 \rangle = \langle J_y^2 \rangle = 6 \times 10^6 \pm 0.6 \times 10^6$$

- For that large system, we start from **small ones** similar to the spin-squeezed states.

- Numerical lower bound.



Similarly to the spin-squeezed states, the bound *converges quickly*.

[I.A., M. Kleinmann, O. Güne & G. Tóth, arXiv:1511.05203]

Conclusion and Outlook

- 1 We prove that for realistic cases *the optimisation is feasible*.

Conclusion and Outlook

- ① We prove that for realistic cases *the optimisation is feasible*.
- ② We used *our approach to verify* that the P-S bound is tight.

Conclusion and Outlook

- ① We prove that for realistic cases *the optimisation is feasible*.
- ② We used *our approach to verify* that the P-S bound is tight.
- ③ We have shown that the lower bounds can be *improved with extra constraints*.

Conclusion and Outlook

- ① We prove that for realistic cases *the optimisation is feasible*.
- ② We used *our approach to verify* that the P-S bound is tight.
- ③ We have shown that the lower bounds can be *improved with extra constraints*.
- ④ For large systems the *optimisation method can be complemented* with scaling considerations.

Conclusion and Outlook

- 1 We prove that for realistic cases *the optimisation is feasible*.
- 2 We used *our approach to verify* that the P-S bound is tight.
- 3 We have shown that the lower bounds can be *improved with extra constraints*.
- 4 For large systems the *optimisation method can be complemented* with scaling considerations.
- 5 The *method very versatile* and it can be used in many other situations.

Thank you for your attention!

Preprint → arXiv:1511.05203

Groups' home pages

→ <https://sites.google.com/site/gedentqopt>

→ <http://www.physik.uni-siegen.de/tqo/>



iagoba



matthias



otfried



géza