

# Optimal bound on the quantum Fisher Information

*Based on few initial expectation values of the prove state.*

**Iagoba Apellaniz**<sup>1</sup>, Matthias Kleinmann<sup>1</sup>, Otfried Gühne<sup>2</sup>,  
& Géza Tóth<sup>1,3,4</sup>

**iagoba.apellaniz@gmail.com**

<sup>1</sup>Department of Theoretical Physics, University of the Basque Country, Spain

<sup>2</sup>Naturwissenschaftlich-Technische Fakultät, Universität Siegen, Germany

<sup>3</sup>IKERBASQUE, Basque Foundation for Science, Spain

<sup>4</sup>Wigner Research Centre for Physics, Hungarian Academy of Sciences, Hungary

Recent Advances in Quantum Metrology; Warsaw - 2016

# Outline

- 1 Introduction and Motivation
- 2 QFI based on expectation values: Are they optimal?
  - Optimisation problem
- 3 Case study
  - Spin squeezed states
  - Unpolarised Dicke states
- 4 Conclusion and outlook

- ① **Many inequalities** have been proposed to lower bound the quantum Fisher Information.

### Bounds for qFI

$$F_Q[\varrho, J_z] \geq \frac{\langle J_x \rangle^2}{(\Delta J_y)^2}, \quad F_Q[\varrho, J_y] \geq \beta^{-2} \frac{\langle J_x^2 + J_z^2 \rangle}{(\Delta J_z)^2 + \frac{1}{4}},$$


$$F_Q[\varrho, J_z] \geq \frac{4(\langle J_x^2 + J_y^2 \rangle)^2}{2\sqrt{(\Delta J_x^2)^2 (\Delta J_y^2)^2 + \langle J_x^2 \rangle - 2\langle J_y^2 \rangle (1 + \langle J_x^2 \rangle)} + 6}$$

[ I.A., B. Lücke, J. Peise, C. Klempt & G. Toth, NJP **17**, 083027 (2015) ]

[ L. Pezzé & A. Smerzi, PRL **102**, 100401 (2009) ]

[ Z. Zhang & L.-M. Duan, NJP **16**, 103037 (2014) ]

- 1 **Many inequalities** have been proposed to lower bound the quantum Fisher Information.
- 2 Typically, we only have *a couple of expectation values* to characterise the state.



- 1 **Many inequalities** have been proposed to lower bound the quantum Fisher Information.
- 2 Typically, we only have *a couple of expectation values* to characterise the state.



- 1 *Many inequalities* have been proposed to lower bound the quantum Fisher Information.
- 2 Typically, we only have *a couple of expectation values* to characterise the state.
- 3 The archetypical criteria that demonstrates *metrologically useful entanglement* on the state.

$$F_Q[\varrho, J_z] \geq \frac{\langle J_x \rangle}{(\Delta J_z)^2}$$

[ L. Pezzé & A. Smerzi, PRL **102**, 100401 (2009) ]

- ① *Many inequalities* have been proposed to lower bound the quantum Fisher Information.
- ② Typically, we only have *a couple of expectation values* to characterise the state.
- ③ The archetypical criteria that demonstrates *metrologically useful entanglement* on the state.
- ④ It is essential either to *verify them or find new ones* for different set of expectation values.

- 1 Introduction and Motivation
- 2 QFI based on expectation values: Are they optimal?
  - Optimisation problem
- 3 Case study
  - Spin squeezed states
  - Unpolarised Dicke states
- 4 Conclusion and outlook



# The non-trivial exercise of computing the qFI

- Different forms of the qFI

$$F_Q[\varrho, J_z] = 2 \sum_{\lambda, \gamma} \frac{(p_\lambda - p_\gamma)^2}{p_\lambda + p_\gamma} |\langle \lambda | J_z | \gamma \rangle|^2$$

$$F_Q[\varrho, J_z] = \min_{\{p_k, |\Psi_k\rangle\}} 4 \sum_k p_k (\Delta J_z)_{|\Psi_k\rangle}^2$$

[ M.G.A. Paris, Int. J. Quant. Inf. **7**, 125 (2009) ]

[ G. Tóth & D. Petz, PRA **87**, 032324 (2013) ]

[ S. Yu, arXiv:1302.5311 ]

# The non-trivial exercise of computing the qFI

- Different forms of the qFI

$$F_Q[\varrho, J_z] = 2 \sum_{\lambda, \gamma} \frac{(p_\lambda - p_\gamma)^2}{p_\lambda + p_\gamma} |\langle \lambda | J_z | \gamma \rangle|^2$$

$$F_Q[\varrho, J_z] = \min_{\{p_k, |\Psi_k\rangle\}} 4 \sum_k p_k (\Delta J_z)_{|\Psi_k\rangle}^2$$

[ M.G.A. Paris, Int. J. Quant. Inf. **7**, 125 (2009) ]

[ G. Tóth & D. Petz, PRA **87**, 032324 (2013) ]

[ S. Yu, arXiv:1302.5311 ]

- For pure states it's extremely simple

$$F_Q[\varrho, J_z] = 4 (\Delta J_z)^2$$

# The non-trivial exercise of computing the qFI

- Different forms of the qFI

$$F_Q[\varrho, J_z] = 2 \sum_{\lambda, \gamma} \frac{(p_\lambda - p_\gamma)^2}{p_\lambda + p_\gamma} |\langle \lambda | J_z | \gamma \rangle|^2$$

$$F_Q[\varrho, J_z] = \min_{\{p_k, |\Psi_k\rangle\}} 4 \sum_k p_k (\Delta J_z)_{|\Psi_k\rangle}^2$$

[ M.G.A. Paris, Int. J. Quant. Inf. **7**, 125 (2009) ]

[ G. Tóth & D. Petz, PRA **87**, 032324 (2013) ]

[ S. Yu, arXiv:1302.5311 ]

- For pure states it's extremely simple

$$F_Q[\varrho, J_z] = 4 (\Delta J_z)^2$$

- In the general case, usually *lower bounded* by its "classical" counterparts.

# Optimisation: Legendre Transform

- For a convex function of the state, we construct a *tight lower bound* as follows,

$$g(\varrho) \geq \mathcal{B}(\{w_k := \langle W_k \rangle\}) = \sup_{\{r_k\}} (r \cdot w - \sup_{\varrho} [r \cdot \langle W \rangle_{\varrho} - g(\varrho)]).$$

The scalar product reads as follows  $r \cdot w \equiv \sum_k r_k w_k$   
 and  $r \cdot \langle W \rangle_{\varrho} \equiv \sum_k r_k \langle W_k \rangle_{\varrho}$ .

# Optimisation: Legendre Transform

- For a convex function of the state, we construct a *tight lower bound* as follows,

$$g(\varrho) \geq \mathcal{B}(\{w_k := \langle W_k \rangle\}) = \sup_{\{r_k\}} (r \cdot w - \sup_{\varrho} [r \cdot \langle W \rangle_{\varrho} - g(\varrho)]).$$

- When  $g(\varrho)$  is defined as *infimum over the convex roof*, the 2<sup>nd</sup> optimisation simplified to pure states only,

$$\mathcal{B}(\{w_k\}) = \sup_{\{r_k\}} (r \cdot w - \sup_{|\psi\rangle} [r \cdot \langle W \rangle - g(|\psi\rangle)]).$$

[ O. Gühne, M. Reimpell & R.F. Werner, PRL **98**, 110502 (2007) ]

[ J. Eisert, F.G.S.L. Brandão & K.M.R. Audenaert, NJP **9**, 46 (2007) ]

## Optimisation for the qFI

The *simplicity* of qFI for pure states allows to rewrite the problem:

$$\mathcal{F}(\{w_k\}) = \sup_{\{r_k\}} (r \cdot w - \sup_{\mu} [\lambda_{\max}(r \cdot W - 4(J_z - \mu)^2)]).$$

- Therefore, we have parametrised the optimisation, which leads to a *more efficient finding* of the solution.



[ I.A., M. Kleinmann, O. Güne & G. Tóth, arXiv:1511.05203 ]

- 1 Introduction and Motivation
- 2 QFI based on expectation values: Are they optimal?
  - Optimisation problem
- 3 Case study
  - Spin squeezed states
  - Unpolarised Dicke states
- 4 Conclusion and outlook

- We'll present *2 main cases*, spin-squeezed states and unpolarized Dicke states.
- Though, we apply our method to projectors with great success, we will focus on collective  $J_n$  operators.



- We'll present *2 main cases*, spin-squeezed states and unpolarized Dicke states.
- Though, we apply our method to projectors with great success, we will focus on collective  $J_n$  operators.



- One of the cases using projector operators, *i.e.*, using the *fidelity*, leads to *analytic solution*!

# Measuring $\langle J_z \rangle$ and $(\Delta J_x)^2$ for Spin Squeezed States

- 1 We use the following 3 operators  $\{J_z, J_x, J_x^2\}$  to characterize the input state with their respective expectation values.

# Measuring $\langle J_z \rangle$ and $(\Delta J_x)^2$ for Spin Squeezed States

- 1 We use the following 3 operators  $\{J_z, J_x, J_x^2\}$  to characterize the input state with their respective expectation values.
- 2 In the direction of  $\langle J_x \rangle$  the worst case is when  $\langle J_x \rangle$  takes the *value zero*.


# Measuring $\langle J_z \rangle$ and $(\Delta J_x)^2$ for Spin Squeezed States

- 1 We use the following 3 operators  $\{J_z, J_x, J_x^2\}$  to characterize the input state with their respective expectation values.
- 2 In the direction of  $\langle J_x \rangle$  the worst case is when  $\langle J_x \rangle$  takes the *value zero*.
- 3 Hence, the optimisation can be accomplished *only using 2 operators*  $\{J_z, J_x^2\}$  while the resulting bound is mapped directly to  $\langle J_z \rangle, (\Delta J_x)^2$ .

# Measuring $\langle J_z \rangle$ and $(\Delta J_x)^2$ for Spin Squeezed States

- 1 We use the following 3 operators  $\{J_z, J_x, J_x^2\}$  to characterize the input state with their respective expectation values.
- 2 In the direction of  $\langle J_x \rangle$  the worst case is when  $\langle J_x \rangle$  takes the *value zero*.
- 3 Hence, the optimisation can be accomplished *only using 2 operators*  $\{J_z, J_x^2\}$  while the resulting bound is mapped directly to  $\langle J_z \rangle, (\Delta J_x)^2$ .
- 4 Since  $F_Q \geq \langle J_z \rangle^2 / (\Delta J_x)^2$  is a valid lower bound, we compare it with our numerical result.

- We numerically optimise the lower bound of qFI for a 4 particle system for all possible values of  $\langle J_z \rangle$  and  $(\Delta J_x)^2$ .



**Figure:** Below the dashed line,  $\mathcal{F}$  surpasses the shot noise limit. Cross point enhanced with extra parameter. The result shows an extreme similarity with respect to  $F_Q \geq \langle J_z \rangle^2 / (\Delta J_x)^2$ .


[ I.A., M. Kleinmann, O. Güne & G. Tóth, arXiv:1511.05203 ]

# Metrology with unpolarised Dicke states

- 1 In order to characterize the unpolarised Dicke state, we use these 3 operators,  $\{J_x^2, J_y^2, J_z^2\}$ , with the following constraint,  $\langle J_x^2 \rangle = \langle J_y^2 \rangle$ .

# Metrology with unpolarised Dicke states

- 1 In order to characterize the unpolarised Dicke state, we use these 3 operators,  $\{J_x^2, J_y^2, J_z^2\}$ , with the following constraint,  $\langle J_x^2 \rangle = \langle J_y^2 \rangle$ .
- 2 For  $\sum_I \langle J_I^2 \rangle = \frac{N}{2}(\frac{N}{2} + 1)$ , *i.e.* maximal, the following figure shows an illustrative result for 6 particles.





# Realistic characterisation of Dicke state

Experiment  $\rightarrow$  [ B. Lücke *et al.*, PRL **112**, 155304 (2014) ]

- We now consider an interesting experimental case with BECs where the following initial values are measured.

## Experimental details of unpolarized Dicke state with BEC

$$\begin{aligned} N &= 7900 & \langle J_z^2 \rangle &= 112 \pm 31 \\ \langle J_x^2 \rangle &= \langle J_y^2 \rangle &= 6 \times 10^6 \pm 0.6 \times 10^6 \end{aligned}$$

- For that number of particles we developed a *powerful extrapolation* tool.

# Extrapolation

Procedure:

- 1 First, we equivalently increase the expectation values and the unknown bound *to the symmetric subspace*.

$$\langle J_z^2 \rangle_{\text{sym}, N} = 145.69 \quad \langle J_x^2 \rangle_{\text{sym}, N} = 7.8 \times 10^6$$

# Extrapolation

Procedure:

- 1 First, we equivalently increase the expectation values and the unknown bound *to the symmetric subspace*.

$$\langle J_z^2 \rangle_{\text{sym}, N} = 145.69 \quad \langle J_x^2 \rangle_{\text{sym}, N} = 7.8 \times 10^6$$

- 2 We freeze  $\langle J_z^2 \rangle$  while we reduce the particle number of an *auxiliary system*. Here, we perform the optimisation.

$$2\langle J_x^2 \rangle_{\text{sym}, N'} = \frac{N'}{2} \left( \frac{N'}{2} + 1 \right) - \langle J_z^2 \rangle_{\text{sym}, N'}$$

# Extrapolation

Procedure:

- 1 First, we equivalently increase the expectation values and the unknown bound *to the symmetric subspace*.

$$\langle J_z^2 \rangle_{\text{sym}, N} = 145.69 \quad \langle J_x^2 \rangle_{\text{sym}, N} = 7.8 \times 10^6$$


- 2 We freeze  $\langle J_z^2 \rangle$  while we reduce the particle number of an *auxiliary system*. Here, we perform the optimisation.

$$2\langle J_x^2 \rangle_{\text{sym}, N'} = \frac{N'}{2} \left( \frac{N'}{2} + 1 \right) - \langle J_z^2 \rangle_{\text{sym}, N'}$$

- 3 The extrapolation is directly obtained by

$$\frac{\mathcal{F}}{\sum_l \langle J_l^2 \rangle} \approx \frac{\mathcal{F}_{\text{sym}, N}}{N^2/4} \approx \frac{\mathcal{F}_{\text{sym}, N'}}{N'^2/4}$$

- We are ready now to find numerically the lower bound for the experimental case.



**Figure:** The extrapolated lower bound approaches rapidly to the desired value. The points over the line indicate the systems used for optimisation.

[ I.A., M. Kleinmann, O. Güne & G. Tóth, arXiv:1511.05203 ]

# Conclusion and Outlook

- ① We have found that for very interesting and realistic cases *the optimisation is feasible*.

# Conclusion and Outlook

- ① We have found that for very interesting and realistic cases *the optimisation is feasible*.
- ② We used *our approach to verify* the tight bounding of one of the inequalities.

## Conclusion and Outlook

- ① We have found that for very interesting and realistic cases *the optimisation is feasible*.
- ② We used *our approach to verify* the tight bounding of one of the inequalities.
- ③ We have shown that the lower bounds can be *improved with few extra considerations*.



## Conclusion and Outlook

- ① We have found that for very interesting and realistic cases *the optimisation is feasible*.
- ② We used *our approach to verify* the tight bounding of one of the inequalities.
- ③ We have shown that the lower bounds can be *improved with few extra considerations*.
- ④ It has been shown that the *optimisation method can be complemented* with scaling considerations such as the extrapolation for the Dicke states case.

## Conclusion and Outlook

- 1 We have found that for very interesting and realistic cases *the optimisation is feasible*.
- 2 We used *our approach to verify* the tight bounding of one of the inequalities.
- 3 We have shown that the lower bounds can be *improved with few extra considerations*.
- 4 It has been shown that the *optimisation method can be complemented* with scaling considerations such as the extrapolation for the Dicke states case.
- 5 This *method is constructed to be universal* and it could be used in many other situations.

# *Thank you for your attention!*

Groups' home pages

→ <https://sites.google.com/site/gedentqopt>

→ <http://www.physik.uni-siegen.de/tqo/>



iagoba



matthias



ottfried



géza