

# Optimal bound on the quantum Fisher Information

*Based on few initial expectation values of the prove state.*

**Iagoba Apellaniz**<sup>1</sup>, Matthias Kleinmann<sup>1</sup>, Otfried Gühne<sup>2</sup>,  
& Géza Tóth<sup>1,3,4</sup>

**iagoba.apellaniz@gmail.com**

<sup>1</sup>Department of Theoretical Physics, University of the Basque Country, Spain

<sup>2</sup>Naturwissenschaftlich-Technische Fakultät, Universität Siegen, Germany

<sup>3</sup>IKERBASQUE, Basque Foundation for Science, Spain

<sup>4</sup>Wigner Research Centre for Physics, Hungarian Academy of Sciences, Hungary

Recent Advances in Quantum Metrology; Warsaw - 2016

# Outline

- 1 Introduction and Motivation
- 2 QFI based on expectation values: Are they optimal?
  - Optimization problem
- 3 Case study
  - Spin squeezed states
  - Unpolarized Dicke states
- 4 Conclusion and outlook

- ① **Many inequalities** have been proposed to lower bound the quantum Fisher Information.

### Bounds for qFI

$$\mathcal{F}[\varrho, J_z] \geq \frac{\langle J_x \rangle^2}{(\Delta J_y)^2}, \quad \mathcal{F}[\varrho, J_y] \geq \beta^{-2} \frac{\langle J_x^2 + J_z^2 \rangle}{(\Delta J_z)^2 + \frac{1}{4}},$$


$$\mathcal{F}[\varrho, J_z] \geq \frac{4(\langle J_x^2 + J_y^2 \rangle)^2}{2\sqrt{(\Delta J_x^2)^2 (\Delta J_y^2)^2 + \langle J_x^2 \rangle - 2\langle J_y^2 \rangle (1 + \langle J_x^2 \rangle)} + 6}$$

[ I.A., B. Lücke, J. Peise, C. Klempt & G. Toth, New J. Phys. **17**, 083027 (2015) ]

[ L. Pezzé & A. Smerzi, Phys. Rev. Lett. **102**, 100401 (2009) ]

[ Z. Zhang & L.-M. Duan, 2014 New J. Phys. **16** 103037 (2014) ]

- 1 **Many inequalities** have been proposed to lower bound the quantum Fisher Information.
- 2 Typically, we only have *a couple of expectation values* to characterize the state.



- 1 *Many inequalities* have been proposed to lower bound the quantum Fisher Information.
- 2 Typically, we only have *a couple of expectation values* to characterize the state.



- ① *Many inequalities* have been proposed to lower bound the quantum Fisher Information.
- ② Typically, we only have *a couple of expectation values* to characterize the state.
- ③ The archetypical criteria that demonstrates *useful entanglement* on the state.

$$\mathcal{F}[\varrho, J_z] \geq \frac{\langle J_x \rangle}{(\Delta J_z)^2}$$

[ L. Pezzé & A. Smerzi, Phys. Rev. Lett. **102**, 100401 (2009) ]

- ① *Many inequalities* have been proposed to lower bound the quantum Fisher Information.
- ② Typically, we only have *a couple of expectation values* to characterize the state.
- ③ The archetypical criteria that demonstrates *useful entanglement* on the state.
- ④ It is essential either to *verify them or find new ones* for different set of expectation values.

- 1 Introduction and Motivation
- 2 QFI based on expectation values: Are they optimal?
  - Optimization problem
- 3 Case study
  - Spin squeezed states
  - Unpolarized Dicke states
- 4 Conclusion and outlook



# The non trivial exercise of computing the qFI

## 1 Different forms of the qFI

$$\mathcal{F}[\varrho, J_z] = 2 \sum_{\lambda, \gamma} \frac{(p_\lambda - p_\gamma)^2}{p_\lambda + p_\gamma} |\langle \lambda | J_z | \gamma \rangle|^2$$

$$\mathcal{F}[\varrho, J_z] = \min_{\{p_k, |\psi_k\rangle\}} 4 \sum_k p_k (\Delta J_z)_{|\psi_k\rangle}^2$$

# The non trivial exercise of computing the qFI

## 1 Different forms of the qFI

$$\mathcal{F}[\varrho, J_z] = 2 \sum_{\lambda, \gamma} \frac{(p_\lambda - p_\gamma)^2}{p_\lambda + p_\gamma} |\langle \lambda | J_z | \gamma \rangle|^2$$

$$\mathcal{F}[\varrho, J_z] = \min_{\{p_k, |\psi_k\rangle\}} 4 \sum_k p_k (\Delta J_z)_{|\psi_k\rangle}^2$$

## 2 For pure states it's extremely simple

$$\mathcal{F}[\varrho, J_z] = 4 (\Delta J_z)^2$$

# The non trivial exercise of computing the qFI

## 1 Different forms of the qFI

$$\mathcal{F}[\varrho, J_z] = 2 \sum_{\lambda, \gamma} \frac{(p_\lambda - p_\gamma)^2}{p_\lambda + p_\gamma} |\langle \lambda | J_z | \gamma \rangle|^2$$

$$\mathcal{F}[\varrho, J_z] = \min_{\{p_k, |\psi_k\rangle\}} 4 \sum_k p_k (\Delta J_z)_{|\psi_k\rangle}^2$$

## 2 For pure states it's extremely simple

$$\mathcal{F}[\varrho, J_z] = 4 (\Delta J_z)^2$$

## 3 In the general case, usually *lower bounded* by its "classical" counterparts.



- 1 Introduction and Motivation
- 2 QFI based on expectation values: Are they optimal?
  - Optimization problem
- 3 Case study
  - Spin squeezed states
  - Unpolarized Dicke states
- 4 Conclusion and outlook











*Thank you for your attention!*

Group's home page → <https://sites.google.com/site/gedentqopt>