RAFFLES INSTITUTION
H2 Mathematics 9758
2026 Year 6 Term 3 Revision 2 (Zeta)
Topic(s): Complex Numbers and Sequences and of Series (Questions)

1 A sequence $u_1, u_2, u_3, \ldots$ is given by

$$u_1 = 2 \text{ and } u_{n+1} = \frac{3u_n - 1}{6} \text{ for } n \geq 1.$$

(i) Find the exact values of u_2 and u_3, and the value of u_{10} to 5 decimal places. [3]

(ii) It is given that $u_n \to l$ as $n \to \infty$. Showing your working, find the exact value of l. [2]

(iii) Describe the behaviour of the sequence. [1]

2 A sequence $a_1, a_2, a_3, \ldots$ is such that $a_n = 2a_{n-1} - 3n + K$, where K is a constant, and $n \geq 2$.

(i) Given that $a_1 = 2$ and $a_2 = 4$, find K, a_3 and a_{18}. [2]

It is known that the nth term of this sequence is given by $a_n = p(2^n) + qn + r$, where p, q and r are constants.

(ii) Find p, q and r. [3]

(iii) Find $\sum_{n=1}^{N} a_n$ in terms of N. [2]

3 Given that all terms in a geometric progression $\{u_n\}$, $n = 1, 2, 3, \ldots$ are positive with first term a and common ratio r, where $r \neq 1$, the sums H and C are defined as follows:

$$H = u_1 + u_2 + \ldots + u_n, \qquad C = \frac{1}{u_1} + \frac{1}{u_2} + \ldots + \frac{1}{u_n}.$$

(i) By expressing H and C in terms of a and r, prove that $\frac{H}{C} = u_1 u_n$. [3]

(ii) Express the product $u_1 u_2 u_3 \ldots u_n$ in the form of $\left(\frac{H}{C}\right)^{\mathrm{f}(n)}$, where f($n$) is a function of n. [2]

4 A finite sequence $\{a_n\}$ has 50 terms and is such that $a_{n+1} = a_n + 0.15$ for $n = 1, 2, 3, \ldots, 49$.

(i) Given that $a_{50} = 99a_1$, show that $a_1 = 0.075$. [2]

(ii) Find, without using a calculator, the value of $\sum_{n=1}^{50} a_n$. [2]


Another infinite sequence $\{b_m\}$ is such that $b_1 = a_{50}$ and $\frac{b_m}{b_{m-1}} = 0.98$ for $m \geq 2$.

(iii) Determine the smallest value of k such that $b_k < a_{25}$. [2]

(iv) Find the least value of h such that the sum of the first h terms of $\{b_m\}$ is more than 99% of its sum to infinity. [2]

(v) If $\frac{b_m}{b_{m-1}} = -0.98$ instead, find $\sum_{m=0}^{\infty} b_{1+3m}$. [2]

5



The line l with equation $y = 3x$ and the curve C with equation $y = x^2$ are shown in the diagram above. The x-coordinates of the point Q_1 on C is a_1, where $0 < a_1 < 3$.

A line parallel to the x-axis passing through the point Q_k, $(k \geq 1)$ on the curve C intersects the line l at P_{k+1}, and another line parallel to the y-axis passing through P_{k+1} intersects C at Q_{k+1}.

The sequence $a_1, a_2, \ldots,$ is defined by the x-coordinates of the points $Q_1, Q_2, \ldots,$ respectively.

(i) Show that the sequence satisfies $a_{k+1} = \dfrac{a_k^2}{3}$ for all $k \geq 1$. [2]

It is given that the k-th term of the sequence is given by $a_k = 3\left(\dfrac{a_1}{3}\right)^{2^{k-1}}$ for all $k \geq 1$.

For the rest of the question, assume $a_1 = 1$.

(ii) Find a_3. [1]

(iii) Find the area of the shaded rectangle in terms of a_k and a_{k+1}. By considering the area of the region bounded by C, the lines $x = 0$, $x = 1$ and the x-axis, give a geometrical reason why $\sum_{k=1}^{n}(a_k - a_{k+1})a_{k+2} < \dfrac{1}{9}$ for all $n \geq 1$. [3]

(iv) Given that $\sum_{k=1}^{n}(a_k - a_{k+1}) = 1 - \left(\dfrac{1}{3}\right)^{2^n - 1}$, show that $\sum_{k=1}^{n}(a_k - a_{k+1})a_{k+2} < \dfrac{1}{27}$ for all $n \geq 1$. [2]

6 **(a)** Find the sum of all integers between 200 and 1000 (both inclusive) that are not divisible by 7. [4]

(b)




Fig. 1




Fig. 2




Fig. 3




Fig. 4

Snowflakes can be constructed by starting with an equilateral triangle (Fig. 1), then repeatedly altering each line segment of the resulting polygon as follows:

1. Divide each outer line segments into three segments of equal length.
2. Add an equilateral triangle that has the middle segment from step 1 as its base.
3. Remove the line segment that is the base of the triangle from step 2.
4. Repeat the above steps for a number of iterations, n.

The 1st, 2nd and 3rd iteration produces the snowflakes in Fig. 2, Fig. 3 and Fig. 4 respectively.

(i) If a_0 denotes the area of the original triangle and the area of each new triangle added in the nth iteration is denoted by a_n, show that $a_n = \frac{1}{9}a_{n-1}$ for all positive integers n. [2]

(ii) Write down the number of sides in the polygons in Fig.1, Fig. 2 and Fig. 3, and deduce, with clear explanations, that the number of new triangles added in the nth iteration is $T_n = k\left(4^n\right)$ where k is a constant to be determined. [2]

(iii) Find the total area of triangles added in the nth iteration, A_n in terms of a_0 and n. [2]

(iv) Show that the total area of the snowflake produced after the nth iteration is $a_0\left[\frac{8}{5}-\frac{3}{5}\left(\frac{4}{9}\right)^n\right]$ units^2. [3]

The snowflake formed when the above steps are followed indefinitely is called the Koch snowflake.

(v) Determine the least number of iterations needed for the area of the snowflake to exceed 99% of the area of a Koch snowflake. [3]

7 **(a)** Meredith owns a set of screwdrivers numbered 1 to 17 in decreasing lengths. The lengths of the screwdrivers form a geometric progression. It is given that the total length of the longest 3 screwdrivers is equal to three times the total length of the 5 shortest screwdrivers. It is also given that the total length of all the odd-numbered screwdrivers is 120 cm. Find the total length of all the screwdrivers, giving your answer correct to 2 decimal places. **[4]**


(b) Meredith is building a DIY workbench, and she needs to secure several screws by twisting them with a screwdriver drill. Each time Meredith presses the button on the drill, the screw is rotated clockwise by u_n radians, where n is the number of times the button is pressed. Each press rotates the screw more than the previous twist, and on the first press, the screw is rotated by $\frac{2\pi}{3}$ radians. It is given that $\cos u_{n+1} = \frac{1}{2}\cos u_n - \frac{\sqrt{3}}{2}\sin u_n$ and $\sin u_{n+1} = \frac{1}{2}\sin u_n + \frac{\sqrt{3}}{2}\cos u_n$ for all $n \geq 1$.

(i) By considering $\cos(u_{n+1} - u_n)$ or otherwise, and assuming that the increase in rotation in successive twists is less than π radians, prove that $\{u_n\}$ is an arithmetic progression with common difference $\frac{\pi}{3}$ radians. **[3]**

(ii) Each screw requires at least 25 complete revolutions to ensure that it does not fall out. Find the minimum number of times Meredith has to press the drill button to ensure the screw is fixed in place. **[3]**

(iii) The distance the screw is driven into the workbench on the nth press of the drill, d_n, is proportional to the angle of rotation u_n. If the total distance the screw is driven into the workbench after 21 presses is 144mm, find the distance the screw is driven into the workbench on the first press. **[2]**

8 A company produces festive decorative Light Emitting Diode (LED) string lights, where micro LEDs are placed at intervals along a thin wire. In a particular design, the first LED (LED 1) is placed on one end of a wire with the second LED (LED 2) placed at a distance of d cm from LED 1, and each subsequent LED is placed at a distance $\frac{4}{5}$ times the preceding distance as shown.



(i) If the distance between LED 8 and LED 9 is 56.2 cm, find the value of d correct to 1 decimal place. [2]

(ii) Find the theoretical maximum length of the wire, giving your answer in centimetres correct to 1 decimal place. [2]

The LEDs consisting of three colours red, orange and yellow, are arranged in that order in a repeated manner, that is, LED 1 is red, LED 2 is orange, LED 3 is yellow, LED 4 is red, LED 5 is orange, LED 6 is yellow, and so on.

(iii) Find the colour of the LED nearest to a point on the wire 12.9 m from LED 1. [3]

(iv) If the minimum distance between any two consecutive LEDs is 1 cm so that they can be mounted on the wire, find the colour of the last LED. [3]

9 Bank *A* offers a study loan to students enrolled in an undergraduate course of study. The key features of the loan are:

- interest-free during the course of study,
- fixed monthly interest of 0.3% upon graduation,
- minimum monthly repayment of \$100.

Ali decides to take a study loan of \$50000 from Bank *A* on 1 January 2021 at the start of his 3-year undergraduate course.

(a) During his course of study, Ali pays \$200 at the end of January 2021 and on the last day of each subsequent month, he pays \$10 more than in the previous month. Thus on 28 February 2021, he pays \$210 and on 31 March 2021, he pays \$220, and so on. How much does Ali owe the bank at the end of his course of study? [2]

(b) Bank *A* charges interest on any outstanding amount of the loan on the first day of each month, starting on the month right after a student graduates.

(i) Upon graduation, Ali immediately found a job and decides to pay \$900 to bank *A* on the last day of each month, starting on the month right after he graduated. Show that Ali owes the bank \$34121 (to the nearest dollar) on the last day of the 3rd month after he graduated. [2]

(ii) Use the formula for the sum of a geometric progression to find an expression for the amount owed by Ali on the last day of the *n*th month after he graduated. Hence find in which month Ali pays off his study loan. [5]

(iii) Find the total amount of interest that Ali paid. [2]

(iv) If Ali decides to pay off his study loan within 3 years upon graduation, i.e. at the end of December 2026, what is the minimum amount, to the nearest dollar, that he needs to pay per month after graduation? [3]

10 **Do not use a graphing calculator for this question.**

(i) The complex number $a+\mathrm{i}$ is a root of the equation $z^2-2\sqrt{2}z+b=0$ where a and b are positive real constants. Find the exact values of a and b. [3]

It is given that $\mathrm{f}(z)=z^6-3z^4+11z^2-9$.

(ii) Show that $\mathrm{f}(z)=\mathrm{f}(-z)$. [1]

(iii) It is given that $a+\mathrm{i}$ is a root of the equation $\mathrm{f}(z)=0$. Using the results found in parts **(i)** and **(ii)**, show that $\mathrm{f}(z)$ can be factorised into 3 quadratic factors with real coefficients, leaving your answer in exact form. [4]

11 **(i)** Show that $\tan\frac{\alpha}{2}=\frac{1-\cos\alpha}{\sin\alpha}$. [2]

It is given that $w=-\frac{1}{2}+\mathrm{i}$ and $z=\frac{\sqrt{3}}{2}\mathrm{i}$.

(ii) Show that $|w-z|=\sqrt{2-\sqrt{3}}$. [2]

(iii) Using part **(i)**, find the exact value of $\arg(w-z)$. [3]

12 The complex number z is given by $z=\sqrt{3}-\mathrm{i}$. In an Argand diagram, mark and label clearly the points *Z, X* and *Y* representing the complex numbers z, $\mathrm{i}z$ and $z+\mathrm{i}z$ respectively. [2]

(i) State, with a reason, the geometrical representation of *OXYZ*. [1]

(ii) Using your Argand diagram, determine the argument of $z+\mathrm{i}z$. Hence deduce that $\tan\left(\frac{\pi}{12}\right)=2-\sqrt{3}$. [5]

13 **(a)** **(i)** Showing your working, find the complex numbers v and w which satisfy the following simultaneous equations.

$$v+2w=5$$
$$3v-w^*=1+5\mathrm{i}$$ [4]

(ii) Points *O*, *V* and *W* on an Argand diagram represent the complex numbers 0, v and w respectively.
Plot these points on an Argand diagram and state the transformation that maps the point *W* onto *V*. [2]

(b) The roots of the equation $z^4-2z^3-3z^2+az+b=0$, where a and b are real numbers, are z_1, z_2, z_3 and z_4. It is given that $z_1^2+z_2^2+z_3^2+z_4^2<0$. Explain why at most two of z_1, z_2, z_3 and z_4 are real. [2]

14 A complex number z varies with t such that

$$z = 2\cos t + \mathrm{i}(3\sin t), \text{ where } 0 \le t < 2\pi.$$

(a) By taking $x = \mathrm{Re}(z)$ and $y = \mathrm{Im}(z)$, sketch on an Argand diagram the curve that shows the positions of the points representing the complex number z. Find the cartesian equation of this curve. [3]

Two complex numbers z_1 and z_2 for two distinct values of t are such that $|z_1| = |z_2|$ and $0 < \arg(z_1) < \frac{\pi}{2}$.

(b) By referring to the Argand diagram in part **(a)**, find the possible values of $\arg(z_1 + z_2)$. [2]

It is given further that z_1 and z_2 are roots to the quadratic equation $z^2 + \alpha z + \beta = 0$.

(c) Explain whether it is necessary for α to be real. [2]

(d) Given that α is not real and $|z_1| = |z_2| = \frac{\sqrt{26}}{2}$, find the values for α and β. [4]

15 **(a)** The complex numbers z and w satisfy the following equations.

$$2z + |w| = 2 - 4\mathrm{i}$$
$$\mathrm{i}z - w = 2\mathrm{i}$$

Find z and w, giving your answers in the form $a + \mathrm{i}b$, where a and b are real numbers. [5]

(b) It is given that $z_1 = 2 - \mathrm{i}$ and $z_3 = -3 + 2\mathrm{i}$. On an Argand diagram, mark the points A and C representing z_1 and z_3 respectively. [1]

The point B and D on the Argand diagram represent complex numbers z_2 and z_4 respectively. Given that ABCD is a square, labelled in an anti-clockwise direction, find z_2 and z_4. [4]