



Basic Tooth Numbering


community Lab 4




Teeth names and numbers:

Teeth names are created by a combination of 4 different characteristics:

1. The arch where they are located: upper and lower or (maxillary and mandibular).




2. The side of the face: left and right.



3. The type of tooth: incisor, canine, premolar, molar



4. Set (primary or secondary).





Dentists refer to a specific tooth using a number or coding more usually than using teeth names.

There are several teeth numbering systems:

1_ Universal Tooth Numbering System :


The Universal Numbering System is a simplified method of identifying teeth that is approved and adopted by the American Dental Association.

Permanent teeth:

In the universal tooth numbering system,

The tooth number 1 is the patient's upper right (maxillary) third molar,


Numbering of teeth continues along the upper teeth and across to the last molar tooth back on the left side (**number 16**).



The tooth numbering continues ➡

By descending to the lower left third molar (number17) And numbering of teeth continues along the lower teeth and across to the last molar back on the right side of the mouth (number32).


All teeth that should be there are numbered, including those teeth that have been removed for any reason or have not erupted yet (e.g. wisdom teeth).



Deciduous (primary) teeth:

Universal Numbering System for children use letters instead of teeth numbers. The primary teeth are designated by upper case letters A through T,



with **A** being the patient's **upper right second primary** molar and the **T** being **lower right second primary** molar.



2_Fédération Dentaire Internationale (FDI):

Permanent teeth: For adults, the mouth is divided into quadrants numbered from 1 to 4 clockwise starting from the upper-right from the dentist's view.

The teeth are numbered from 1 to 8, going from the central incisor, canines, premolars to the third molar.



Deciduous (primary) teeth: For children, the mouth is divided into quadrants numbered from 5 to 8 clockwise starting from the upper-right from the dentist's view.

The children's deciduous teeth are numbered from 1 to 5, going from the central incisor, canines to the molars.


Maxillary Right

Maxillary Left

5	6
8	7

Mandibular Right

Mandibular Left



3- Palmer Notation (Chevron System, Quadrant System):

Permanent teeth:


In this system, the mouth is divided into four sections called quadrants. The numbers 1 through 8 and a unique **symbol** are used to identify the teeth in each quadrant

Deciduous (primary) teeth:

In children, this system use suppercase letters instead of numbers.

Upper Right	EDCBA	ABCDE	Upper Left
	87654321	12345678	
	87654321	12345678	
Lower Right	EDCBA	ABCDE	Lower Left

symbol



Upper Right			Upper Left
E D C B A		A B C D E	
8 7 6 5 4 3 2 1		1 2 3 4 5 6 7 8	
8 7 6 5 4 3 2 1		1 2 3 4 5 6 7 8	
E D C B A		A B C D E	
Lower Right			Lower Left



4-European System (Plus/Minus System)

For the permanent teeth, this system uses (+) for the maxillary teeth and (-) for the mandibular teeth.

The teeth of each quadrant are numbered from 1-8, beginning with the central incisor.

In primary teeth, the teeth of each quadrant are numbered from 01,02 to....05

Upper Right		Upper Left
50+40+30+20+10+		+01+02+03+04+05
8+7+6+5+4+3+2+1+		+1+2+3+4+5+6+7+8
8- 7- 6- 5- 4- 3- 2- 1-		-1 -2 -3 -4 -5 -6 -7 -8
50- 40- 30- 20- 10-		-01 -02 -03 -04 -05
Lower Right		Lower Left

Dental caries indices

- Index A numerical value describing the relative status of a population on a graduated scale with definite upper and lower limits, which is designed to permit and facilitate comparison with other populations classified by the same criteria and methods.

Indices used for dental caries assessment:

A- Permanent teeth index

B- Primary teeth index

Decayed-Missing-Filled Index (DMF) This index is of two types:


- 1- DMF teeth index (DMFT) which describe the amount (the prevalence) of dental caries in an individual. DMFT numerically expresses the caries prevalence and is obtained by calculating the number of teeth.
- 2- DMF surfaces index (DMFS) which measures the severity of dental caries and is obtained by calculating the number of surfaces

• D,d component:

Used to describe (Decayed teeth) which include:

1. Carious tooth.
2. Filled tooth with recurrent decay.
3. Only the roots are left.
4. Defect filling.
5. Temporary filling.
6. Filled tooth surface with another surface decayed (DMFT)


كل صورة ممكن تجيبها
وتسأل D M. F




Carious tooth



Only the roots are left



recurrent decay



Temporary filling.



Filled tooth surface with another surface with caries

M,m component:

Used to describe (Missing teeth due to caries only).

Other cases should be excluded these are:

1. Missing due Orthodontic treatment.
2. Impaction.
3. Missing due Periodontal disease.
4. Uneruptedteeth.
5. Congenitally missing.
6. Avulsion of teeth due to trauma or accident.



Avulsion tooth

F,f component:

All the following points should be present:

A. Filled teeth due to caries.

B. Permanent restorations.

C. There was no secondary (recurrent) caries or other area of the tooth with primary caries.

D. Undefected filling.

Teeth restored for reason other than dental caries should be excluded,
which include:



1. Trauma (fracture).
2. Hypoplasia (cosmetic purposes).
3. Bridge abutment (retention).
4. Preventive filling.



Trauma



Preventive filling



Hypoplasia

The excluded teeth in the DMF index are:


1-Supernumerary teeth.

2-The third molar





Supernumerary teeth

The examination should proceed in an orderly manner from one tooth or tooth space to the adjacent tooth or tooth space starting from **upper right second molar** and ending with **lower right second molar**.




Each tooth recorded as 4 surfaces for anterior teeth and 5 surfaces for posterior teeth, retained root (only the roots are left) was recorded as 4 D for anterior teeth, 5 D for posterior teeth. Missing tooth was recorded as 4 M for anterior teeth, 5 M for posterior teeth.



Differentiation between tooth missing due to caries and due to exfoliation:


هذه النقاط الثلاثة مهمة

1-By **age** of the patient if it is near to exfoliation time or not.



2-The **shape of ridge** is concave in carious missing tooth and straight in exfoliated one and permanent successor may be seen.

3-**Bad oral hygiene** mainly associated with carious teeth



concave in carious missing tooth



straight in normal exfoliated teeth

Differentiation between tooth missing due to caries and due to orthodontic treatment:

- 1-By type of teeth, in orthodontic treatment most teeth should be extracted are 4,5/c, d while in carious missing teeth any teeth may be involved.
- 2-Bilateral and /or opposing missing generally associated with orthodontic treatment, while in carious missing teeth it is not necessary.
- 3-Crowding or appliance may be seen in Orthodontic treatment.

Dentition status (DMFS / dmfs Index)

Right		Upper				
		M	O	D	B	L
	7					
	6					
E	5					
D	4					
C	3					
B	2					
A	1					
A	1					
B	2					
C	3					
D	4					
E	5					
	6					
	7					

DS:

MS:

FS:

DMFS:

DMFT:

Left		Lower				
		M	O	D	B	L
	7					
	6					
E	5					
D	4					
C	3					
B	2					
A	1					
A	1					
B	2					
C	3					
D	4					
E	5					
	6					
	7					

ds:

ms:

fs:

dmfs:

dmft:

D/d= Decay

M/m= Missing due to caries

F/f= Permanent filling due to caries

Permanent teeth index

Calculation of DMFT\DMFS:

$$DMF = D + M + F$$

Maximum score:

$$1-DMFT = 32 \text{ or } 28$$

$$2-DMFS = 12 * 4 + 20 * 5$$

$$= 48 + 100 = 148 \text{ or } 128$$

Minimum score = 0

Primary teeth index

dmft/ dmfs

Maximum scores: dmft= 20,

dmfs= 88

Minimum score = 0



Thank You

