

# STOAViz: Visualizing Saturated Thickness of Ogallala Aquifer

Tommy Dang, Long Hoang Nguyen

## Abstract

In this paper, we introduce STOAViz, a visual analytics tool for analyzing the saturated thickness of the Ogallala aquifer. The saturated thicknesses are monitored by sensors integrated on wells distributed on a vast geographic area. Our analytics application also captures the trends and patterns (such as average/standard deviation over time, sudden increase/decrease of saturated thicknesses) of water on individual well and group of wells based on their geographic locations. To highlight the usefulness and effectiveness of STOAViz, we demonstrate it on the Southern High Plains Aquifer of Texas. The work was developed using feedback from experts at the water resource center at a university. Moreover, our technique can be applied on any geographic areas where wells and their measurements are available.

## 1. Introduction

Water is the basic element that human relies on for all living and manufacturing activities. According to the National Ground Water Association report in 2016 [NGW16], we extract around 982 km<sup>3</sup> of water per year from the ground. This number serves as a good indicator of the need of water in daily life. Therefore, monitoring ground water, sustaining aquifer capability, and analyzing its changes are highly desirable by decision makers. Venki et al. [VUS17] discussed two types of errors in well management: improper removal of wells and redundancy of insensitive wells. Moreover, as the number of wells increases, it becomes more and more challenging for specialists to visualize and analyze such large amount of well monitoring data which comes in real time.

In this paper, we propose a visual analytics tool for analyzing the saturated thickness of the Ogallala aquifer, called STOAViz. Saturated thickness is the vertical thickness of the hydro-geologically defined aquifer in which the pore spaces of the rock forming the aquifer are filled with water [JAS00]. The main contributions of this paper are:

- We provide a data analytics tool for visualizing and analyzing a large number of well data distributed on a vast geographic area. Our approach is implemented as a web application which can track user location in real time.
- We integrate time series feature for detecting trends/patterns of monitored data at wells. This provides policy makers to come up with quick decisions.
- We demonstrate our application in the Southern High Plains Aquifer of Texas and conduct an informal user study at a water resource center.

The rest of this paper is organized as follows: We discuss the related work in the next section. Then we provide an overview of visualization tasks and describe implemented components in our system. We discuss our implementation and availability of our tool

in Section 4. We provide the expert feedback from our user study and discussion in Section 5. Finally, we conclude our paper with future plans.


## 2. Related Work

This section does not intend to survey all visualization tools for geospatial temporal visualizations [BDA\*14]. Instead, we discuss some related tools. One of the earliest spatial temporal illustration is the Napoleon's march towards Moscow. The illustration uses position, color, size, and text annotations to captures different states of the Napoleon's army during the Russian campaign in 1812 [Tuf86] within a single snapshot.

A simple approach to visualize temporal data on a 2D map is to attach the time series graphs directly on top of each geolocation [AA04]. This easily becomes too clustered for larger time series, especially when geolocations are not equally distributed. Showing summary statistics [DW13], such as average, standard deviation, or trends [SRd05] on a choropleth map resolves this concern. However, this comes back to the trade-off between details and simplifications.

There are numerous recent visualization ideas have been proposed to address the challenges of big spatial and temporal data analytics. Keim et al. [KPS04] suggest three step process for information visualization: overview first, zoom and filter, and then details-on-demand. CrimeViz [RRF\*10] utilizes this framework but also combines with mashup techniques to integrate additional visual elements on top of google map for crime analysis. Similarly, Ramakrishna et al. [RCM13] present another software mashup which can handle a large spatio-temporal data set with 2.5 million records using hexagon binning.

Keim et al. [Kei02] propose a classification of information visualization and visual data mining techniques which is based on



**Figure 1:** Major stages in STOAViz visualization: data cleansing and processing, visualization, and interaction.

the data type, the visualization technique, and the interaction technique. Soon after, Andrienko et al. [AAG03] provide an in-depth analytical review of exploratory spatio-temporal visualizations. The authors also categorize existing visualization-based techniques into two different types based on: (1) what types of spatio-temporal data they are applicable to; (2) what exploratory tasks they can potentially support. The work further defines a mapping mechanism between exploratory techniques and typologies of data exploration tasks. Even though many techniques use spatial location, color, shape, and size to encode information, they do not generate useful insights by just using simple statistical summaries, such as average or standard deviation [AA04].

Mohamed et al. [MLA13] discuss the use of interactive visualizations for knowledge discovery in order to support decision making in medical domain. In the water resources domain, Venki et al. [VUS17] combine ArgGIS 10.5 [Env16] with R integration [R C13] for well data visualization and analysis. This requires a strong technical skills from analysts and present limitations on interactive capabilities while analyzing multi-dimensional data.

### 3. STOAViz Stages

STOAViz aims to provide a high-level overview of saturated thickness of Ogallala aquifer. This tool also stretches out unusual patterns of water level such as sudden monthly increase/decrease [HS04, DW13] and overall trends [BPS\*07]. This section explains STOAViz stages in detail. Figure 1 shows a schematic overview of STOAViz:

- Processing:** We first clean and preprocess the monthly data collected from well integrated sensors from 1995 to 2016. At the end of the Processing stage, we have time series data, spatial interpolated data, and well metadata. (see Section 3.1)
- Visualization:** There are multiple linked view in our STOAViz visualization: the saturated thickness contour map, the google map, the standard time series graph, and the horizon graphs [HR08] (see Section 3.2).
- Interaction:** Users can click on a well or location on the map.

The selected well (and other neighboring wells) are plotted in the time series view. Users can further request a comparison chart to visualize individual well data and compare to its county average time series (see Section 3.3).

The STOAViz implements five low-level analysis tasks that largely capture users' activities when working with a given dataset [AES05, KPS04, AAG03]:

- T1: Provide an overview of large number of well data.
- T2: Retrieve and display well details on demand.
- T3: Group wells based on their geolocations.
- T4: Filter and sort wells by their time series features [HS04].
- T5: Detect suspicious or abnormal wells.

#### 3.1. Input datasets

We use the saturated thicknesses calculated at over 5,200 wells between the years 1995 and 2016 in the Southern High Plains Aquifer of Texas for visualization purpose. Saturated thickness in an unconfined aquifer is calculated using the equation: Saturated thickness (ST) = Land surface datum (LSD) - Water level (WL) - Bottom elevation of the aquifer (BT). LSD and BT are measured with respect to mean sea level (MSL), and WL is below ground surface.


#### 3.2. STOAViz overview

Main visualization components of STOAViz include the saturated thickness contour map, the google map, the standard time series chart, the horizon graphs, and the average comparison chart. All of these components share the same encodings: wells are drawn as filled circles whose radius is computed based on user selection strategy/feature. Selected wells are highlighted in different colors while other wells are in black with lower opacity.

**The Contour map** This contour map is plotted based on the well measurements and their interpolated data for areas where these measurements are not available. This map provides an high-level overview of water supply capability for a given geographic area (visualization task T1) as depicted in Figure 2. In particular, the

darker blue regions are areas with high volume of water supply while the lighter brown regions have thinner saturated thicknesses. Wells are overlaid on top of this contour map. Colored circles are wells with “highest monthly drop” on their time series.

Moreover, if user has location tracking enabled via browser or mobile phone, a flickering filled circle is displayed to represent current position of the user. This will help user to figure out how much water capability is existing in that area.



**Figure 2:** Saturated thickness contour map of Ogallala in 2013.


**The google map** The contour map serves as a starting point. As users select a well/location on the contour map, all of the neighboring wells (or all wells in the selected county) are plotted (visualization task **T3**) in the google map for further inspections. All google-map navigation capabilities, such as zooming in/out and panning, are also supported within this view. Figure 3 shows an example when a user mouse over a well (at the end of red arrow). All information (well id, GPS location, county, and measured saturated thickness over time) are displayed in a popup window (visualization task **T2**). This view is also linked to time series visualizations (the standard line graph or horizon graphs, based on user choice).

#### The time series line graphs


The time series line graphs for selected wells (or wells with “highest monthly drop” in this case) is depicted in Figure 4. We use the same color encoding across different views in *STOAViz*. When user has mouse over any well at the bottom legend, the selected well’s line chart is highlighted while other are faded out.

#### The horizon graphs

Users can switch to horizon graphs (instead of standard line graph) as depicted in Figure 5. This graph provides a different perspective in viewing well time series [JME10]. In particular, the time series is split into bands using a uniform interval. The interval and



**Figure 3:** Visualizing wells on google map (satellite view). The popup window displays well details on mouse over.




**Figure 4:** The standard line graphs for wells with “highest monthly drop” on their time series.

band colors are the same as on the contour map (see the right color legend of Figure 2). The split bands are now collapsed so that the higher bands layer on top [HR08]. The filled bacon time series chart becomes a horizon graph that takes up much less space [YAU] (how much less depends on how many split bands that we want).

The blank time intervals on the horizon graph are time periods when well measurements are not available. Blue bands represent high volumes of water supply. The wells in Figure 5 have been ordered base on how much water they lost within two consecutive months (visualization task **T4**). Upon mousing over any wells, a comparison graph of this well to the average time series is displayed. The comparison graph is described in the next section.


#### 3.3. Interaction

All views in *STOAViz* are linked. In other words, as users make a different selection on one view, others are updated accordingly. Moreover, users can request to show the comparison graph of a selected well to county/neighbor average time series. A bipolar color band is used to highlight differences: green for above the county



**Figure 5:** The horizon graphs of the same 20 wells as in Figure 4. Blue are thicker saturated thicknesses.

average and orange for below the county average. As depicted in Figure 6, the saturated thickness of well 712401 in Hartley county is thinner and thinner (after 2008, the saturated thickness is constantly below the county average). Policy makers in Hartley county may want to take actions to stop this decreasing trend.



**Figure 6:** Well 712401 in Hartley county is compared to its county average: green for above the county average and orange for below the county average. The light blue circles are monthly measurements of well 712401 from 1996 to 2016.

#### 4. Implementation

STOAViz is implemented as a web application utilizing the D3.js library [BOH11] and the Google map API [Inc]. This is an open source project. Source code, video, web demo, and project documentation are available on our Github repository at <https://github.com/iDataVisualizationLab/SaturatedThickness>.

#### 5. Evaluation and discussion

We solicited qualitative responses about STOAViz from two experts at a water resource center: one post-doc researcher and one associate professor. Both have at least nine-year of experiences in this research domain. The informal study started with a quick description (around 10 minutes) of the main components and their functionalities to familiarize users to the main GUI of STOAViz. Then the experts are free to use STOAViz before providing feedback. Both of them agree that the contour map provides an high level overview and can serve as the starting point for their well selection while the google map provides details of well (they can turn on the satellite images to inspect well location and county, see Figure ref). Moreover, the average graph is useful to see how the saturated thickness at a particular well behaves compared to others in the same county.

For the time series visualizations, they indicated that they are familiar with the line graphs but not the horizon graph. However, they both got the main idea of horizon graphs after a short explanation. Each expert has different interests and selected different features to inspect the time series data of wells. One selected wells with high variance of saturated thickness over time (depicted in Figure 3). She indicated that with horizon graphs, she can quickly compare patterns of a large number of wells. It makes sense that neighboring wells on the map have similar patterns. Moreover, missing data (well measurements are not available for certain period of time) can be discerned quickly.

The other expert selected “highest monthly drop” and detected an suspicious drop of water within a month (visualization task T5). He then used the Google map and grasp all wells within that county and many of them have similar drop on that month. He also commented that “for a smaller selection of wells (fewer than 5), I would like to use standard line graph since I can read the monthly values easily. For larger number of wells, line graph becomes too cluttered. Therefore, horizon patterns are much easier to compare, especially for wells with higher saturated thicknesses (darker blue areas).”

Besides the positive feedback, the experts also pointed out some limitations of STOAViz. For example, they suggested that the Google map and contour map should be automatically updated via interactions with the time series visualizations. These features have been incorporated in the final product on our Github repository.

#### 6. Conclusion and Future work

In this paper, we present STOAViz, a visual analytics tool with a case study of the Southern High Plains Aquifer of Texas. The tool utilize state of art spatial-temporal visualization techniques to support various linked views of saturated thicknesses monitored by a large number of wells distributed on a vast geographic area.

For future work, we will extend the application with Nanocubes integration [LKS13]. Nanocubes provides real-time visualization of billions of multidimensional spatial-temporal data expand zooming and other map viewing experiences. While the current version of STOAViz is limited to one layer of underground water (the saturated thickness contour map in Figure 2), we eventually target a 3D STOAViz which allows users to visualize and interact with different dynamic layers of underground water in a virtual reality environment.

## References

- [AA04] ANDRIENKO N., ANDRIENKO G.: Interactive visual tools to explore spatio-temporal variation. In *Proceedings of the working conference on Advanced visual interfaces* (2004), ACM, pp. 417–420. 1, 2
- [AAG03] ANDRIENKO N., ANDRIENKO G., GATALSKY P.: Exploratory spatio-temporal visualization: an analytical review. *Journal of Visual Languages & Computing* 14, 6 (2003), 503–541. 2
- [AES05] AMAR R., EAGAN J., STASKO J.: Low-level components of analytic activity in information visualization. In *Proc. of the IEEE Symposium on Information Visualization* (2005), pp. 15–24. 2
- [BDA\*14] BACH B., DRAGICEVIC P., ARCHAMBAULT D., HURTER C., SHEELAGH C.: A review of temporal data visualizations based on space-time cube operations. In *Eurographics Conference on Visualization* (2014), Eurographics. 1
- [BOH11] BOSTOCK M., OGIEVETSKY V., HEER J.: D3 data-driven documents. *IEEE Trans. Vis. Comput. Graph.* 17, 12 (2011), 2301–2309. 4
- [BPS\*07] BUONO P., PLAISANT C., SIMEONE A., ARIS A., SHMUEL G., JANK W.: Similarity-based forecasting with simultaneous previews: A river plot interface for time series forecasting. In *Information Visualization, 2007. IV '07. 11th International Conference* (July 2007), pp. 191–196. 2
- [DW13] DANG T., WILKINSON L.: TimeExplorer: Similarity search time series by their signatures. In *Proc. International Symp. on Visual Computing* (2013), pp. 280–289. 1, 2
- [Env16] ENVIRONMENTAL SYSTEMS RESEARCH INSTITUTE: *ArcGIS Desktop: Release 10.5*. ESRI, Redlands, California, CA, 2016. 2
- [HR08] HANNES REIJNER R. W. B.: The development of the horizon graph. <http://www.stonesc.com/Vis08workshop/DVD/Reijner,submission.pdf>, 2008. 2, 3
- [HS04] HOCHHEISER H., SHNEIDERMAN B.: Dynamic query tools for time series data sets: Timebox widgets for interactive exploration. *Information Visualization* 3, 1 (Mar. 2004), 1–18. 2
- [Inc] INC. A.: Google map. <http://maps.google.com>. 4
- [JAS00] J. A. SCHLOSS R. W. B.: Current saturated thickness. <http://www.kgs.ku.edu/HighPlains/atlas/atcst.htm>, last revision 2000. 1
- [JME10] JAVED W., McDONNEL B., ELMQVIST N.: Graphical perception of multiple time series. *IEEE Trans. Vis. Comput. Graph.* 16, 6 (Nov. 2010), 927–934. 3
- [Kei02] KEIM D. A.: Information visualization and visual data mining. *IEEE transactions on Visualization and Computer Graphics* 8, 1 (2002), 1–8. 1
- [KPS04] KEIM D. A., PANSE C., SIPS M.: Information visualization : Scope, techniques and opportunities for geovisualization. In *Exploring Geovisualization*, Dykes J., (Ed.). Elsevier, Oxford, 2004, pp. 1–17. 1, 2
- [LKS13] LINS L., KŁOSOWSKI J. T., SCHEIDECKER C.: Nanocubes for real-time exploration of spatiotemporal datasets. *IEEE Transactions on Visualization and Computer Graphics* 19, 12 (2013), 2456–2465. 4
- [MLA13] MOHAMED E. B., LTIFI H., AYED M. B.: Using visualization techniques in knowledge discovery process for decision making. In *Hybrid Intelligent Systems (HIS), 2013 13th International Conference on* (2013), IEEE, pp. 93–98. 2
- [NGW16] NGWA: Facts about global groundwater usage. Westerville, Ohio 43081-8978 USA. 1
- [R C13] R CORE TEAM: *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2013. ISBN 3-900051-07-0. 2
- [RCM13] RAMAKRISHNA A., CHANG Y.-H., MAHESWARAN R.: An interactive web based spatio-temporal visualization system. In *International Symposium on Visual Computing* (2013), Springer, pp. 673–680. 1
- [RRF\*10] ROTH R. E., ROSS K. S., FINCH B. G., LUO W., MAC EACHREN A. M.: A user-centered approach for designing and developing spatiotemporal crime analysis tools. In *Proceedings of GIScience* (2010), vol. 15. 1
- [SRd05] SHANBHAG P., RHEINGANS P., DESJARDINS M.: Temporal visualization of planning polygons for efficient partitioning of geo-spatial data. In *IEEE Symposium on Information Visualization, 2005. INFOVIS 2005*. (Oct 2005), pp. 211–218. 1
- [Tuf86] TUFTE E. R.: *The Visual Display of Quantitative Information*. Graphics Press, Cheshire, CT, USA, 1986. 1
- [VUS17] VENKI UDDAMERI ABDULLAH KARIM E. U., SRIVASTAVA P.: Sensitivity of wells in a large groundwater monitoring network and its evaluation using grace satellite derived information. In *Sensitivity Analysis in Earth Observation* (2017), Elsevier, pp. 235–256. 1, 2
- [YAU] YAU N.: Horizon graphs, with a food pricing example. <https://flowingdata.com/2015/07/02/changing-price-of-food-items-and-horizon-graphs/>. 3