



Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Lecture with Computer Exercises: Modelling and Simulating Social Systems with MATLAB

Project Report

Insert Title Here
...

Joël Mathys & Marc Rauch

Zurich
December 2017

Agreement for free-download

We hereby agree to make our source code for this project freely available for download from the web pages of the SOMS chair. Furthermore, we assure that all source code is written by ourselves and is not violating any copyright restrictions.

Joël Mathys

Marc Rauch

Contents

1	Abstract	4
2	Individual contributions	4
3	Introduction and Motivations	4
4	Description of the Model	5
4.1	Trail laying	5
4.2	Binary choices	5
4.3	U-Turns	6
4.4	Extensions	6
5	Implementation	7
6	Simulation Results and Discussion	7
7	Summary and Outlook	7

1 Abstract

The amount of people transported by plane in Europe has been rising in the last years and is predicted to grow further in the next years. This growth brings many questions and challenges. Many airports already operate at the limit of their capacity and the environment is suffering under the increasing air traffic. [2] In this paper we discuss the approach of maximizing the amount of people transported in the European air network. In an optimised scenario planes would be able to fill their capacity and more people could be transported with the same resources. This increase of efficiency of the planes would help with counteracting the pressure on the European flight network. We will discuss this problem with a model, which is inspired by the path finding of ant colonies. Ants have been found to be very efficient in finding paths, which lead to food sources. Those paths are optimised in terms of travel time to the source and the quality of the food.

2 Individual contributions

The paper is an team effort of Joël Mathys and Marc Rauch. There was no strict division of the work, but both of us had a focus on a certain part of the paper.

- Joël Mathys: Simulation, visualisation, graph generation, writing of report
- Marc Rauch: Behaviour of single ants, writing of report

3 Introduction and Motivations

Every year there are more flights starting from the airports of Europe. Therefore arises the question, whether the passengers can be transported more efficiently. In this paper we describe the efficiency as passengers transported per time. The goal is to maximize this efficiency and transport as many passengers as possible in a given time period. For this optimization the resources should only be redistributed, but no additional planes or airports have to be added to the system. We assume, that the most efficient way to transport these people is a compromise between distance to the destination and the size of the target airport. If the flight is long distance, then the destination airport has to be large, so that enough passengers have the desire to take this flight. If the distance is short, on the other hand, the flight is still worthwhile. Because ants use their trail laying to

4 Description of the Model

The model is based on research on the ant *Lasius niger*, in the book Self-organization in biological systems [1]. This paper describes the process of trail formation in ants. This phenomenon can easily be observed in the wild, by placing some sugar solution. After a short while the food will be discovered and shortly after the number of ants at the food source will increase rapidly, until eventually it stabilizes and a well established trail of ants can be observed from the nest to the sugar. To study this behaviour, several experiments have been conducted on a colony of *Lasius niger*. However the research was done under Laboratory conditions, so that the terrain could be controlled and therefore the experiments could be repeated. The terrain was constructed in such a way, that there was one path leading away from the nest, which then split into two paths with a food source at the end of both paths. The length, as well as the quality of the food sources could be controlled. Then the behaviour of the ants was observed and formulated as a mathematical model.

4.1 Trail laying

It was found, that the ants would place chemicals, so called pheromones, which brings other ants to follow the laid path. When an ant finds a food source it will lay pheromones on the way back to the nest and on the subsequent trips to the source. This acts as a positive feedback for the other ants, which are likely to follow that trail. The behaviour of trail laying is only observed by ants, who found food. The intensity of trail laying is dependant on the quality of the found food. The better the food source, the more pheromones will be deposited. The frequency of laying pheromones is also reduced based on the current direction of the ant. If the ant is walking in a trajectory with a bad angle to the direct route to the nest less pheromones will be placed. The placed chemicals also evaporate over time. They undergo an exponential decay given by formula 1.

$$\frac{dC}{dt} = -\frac{1}{l}C \quad (1)$$

where l is the lifetime of the pheromones and C the concentration.

4.2 Binary choices

In the experiment there was a choice the ants had to make, when the path split into two and they have to decide, whether to go left or right. This decision is made based on the concentration of pheromones on each branch. SO that the branch with a higher concentration of pheromones on it has a better chance of being chosen. For

that decision a formula for the probability of choosing one of the two branches was found.

$$P_L = \frac{(k + C_L)^n}{(k + C_L)^n + (k + C_R)^n} \text{ and } P_R = 1 - P_L \quad (2)$$

n is the non-linearity factor. with a high value of n a branch with little more pheromones then the other will be strongly favoured. With experiments it is approximately found that $n = 2$. k stands for the likelihood of the ant taking a path with no pheromones on it. P is the probability of choosing the left or right path and C the concentration of pheromones on each path.

4.3 U-Turns

It is often observed, that ants on a trail make an U-turn, return to the common branch point and follow the other path. After a U-turn the ant does not engage in trail laying, until it has gone back to the branch and follows the new path. There are two causes of this behaviour. If the amount of pheromones on the trail is low the ant may turn around. The second reason is direction based. If the orientation to the target is not good the probability of an U-turn is high. This probability can be modelled by equation 3.

$$P = \frac{P_0}{1 + \alpha C} \quad (3)$$

P_0 stands for the probability of an direction based U-turn. α describes the importance of the pheromone based U-turn.

4.4 Extensions

For our project we could not use the model as given in the paper. For that reason the model was changed in some aspects to ease the implementation and add additional functionality. Because the goal of this paper is to model more complicated networks then the one with only one binary decision, formula 2 has to be adapted to support a choice between more than just two paths.

$$P_i = \frac{(k + C_i)^n}{\sum_{j=1}^{nr} (k + C_j)^n} \quad (4)$$

[1]

5 Implementation

6 Simulation Results and Discussion

7 Summary and Outlook

References

- [1] Camazine, S. (2003). *Self-organization in biological systems*. Princeton University Press.
- [2] Eurocontrol (2009). Five major challenges of long-term air traffic growth. Retrieved from https://www.eurocontrol.int/eec/public/standard_page/ETN_2009_1_Growth.html.