

A mathematical understanding of deep learning

Jiang J.

Data Engineer, Data Scientist

ENSEEIH Computer Science Engineering Degree, INP Toulouse Dual MSc Research Degree in AI / Big Data / Ops
France

1 – Introduction

Frameworks such as TensorFlow or PyTorch make deep learning developments easy. They have made this field wide spread for every enthusiast. Implementations only needs an instinctive understanding of deep learning. The proper math aspect is little by little forgotten.

The objective is to do a summary of the important propositions. These propositions will be mathematically proven. The subject tackled is a multi-class classification problem with – dense layers, ReLU and SoftMax activation layers, Categorical cross-entropy loss, Stochastic gradient descent optimizer. However, all the elements below can be re-used or easily modified to cover a regression problem.

2 – Notation and Nomenclature

Definition – Let Ω a non-empty open subset of \mathbb{R}^m , $\|\cdot\|_m$ a norm on \mathbb{R}^m , $\|\cdot\|_{m'}$ a norm on $\mathbb{R}^{m'}$, and $f: \Omega \rightarrow \mathbb{R}^{m'}$. Then f continuous function is equivalent to

$$\forall a \in \Omega,$$

$$\forall \epsilon > 0, \exists \eta > 0, \forall x \in \Omega, \|x - a\|_m \leq \eta \Rightarrow \|f(x) - f(a)\|_{m'} \leq \epsilon$$

The notation $\mathcal{C}(\Omega, \mathbb{R}^{m'})$ means the set of continuous functions from Ω to $\mathbb{R}^{m'}$.

The notation $\mathcal{C}(\mathbb{R}^m)$ means the set of continuous functions from \mathbb{R}^m to $\mathbb{R}^{m'}$.

Definition – Let Ω a non-empty open subset of \mathbb{R}^m , $\|\cdot\|_m$ a norm on \mathbb{R}^m , $\|\cdot\|_{m'}$ a norm on $\mathbb{R}^{m'}$, and $f: \Omega \rightarrow \mathbb{R}^{m'}$. Then f derivable is equivalent to

$$\forall a \in \Omega, \exists f'(a) \in \mathbb{R}^{m'},$$

$$\forall \epsilon > 0, \exists \eta > 0, \forall x \in \Omega, \|x - a\|_m \leq \eta \Rightarrow \left\| \frac{f(x) - f(a)}{\|x - a\|_m} - f'(a) \right\|_{m'} \leq \epsilon$$

The notation $D(\Omega, \mathfrak{R}^{m'})$ means the set of derivable functions from Ω to $\mathfrak{R}^{m'}$.

The notation $D(\mathfrak{R}^m)$ means the set of derivable functions from \mathfrak{R}^m to \mathfrak{R}^m .

Definition – Let Ω a non-empty open subset of \mathfrak{R}^m , $\|\cdot\|_m$ a norm on \mathfrak{R}^m , $\|\cdot\|_{m'}$ a norm on $\mathfrak{R}^{m'}$, and $f: \Omega \rightarrow \mathfrak{R}^{m'}$. Then f piece-wise derivable is equivalent to

$\forall K \subset \Omega$ such as K compact and bounded,

$$\exists (K_i)_{i \in \llbracket 0, n \rrbracket} \text{ non-empty open subsets such as } \bigcup_{i=0}^n \overline{K_i} = K \text{ and } \forall (i, i') \in \llbracket 0, n \rrbracket^2, \\ i \neq i' \Rightarrow K_i \cap K_{i'} = \emptyset,$$

$$\forall i \in \llbracket 0, n \rrbracket, \exists f_i \in D^0(\overline{K_i}, \mathfrak{R}^{m'}), \forall x \in K_i, f_i(x) = f(x)$$

The notation $D_{pw}(\Omega, \mathfrak{R}^{m'})$ means the set of piece-wise derivable functions from Ω to $\mathfrak{R}^{m'}$.

The notation $D_{pw}(\mathfrak{R}^m)$ means the set of piece-wise derivable functions from \mathfrak{R}^m to \mathfrak{R}^m .

Definition – Let $(a_{i,j})_{i \in \llbracket 1, n \rrbracket, j \in \llbracket 1, m \rrbracket} \in \mathfrak{R}^{n \times m}$. Then the ordered rectangular array

$$A = \begin{bmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,m} \\ a_{2,1} & a_{2,2} & \cdots & a_{2,m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n,1} & a_{n,2} & \cdots & a_{n,m} \end{bmatrix}$$

is a real matrix of dimension $n \times m$.

The following notations are considered

$$\forall i \in \llbracket 1, n \rrbracket, \forall j \in \llbracket 1, m \rrbracket, A_{i,j} = a_{i,j}$$

$$\forall j \in \llbracket 1, m \rrbracket, A_{:,j} = \begin{bmatrix} a_{1,j} \\ a_{2,j} \\ \vdots \\ a_{n,j} \end{bmatrix}$$

$$\forall i \in \llbracket 1, n \rrbracket, A_{i,:} = [a_{i,1} \quad a_{i,2} \quad \cdots \quad a_{i,m}]$$

The notation $M_{n,m}$ means the matrix set of dimension $n \times m$ with coefficients in \mathfrak{R} .

The notation $M_{n,m}(E)$ means the matrix set of dimension $n \times m$ with coefficients in $E \subseteq \mathfrak{R}$.

Convention – A vector is a matrix with only one row. Thus, the real vector set \mathfrak{R}^m is equivalent to $M_{1,m}$.

Notation – The matrix transpose operation will be noted as A^T .

Definition – Let $A \in M_{n,m}$, and $B \in M_{m,p}$, and let the product noted $A \times B$ or AB be

$$C = A \times B = AB$$

where C is a $m \times p$ matrix with

$$\forall i \in [1, n] \quad , \quad \forall j \in [1, p] \quad , \quad C_{i,j} = \sum_{k=1}^m A_{i,k} \times B_{k,j}$$

3 – Activation functions

Definition – Let $z \in M_{n,m}$, and $f \in \zeta(\mathbb{R}^m) \cap D_{pw}(\mathbb{R}^m)$. Then the vector wise application

$$\forall i \in [1, n] \quad , \quad f : \begin{cases} \mathbb{R}^m \rightarrow \mathbb{R}^m \\ z_{i,:} \mapsto f(z_{i,:}) \end{cases}$$

is an activation function.

Proposition – Let $z \in M_{n,m}$. Then the following vector wise applications are activation functions

$$\forall i \in [1, n] \quad , \quad ReLU : \begin{cases} \mathbb{R}^m \rightarrow \mathbb{R}^m \\ z_{i,:} \mapsto \max(0_{\mathbb{R}^m}, z_{i,:}) \end{cases}$$

with \max the element-wise maximum operation between two vectors.

$$\forall i \in [1, n] \quad , \quad SoftMax : \begin{cases} \mathbb{R}^m \rightarrow \mathbb{R}^m \\ z_{i,:} \mapsto \frac{e^{z_{i,:}}}{\sum_{j'=1}^m e^{z_{i,j'}}} \end{cases}$$

with e the element-wise exponential operation.

The *SoftMax* function will be denoted as S for simplicity.

The Jacobian matrices are the following

$$\forall i \in \llbracket 1, n \rrbracket, J_{ReLU} : \left\{ \begin{array}{c} \mathfrak{R}_{\setminus \{0\}}^m \rightarrow M_{m,m} \\ z_{i,:} \mapsto \begin{bmatrix} 1_{\mathfrak{R}_{\setminus \{0\}}^+}(z_{i,1}) & 0 & \dots & 0 \\ 0 & 1_{\mathfrak{R}_{\setminus \{0\}}^+}(z_{i,2}) & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & 1_{\mathfrak{R}_{\setminus \{0\}}^+}(z_{i,m}) \end{bmatrix} \end{array} \right.$$

with $1_{\mathfrak{R}_{\setminus \{0\}}^+}$ the $\mathfrak{R}_{\setminus \{0\}}^+$ indicator function on $\mathfrak{R}_{\setminus \{0\}}$.

$$\forall i \in \llbracket 1, n \rrbracket, J_S : \left\{ \begin{array}{c} \mathfrak{R}^m \rightarrow M_{m,m} \\ z_{i,:} \mapsto J_S(z_{i,:}) \end{array} \right.$$

where $\forall (j, j') \in \{1, 2, \dots, m\}^2$, $[J_S(z_{i,:})]_{j,j'} = S(z_{i,:})_{i,j} \times (\delta_{j,j'} - S(z_{i,:})_{i,j'})$

with $\delta_{j,j'}$ the Kronecker delta.

Proof: TO DO.

3 – Categorical cross-entropy loss

Definition – Let $\hat{\Omega}$ and Ω non-empty \mathfrak{R} subsets with $\Omega =]\min_{\hat{\Omega}}, \max_{\hat{\Omega}}[$. Let $\hat{y} \in M_{n,m}(\hat{\Omega})$, $y \in M_{n,m}(\Omega)$, and $f \in \zeta(\Omega^m, \mathfrak{R}) \cap D_{pw}(\Omega^m, \mathfrak{R})$. Let $\|\cdot\|$ a norm on \mathfrak{R}^m . Then f is a loss function is equivalent to the application

$$f \circ g : \left\{ \begin{array}{c}]0, \max_{\hat{\Omega}} - \min_{\hat{\Omega}}[{}^m \rightarrow \mathfrak{R} \\ \epsilon \mapsto (f \circ g)(\epsilon) = f(\hat{y} + \epsilon) \end{array} \right.$$

is an increasing function according each component.

Proposition – Let $\hat{y} \in M_{n,m}(\{0,1\})$, and $y \in M_{n,m}(]0,1[)$. Then the application

$$\forall i \in \llbracket 1, n \rrbracket, \xi_{entropy} : \left\{ \begin{array}{c}]0,1[{}^m \rightarrow \mathfrak{R} \\ y_{i,:} \mapsto - \sum_{j=1}^m \hat{y}_{i,j} \log(y_{i,j}) \end{array} \right.$$

is a loss function.

The Gradient matrix is the following

$$\forall i \in \llbracket 1, n \rrbracket, \nabla_{\xi_{entropy}} : \left\{ \begin{array}{l}]0, 1[^m \rightarrow \Re^m \\ y_{i,:} \mapsto \begin{bmatrix} \frac{\hat{y}_{i,1}}{y_{i,1}} & \dots & \frac{\hat{y}_{i,m}}{y_{i,m}} \end{bmatrix} \end{array} \right.$$

Proof: TO DO.

4 – Dense layers

Definition – Let $y \in M_{n,m}$ the input vector, and $W \in M_{n,m}$ the weight matrix. Then the simple perceptron is defined as the operation

5 – Neural Network

Definition – Suppose a data set with n samples. Each sample have m features and a corresponding one-hot encoded label among l possible labels.

Let $X \in M_{n,m}$, and $Y \in M_{n,l}$ the matrices defining the features and the labels respectively for each sample.

Suppose a neural network with k layers.

6 – References