

Name: .....

Maths Class: .....

# SYDNEY TECHNICAL HIGH SCHOOL



## Year 12 Mathematics TRIAL HSC

August, 2015

*Time allowed: 3 hours plus 5 minutes reading time*

### ***General Instructions:***

- Reading time – 5 minutes
- Working time – 3 hours
- Marks for each question are indicated on the question.
- Approved calculators may be used
- All necessary working should be shown
- Full marks may not be awarded for careless work or illegible writing
- ***Begin each question on a new page***
- Write using black or blue pen
- All answers are to be in the writing booklet provided
- A set of Standard Integrals is provided at the rear of this Question Booklet, and may be removed at any time.

Total marks - 100

### **Section 1**

**10 Marks**

- \* Attempt Questions 1-10 on the sheet provided
- \* Allow about 15 minutes for this section

### **Section II**

**90 marks**

- Attempt Questions 11 – 16
- Allow about 2 hours and 45 minutes for this section

## Section I

10 marks

Attempt Questions 1-10

Allow about 15 minutes for this section.





Use the multiple-choice answer sheet for Questions 1-10

---

1. What is the value of  $\frac{\sqrt[3]{3}}{2\pi}$ , correct to 3 significant figures?

- A. 0.23
- B. 0.230
- C. 0.229
- D. 0.22

2. Which graph best represents  $y = x^2 + 2x + 1$ ?



3. What is the solution to the equation  $\log_3(x + 1) = 4$ ?

- A. 11
- B. 81
- C. 80
- D. 12

4. Which equation represents the line parallel to  $2x - 3y = 8$ , passing through the point  $(-1, 2)$ ?

- A.  $3x + 2y - 1 = 0$
- B.  $3x + 2y - 8 = 0$
- C.  $2x - 3y - 8 = 0$
- D.  $2x - 3y + 8 = 0$

5. Which expression is a factorisation of  $8x^3 - 27$  ?
- A.  $(2x - 3)(4x^2 + 12x - 9)$
- B.  $(2x + 3)(4x^2 - 12x + 9)$
- C.  $(2x - 3)(4x^2 + 6x + 9)$
- D.  $(2x + 3)(4x^2 - 6x + 9)$

6. The correct solutions to the equation  $2\sin^2 x - 1 = 0$  for  $-\pi \leq x \leq \pi$  are ?

A.  $\frac{\pi}{4}, \frac{3\pi}{4}$

B.  $\frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$

C.  $\pm \frac{\pi}{4}, \pm \frac{3\pi}{4}$

D.  $\pm \frac{\pi}{6}, \pm \frac{5\pi}{6}$

7. The value of  $\sum_{n=0}^{\infty} 2 \times \left(\frac{3}{5}\right)^n$  is ?


A.  $\frac{6}{5}$

B. 2

C. 5

D. 3

8. The length of  $PS$  in the following diagram is :



- A. 2
- B. 24
- C. 32
- D. 16

9. A parabola with a directrix  $x = 2$  has a focus at  $(-4, 3)$ . The focal length of this parabola is?


A. -6

B. -3

C. 6

D. 3

10. For the curve  $y = f(x)$ , which of the following statements is correct?



- A.  $f'(x) > 0$  at A and  $f''(x) < 0$  at B
- B.  $f'(x) < 0$  at A and  $f''(x) < 0$  at B
- C.  $f'(x) > 0$  at A and  $f''(x) > 0$  at B
- D.  $f'(x) < 0$  at A and  $f''(x) > 0$  at B

End of Section 1

## Section II

90 marks

Attempt Questions 11-16

Allow about 2 hours and 45 minutes for this section.


Answer each question on the appropriate writing page. Extra pages are available.

In Questions 11-16, your responses should include relevant mathematical reasoning and/or calculations.

---

### Question 11 (15 marks)

- a) Rationalise the denominator of  $\frac{4}{\sqrt{6}-2}$  2
- b) Factorise  $9x^2 - 37x + 4$  2
- c) Differentiate  $\frac{x+1}{x^3}$  2
- d) Find  $\int \frac{dx}{(2x+1)^3}$  2
- e) Find  $\int \cos\left(\frac{x}{2}\right) dx$  2
- f) Find the equation of the normal to the curve  $y = 2\sqrt{x}$  at the point  $x = 9$ . 3
- g) The perimeter of the sector AOB is 15cm. Calculate the size of angle AOB, correct to the nearest degree. 2






**Question 12 (15 marks)** *Start a new page.*

- a) Consider the series  $106 + 97 + 88 + \dots$

Find the sum of all the positive terms belonging to this series.

3


b)



Consider the points  $A(3,5)$ ,  $O(0,0)$  and  $B(6,2)$  in the diagram above.

- I. Find the equation of BO. 2
- II. Show that the distance of the interval BO is  $2\sqrt{10}$ . 1
- III. Show that the area of triangle AOB is 12 square units. 2
- IV. Hence, or otherwise, find perpendicular distance from the point O to the line AB. 2

- c) The parabola  $y = x^2 - 4x$  and the line  $y = x + 6$  intersect at the points A and B.



- I. Find the  $x$  co-ordinate of the points A and B. 2
- II. Calculate the area enclosed by the parabola  $y = x^2 - 4x$  and the line  $y = x + 6$ . 3

Question 13 (15 marks) *Start a new page.*

a) i) Write down the exact value of  $\tan 2x$  when  $x = \frac{\pi}{6}$  1

ii) Give the exact value of  $\int_0^{\frac{\pi}{6}} \sec^2 2x \, dx$  2

b) Simplify the expression  $\frac{\cos(\frac{\pi}{2} - \theta)}{\sin(\pi + \theta)}$  2

c) Consider the curve  $y = 2x^3 + 3x^2 - 12x - 9$

i. Find the co-ordinates of any stationary points and determine their nature. 3

ii. Show that a point of inflexion exists and state its co-ordinates. 2

iii. Sketch the curve  $y = f(x)$  in the domain  $-3 \leq x \leq 3$ , showing the  $y$ -intercept. 2

iv. For what values of  $x$ , in the domain given in part (iii), is the curve both increasing and concave down? 2

v. Write down the minimum value for  $y = f(x)$  in the interval  $-3 \leq x \leq 3$ . 1



Question 14 (15 marks) *Start a new page.*

a) i) Differentiate  $2xe^{-x}$  2

ii) Hence find  $\int_0^1 xe^{-x} dx$  2

b) The roots of the quadratic equation  $3x^2 - kx + 18 = 0$  are  $\alpha$  and  $\beta$

i) Find the value of  $\alpha\beta$ . 1

ii) Given that  $\alpha^2 + \beta^2 = 4$ , find the value/s of  $k$ . 2

c) The region bounded by the curve  $y = 1 + x^2$  and the  $x$ -axis between

$x = 0$  and  $x = 3$  is rotated about the  $x$ -axis to form a solid. Find the

volume of the solid. 3

d) The quantity of bacteria in a culture is growing according to the equation

$$\frac{dM}{dt} = kM$$

where  $M$  is the mass of the bacteria present in the culture in mg,  $t$  is the time in

hours and  $k$  is a constant.

i) Show that  $M = Ae^{kt}$  is a solution to the equation, where  $A$  is a constant. 1

ii) The time for the bacteria to double in mass is calculated to be 12 minutes.

Write down the value of  $k$ , correct to 4 significant figures. 2

iii) If the initial amount of bacteria is 8mg, after how many minutes does the


bacteria reach 1 gram? ( Answer to one decimal place ). 2

Question 15 (15 marks) *Start a new page.*

a) Solve  $2 \log_5 x - \log_5(x + 2) = \frac{2}{3} \log_5 125$

3

b)



In the diagram BEF is a triangle with  $BE = EF$ . BF is perpendicular to BD and the line AC through B is parallel to EF.

i) Copy the diagram into your answer booklet.

ii) Prove that BF bisects angle ABE.

2

iii) Prove that BD bisects angle EBC.

2

c) Water is being released from a rainwater tank. The rate of flow,  $R$  litres per minute is given by  $R = t(t - 12)^2$ , where  $t$  is the number of minutes since the water began to flow.

i) For how long does the water flow?

1

ii) Find the maximum rate of flow.

2

iv) What is the total volume of water released from the tank?

3

d) Given  $y = e^{kx}$ , find the value of  $k$  such that  $y = 2 \frac{dy}{dx} - \frac{d^2y}{dx^2}$

2

Question 16 (15 marks) *Start a new page.*

- a) Allied Lending is offering a special on loans of \$50,000.00 or more. The terms offered are a reducible interest rate of only  $\frac{1}{2}\%$  per month with the first six months interest free.

Paddington takes out a loan of \$80,000.00 to start his marmalade shop and agrees to the terms set out by Allied Lending.

Paddington agrees to repay the loan in equal monthly instalments of \$M, over 10 years with the first repayment due at the end of the first month. Let  $A_n$  be the amount owing at the end of the nth repayment.

- i) Write down an expression for the amount Paddington owes at the end of the first six months. 1

- ii) If the interest is calculated immediately before each repayment is made, show that the amount owing at the end of 8 months is,

$$A_8 = (80000 - 6M)(1.005)^2 - M(1.005 + 1) \quad 2$$

- iii) Hence show that  $A_{120} = (80000 - 6M)(1.005)^{114} - 200M(1.005^{114} - 1)$  2


- iv) Find the value of each monthly repayment correct to the nearest dollar. 2

b) ABCD is a rhombus of side 2cm.

P and Q are points on AC and AB respectively such that

$CP = AQ = x \text{ cm}$ .  $\angle DAP = \theta$  (where  $0 < \theta < \frac{\pi}{2}$ ) and  $\theta$  is a constant.

Let the area of the shaded area PDAQ be  $S \text{ cm}^2$ .



- |       |  |   |
|-------|--|---|
| (i)   | Show that $S = \frac{\sin \theta}{2} (4 \cos \theta - x)(2 + x)$               | 3 |
| (ii)  | If $\frac{dS}{dx} = 0$ , find $x$ in terms of $\theta$                         | 2 |
| (iii) | Find $\frac{d^2S}{dx^2}$ in terms of $\theta$                                  | 1 |
| (iv)  | Suppose that $\theta = \frac{\pi}{6}$ , show that $S$ attains its maximum when |   |
|       | $\frac{PC}{AC} = \frac{\sqrt{3}-1}{2\sqrt{3}}$                                 | 2 |

End of Assessment task

# STHS 2unit Trial

## Solutions

1. B

2. A

3. C

4. D

5. C

6. C

7. C

8. D

9. D

10. B

$$e) \int \cos\left(\frac{1}{2}x\right) dx$$

$$= 2\sin\left(\frac{x}{2}\right) + C$$

$$f) x = 9 \quad y = 2\sqrt{9} \quad \therefore (9, 6)$$

$$= 6$$

$$y' = x^{-1/2}$$

$$m_T = \frac{1}{\sqrt{9}}$$

$$= \frac{1}{3} \quad \therefore M_N = -3$$

$$\text{Eq: } y - 6 = -3(x - 9)$$

$$y - 6 = -3x + 27$$

$$3x + y - 33 = 0$$

## Question 11

$$a) \frac{4}{\sqrt{6}-2} \times \frac{\sqrt{6}+2}{\sqrt{6}+2}$$

$$= \frac{4(\sqrt{6}+2)}{6-4}$$

$$= 2(\sqrt{6}+2)$$

$$= 2\sqrt{6} + 4$$

either

$$g) \rho = r\theta + 12$$

$$r\theta = 3$$

$$6\theta = 3$$

$$\theta = \frac{1}{2}$$

$$\therefore \theta = \frac{1}{2} \times \frac{180}{\pi}$$

$$\approx 29^\circ$$

$$b) 9x^2 - 37x + 4$$

$$= (9x-1)(x-4)$$

$$c) \frac{x+1}{x^3} = x^{-2} + x^{-3}$$

$$\frac{d}{dx} = -2x^{-3} - 3x^{-4}$$

$$= \frac{-2}{x^3} - \frac{3}{x^4}$$

$$d) \int \frac{dx}{(2x+1)^3} = \int (2x+1)^{-3} dx$$

$$= \frac{(2x+1)^{-2}}{-2 \times 2} + C$$

$$= \frac{-1}{4(2x+1)^2} + C$$

## Question 12

a)  $106 + 97 + 88 + \dots$

AP  $a = 106$

$d = -9$

$T_n < 0$

$106 + (n-1)(-9) < 0$

$-9n < -715$

$n > 12.7$


$\therefore n = 13$

$T_{12} = 7$

$S_n = \frac{n}{2}(a+1) \text{ or } \frac{n}{2}(2a+(n-1)d)$

$= \frac{12}{2}(106+7)$

$= 678$



AB  $(3, 5)$   $(6, 2)$

$d_{AB} = \sqrt{(6-3)^2 + (2-5)^2}$   
 $= \sqrt{9+9}$

$= \sqrt{18}$

$= 3\sqrt{2}$

$\therefore \frac{1}{2}hb = \frac{1}{2} \times h \times 3\sqrt{2}$

$12 = \frac{1}{2} \times h \times 3\sqrt{2}$

$8 = h\sqrt{2}$

$h = \frac{8}{\sqrt{2}} \quad (4\sqrt{2})$

c) Solving

$x^2 - 4x = x + 6$

$x^2 - 5x - 6 = 0$

$(x-6)(x+1) = 0$

$x = 6, x = -1$

ii)  $A = \int_{-1}^6 (x+6) - (x^2-4x) dx$

$= \int_{-1}^6 (5x+6-x^2) dx$

$= \left[ \frac{5x^2}{2} + 6x - \frac{x^3}{3} \right]_{-1}^6$

$= \frac{5}{2}(36) + 36 - \frac{6^3}{3} - \left( \frac{5}{2} - 6 + \frac{1}{3} \right)$

$= 57 \frac{1}{6} u^2$

i) Equation  $M_{B0} = \frac{0-2}{0-6}$

$= \frac{1}{3}$

$\therefore y = \frac{1}{3}x \rightarrow 3y = x$

iii)  $B(6, 2)$   $O(0, 0)$

$d_{B0} = \sqrt{36+4}$

$= \sqrt{40}$

$= 2\sqrt{10}$

ii)  $A(3, 5)$   $d_1 = 12/\sqrt{10}$

$A = \frac{1}{2}hb$

$= \frac{1}{2} \times 12/\sqrt{10} \times 2\sqrt{10}$

$= 12 u^2$

# Question 13

$$\begin{aligned} i) \tan 2x &= \tan 2(\pi/6) \\ &= \tan \pi/3 \\ &= \sqrt{3} \end{aligned}$$

$$\begin{aligned} ii. \text{ inflexion } y'' &= 0 \\ 12x + 6 &= 0 \\ x &= -1/2 \\ y &= -5/2 \end{aligned}$$

$$\begin{aligned} ii) \int_0^{\pi/6} \sec^2 2x \, dx &= \left[ \frac{1}{2} \tan 2x \right]_0^{\pi/6} \\ &= \frac{1}{2} [\tan \pi/3 - \tan 0] \\ &= \frac{1}{2} [\sqrt{3} - 0] \\ &= \sqrt{3}/2 \end{aligned}$$

$$\begin{aligned} \text{concavity check} \\ x = -1 \quad x = -1/2 \quad x = 0 \\ y'' = -6 \quad 0 \quad y'' = 6 \end{aligned}$$

$\therefore$  change in concavity @  $(-1/2, -5/2)$


iii. end pts  $(-3, 0) \& (3, 36)$

$$\begin{aligned} b) \frac{\cos(\pi/2 - \theta)}{\sin(\pi + \theta)} &= \frac{\sin \theta}{-\sin \theta} \\ &= -1 \end{aligned}$$

$$c) y = 2x^3 + 3x^2 - 12x - 9$$

$$dy/dx = 6x^2 + 6x - 12$$

$$d^2y/dx^2 = 12x + 6$$



iv)  $\nearrow$  and  $\searrow$   $-3 \leq x < -2$

accept  $-3 < x < -2$

donot accept  $-3 \leq x \leq -2$

i) Stat pts

$$6(x^2 + x - 2) = 0$$

$$6(x+2)(x-1) = 0$$

$$\begin{array}{l|l} x = -2 & x = 1 \\ y = 11 & y = -16 \end{array}$$

$\therefore (-2, 11) \& (1, -16)$

v) min value is  $-16$

Nature

donot accept  $(1, -16)$

$x$	-3	-2	-1
$y'$	18	0	-12

$x$	-1	1	2
$y'$	-12	0	24

$\therefore$  MAX TP

$\&$

MIN TP

$(-2, 11)$

$(1, -16)$

### Question 14

a)  $\frac{d}{dx}(2xe^{-x})$

$$= 2e^{-x} - 2xe^{-x}$$

$$= \pi \left[ x + \frac{2x^3}{3} + \frac{x^5}{5} \right]_0^3$$

$$= \pi \left[ 3 + \frac{2 \cdot 27}{3} + \frac{3^5}{5} - (0) \right]$$

ii)  $\frac{1}{2} \frac{d}{dx}(2xe^{-x}) = e^{-x} - xe^{-x}$

$$\therefore \int xe^{-x} dx = \left[ \int e^{-x} - \int \frac{1}{2} \frac{d}{dx}(2xe^{-x}) \right] dx = \frac{348\pi}{5} u^3$$

$$= \int e^{-x} dx - xe^{-x}$$

d)  $M = Ae^{kt}$

$$= -e^{-x} - xe^{-x}$$

$$\frac{dM}{dt} = k \cdot Ae^{kt}$$

$$= k \cdot M \therefore \text{a sol}^n$$

$$\therefore \int_0^1 xe^{-x} dx = \left[ -e^{-x} - xe^{-x} \right]_0^1$$

$$= -e^{-1} - 1e^{-1} - (-e^0 - 0)$$

ii.  $2A = Ae^{k(12)}$

$$2 = e^{\frac{1}{5}k}$$

$$= -1/e - 1/e + 1$$

$$2 = e^{12k}$$

$$\ln 2 = \frac{1}{5}k$$

$$= 1 - 2/e$$

$$\ln 2 = 12k$$

$$k = 3.466$$

$$= \boxed{1 - \frac{2}{e}}$$

$$2 \frac{e^{-2}}{e}$$

$$k = \frac{1}{12} \ln 2$$

$$\approx 0.05776 \text{ (4 sig fig)}$$

b) i.  $\alpha\beta = c/a$

$$= 6$$

iii.  $1 \text{ gram} = 1000 \text{ mg}$

$$1000 = 8e^{kt}$$

using  $k = 0.0577$

$$125 = e^{kt}$$

ii.  $\alpha^2 + \beta^2 = 4$

$$\alpha + \beta = \frac{k}{3}$$

$$(\alpha + \beta)^2 - 2\alpha\beta = 4$$

$$\ln 125 = kt$$

$$(\alpha + \beta)^2 - 12 = 4$$

$$\left(\frac{k}{3}\right)^2 = 16$$

$$t = \ln 125 \div k$$

$$\frac{k}{3} = 4 \text{ or } -4$$

$$= 83.592 \dots$$

$$\therefore k = \pm 12$$

$$\therefore 83.6 \text{ minutes.}$$

c)  $V_x = \pi \int y^2 dx$

$$= \pi \int_0^3 (1+x^2)^2 dx$$

$$= \pi \int_0^3 1 + 2x^2 + x^4 dx$$



# Question 15

$$a) 2 \log_5 x - \log_5 (x+2) = \frac{2}{3} \log_5 125$$

$$\log_5 \left( \frac{x^2}{x+2} \right) = \log_5 (5^3)^{2/3}$$

$$\log_5 \left( \frac{x^2}{x+2} \right) = \log_5 5^2$$


$$\frac{x^2}{x+2} = 25$$

$$x^2 - 25x - 50 = 0$$

$$x = \frac{25 \pm \sqrt{825}}{2}$$

$$\text{as } x > 0$$

$$x = \frac{25 + \sqrt{825}}{2} \text{ only}$$



$$b) \text{ let } \angle ABF = \alpha$$

$$\angle BFE = \angle ABF \text{ (alternate angles } AC \parallel EF)$$

$$\angle FBE = \angle BFE \text{ (equal angles opposite equal sides } FE = BE \text{ given)}$$

$$\therefore \angle ABF = \angle FBE (= \alpha)$$

$$\therefore BF \text{ bisects angle ABE}$$

$$\angle EBD = 90 - \alpha \text{ (adjacent complementary with } \angle FBE)$$

$$\angle ABE + \angle EBD + \angle CBD = 180 \text{ (straight line)}$$

$$\therefore \angle CBD = 90 - \alpha = \angle EBD$$

and BD bisects  $\angle EBC$

$$c) R = \frac{dv}{dt} = t(t-12)^2$$

$$1) R > 0 \therefore 0 \leq t \leq 12$$

$\therefore$  flows for 12 minutes

$$ii) \text{ MAX flow rate } \frac{dR}{dt} = 0$$

$$R = t(t^2 - 24t + 144)$$

$$= t^3 - 24t^2 + 144t$$

$$\frac{dR}{dt} = 3t^2 - 48t + 144 = 0$$

$$t^2 - 16t + 48 = 0$$

$$(t-12)(t-4) = 0$$

$$\text{as } 0 \leq t \leq 12$$

test	$t = 4$	$t = 3$	$t = 4$	$t = 5$
$\frac{dR}{dt}$	27	0	0	-21

$$\text{max flow rate is } R = 4(4-12)^2 = 256 \text{ L/min}$$

$$iii) V = \int_0^{12} t^3 - 24t^2 + 144t \, dt$$

$$= \left[ \frac{t^4}{4} - 8t^3 + 72t^2 \right]_0^{12}$$

$$= \left[ \frac{12^4}{4} - 8(12)^3 + 72(12)^2 - 0 \right]$$

$$= 1728 \text{ L}$$

$$d) y = e^{kx} \quad y' = k e^{kx} \quad y'' = k^2 e^{kx}$$

$$y = 2y' - y''$$

$$e^{kx} = 2(ke^{kx}) - (k^2 e^{kx})$$

$$1 = 2k - k^2$$

$$k^2 - 2k + 1 = 0$$

$$(k-1)^2 = 0$$

$$\therefore k = 1$$

# Question 6

a)

$$(i) A_6 = 80000 - 6M$$

$$(ii) A_7 = (80000 - 6M)(1.005) - M$$

$$A_8 = [(80000 - 6M)(1.005) - M](1.005) - M$$

$$= (80000 - 6M)(1.005)^2 - M(1.005) - M$$

$$= (80000 - 6M)(1.005)^2 - M[1.005 + 1]$$

$$= (80000 - 6M)(1.005)^2 - M(2.005)$$

$$(iii) A_{120} = (80000 - 6M)(1.005)^{114} - M[1 + 1.005 + \dots + 1.005^{113}]$$

G.P.  $a=1$

$$r = 1.005$$

$$n = 114$$

$$= (80000 - 6M)(1.005)^{114} - M \left[ \frac{a(r^n - 1)}{r - 1} \right]$$

$$= (80000 - 6M)(1.005)^{114} - M \left[ \frac{1.005^{114} - 1}{0.005} \right] \quad (ii)$$

$$= (80000 - 6M)(1.005)^{114} - 200M(1.005^{114} - 1)$$

$$A_{120} = 0 \text{ Find } M$$

$$0 = 80000(1.005)^{114} - M[6(1.005) + 200(1.005^{114} - 1)] \quad (iii)$$

$$M = \frac{80000(1.005)^{114}}{6(1.005) + 200(1.005^{114} - 1)}$$

$$= \$862.681 \dots$$

∴ repayment is \$863.

(1)

Let  $A$  be the intersection of the diagonals.

$\angle XAD = \angle XAC = \theta$  [property of rhombi]

$$AX = 2\cos\theta \Rightarrow AC = 4\cos\theta$$


$$\therefore AP = 4\cos\theta - x$$

The shaded area is the sum of triangles  $ADP$  and

$$S = \frac{1}{2} \times 2 \times (4\cos\theta - x)\sin\theta + \frac{1}{2} \times (4\cos\theta - x)x$$

$$= \frac{\sin\theta}{2} (4\cos\theta - x)(x + 2)$$

[NB  $S$  is a concave down parabola in  $x$ ]



$$S = \frac{\sin\theta}{2} [8\cos\theta + (4\cos\theta - 2)x - x^2]$$

$$\frac{dS}{dx} = \frac{\sin\theta}{2} [(4\cos\theta - 2) - 2x] = \sin\theta(2\cos\theta - 1 - x)$$

$$\therefore \frac{dS}{dx} = 0 \Rightarrow x = 2\cos\theta - 1 \quad [\because \sin\theta \neq 0]$$

$$\frac{dS}{dx} = \sin\theta(2\cos\theta - 1 - x)$$

$$\therefore \frac{d^2S}{dx^2} = -\sin\theta \quad \left[ < 0 \text{ for } 0 < \theta < \frac{\pi}{2} \right]$$

$$(iv) \theta = \frac{\pi}{6}$$

$$\frac{dS}{dx} = 0 \Rightarrow x = 2\cos\left(\frac{\pi}{6}\right) - 1 = \sqrt{3} - 1$$

$$\frac{d^2S}{dx^2} < 0 \Rightarrow S \text{ is a maximum}$$

$$AC = 4\cos\left(\frac{\pi}{6}\right) = 2\sqrt{3}$$

$$\therefore \frac{PC}{AC} = \frac{\sqrt{3} - 1}{2\sqrt{3}}$$