

# SYDNEY TECHNICAL HIGH SCHOOL



## MATHEMATICS

Year 11

2 Unit

May 2006

Common Test

Time allowed: 70 mins

Name: \_\_\_\_\_

Teacher: \_\_\_\_\_

### Instructions:

- Begin each question on a new page
- Marks shown are approximate and may be varied
- Show necessary working
- Full marks may not be awarded if working is poorly set out or difficult to read

Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	TOTAL
/8	/7	/7	/7	/8	/7	/8	/6	/58

### Question 1

- a) Evaluate  $\frac{1.3 \times 10^{-3}}{8.6 - 4.2}$ . Give your answer in scientific notation using 2 significant figures. 2
- b) Express  $0.\dot{4}\dot{2}$  as a simplified fraction. 2
- c) Simplify  $\sqrt{27} - \sqrt{12}$  2
- d) Evaluate  $x^2 - 3x + 1$  when  $x = 3\sqrt{5}$ . Leave your answer in simplest form 2

### Question 2

- a) Simplify: (i)  $3x - 4x(3 - x)$  1
- (ii)  $(4a^3b)^2 \times \frac{a^2}{6b^3}$  2
- b) Express the answer to  $\sqrt{x} \div \frac{1}{x}$  in simplified index form 2
- c) Express  $8^a \times 4^{a+1}$  as a power of 2 2

### Question 3

- a) Factorise (i)  $x^2 - 5x - 6$  1
- (ii)  $x^3 - 8$  1
- (iii)  $x^3 + x^2 + x + 1$  1
- b) Simplify (i)  $\frac{x^2 - x}{x^2 - 1}$  2
- (ii)  $\frac{4}{x+2} - \frac{1}{x}$  2

### Question 4

- a) Evaluate  $|-3| - |-4|^2$  1
- b) Solve (i)  $\frac{2x-4}{3} = x+2$  2
- (ii)  $2x^2 + 9x - 5 = 0$  2
- (iii)  $|1 - 2x| = 6$  2

**Question 5**

- a) Solve simultaneously  $3x - y = 0$  and  $4x - 2y = 3$  3
- b) Solve  $\frac{4}{x+2} + x = 3$  2
- c) Given  $f(x) = 3x^2 - 2x$ , find and simplify:
- (i)  $f(-3)$  1
- (ii)  $f(x+h) + f(x)$  2

**Question 6**

a) Sketch each function. Use a ruler for all straight lines and show any  $x, y$  intercepts:

(i)  $y = \frac{1}{x+1}$  1

(ii)  $y = -\sqrt{4-x^2}$  1

(iii)  $y = |3x-6|$  1


- b) (i) Sketch  $y = x^2 + 2x - 3$ . Show all intercepts 2
- (ii) Solve  $x^2 + 2x - 3 \leq 0$ . You may use the sketch above to help 1
- (iii) What is the range of the function in (i) 1

**Question 7**

- a) State the domain and range of the function  $y = \sqrt{x-4}$  2
- b) Find the centre of the circle  $x^2 - 4x + 4 + y^2 + 6y = 0$  2
- c) (i) Sketch  $y = 2^x$  and  $y = |x-1|$  on the same axes 2
- (ii) Hence or otherwise, solve  $|x-1| > 2^x$  1
- (iii) On your graph in (i), neatly shade the intersected region of  $x \geq 0$  and  $y \geq 2^x$  1

**Question 8**

a)




O is the centre of the circle. If  $\angle AOC = x^\circ$ , prove

that  $\angle B = \frac{x^\circ}{2}$

3

b)



ABCDE is a regular pentagon.  $AF \parallel CD$  and DE is produced to F.

Set out a geometrical proof to find the size of  $\angle FAE$ . 3

# SOLUTIONS

① a)  $3.0 \times 10^{-4} \leftarrow \textcircled{1}$

b)  $x = 0.4242 \dots$

$100x = 42.4242 \dots$

$99x = 42 \quad \textcircled{1} \text{ for correct method}$

$x = \frac{42}{99}$

$= \frac{14}{33} \leftarrow \textcircled{1}$

① ↘

c)  $3\sqrt{3} - 2\sqrt{3} = \sqrt{3} \leftarrow \textcircled{1}$

d)  $(3\sqrt{5})^2 - 3(3\sqrt{5}) + 1$

$= 9 \times 5 - 9\sqrt{5} + 1 \quad \textcircled{1} \text{ for substitution}$

$= 46 - 9\sqrt{5} \leftarrow \textcircled{1}$

② a) (i)  $3x - 12x + 4x^2$   
 $= 4x^2 - 9x \quad \textcircled{1}$

(ii)  $16a^6b^2 \times \frac{a^2}{6b^3}$   
 $\textcircled{1}$

$= \frac{8a^8}{3b} \leftarrow \textcircled{1}$

b)  $x^{\frac{1}{2}} \div x^{-1} = x^{\frac{3}{2}} \quad \textcircled{1}$

c)  $(2^3)^n \times (2^2)^{n+1} \leftarrow \textcircled{1}$

$= 2^{3n} \times 2^{2n+2}$

③ a) (i)  $(x-6)(x+1) \quad \textcircled{1}$

(ii)  $(x-2)(x^2+2x+4) \quad \textcircled{1}$

(iii)  $x^2(x+1) + 1(x+1)$

$= (x+1)(x^2+1) \quad \textcircled{1}$

b) (i)  $\frac{x(\cancel{x-1})}{(x+1)(\cancel{x-1})} = \frac{x}{x+1} \quad \textcircled{1}$

(iii)  $\frac{4x}{x(x+2)} - \frac{1(x+2)}{x(x+2)} \quad \textcircled{1}$

$= \frac{4x - x - 2}{x(x+2)}$

$= \frac{3x - 2}{x(x+2)} \quad \textcircled{1}$

④ a)  $3 - 16 = -13 \quad \textcircled{1}$

b) (i)  $2x - 4 = 3x + 6 \quad \textcircled{1}$

$-x = 10$

$x = -10 \quad \textcircled{1}$

(ii)  $(2x-1)(x+5) = 0 \quad \textcircled{1}$

$\therefore x = \frac{1}{2}, -5 \quad \textcircled{1}$

(iii)  $1 - 2x = 6 \text{ or } -(1 - 2x) = 6 \quad \textcircled{1}$

$-2x = 5 \text{ or } -1 + 2x = 6$

$x = -2\frac{1}{2}$

$2x = 7$

5 a)  $3x - y = 0$  — (1)  
 $4x - 2y = 3$  — (2)

(1)  $\times -2$ :  $-6x + 2y = 0$  1 mark for clear method

(2) + (3):  $-2x = 3$

$x = -1\frac{1}{2}$  1 mark

Sub in (1):  $-4\frac{1}{2} - y = 0$

$y = -4\frac{1}{2}$  1 mark

b)  $4 + x(x+2) = 3(x+2)$

$4 + x^2 + 2x = 3x + 6$

$x^2 - x - 2 = 0$  ← (1)

$(x-2)(x+1) = 0$

$x = 2$  or  $-1$  ← (1)

c) (i)  $f(-3) = 27 + 6$

$= 33$  ← (1)

(ii)  $3(x+h)^2 - 2(x+h) + 3x^2 - 2x$  ← (1)


$= 3x^2 + 6xh + 3h^2 - 2x - 2h + 3x^2 - 2x$

$= 6x^2 + 6xh + 3h^2 - 4x - 2h$


← (1)

(6) a) (i)


must show marked info




(ii)



(iii)



b) (i)  $y = (x-1)(x+3)$



(ii)  $-3 \leq x \leq 1$  (1)

(iii) min. value when  $x = -1$   
 $\therefore y = -4$

$\therefore$  range is  $y \geq -4$

⑦ a)  $x - 4 \geq 0$

$\therefore D: x \geq 4$  ①


$R: y \geq 0$  ①

b)  $(x-2)^2 + y^2 + 6y + 9 = 9 \leftarrow$  ①

$(x-2)^2 + (y+3)^2 = 9$

$\therefore$  centre at  $(2, -3)$   $\leftarrow$  ①

c) (i)



(ii)  $x < 0$  ①

(iii) above ①

⑧ a)  $x = \angle B + \angle C$  (exterior angle of  $\triangle OBC$ )

$\triangle OBC$  is isosceles ( $OC = OB$ , equal radii)

$\therefore \angle B = \angle C$  (base angles)

$\therefore \angle B = \angle C = \frac{x}{2}$

$\leftarrow$  1 mark deducted for each error of reasoning

b)  $\angle D = 108^\circ$  (angle sum for reg. pentagon is  $540^\circ$ )

$\therefore \angle F = 72^\circ$  (co-interior angles,  $AF \parallel CD$ )

Similarly,  $\angle DEA = 108^\circ$

Now,  $\angle DEA = \angle F + \angle FAE$  (exterior angle rule for  $\triangle FAE$ )

$\therefore \angle FAE = 108^\circ - 72^\circ$   
 $= 36^\circ$