

IAO 2011


QUESTION PAPERS



THEORY

QUESTION PAPERS





XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

| | |
|----------|-----------------------|
| ЯЗЫК | <i>English</i> |
| language | |

Theoretical round. Problems to solve

General note. Maybe not all problems have correct questions. Some questions (maybe the main question of the problem, maybe one of the subquestions) may make no real sense. In this case you have to write in your answer (in English or Russian): «**impossible situation** – **ситуация невозможна**». Of course, this answer has to be explained numerically or logically.

Data from the tables (Planetary data, stars, constants, etc.) may be used for solving every problem.


The answers «Да-Yes» or «Нет-No» has to be written in English or Russian.

- 1. Observation of a star.** Observations were done by the naked eye on June 16, 2008, Universal time was used. An observer has registered that a star passed zenith at $0^{\text{h}}18^{\text{m}}$, and at $8^{\text{h}}17^{\text{m}}$ its height above the horizon was $87^{\circ}12'$. Find the latitude of the observations.
- 2. Planetarium.** Classical devices "planetaria" are arranged so that each group of stars is projected on a dome by a small optical system. Foils with hole-stars of the corresponding sizes are often used as these "slides" of constellations which are projected on the dome, so that most of the light is blocked by the foil (the black sky appears), and only light through the holes are transmitted (so stars appear). For example, images of 0^{m} stars on the foil have the size $l_0 = 0.1 \text{ mm}$, and stars up to 6^{m} are demonstrated, the focal length of the projecting system is $f = 25 \text{ cm}$, and the device has 16 separate projecting systems for every hemisphere. The dome of the planetarium of the observatory "Bobek" has a diameter of $2R = 10 \text{ m}$.
Let's suppose that all slides have been removed for replacement by more perfect ones, and the whole light began to be projected on the dome. What would the total stellar magnitude of the illuminated dome be (the artificial gray sky)? Would it be possible to read a newspaper in such an illuminance?
The answer has to include a list of the necessary parameters with formulae and numerical values.
- 3. Sunrise on Mars.** The Polar Bear (whom was already met in the texts of many International Astronomy Olympiads) was tired to make astronomical observations on Earth. He made a fascinating journey to the North Pole of Mars, and decided to observe a sunrise there. Calculate how long this sunrise lasts. The solution has to include a picture with an image of the Bear on the North Pole of Mars. Necessary sizes or angular sizes should be in the picture. Assume that Mars is spherical and its orbit may be considered circular. Recollect for yourself the necessary information about the Polar Bear.
- 4. Photo of Jupiter.** In the photo of Jupiter that was taken on October 19, 2009, one of the Galilean moons and its shadow on the disc of the planet can be seen. Jupiter was near the middle of Capricornus constellation when the photo was taken.
Find the orbital distance from the moon to the surface of the planet. Determine the name of the moon. The solution should be illustrated by drawings. The name of the moon in the solution and in the drawings should be written (or duplicated) in English.
- 5. Jupiter disappeared.** Let us suppose that Jupiter suddenly disappeared. The moons of Jupiter became independent bodies.

5.1. Which former Galileo moon(s) and in what case may leave the Solar system?

5.2. Which former Galileo moon(s) and in what case may fall into the Sun?

The answers "which moon" and "in what case" (configurations at the moment of Jupiter disappearance) have to be given in the form of drawings, and calculations should be the base for the answers. The names of the moons in the solution and in the drawings should be written in English. Consider the orbit of Jupiter (before its disappearance) to be circular.



XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

Казахстан, Алма-Ата

22 – 30. IX. 2011

Almaty, Kazakhstan

| | |
|----------|-----------------------|
| ЯЗЫК | <i><u>Русский</u></i> |
| language | |
| ЯЗЫК | <i><u>English</u></i> |
| language | |


Задачи теоретического тура. Рисунок

Theoretical round. Problems to solve. Picture

4. Снимок Юпитера.

4. Photo of Jupiter.





XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

Казахстан, Алма-Ата

22 – 30. IX. 2011

Almaty, Kazakhstan

Groups

α

β

| | |
|----------|-----------------------|
| ЯЗЫК | <i>Русский</i> |
| language | |
| ЯЗЫК | <i>English</i> |
| language | |


Элементы орбит. Физические характеристики Солнца, некоторых планет,
Луны и Галилеевых спутников Юпитера

Parameters of orbits. Physical characteristics of Sun, some planets,
Moon and Galilean moons of Jupiter

| Небесное тело, планета | Среднее расстояние от центрального тела | | Сидерический (или аналогичный) период обращения | | Экс- цен- три- тет, <i>e</i> | Эквато- риальн. диаметр <i>км</i> | Масса <i>10²⁴ кг</i> | Сред- няя плот- ность <i>г/см³</i> | Ускор. свооб. пад. у пов. <i>м/с²</i> | На- клон оси | Аль- bedo |
|------------------------------|---|--------------------------|--|------------------------|--|--|--|---|--|--------------------|--------------|
| | в <i>астр. ед.</i> | в <i>млн. км</i> | в тропич. годах | в средних сутках | | | | | | | |
| Body, planet | Average distance to central body in <i>astr. units</i> | in <i>mln. km</i> | Sidereal (or analogous) period in troph. years | in days | Ec- centri- city <i>e</i> | Equat. diameter <i>km</i> | Mass <i>10²⁴ kg</i> | Av. den- sity <i>g/cm³</i> | Grav. acceler. at surf. <i>m/s²</i> | Axial tilt | Al- bedo |
| Солнце Sun | 1,6·10 ⁹ | 2,5·10 ¹¹ | 2,2·10 ⁸ | 8·10 ¹⁰ | | 1392000 | 1989000 | 1,409 | | | |
| Меркурий Mercury | 0,387 | 57,9 | 0,241 | 87,97 | 0,206 | 4 879 | 0,3302 | 5,43 | 3,70 | 0,01° | 0,06 |
| Венера Venus | 0,723 | 108,2 | 0,615 | 224,70 | 0,007 | 12 104 | 4,8690 | 5,24 | 8,87 | 177,36 | 0,78 |
| Земля Earth | 1,000 | 149,6 | 1,000 | 365,26 | 0,017 | 12 756 | 5,9742 | 5,515 | 9,81 | 23,44 | 0,36 |
| Луна Moon | 0,00257 | 0,38440 | 0,0748 | 27,3217 | 0,055 | 3 475 | 0,0735 | 3,34 | 1,62 | 6,7 | 0,07 |
| Марс Mars | 1,524 | 227,9 | 1,880 | 686,98 | 0,093 | 6 794 | 0,6419 | 3,94 | 3,71 | 25,19 | 0,15 |
| Юпитер Jupiter | 5,204 | 778,6 | 11,862 | 4 332,59 | 0,048 | 142 984 | 1899,8 | 1,33 | 24,86 | 3,13 | 0,66 |
| Сатурн Saturn | 9,584 | 1433,7 | 29,458 | 10 759,20 | 0,054 | 120 536 | 568,50 | 0,70 | 10,41 | 26,73 | 0,68 |

| Спутник | Среднее расстояние от центра планеты | | Сидерич. период обращения в средних сутках | Накло- нение орбиты (°) | Экс- цен- три- тет, <i>e</i> | Диаметр <i>км</i> | Масса <i>10²¹ кг</i> | Сред- няя плот- ность <i>г/см³</i> | Ускор. свооб. пад. у пов. <i>м/с²</i> | Макс. блеск. вид. с Земли (*) | Геом Аль- bedo |
|----------------------|--|----------------------------|---|----------------------------------|--|---------------------------|--|---|--|---|----------------------|
| | в <i>астр. ед.</i> | в <i>тыс. км</i> | | | | | | | | | |
| Moon | Average distance to planet center in <i>astr. units</i> | in <i>thous. km</i> | Sidereal period in days | Orbit inclination (°) | Ec- centri- city <i>e</i> | Diameter <i>km</i> | Mass <i>10²¹ kg</i> | Av. den- sity <i>g/cm³</i> | Grav. acceler. at surf. <i>m/s²</i> | Max. magn. From Earth (*) | Geom Al- bedo |
| Ио Io | 0,00282 | 421,70 | 1,769137 | 0,050° | 0,0041 | 3 643 | 89,31 | 3,53 | 1,80 | 5,02 ^m | 0,63 |
| Европа Europa | 0,00449 | 671,03 | 3,551181 | 0,471° | 0,0094 | 3 122 | 48,00 | 3,01 | 1,32 | 5,29 ^m | 0,67 |
| Ганимед Ganymede | 0,00716 | 1070,41 | 7,154553 | 0,204° | 0,0011 | 5 262 | 148,19 | 1,94 | 1,43 | 4,61 ^m | 0,43 |
| Каллисто Callisto | 0,01259 | 1882,71 | 16,689018 | 0,205° | 0,0074 | 4 821 | 107,59 | 1,83 | 1,23 | 5,65 ^m | 0,20 |

*) В среднем противостоянии. *) In mean opposition.



XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

Казахстан, Алма-Ата

22 – 30. IX. 2011


Almaty, Kazakhstan

| | |
|------------------|-----------------------|
| ЯЗЫК language | <i>Русский</i> |
| ЯЗЫК language | <i>English</i> |

Некоторые константы и формулы

Some constants and formulae

| | | |
|---|---|---|
| Скорость света в вакууме, c (м/с) | 299 792 458 | Speed of light in vacuum, c (m/s) |
| Гравитационная постоянная, G (Н·м ² /кг ²) | $6.674 \cdot 10^{-11}$ | Constant of gravitation, G (N·m ² /kg ²) |
| Солнечная постоянная, A (Вт/м ²) | 1367 | Solar constant, A (W/m ²) |
| Параметр Хаббла, H_0 (км/с/Мпк) | 71 среднее значение диапазон значений | mean value Hubble parameter, diapason of values H_0 (km/s/Mpc) |
| Постоянная Планка, h (Дж·с) | $6.626 \cdot 10^{-34}$ | Plank constant, h (J·s) |
| Заряд электрона, e (Кл) | $1.602 \cdot 10^{-19}$ | Charge of electron, e (C) |
| Масса электрона, m_e (кг) | $9.109 \cdot 10^{-31}$ | Mass of electron, m_e (kg) |
| Соотношение масс протона и электрона | 1836.15 | Proton-to-electron ratio |
| Постоянная Фарадея, F (Кл/моль) | 96 485 | Faraday constant, F (C/mol) |
| Магнитная постоянная, μ_0 (Гн/м) | $1.257 \cdot 10^{-6}$ | Magnetic constant, μ_0 (H/m) |
| Универсальная газовая постоянная, R (Дж/моль/К) | 8.314 | Universal gas constant, R (J/mol/K) |
| Постоянная Больцмана, k (Дж/К) | $1.381 \cdot 10^{-23}$ | Boltzmann constant, k (J/K) |
| Стандартная атмосфера (Па) | 101 325 | Standard atmosphere (Pa) |
| Постоянная Стефана-Больцмана, σ (Вт/м ² /К ⁴) | $5.670 \cdot 10^{-8}$ | Stefan-Boltzmann constant, σ (W/m ² /K ⁴) |
| Константа смещения Вина, b (м·К) | 0.002897 | Wien's displacement constant, b (m·K) |
| Лабораторная длина волны $H\alpha$ (Å) | 6562.81 | Laboratory wavelength of $H\alpha$ (Å) |
| Длина тропического года, T (сут) | 365.242199 | Tropical year length, T (days) |
| Показатель преломления воды при 20°C, n | 1.334 | Refractive index of water for 20°C, n |
| Момент инерции шара | $I = \frac{2}{5} MR^2$ | Moment of inertia of a solid ball |
| Площадь сферы | $S = 4\pi R^2$ | Area of sphere |
| π | 3.14159265 | π |
| e | 2.71828183 | e |
| Золотое сечение, ϕ | 1.61803399 | Golden ratio, ϕ |




Theoretical round. Problems to solve

General note. Maybe not all problems have correct questions. Some questions (maybe the main question of the problem, maybe one of the subquestions) may make no real sense. In this case you have to write in your answer (in English or Russian): «**impossible situation – ситуация невозможна**». Of course, this answer has to be explained numerically or logically.

Data from the tables (Planetary data, stars, constants, etc.) may be used for solving every problem.

The answers «Да-Yes» or «Нет-No» has to be written in English or Russian.

- Solar radiation.** Find, with how many percents the mass of the Sun is diminishing per year due to its radiation.
- Planetarium.** Classical devices "planetaria" are arranged so that each group of stars is projected on a dome by a small optical system. Consider the planetarium of observatory "Bobek" which diameter of the hall (dome) is $2R = 10$ m. On slides of the constellations projected on the dome images of 0^m stars have the size $l_0 = 0.1$ mm (foils with hole-stars of the mentioned size are often used as these "slides").
 - Estimate what the parameters should be (decide yourself, which parameters are important here) of the objective of this optical system, so that the visitors sitting in the centre of the hall of the planetarium would perceive the "stars" as points (not as circles or nebulae).
 - Let's suppose that all slides have been removed for replacement by more perfect ones, and all the light began to be projected on the dome. What would the total stellar magnitude of the illuminated dome be (the artificial gray sky)? Would it be possible to read a newspaper in such an illuminance?
The answers has to include a list of the necessary parameters with formulae and numerical values.
- Sunrise on Mars.** The Polar Bear (whom was already met in the texts of many International Astronomy Olympiads) was tired to make astronomical observations on Earth. He made a fascinating journey to the North Pole of Mars, and decided to observe a sunrise there. Calculate how long this sunrise lasts. The solution has to include a picture with an image of the Bear on the North Pole of Mars. Necessary sizes or angular sizes should be in the picture. Assume that Mars is spherical and its orbit may be considered circular. Recollect for yourself the necessary information about the Polar Bear.
- Photo of Jupiter.** In the photo of Jupiter that was taken on October 19, 2009, one of the Galilean moons and its shadow on the disc of the planet can be seen. Jupiter was near the middle of Capricornus constellation when the photo was taken.
Find the orbital distance from the moon to the surface of the planet. Determine the name of the moon. The solution should be illustrated by drawings. The name of the moon in the solution and in the drawings should be written (or duplicated) in English.
- Jupiter disappeared.** Let us suppose that Jupiter suddenly disappeared. The moons of Jupiter became independent bodies.
 - Which former Galileo moon(s) and in what case may leave the Solar system?
 - Which former Galileo moon(s) and in what case may fall into the Sun?
 The answers "which moon" and "in what case" (configurations at the moment of Jupiter disappearance) have to be given in the form of drawings, and calculations should be the base for the answers. The names of the moons in the solution and in the drawings should be written in English. Consider the orbit of Jupiter (before its disappearance) to be circular.



XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

Казахстан, Алма-Ата

22 – 30. IX. 2011

Almaty, Kazakhstan

| | |
|----------|-----------------------|
| ЯЗЫК | <i><u>Русский</u></i> |
| language | |
| ЯЗЫК | <i><u>English</u></i> |
| language | |


Задачи теоретического тура. Рисунок

Theoretical round. Problems to solve. Picture

4. Снимок Юпитера.

4. Photo of Jupiter.





XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

Казахстан, Алма-Ата

22 – 30. IX. 2011

Almaty, Kazakhstan

Groups

α

β

| | |
|----------|-----------------------|
| ЯЗЫК | <i>Русский</i> |
| language | |
| ЯЗЫК | <i>English</i> |
| language | |


Элементы орбит. Физические характеристики Солнца, некоторых планет, Луны и Галилеевых спутников Юпитера

Parameters of orbits. Physical characteristics of Sun, some planets, Moon and Galilean moons of Jupiter

| Небесное тело, планета | Среднее расстояние от центрального тела | | Сидерический (или аналогичный) период обращения | | Эксцентриситет, e | Экваториальный диаметр, км | Масса, 10^{24} кг | Средняя плотность, г/см ³ | Ускор. своб. пад. у пов., м/с ² | Наклон оси | Альбедо |
|------------------------|---|---------------------|---|-------------------|---------------------|----------------------------|---------------------|--------------------------------------|--|------------|---------|
| | в астр. ед. | в млн. км | в тропич. годах | в средних сутках | | | | | | | |
| Body, planet | Average distance to central body in astr. units | in mln. km | Sidereal (or analogous) period in troph. years | in days | Ec-centri-city e | Equat. diameter km | Mass 10^{24} kg | Av. den-sity g/cm ³ | Grav. acceler. at surf. m/s ² | Axial tilt | Al-bedo |
| Солнце Sun | $1,6 \cdot 10^9$ | $2,5 \cdot 10^{11}$ | $2,2 \cdot 10^8$ | $8 \cdot 10^{10}$ | | 1392000 | 1989000 | 1,409 | | | |
| Меркурий Mercury | 0,387 | 57,9 | 0,241 | 87,97 | 0,206 | 4 879 | 0,3302 | 5,43 | 3,70 | 0,01° | 0,06 |
| Венера Venus | 0,723 | 108,2 | 0,615 | 224,70 | 0,007 | 12 104 | 4,8690 | 5,24 | 8,87 | 177,36 | 0,78 |
| Земля Earth | 1,000 | 149,6 | 1,000 | 365,26 | 0,017 | 12 756 | 5,9742 | 5,515 | 9,81 | 23,44 | 0,36 |
| Луна Moon | 0,00257 | 0,38440 | 0,0748 | 27,3217 | 0,055 | 3 475 | 0,0735 | 3,34 | 1,62 | 6,7 | 0,07 |
| Марс Mars | 1,524 | 227,9 | 1,880 | 686,98 | 0,093 | 6 794 | 0,6419 | 3,94 | 3,71 | 25,19 | 0,15 |
| Юпитер Jupiter | 5,204 | 778,6 | 11,862 | 4 332,59 | 0,048 | 142 984 | 1899,8 | 1,33 | 24,86 | 3,13 | 0,66 |
| Сатурн Saturn | 9,584 | 1433,7 | 29,458 | 10 759,20 | 0,054 | 120 536 | 568,50 | 0,70 | 10,41 | 26,73 | 0,68 |

| Спутник | Среднее расстояние от центра планеты | | Сидерич. период обращения в средних сутках | Накло-нение орбиты (°) | Экс-центриситет, e | Диаметр, км | Масса, 10^{21} кг | Сред-няя плотность, г/см ³ | Ускор. своб. пад. у пов., м/с ² | Макс. блеск, вид. с Земли *) | Геом Альбедо |
|-------------------|--|--------------|--|------------------------|----------------------|-------------|---------------------|---------------------------------------|--|------------------------------|--------------|
| | в астр. ед. | в тыс. км | | | | | | | | | |
| Moon | Average distance to planet center in astr. units | in thous. km | Sidereal period in days | Orbit inclination (°) | Ec-centri-city e | Diameter km | Mass 10^{21} kg | Av. den-sity g/cm ³ | Grav. acceler. at surf. m/s ² | Max. magn. From Earth *) | Geom Al-bedo |
| Ио Io | 0,00282 | 421,70 | 1,769137 | 0,050° | 0,0041 | 3 643 | 89,31 | 3,53 | 1,80 | 5,02 ^m | 0,63 |
| Европа Europa | 0,00449 | 671,03 | 3,551181 | 0,471° | 0,0094 | 3 122 | 48,00 | 3,01 | 1,32 | 5,29 ^m | 0,67 |
| Ганимед Ganymede | 0,00716 | 1070,41 | 7,154553 | 0,204° | 0,0011 | 5 262 | 148,19 | 1,94 | 1,43 | 4,61 ^m | 0,43 |
| Каллисто Callisto | 0,01259 | 1882,71 | 16,689018 | 0,205° | 0,0074 | 4 821 | 107,59 | 1,83 | 1,23 | 5,65 ^m | 0,20 |

*) В среднем противостоянии. *) In mean opposition.



XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

Казахстан, Алма-Ата

22 – 30. IX. 2011

Almaty, Kazakhstan

| | |
|------------------|-----------------------|
| ЯЗЫК language | <i>Русский</i> |
| ЯЗЫК language | <i>English</i> |

Некоторые константы и формулы


Some constants and formulae

| | | |
|---|---|---|
| Скорость света в вакууме, c (м/с) | 299 792 458 | Speed of light in vacuum, c (m/s) |
| Гравитационная постоянная, G (Н·м ² /кг ²) | $6.674 \cdot 10^{-11}$ | Constant of gravitation, G (N·m ² /kg ²) |
| Солнечная постоянная, A (Вт/м ²) | 1367 | Solar constant, A (W/m ²) |
| Параметр Хаббла, H_0 (км/с/Мпк) | 71 среднее значение диапазон значений | mean value Hubble parameter, diapason of values H_0 (km/s/Mpc) |
| Постоянная Планка, h (Дж·с) | $6.626 \cdot 10^{-34}$ | Plank constant, h (J·s) |
| Заряд электрона, e (Кл) | $1.602 \cdot 10^{-19}$ | Charge of electron, e (C) |
| Масса электрона, m_e (кг) | $9.109 \cdot 10^{-31}$ | Mass of electron, m_e (kg) |
| Соотношение масс протона и электрона | 1836.15 | Proton-to-electron ratio |
| Постоянная Фарадея, F (Кл/моль) | 96 485 | Faraday constant, F (C/mol) |
| Магнитная постоянная, μ_0 (Гн/м) | $1.257 \cdot 10^{-6}$ | Magnetic constant, μ_0 (H/m) |
| Универсальная газовая постоянная, R (Дж/моль/К) | 8.314 | Universal gas constant, R (J/mol/K) |
| Постоянная Больцмана, k (Дж/К) | $1.381 \cdot 10^{-23}$ | Boltzmann constant, k (J/K) |
| Стандартная атмосфера (Па) | 101 325 | Standard atmosphere (Pa) |
| Постоянная Стефана-Больцмана, σ (Вт/м ² /К ⁴) | $5.670 \cdot 10^{-8}$ | Stefan-Boltzmann constant, σ (W/m ² /K ⁴) |
| Константа смещения Вина, b (м·К) | 0.002897 | Wien's displacement constant, b (m·K) |
| Лабораторная длина волны $H\alpha$ (Å) | 6562.81 | Laboratory wavelength of $H\alpha$ (Å) |
| Длина тропического года, T (сут) | 365.242199 | Tropical year length, T (days) |
| Показатель преломления воды при 20°C, n | 1.334 | Refractive index of water for 20°C, n |
| Момент инерции шара | $I = \frac{2}{5} MR^2$ | Moment of inertia of a solid ball |
| Площадь сферы | $S = 4\pi R^2$ | Area of sphere |
| π | 3.14159265 | π |
| e | 2.71828183 | e |
| Золотое сечение, ϕ | 1.61803399 | Golden ratio, ϕ |

OBSERVATION

QUESTION PAPERS





XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

Казахстан, Алма-Ата

22 – 30. IX. 2011

Almaty, Kazakhstan

Groups

α

β

| | |
|------------------|----------------|
| ЯЗЫК language | <u>Русский</u> |
| ЯЗЫК language | <u>English</u> |

Вопросы наблюдательного тура

Observational round. Questions

| <u>Русский</u> | <u>English</u> |
|---|--|
| В Вашем распоряжении фонарик и часы | You are provided by a light and a watch |
| <ol style="list-style-type: none"> 1. Установите Юпитер в центре поля зрения телескопа. Покажите результат экзаменатору. Зарисуйте расположение спутников Юпитера на момент наблюдения. 2. Определите и напишите на листке часовой угол и горизонтальные координаты α Peg. 3. Зарисуйте и пронумеруйте в порядке убывания яркости (1, 2, ...) восемь основных звёзд созвездия "Cygnus". 4. Определите поле зрения телескопа, стоящего перед Вами. Фокусное расстояние окуляра $F = 10$ мм. | <ol style="list-style-type: none"> 1. Bring Jupiter to the centre of field of view of the telescope. Demonstrate the result to the examiner. Draw a picture with the positions of the moons of Jupiter at the moment of observation. 2. Calculate and write down on the paper the hour angle and horizontal coordinates of α Peg. 3. Draw and enumerate in descending order by brightness (1, 2, ...) the eight main stars of the "Cygnus" constellation. 4. Calculate the field of view of the telescope in front of you. The focal length of the eyepiece is $F = 10$ mm. |
| Максимальное время выполнения задания – 20 минут . | The maximum total time for all tasks is 20 minutes . |




Медведи желают Вам успехов!
The Bears wish you success!



PRACTICAL

QUESTION PAPERS





XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

Казахстан, Алма-Ата

22 – 30. IX. 2011

Almaty, Kazakhstan



ЯЗЫК
language

English

Practical round. Problems to solve

6. Saturn 2010.

САТУРН 2010




You are provided with a series of Saturn images. Find the rotational period for Saturn, and the velocity and the direction of the movement of unique formation in the atmosphere of Saturn – Great Northern Disturbance (GND), which appeared on December 8, 2010.

- 6.1. Use measurements of two images on Saturn on 13 December 2010 to determine the rotational period T for Saturn from the angular change of the GND position during the time interval between two images.
- 6.2. Plot from all measurements of the GND core a graph showing how the core longitude L depends on time t . The time should be expressed in hours and parts of hours. Set the time of the first image as zero time (0.0^h). The inclination of the graph must show the direction and the angular velocity of GND.
- 6.3. Determine the value of radius $R=(a+b)/2$ at the GND latitude in kilometers and determine the linear velocity of the GMD core latitude in kilometers using the graph that shows L vs t .

You may neglect the tilt of the equator to the line of sight (the direction Earth-Saturn). In the images north is up. Saturn's equatorial radius $R_0 = 60266$ km. The planet is rotating counterclockwise if you look from the north pole, or from left to right in the images. The longitude of the central meridian of the planet increases following the planet's rotation. It means that L increases from right to left in the images.

Data on the images: date (2010, December, day), time UT (h, m), and LCM, the longitude of the central meridian (in degrees) in system of coordinates of planet.

On the above drawing you can see the scheme (a view from the North), R is the radius on the GND latitude. Evidently $R < R_0$.



XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

Казахстан, Алма-Ата

22 – 30. IX. 2011

Almaty, Kazakhstan


| | |
|------------------|-----------------------|
| ЯЗЫК language | <u><i>Русский</i></u> |
| ЯЗЫК language | <u><i>English</i></u> |

6. Сатурн 2010.

6. Saturn 2010.







XVI Международная астрономическая олимпиада

XVI International Astronomy Olympiad

Казахстан, Алма-Ата

22 – 30. IX. 2011

Almaty, Kazakhstan

| |
|----------|
| ЯЗЫК |
| language |

English

Practical round. Problems to solve

7. Exoplanet.

Recently astronomers discovered an exoplanet that practically reflects no light and belongs to the class of hot Jupiters. Its mass corresponds to 12 Jupiter's masses, the average orbit radius is 0.036 AU and the period is 2.47 days. The given data about the exoplanet has been obtained with the help of the data from the "Kepler" telescope which continuously monitors the sky region between the constellations Cygnus and Lyra. The device was launched in March 2009 and is capable of seeing not only gas giants, but also planets with the size of Earth. Recently scientists have discovered that exoplanets are capable of causing huge tidal waves in the stars that have a significant influence on the spectral parameters of the emitted light from the star. In fact, with the help of this effect scientists have been able to prove that the planet moves around a star in a circular orbit.

In the graph (see separate sheet) the flux of radiation from the system vs. the phase (phase ϕ given in radians) can be seen, normalized to the radiation flux from the star.

7.1. During the transit there are four "contacts" when the contour of the smaller object touches the contour of the larger object at one point. The contacts take place in the following order:

- First contact: The smaller body is completely outside of the larger and moving inside.
- Second contact: The smaller body is completely inside of the larger and continues to move inwards.
- Third contact: The smaller body is completely inside of the larger and moves outwards.
- Fourth contact: The smaller body is completely outside of the larger and starts to move away.


In the given graph mark the points of the first, second, third and the fourth contacts. Write in the copybook the value of the phase for each contact.

7.2. Determine the duration of the transit (duration of the passage of the disk of planet over the disk of the star).

7.3. Determine the inclination of the orbit (for exoplanets and binary stars the reference plane is assumed to be the "image plane" – the plane that is perpendicular to the line of sight).

7.4. Calculate the radius of the star.

7.5. Calculate the radius of the planet.



XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

Казахстан, Алма-Ата


22 – 30. IX. 2011


Almaty, Kazakhstan

| | |
|----------|-----------------------|
| ЯЗЫК | <u><i>Русский</i></u> |
| language | |
| ЯЗЫК | <u><i>English</i></u> |
| language | |

7. Экзопланета.

7. Exoplanet.





XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

Казахстан, Алма-Ата

22 – 30. IX. 2011

Almaty, Kazakhstan

ЯЗЫК
language


English

Practical round. Problems to solve

6. Planetary nebula.

The table provides the results of the measurements of the wavelength of the spectral line of hydrogen with $\lambda_0 = 6562,81\text{\AA}$ ($H\alpha$) of the planetary nebula NGC 5643 (points shown in the picture on separate sheet). Draw a graph that shows the dependence of the radial velocity of the points on the declination δ . Draw a diagram showing the direction in which each point is moving. Assuming that the extreme points in the table correspond to the boundary of the nebula, calculate the motion velocity (the velocity at which the nebula as a whole approaches or recedes) and the velocity of the nebula expansion.

| | | | | | | | |
|-----------------------|----------|----------|----------|----------|----------|----------|----------|
| δ | +66° 34' | +66° 35' | +66° 36' | +66° 37' | +66° 38' | +66° 39' | +66° 40' |
| $\lambda, \text{\AA}$ | 6563,17 | 6562,80 | 6562,68 | 6562,40 | 6562,65 | 6562,77 | 6563,16 |



XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

Казахстан, Алма-Ата


22 – 30. IX. 2011


Almaty, Kazakhstan

| | |
|----------|-----------------------|
| ЯЗЫК | <i><u>Русский</u></i> |
| language | |
| ЯЗЫК | <i><u>English</u></i> |
| language | |

6. Планетарная туманность.

6. Planetary nebula.





XVI Международная астрономическая олимпиада

XVI International Astronomy Olympiad

Казахстан, Алма-Ата

22 – 30. IX. 2011

Almaty, Kazakhstan

| |
|----------|
| ЯЗЫК |
| language |

English

Practical round. Problems to solve

7. Exoplanet.

Recently astronomers discovered an exoplanet that practically reflects no light and belongs to the class of hot Jupiters. Its mass corresponds to 12 Jupiter's masses, the average orbit radius is 0.036 AU and the period is 2.47 days. The given data about the exoplanet has been obtained with the help of the data from the "Kepler" telescope which continuously monitors the sky region between the constellations Cygnus and Lyra. The device was launched in March 2009 and is capable of seeing not only gas giants, but also planets with the size of Earth. Recently scientists have discovered that exoplanets are capable of causing huge tidal waves in the stars that have a significant influence on the spectral parameters of the emitted light from the star. In fact, with the help of this effect scientists have been able to prove that the planet moves around a star in a circular orbit.

In the graph (see separate sheet) the flux of radiation from the system vs. the phase (phase ϕ given in radians) can be seen, normalized to the radiation flux from the star.

7.1. During the transit there are four "contacts" when the contour of the smaller object touches the contour of the larger object at one point. The contacts take place in the following order:

- First contact: The smaller body is completely outside of the larger and moving inside.
- Second contact: The smaller body is completely inside of the larger and continues to move inwards.
- Third contact: The smaller body is completely inside of the larger and moves outwards.
- Fourth contact: The smaller body is completely outside of the larger and starts to move away.


In the given graph mark the points of the first, second, third and the fourth contacts. Write in the copybook the value of the phase for each contact.

7.2. Determine the duration of the transit (duration of the passage of the disk of planet over the disk of the star).

7.3. Determine the inclination of the orbit (for exoplanets and binary stars the reference plane is assumed to be the "image plane" – the plane that is perpendicular to the line of sight).

7.4. Calculate the radius of the star.

7.5. Calculate the radius of the planet.



XVI Международная астрономическая олимпиада
XVI International Astronomy Olympiad

Казахстан, Алма-Ата

22 – 30. IX. 2011

Almaty, Kazakhstan

| | |
|----------|-----------------------|
| ЯЗЫК | <i><u>Русский</u></i> |
| language | |
| ЯЗЫК | <i><u>English</u></i> |
| language | |

7. Экзопланета.

7. Exoplanet.

