

Two-Year Randomized Controlled Trial of Vitamin K₁ (Phylloquinone) and Vitamin D₃ Plus Calcium on the Bone Health of Older Women

Caroline Bolton-Smith,¹ Marion ET McMurdo,² Colin R Paterson,³ Patricia A Mole,² Julia M Harvey,⁴
Steven T Fenton,¹ Celia J Prynne,⁵ Gita D Mishra,⁵ and Martin J Shearer⁴

ABSTRACT: Dietary supplementation with vitamin K₁, with vitamin D₃ and calcium or their combination, was examined in healthy older women during a 2-year, double-blind, placebo-controlled trial. Combined vitamin K with vitamin D plus calcium was associated with a modest but significant increase in BMC at the ultradistal radius but not at other sites in the hip or radius.

Introduction: The putative beneficial role of high dietary vitamin K₁ (phylloquinone) on BMD and the possibility of interactive benefits with vitamin D were studied in a 2-year double-blind, placebo-controlled trial in healthy Scottish women ≥ 60 years of age.

Materials and Methods: Healthy, nonosteoporotic women ($n = 244$) were randomized to receive either (1) placebo, (2) 200 $\mu\text{g/day}$ vitamin K₁, (3) 10 μg (400 IU) vitamin D₃ plus 1000 mg calcium/day, or (4) combined vitamins K₁ and D₃ plus calcium. Baseline and 6-month measurements included DXA bone mineral scans of the hip and wrist, markers of bone turnover, and vitamin status. Supplementation effects were tested using multivariate general linear modeling, with full adjustment for baseline and potential confounding variables.

Results: Significant bone mineral loss was seen only at the mid-distal radius but with no significant difference between groups. However, women who took combined vitamin K and vitamin D plus calcium showed a significant and sustained increase in both BMD and BMC at the site of the ultradistal radius. Serum status indicators responded significantly to respective supplementation with vitamins K and D. Over 2 years, serum vitamin K₁ increased by 157% ($p < 0.001$), the percentage of undercarboxylated osteocalcin (%GluOC) decreased by 51% ($p < 0.001$), serum 25-hydroxyvitamin D [25(OH)D] increased by 17% ($p < 0.001$), and PTH decreased by 11% ($p = 0.049$).

Conclusions: These results provide evidence of a modest synergy in healthy older women from nutritionally relevant intakes of vitamin K₁ together with supplements of calcium plus moderate vitamin D₃ to enhance BMC at the ultradistal radius, a site consisting of principally trabecular bone. The substantial increase in γ -carboxylation of osteocalcin by vitamin K may have long-term benefits and is potentially achievable by increased dietary intakes of vitamin K rather than by supplementation.

J Bone Miner Res 2007;22:509–519. Published online on January 22, 2007; doi: 10.1359/JBMR.070116

Key words: vitamin K, vitamin D, osteoporosis, bone densitometry, bone turnover markers, osteocalcin, undercarboxylated osteocalcin

INTRODUCTION

POPULATION-BASED PREVENTATIVE MEASURES that reduce the rate of bone mineral loss and therefore extend years of active, good-quality life offer a potential means of improving bone health. Dietary change is one potentially achievable and sustainable approach. Undernutrition is often observed in older people and is likely to be a significant

contributing risk factor to the pathogenesis and outcome of hip fracture in the elderly.^(1,2) Dietary intervention strategies have focused mainly on calcium and vitamin D; supplementation with either or both of these nutrients having been shown in some studies to be effective in maintaining BMD or in reducing the incidence of fractures.^(3–6) Not all studies have shown these benefits,⁽⁷⁾ including three recent fracture prevention trials carried out in the United Kingdom^(8,9) and the United States.⁽¹⁰⁾ Another nutrient that is being increasingly linked to bone health and osteoporosis is vitamin K.^(2,11–13)

The authors state that they have no conflicts of interest.

¹Nutrition Research Group, CVEU, University of Dundee, Ninewells Hospital and Medical School, Dundee, Scotland, United Kingdom; ²Section of Ageing and Health, University of Dundee, Ninewells Hospital and Medical School, Dundee, Scotland, United Kingdom; ³Division of Medicine and Therapeutics, University of Dundee, Ninewells Hospital and Medical School, Dundee, Scotland, United Kingdom; ⁴Vitamin K Research Unit, Centre for Haemostasis and Thrombosis, St Thomas' Hospital, London, United Kingdom; ⁵Medical Research Council-Human Nutrition Research, Elsie Widdowson Laboratory, Cambridge, United Kingdom.




FIG. 1. Study design showing the four treatment arms and subject numbers recruited and retained.

Vitamin K is needed as a co-factor to convert certain peptide-bound glutamate (Glu) residues to γ -carboxyglutamate (Gla) residues in target proteins (Gla proteins) that are synthesized by a variety of tissues. Bone matrix contains several Gla proteins, of which osteocalcin is the most abundant.^(11,12) The synthesis of Gla residues imbues vitamin K-dependent proteins with structural integrity, metal binding characteristics, and functionality; in the case of osteocalcin, the Gla residues confer high affinity binding to bone hydroxyapatite.⁽¹⁴⁾ Although its precise function is unclear, osteocalcin seems to act as a regulator of bone remodeling and/or bone mineral maturation.⁽¹⁵⁾ A clear rationale for ensuring optimal dietary intakes of vitamin K is the evidence that a high circulating level of undercarboxylated osteocalcin (GluOC) is an independent risk predictor of bone fractures^(16–19) and of low BMD.^(20,21) In addition, short-term studies show that the γ -carboxylation status of osteocalcin responds to increased vitamin K intakes in the usual dietary range.^(22,23) The possibility that vitamin D itself may also influence the γ -carboxylation status of osteocalcin has been raised by evidence that the free Gla content of animal bones⁽²⁴⁾ and circulating GluOC in elderly women⁽¹⁶⁾ both reflect vitamin D status. This unsubstantiated effect of vitamin D on posttranslational γ -carboxylation is independent of the well-established role of 1,25-dihydroxyvitamin D as a potent enhancer of osteocalcin expression.⁽²⁵⁾

To date, the only published trials that have assessed the effects of vitamin K on postmenopausal bone loss have used supradietary daily doses of either 1 mg of vitamin K₁⁽²⁶⁾ or 45 mg of vitamin K₂ (menaquinone-4; menatetranone).^(27–29) Our placebo-controlled study was designed to establish the influence of an additional daily intake of 200 µg vitamin K₁ on the bone health of older women. This extra vitamin K₁ intake was equivalent to the 95th percentile of younger adult intake in Scotland⁽³⁰⁾ and equivalent to that contained in a 40- to 60-g portion of leafy green vegetables such as collards and spinach.⁽³¹⁾ This intake could reasonably be considered to be attainable by dietary modification rather than from supplementation. Supple-

mentation at a high dietary level is justified by evidence from population studies in the United States that low consumption of vitamin K in some groups is associated with elevated fracture risk^(32,33) or lower bone mass⁽³⁴⁾ and that an impaired vitamin K status is associated with low bone mass⁽³⁵⁾ or increased bone turnover.⁽³⁶⁾ Our trial was also designed to study the effects, and possible interaction with vitamin K, of vitamin D and calcium at additional daily intakes of 10 µg (400 IU) and 1000 mg, respectively. The vitamin D intake was as recommended for elderly people in the United Kingdom at the time this study was initiated.⁽³⁷⁾

MATERIALS AND METHODS

Participants and intervention protocol

Healthy women ≥ 60 years of age were recruited from the general population to provide approximately equal numbers in the age groups 60–64, 65–69, 70–74, and ≥ 75 years. Exclusion criteria were clinical osteoporosis or chronic disease (e.g., diabetes mellitus, cardiovascular disease, cancer, fat malabsorption syndromes), routine medication that interferes with vitamin K, vitamin D, or bone metabolism (notably warfarin and steroids), and consumption of nutrient supplements that provided in excess of 30 µg vitamin K, 10 µg (400 IU) vitamin D, or 500 mg calcium daily. Hoffmann-La Roche (Basel, Switzerland) provided the supplementation tablets. One set of these tablets was in a chewable form and contained either 1000 mg of elemental calcium as calcium carbonate plus 10 µg of vitamin D₃ or placebo ingredients in a matching form. A second set of smaller, oval-shaped tablets contained 200 µg of vitamin K₁ (or placebo); these were swallowed whole with a drink. The 244 participants were randomly assigned to one of the four study groups: (1) placebo, (2) vitamin K₁ 200 µg/day, (3) calcium 1000 mg plus vitamin D₃ 10 µg (400 IU)/day, or (4) combined supplementation with vitamin K₁ and vitamin D₃ plus calcium at the levels in groups 2 and 3 (Fig. 1). To ensure double-blinding, an independent statistician at Hoffmann-La Roche, who had no other connection to the study,

provided a computer-generated randomization list to the researchers. Each participant was asked to take two tablets every day with their breakfast. The Tayside Committee on Medical Research Ethics approved the study, and all participants gave written informed consent.

Baseline measurements were performed in the winter, and subsequent measurements were performed every 6 months for 2 years. On each occasion, participants visited between 8:30 and 9:30 a.m. after an overnight fast, and a venous blood sample was taken, spun, aliquoted, and frozen for storage at -70°C within 1 h of collection. Weight and height were measured. Smoking habits, sunshine exposure, adherence to protocol by pill counting, and medication changes were noted. Dietary intake was assessed using a validated food frequency questionnaire (FFQ) designed for studies of osteoporosis in Scottish women,⁽³⁸⁾ and nutrient intakes from supplements were added to those derived from diet alone to produce a total energy and nutrient intake profile for participants. Subjects provided a urine sample between 10:00 and 11:00 a.m. after an early morning void.

Bone densitometry

DXA of the femur was carried out every 6 months using the same Lunar (DPX α) machine with software version 1.11 and 1.15 (Lunar Corp., Madison, WI, USA). Changes in software did not affect earlier analyses. The recommended Lunar scanning procedure guidelines were followed. Medium scan mode was used for all subjects, and the current was either 750 or 3000 μA , depending on the body mass index (BMI) of the subject. Previous scans were always examined before and during follow-up scans. The Osteoscan pDXA was used for the scanning of the radius, with software version 2.1x (NIM, Verona, Italy). In the standard analysis, the Osteoscan automatically marks the bone edges and positions two region of interest (ROI) boxes, of fixed size: one at the ultradistal site (just distal to the ulnar styloid) and one at the mid-distal site (starting at one third of the distance between the ulna styloid and the olecranon). The nondominant wrist was measured unless there had been a fracture, and on repeat scans, the analysis ROI boxes were matched to the baseline scan. Bone scan results were expressed as BMC (mg) and BMD (BMC divided by bone area; mg/cm^2). The CVs for BMD and BMC measurements, respectively, were as follows: femoral neck, 1.7% and 2.1%; Ward's, 5.1% and 8.7%; trochanter, 2.2% and 7.1%; ultradistal radius, 1.8% and 2.6%; mid-distal radius, 1.9% and 2.1%.

Biochemical analyses

Serum vitamin K₁ was measured by HPLC as previously described.⁽³⁹⁾ Serum 25-hydroxyvitamin D [25(OH)D] was measured by radioimmunoassay using a ^{125}I radioimmunoassay (RIA) kit (68100E; DiaSorin, Wokingham, UK). PTH was measured with the Immulite intact PTH assay (EURO/DPC, Llanberis, Wales, UK). Serum bone-specific alkaline phosphatase (BALP) analyses were carried out using the Alkphase-B immunoassay kit (Metra Biosystems, Great Haseley, UK). Undercarboxylated and carboxylated

serum osteocalcin (GluOC and GlaOC) were determined by separate immunoassays using the respective ELISA kits from Takara Shuzo (Otsu, Shiga, Japan). The sum of GluOC and GlaOC was used as a measure of total OC. As previously recommended,⁽⁴⁰⁾ the vitamin K status of bone was also evaluated by expressing the GluOC fraction as a percentage of total OC. Midmorning urine samples were analyzed for creatinine with a COBAS Bio multichannel analyser using the Boehringer alkaline picrate colorimetric method (124192; Boehringer Mannheim, Mannheim, Germany). Urinary cross-linked N-telopeptides of type I collagen (NTX) was measured using the NTX ELISA kit Osteomark (Ostex, Seattle, WA, USA).

The above biochemical indices were monitored at baseline and 6-month intervals (BALP, GluOC, GlaOC, NTX) or at baseline and yearly intervals [25(OH)D, vitamin K₁, PTH]. Measurements of each analyte were carried out on samples batched by visit except for BALP and PTH, which were performed after the study completion, with samples for each subject batched and analyzed in the same assay run. Assay performance was monitored against external QA schemes for 25(OH)D, vitamin K₁, and PTH or, when no scheme was available, against in-house quality assurance (QA) schemes.

Statistical methods

Statistical procedures were carried out in SPSS (version 10) and SAS. Descriptive statistics were performed for all variables at baseline and on each 6-month visit. Data were transformed toward normality and parametric tests used, with statistical significance taken as $p < 0.05$. Correlation and linear regression procedures were used to study relationships between variables. ANOVA and χ^2 tests were used to detect differences between groups and distributions.

Separate analyses of the bone mineral variables (BMC and BMD) were carried out to address the possibility that BMD does not sufficiently correct for influences of body size, because its derivation as BMC/bone area indicates an areal rather than a volumetric density. Use of BMC with separate adjustment for bone area has been advocated by Prentice et al.⁽⁴¹⁾ to be more statistically rigorous (because it allows the relationship between these two measures to be unforced) and to provide a better adjustment for body size. In line with this approach, we used BMC as the dependent variable and included bone area, weight, and height as independent variables in all multiple regression models.⁽⁴¹⁾ To study the influence of baseline values on response to supplementation, changes in bone mineral and biochemical markers [e.g., 25(OH)D, GluOC, GlaOC] were expressed as a percentage of baseline values and compared across fourths of the baseline distribution using ANOVA.

Cross-sectional multiple regression analysis was used to reveal factors contributing to variance in BMD and BMC at baseline and other time-points. Repeated-measures ANOVA and general linear model (GLM) procedures (SPSS) were used initially to study the significance of changes in BMC and BMD within and between the intervention groups, with appropriate adjustment for covariates (including baseline age, bone area, weight, height, physical

TABLE 1. BASELINE CHARACTERISTICS AND BASAL NUTRIENT INTAKES OF THE 209 WOMEN WHO COMPLETED THE STUDY

	Intervention group			
	Placebo	Vitamin K ₁	Vitamin D + calcium	Vitamins K and D + calcium
Final <i>n</i> (<i>n</i> at recruitment)	56 (61)	54 (60)	50 (62)	49 (61)
Baseline characteristics				
Age (years)	67.8 ± 6.0	67.7 ± 4.9	69.4 ± 6.4	67.8 ± 5.4
Age at menopause (years)	47.8 ± 4.8	49.8 ± 4.3	48.3 ± 8.4	49.5 ± 4.1
Weight (kg)	66.0 ± 8.5	67.5 ± 9.3	64.9 ± 8.4	64.5 ± 9.0
BMI (kg/m ²)	26.2 ± 3.3	26.4 ± 3.7	25.8 ± 3.4	26.1 ± 3.7
Basal nutrient intakes				
Energy (kJ/day)	7594 ± 1935	7790 ± 1955	8082 ± 1916	7762 ± 1852
Protein (% energy)	16.5 ± 1.9	15.8 ± 1.9	16.2 ± 1.8	16.4 ± 1.9
Fat (%energy)	30.1 ± 4.6	30.7 ± 3.0	31.6 ± 3.6	30.6 ± 4.0
Carbohydrate (% energy)	48.5 ± 4.9	49.1 ± 2.9	47.9 ± 3.9	48.4 ± 4.8
Vitamin D (μg/day)	5.0 ± 2.3	5.4 ± 2.6	5.6 ± 3.2	5.9 ± 3.1
Vitamin K ₁ (μg/day)	83.0 ± 24.5	86.9 ± 26.6	82.0 ± 26.8	87.1 ± 28.8
Calcium (mg/day)	1068 ± 280	1046 ± 244	1078 ± 248	1026 ± 219

Values are mean ± SD. Basal nutrient intakes are from foods plus approved supplements and were derived from five 6-month FFQ assessments for each subject over the 2-year intervention. There were no significant differences between groups for any variable.

activity, total energy intake, calcium and other nutrient intakes, serum markers of vitamin D and vitamin K status, and markers of bone turnover). Separate GLMs, with pairwise comparisons against baseline values, were carried out by a statistician, unrelated to the study, who was blind to the groups (Dr Mishra), using the SAS package.

RESULTS

Subject history, compliance, and dietary intakes

A flow chart with the numbers of subjects randomly assigned and retained in each treatment arm at successive 6-month visits is shown in Fig. 1. Of the 244 eligible women randomized into the study, 209 (85.6%) completed the 2-year study, with good supplement adherence based on pill count (median, 99; interquartile range [IQR], 97.3–99.8%). Reasons for withdrawal were illness unrelated to the study (*n* = 17); volunteer preference, noncompliance, or other violations of the inclusion criteria (*n* = 14); and low BMD necessitating further medical intervention (*n* = 4). Of six subjects who specifically reported dislike of the tablets as a reason for dropping out, the distribution between groups was vitamin K (*n* = 2), vitamin D plus calcium (*n* = 2), and combined vitamins K and D plus calcium (*n* = 2).

There were no between-group differences in age, age at menopause, history of oophorectomy, smoking habit, past use of oral contraceptives, or hormone replacement therapy. Table 1 shows the baseline characteristics of the participants and their basal dietary intakes of relevant nutrients. Energy and macronutrient intakes at baseline did not differ significantly between groups. The mean calcium and phyloquinone intakes were adequate, according to UK guidelines,⁽³⁷⁾ but vitamin D intakes in all but three subjects were below the 10 μg/day recommended in this age group. There were no significant group differences for intakes of the trace elements copper, magnesium, manganese, potassium, selenium, and zinc (data not shown). Regular supplement taking was permitted as part of usual dietary intake if

daily intakes were within the inclusion criteria of <10 μg vitamin D, <500 mg calcium, and <30 μg vitamin K₁. Fifty-one percent of the study population reported consuming at least one supplement on at least one of the five 6-month visits. Food supplement use did not differ by group and had a substantial effect on the recorded intakes of vitamin C (~20%) and of vitamin D (~40%) but not on vitamin K₁ or calcium.

Changes in BMD and BMC

The unadjusted mean changes (95% CI) in BMD and BMC over the 2-year study are shown in Table 2. There were no significant differences between intervention groups at any bone site. However, significant changes from baseline were found at some sites. At the mid-distal radius, BMD fell significantly from baseline in all groups (*p* < 0.001), indicating bone loss of between 1% and 2% per year, irrespective of intervention. At the ultradistal radius, whereas there were no significant differences from baseline in the placebo, vitamin K, or vitamin D plus calcium groups, there was a significant increase in BMD of 0.8% per year (*p* < 0.01) in the combined vitamin K and vitamin D plus calcium group. Unadjusted BMC at the ultradistal radius showed the same significant gain in bone mineral (*p* < 0.01) in women who had received combined supplementation with vitamin K and D. Although there was a slight 2-year rise in both BMD and BMC at the ultradistal radius in the placebo group, this did not achieve significance. At the femoral trochanter, there were no significant differences from baseline in BMD, but there were significant 2-year rises in BMC in the placebo group, vitamin D group, and combined vitamin groups (Table 2).

The DXA data were also analyzed by GLM procedures, with appropriate adjustment of covariates as described in the Materials and Methods section. An essentially similar picture was obtained for both BMD and BMC results at all bone sites except for the femoral trochanter, where there were unexplained significant gains in BMC but not BMD at

TABLE 2. UNADJUSTED TWO-YEAR MEAN CHANGES (95% CI) IN BMD AND BMC IN THE FOUR INTERVENTION GROUPS

Bone site	Interval (months)	Intervention group				ANOVA from baseline
		Placebo	Vitamin K ₁	Vitamin D + calcium	Vitamins K and D + calcium	
Femoral neck BMD (mg/cm ²)	0-24	0.7 (-10.2, 11.6)	-4.2 (-15.6, 7.2)	1.9 (-6.5, 10.3)	1.0 (-8.0, 10.1)	NS
Femoral neck BMC (mg)	0-24	-32.1 (-127, 0.6)	-49.6 (-141, 42.0)	25.8 (-59.7, 111)	9.9 (-86.6, 66.8)	NS
Femoral trochanter BMD (mg/cm ²)	0-24	8.9 (-3.9, 21.6)	4.3 (-9.1, 17.8)	7.8 (-2.3, 17.8)	6.9 (-3.9, 17.7)	NS
Femoral trochanter BMC (mg)	0-24	630* (218, 1040)	346 (-100, 793)	460* (158, 763)	549* (207, 890)	<i>p</i> < 0.01
Femoral Ward's BMD (mg/cm ²)	0-24	0.1 (-19.6, 19.4)	2.2 (-20.1, 16.2)	11.4 (-6.3, 29.2)	6.0 (-21.8, 9.9)	NS
Femoral Ward's BMC (mg)	0-24	-36.7 (-143, 70.3)	-30.9 (-126, 64.5)	39.9 (-60.3, 140)	25.6 (-106, 54.5)	NS
Mid-distal radius BMD (mg/cm ²)	0-24	-20.4* (-26.8, -13.9)	-18.6* (-26.7, -10.5)	-16.6* (-23.3, -9.9)	-16.9* (-23.7, -10.1)	<i>p</i> < 0.001
Mid-distal radius BMC (mg)	0-24	-20.4* (-25.4, -15.3)	-16.3* (-22.1, -10.6)	-14.9* (-20.8, -9.0)	-14.1* (-20.7, -7.6)	<i>p</i> < 0.001
Ultradistal radius BMD (mg/cm ²)	0-24	3.6 (-0.2, 7.4)	-1.9 (-7.5, 3.8)	1.7 (-3.1, 6.5)	6.2* (2.6, 9.8)	<i>p</i> < 0.01
Ultradistal radius BMC (mg)	0-24	7.2 (-4.9, 19.3)	-6.3 (-20.8, 8.3)	-1.5 (-17.4, 14.4)	30.7* (15.2, 46.2)	<i>p</i> < 0.01

Values are the mean changes (95% CI) in BMD and BMC from baseline over the 2-year intervention period for each group. Significant differences between intervention groups and/or from baseline were investigated by GLM, repeated measures with pairwise comparisons. There were no significant differences between intervention groups at any bone site.

* Significant bone loss from baseline.

† Significant bone gain from baseline.

NS, not significant (*p* > 0.05).

some visits in all groups. There were no significant differences for adjusted BMD or BMC between the four treatment groups, but changes from baseline were observed. For the most part, only adjusted BMC results are presented here, in line with previous recommendations for the reporting of bone mineral measurements in population-based research.⁽⁴¹⁾ The 6-month changes from baseline in covariate-adjusted BMC at the three sites of the femur and for the mid-distal radius are shown in Fig. 2. As found for the unadjusted results, there were significant increases from baseline in adjusted BMC at the femoral trochanter after 1 year and beyond in both the vitamin D group and in the combined vitamins K and D group. On the other hand, there was a significant increase in trochanter BMC in the placebo group after 2 years. Adjusted values for BMC at the mid-distal radius confirmed that there was a progressive loss of bone mineral at this site over 24 months.

The only consistent intervention effect on BMD and BMC was seen at the ultradistal radius. GLM-adjusted changes (95% CI) in BMD and BMC are shown in Fig. 3. This rigorous analysis confirmed the findings from the unadjusted data that bone mineral had increased from baseline at the ultradistal radius in the combined intervention group, with significant increases in BMD at 18 and 24 months and in BMC at all visits from 6 months onward. A significant increase in BMD in the placebo group was only evident at the 6-month time-point.

Changes in indicators of vitamin K and vitamin D status

The effect of intervention on tissue and functional markers of vitamin K and vitamin D status during the study are shown in Table 3. With the exception of %GluOC and 25(OH)D, the status indicators showed a skewed distribution; for these indicators, geometric means are given in Table 3 because they represent better average measures for such non-normally distributed data. Fasting vitamin K₁ serum concentrations rose significantly in response to vitamin K such that the geometric mean for all supplemented subjects had risen by 157% (95% CI, 101, 212) after 2 years. Vitamin K supplementation induced a significant 40% decrease in the absolute concentration of GluOC and a significant 105% increase in GlaOC over 2 years (Table 3). The apparent lower influence of vitamin K on the GluOC fraction compared with GlaOC is explained by the finding that both fractions rose with time independently of vitamin K. This non-vitamin K-dependent rise was slightly greater for GluOC than for GlaOC. Over 2 years, the geometric means for GluOC increased in the nonsupplemented groups by 49% (placebo) and 69% (vitamin D plus calcium), respectively, whereas for GlaOC, the equivalent increases were 30% and 24%, respectively. The bone vitamin K status indicator, %GluOC,⁽⁴⁰⁾ responded substantially to vitamin K supplementation, with 2-year decreases of 48% (vitamin K) and 54% (vitamins K and D) and an overall decrease of 51% (95% CI, -47.5, -54.0). These changes in the γ -carboxylation status of osteocalcin were evident, and consistent, from 6 months onward. In contrast, there was a trend for %GluOC to rise over 2 years in the non-vitamin




FIG. 2. GLM-adjusted mean changes (95% CI) in BMC from baseline at the femoral neck, femoral Ward's, femoral trochanter, and mid-distal radius after successive 6-month intervals of supplementation. *Significant change from baseline.




FIG. 3. GLM-adjusted mean changes (95% CI) in BMD and BMC from baseline at the ultradistal radius after successive 6-month intervals of supplementation. *Significant increase from baseline.

K-supplemented groups, but the increases of 7.1% (placebo) and 15.5% (vitamin D plus calcium) only reached significance in the latter. Quartile analysis showed that the greatest decreases in GluOC and %GluOC in the vitamin K supplementation groups occurred in those women with the highest values at baseline (results not shown).

Serum 25(OH)D rose significantly ($p < 0.001$) in response to vitamin D supplementation, with a mean increase of 16.7% (95% CI, 9.0, 24.4) over 2 years. This rise in 25(OH)D was accompanied by a significant fall ($p = 0.002$) in PTH of 10.9% (95% CI, -20.6, -1.2). In contrast, 25(OH)D fell significantly ($p < 0.001$) in the non-vitamin D-supplemented subjects, with a mean decrease of 17.5% (95% CI, -13.5, -21.4) over 2 years, but PTH showed no significant change. Quartile analysis showed that the women showing higher percentage increases in 25(OH)D had lower concentrations at baseline (results not shown).

For every status marker measured, there was no evidence of an additive effect in women who took combined vitamin K and vitamin D plus calcium.

Relationship between serum 25(OH)D and PTH

Baseline 25(OH)D was negatively correlated to PTH before ($r = -0.23$, $p < 0.01$) and after age adjustment ($r = -0.26$, $p < 0.001$), and this relationship was unaffected by vitamin D supplementation. Further analyses using a repeated-measure, multivariate linear regression model were carried out to study the ability of serum 25(OH)D to predict change in serum PTH concentrations over time. To avoid seasonal influence, the chosen time intervals were baseline, 12 months, and 24 months, representing subject visits during October to January. This data has been previously reported in abstract form.⁽⁴²⁾ In brief, change in serum 25(OH)D was found to be a strong independent negative predictor of the change in PTH, with regression coefficient $\times 100$ (\pm SE) = -0.9 (0.2), $p < 0.0001$. Previous PTH also positively predicted the change in PTH, with coefficient $\times 100$ (\pm SE) in the regression model = 94 (3.7), $p < 0.0001$. When "current" 25(OH)D was substituted for "change" in 25(OH)D in the model, it was not a significant independent predictor of change in PTH. Neither age nor calcium intake was a significant predictor of change in PTH.

Changes in indicators of bone turnover

No significant differences were found between groups for the markers of bone formation and resorption represented

TABLE 3. SERUM INDICATORS OF VITAMIN K AND VITAMIN D STATUS IN THE FOUR INTERVENTION GROUPS

	Study month	Intervention group				ANOVA by intervention group
		Placebo	Vitamin K ₁	Vitamin D + calcium	Vitamins K and D + calcium	
Serum vitamin K ₁ (μg/liter)	0	0.25 (0.20, 0.31)	0.22 (0.17, 0.27)	0.21 (0.16, 0.29)	0.21 (0.16, 0.27)	NS
	12	0.23 (0.17, 0.32)	0.66 (0.53, 0.82)	0.22 (0.16, 0.30)	0.55 (0.45, 0.68)	<i>p</i> < 0.001
	24	0.29 (0.24, 0.34)	0.55 (0.46, 0.67)	0.23 (0.21, 0.32)	0.53 (0.44, 0.64)	<i>p</i> < 0.001
ANOVA* (within-group)		NS	<i>p</i> < 0.001	NS	<i>p</i> < 0.001	
Serum GluOC (μg/liter)	0	5.03 (4.15, 6.10)	5.52 (4.66, 6.53)	4.81 (3.95, 5.87)	5.77 (4.67, 7.14)	NS
	12	7.02 (6.19, 7.95)	4.04 (3.45, 4.73)	6.77 (5.98, 7.66)	3.67 (3.02, 4.45)	<i>p</i> < 0.001
	24	7.48 (6.51, 8.58)	3.77 (3.19, 4.46)	8.13 (7.04, 9.38)	3.15 (2.55, 3.88)	<i>p</i> < 0.001
ANOVA* (within-group)		<i>p</i> < 0.001	<i>p</i> < 0.001	<i>p</i> < 0.001	<i>p</i> < 0.001	
Serum GlaOC (μg/liter)	0	5.80 (5.04, 6.68)	6.04 (5.34, 6.82)	5.81 (4.89, 6.90)	5.66 (4.89, 6.54)	NS
	12	6.99 (6.2, 7.98)	10.61 (9.92, 11.36)	6.80 (5.99, 7.71)	9.90 (9.08, 10.79)	<i>p</i> < 0.001
	24	7.52 (6.86, 8.25)	12.11 (11.24, 13.05)	7.20 (6.40, 8.08)	11.88 (11.12, 12.69)	<i>p</i> < 0.001
ANOVA* (within-group)		<i>p</i> < 0.001	<i>p</i> < 0.001	<i>p</i> = 0.001	<i>p</i> < 0.001	
Percent GluOC	0	46.4 ± 18.0	48.0 ± 15.3	45.8 ± 17.7	50.0 ± 17.8	NS
	12	49.9 ± 14.3	28.8 ± 10.8	50.1 ± 12.2	28.6 ± 13.5	<i>p</i> < 0.001
	24	49.7 ± 13.8	25.2 ± 10.9	52.9 ± 14.4	23.0 ± 13.8	<i>p</i> < 0.001
ANOVA (within-group)		NS	<i>p</i> < 0.001	<i>p</i> = 0.001	<i>p</i> < 0.0001	
Serum 25(OH)D (μg/liter)	0	22.8 ± 6.1	23.1 ± 6.7	25.0 ± 6.2	24.7 ± 7.0	NS
	12	19.1 ± 6.9	18.4 ± 6.8	29.7 ± 6.1	29.2 ± 6.5	<i>p</i> < 0.001
	24	19.5 ± 5.3	18.3 ± 6.4	29.8 ± 6.0	28.3 ± 6.6	<i>p</i> < 0.001
ANOVA (within-group)		<i>p</i> < 0.001	<i>p</i> < 0.001	<i>p</i> < 0.001	<i>p</i> < 0.001	
Serum PTH (pM)	0	4.38 (3.99, 4.81)	4.69 (4.25, 5.17)	4.86 (4.37, 5.40)	4.30 (3.84, 4.81)	NS
	12	4.75 (4.28, 5.28)	4.91 (4.41, 5.47)	4.31 (3.85, 4.83)	4.00 (3.58, 4.48)	<i>p</i> = 0.037
	24	4.42 (3.98, 4.90)	4.64 (4.16, 5.18)	4.47 (4.03, 4.96)	3.71 (3.22, 4.27)	<i>p</i> = 0.036
ANOVA* (within-group)		NS	NS	<i>p</i> = 0.003	<i>p</i> < 0.001	

Values are geometric means (95% CI) for (log-transformed) skewed distributions and arithmetic means ± SD for normal distributions. Significance differences within each intervention group with time were investigated by ANOVA for repeat measures using GLM. Between-group significances at each subject visit were investigated by ANOVA.

* ANOVA on log-transformed data.

NS, not significant (*p* > 0.05).

by serum BALP and urinary NTX, respectively. Total OC represented by the sum of the GluOC and GlaOC fractions also showed no significant group differences. Significant increases with time were seen for total OC in all groups (*p* < 0.01) and for urinary NTX in all but the placebo group (*p* < 0.05).

DISCUSSION

This was a novel study, designed to address the specific question of whether taking regular, high-dietary amounts of vitamin K₁, either alone, or in combination with moderate supplements of vitamin D₃, could reduce the rate of bone mineral loss in healthy, nonosteoporotic older women. Calcium was included in the vitamin D treatment group to ensure adequacy but was not the nutrient of primary interest. The only comparable vitamin K₁ supplementation trial published thus far is the Maastricht study, which differed in design and had a 5-fold higher vitamin K₁ dose.⁽²⁶⁾

The power of our study to detect changes in bone mineral accretion was diminished both by the wide interindividual variation in BMD at baseline and by the lower than expected rate of bone loss in the placebo group. However, it is recognized that the rate of bone loss in women of this age varies greatly, depending on the bone site, nutritional sta-

tus, and other factors.^(43,44) For example, in the recent Women's Health Initiative (WHI) trial in women 50–79 years of age,⁽¹⁰⁾ the placebo group showed a steady 9-year increase in BMD for the whole body and total spine, whereas BMD for the total hip declined by only 1% over the same period. Lesser⁽⁴⁵⁾ suggested that this may have been attributable to the “therapeutic” background intakes of calcium (1154 mg/day) and vitamin D (9.2 μg/day) in that trial. In our study, the mean calcium intake at baseline of 1055 mg/day was similar to that in the WHI trial but considerably higher than the 690 mg/day found in the British National Diet and Nutrition Survey (NDNS) of 1994/5 in women 65–84 years of age.⁽⁴⁶⁾ The mean background vitamin D intake in our study of 5.4 μg/day was less than that of the WHI trial but greater than the nationally representative intake of 3.5 μg/day in British women 65–84 years of age.⁽⁴⁶⁾ In our study, the only bone site showing a consistently significant bone loss was the mid-distal radius, but the loss was not significantly different between placebo and supplementation groups. On the other hand, at the site of the ultradistal radius, we did observe a significant increase in BMC and BMD from baseline in those women who had received combined calcium, vitamin D, and vitamin K. This increase was evident for unadjusted BMD and BMC and after adjustment for covariates. With its high content of

trabecular bone, the ultradistal forearm has a higher metabolic turnover rate than predominantly cortical bone and may be more responsive to intervention effects. A study of healthy, postmenopausal women in Massachusetts suggested that vitamin D supplementation at 10 $\mu\text{g}/\text{day}$ was more effective in preventing bone loss of trabecular bone than of cortical bone.⁽⁴⁷⁾ The ultradistal forearm (but not the mid-distal forearm or lumbar spine) was previously shown by some of us to be similarly responsive to a 2-year combined exercise plus calcium intervention in older women.⁽⁴⁸⁾ The possible greater sensitivity of the ultradistal radius to a vitamin K effect is supported by the findings from a meta-analysis of nine studies in which BMD changes had been measured after exposure to oral anticoagulants.⁽⁴⁹⁾ Although this meta-analysis showed a trend to bone loss at most sites, a significant decrease in BMD was only observed at the ultradistal radius.⁽⁴⁹⁾ Thus, the effect of vitamin K antagonists seems to mirror our findings of vitamin K (with vitamin D) supplementation but in reverse.

We did not measure vertebral BMD because of the confounding effect of vascular calcification in older people and because matrix Gla protein is a vitamin K-dependent inhibitor of vascular calcification.⁽¹³⁾ This decision seems justified in view of our previous exercise study⁽⁴⁸⁾ and the results of a 3-year intervention study from Maastricht in which a supplementation effect of combined vitamin K, vitamin D (with calcium), and additional minerals was found at the femoral neck but not at the lumbar spine.⁽²⁶⁾ In the Maastricht study, the rate of bone loss at the femoral neck was significantly slowed over 3 years in this combined vitamin K group but not in subjects receiving vitamin D plus minerals alone.⁽²⁶⁾ However, at 1 mg/day, the supplementary vitamin K₁ dose in the Maastricht study was 5-fold higher than in our study and would be difficult to achieve from dietary sources alone.

The lack of a significant effect on BMD in the vitamin D plus calcium-supplemented group suggests that the improvement in vitamin D status alone provided no additional benefit in these healthy older women who, on the basis of calcium intake, were already calcium replete. The efficacy of vitamin D given alone or with calcium in the prevention of bone loss or fractures is controversial, with some trials showing significant benefit^(3,5,6) and some not.^(7–10) A variety of factors could account for these differences, but important variables include vitamin D status at baseline⁽⁵⁰⁾ and habitual calcium intake of the population under study. Our participants were younger, less frail, and were less likely to have vitamin D insufficiency and secondary hyperparathyroidism than participants in some trials where a vitamin D/calcium intervention effect was seen; this may have limited any effect on bone mass. Evidence of a supposedly satisfactory vitamin D status in most of our subjects was supported by their mean baseline 25(OH)D concentration of 24 $\mu\text{g}/\text{liter}$ (60 nM) and by the fact that none fell below the level considered adequate (10 $\mu\text{g}/\text{liter}$; 25 nM) by a UK working group.⁽⁴⁶⁾ Despite this, supplementation with 10 $\mu\text{g}/\text{day}$ vitamin D₃ not only significantly raised 25(OH)D levels but also significantly lowered PTH levels. The suppression of PTH, in particular, suggests that suboptimal vitamin D status was common in our participants. As in a

large, global study,⁽⁵⁰⁾ we found that the increase in 25(OH)D after vitamin D supplementation is strongly dependent on the baseline value, with maximal increases being seen in those women with an initially low serum 25(OH)D. Our analyses also revealed that the change in 25(OH)D is predictive of the inverse change in PTH. These data support the concept that pre-existing vitamin D status is a major factor in determining the change in 25(OH)D in response to vitamin D supplementation and therefore of the degree of lowering in PTH levels.⁽⁵⁰⁾ Although methodological differences usually preclude comparisons of absolute values, the overall 2-year changes of 17% for 25(OH)D and 11% for PTH were slightly lower than in other studies for an equivalent vitamin D dose.^(6,50,51) This is again likely to reflect the better initial vitamin D status of our subjects. This is supported by comparing our results with a recent intervention study in free-living elderly in Randers, Denmark.⁽⁶⁾ Comparison with this Danish study is especially valid because the doses of vitamin D (and calcium), the 2-year sampling period, and the methodology for 25(OH)D and PTH measurements were exactly the same in both studies. In a small subset of the Danish participants, the equivalent 2-year changes for 25(OH)D and PTH were 27% and 15%, respectively. These greater changes probably reflected the lower baseline 25(OH)D of 15 $\mu\text{g}/\text{liter}$ in the Danish subjects compared with 24 $\mu\text{g}/\text{liter}$ in our Scottish subjects. Coincidentally, Randers and Dundee share exactly the same latitude (56°28').

The basal intakes of vitamin K₁ (mean, 85 $\mu\text{g}/\text{day}$) are representative of intakes found in a UK National Survey of people ≥ 65 years of age,⁽⁵²⁾ except that a lower proportion of our study population (27% versus 59%) failed to meet the current UK guideline of 1 $\mu\text{g}/\text{kg}$ body weight/day.⁽³⁷⁾ However, nearly two thirds of the women in our study failed to meet the more recent U.S. adequate intake of 90 $\mu\text{g}/\text{day}$.⁽⁵³⁾

Our study supports the concept^(12,22,54,55) that much higher dietary intakes of vitamin K are needed for γ -carboxylation of bone OC compared with the hepatic coagulation proteins. This is the first study to show that regular, long-term supplementation with vitamin K₁ at a high dietary level can substantially improve and maintain vitamin K status as evidenced by increases in both tissue stores (serum vitamin K₁) and a functional marker (OC carboxylation). Although absolute values for %GluOC are assay dependent,⁽⁴⁰⁾ the reduction of ~50% is consistent with that seen in short-term repletion studies after supplementation with ~300 μg of vitamin K₁^(22,23,55) but smaller than that after a supradietary dose of 1000 $\mu\text{g}/\text{day}$.⁽⁵⁵⁾ In our study, maximal OC carboxylation was attained by 6 months and did not change significantly thereafter. It is noteworthy that those women with the most impaired vitamin K status at baseline (as represented by the highest quartiles of GluOC and %GluOC and lowest quartile of GlaOC, respectively) showed the greatest changes in OC carboxylation after vitamin K supplementation.

We chose vitamin K₁ for this study because this molecular form accounts for 90% of dietary vitamin K intakes.⁽⁵⁶⁾ A proprietary preparation of menaquinone-4 (menatetranone) is used as an anti-osteoporotic agent in Japan, and a

recent meta-analysis of Japanese trials showed an association of menatetranone supplementation with reduced fracture incidence.⁽⁵⁷⁾ These menatetranone trials differed from our study in that the administered doses were extremely large (generally 45 mg/day) and the target groups already had pre-existing involutional^(27–29) or secondary^(58,59) osteoporosis or osteopenia. Enhanced effects on BMD have been found when menatetranone is used in combination with 1 α -hydroxycholecalciferol.^(28,29) There is also evidence that the effectiveness of menatetranone in inhibiting bone loss is related to the specific geranyl-geranyl side chain of this form.^(60,61) The existence of an unusual metabolic pathway that converts a fraction of dietary vitamin K₁ to menaquinone-4⁽⁶²⁾ does not exclude the possibility that any benefit of supplementary vitamin K₁ might be attributable to the menaquinone-4 thus formed.

Our study provided no evidence that vitamin K₁ taken alone at high dietary intakes had any influence on maintaining bone mass despite the improvement in OC carboxylation. However, the Japanese experience suggests that the effectiveness of vitamin K₂ in preventing fractures is greater than that predicted from its ability to maintain bone mass.⁽²⁷⁾ This is supported by the finding in older American women that serum GluOC is more strongly related to ultrasonic transmitted velocity than to femoral neck density⁽⁶³⁾; this relationship prompted the authors to suggest that vitamin K insufficiency might have a greater effect on bone quality than on BMD. Our findings showing no demonstrable effect of supplementation on markers of bone formation (BALP) and resorption (NTX) are in agreement with the lack of consistent effect on equivalent bone markers after long-term supplementation with 1 mg vitamin K₁⁽²⁶⁾ or 45 mg menatetranone.⁽²⁷⁾ The opposite findings of a rise⁽²⁷⁾ or fall⁽²⁶⁾ in OC may be artefactual (e.g., a consequence of the varying affinities of different OC antibodies to changing carboxylation status); this variability in OC response after vitamin K supplementation has been noted previously.^(54,64) In this study, OC, as represented by the sum of the GluOC and GlaOC fractions, showed no differences between groups. Subsequent work using a standard RIA for the intact OC (and 1–43 fragment) also showed no differences in OC between groups (MJ Shearer and CM Gundberg, unpublished observations).

Although several large population studies have revealed significant associations of dietary intakes of vitamin K with fracture rates and/or BMD,^(32–34) the results are not always consistent or uniform across the sexes or age groups. The same is true of associations with indicators of vitamin K status.⁽³⁵⁾ Clearly it is difficult to distinguish vitamin K status from all the other factors influencing changes in BMD, and one consistent criticism of these associations has been the difficulty in excluding the confounding effect of overall poor nutrition.^(11,33) Nevertheless, in the Framingham study,⁽³³⁾ it is noteworthy that the top quartile of vitamin K₁ intake (median, 250 μ g/day), which was associated with reduced fracture risk, was comparable with the mean total dietary intake achieved in this study of 285 μ g/day.

This study lends support to the concept that vitamin K may act in concert with vitamin D to reduce bone loss. Three other studies have shown an enhanced effect of com-

bined vitamin K and vitamin D supplementation,^(26,28,29) but ours is the first in which the extra vitamin K intake can be considered to be within the dietary range. The mechanism(s) whereby parallel improvements in the status of both vitamins D and K might have an enhanced effect over either vitamin alone is unclear. It is feasible that any synergy between vitamins D and K derives from separate effects exerted independently or alternatively from their concerted action through common proteins or pathways. It is known that 1,25-dihydroxyvitamin D enhances the transcription of vitamin K-dependent bone proteins (e.g., osteocalcin, matrix Gla protein, and Gas6). Other evidence has led to the suggestion that vitamin D might directly influence the γ -carboxylation reaction of bone Gla proteins.^(16,24) This hypothesis was not supported by our findings because we showed that γ -carboxylation of osteocalcin was neither enhanced by vitamin D (relative to the placebo group) nor by its combination with vitamin K (relative to the vitamin K group).

In summary, dietary supplementation with a combination of nutritionally relevant amounts of vitamin K with vitamin D and calcium in healthy older women was associated with a modest but significant increase in BMC at one site, consisting predominantly of trabecular bone. Similar changes were not observed in either the vitamin K group alone or in the calcium plus vitamin D group, suggesting a synergistic role of the combination as suggested by previous reports.

ACKNOWLEDGMENTS

This study was supported by a contract (N05001) from the UK Food Standards Agency. The authors thank Hoffmann-La Roche, Vitamins Division (now DSM Nutritional Products), Basel, Switzerland, for supplying the tablets for supplementation, and particularly to Jürg Haller for his keen interest in the project. We thank the participants, Wendy Cassells (the project research nurse), Rosemary Price (the project research dietician), Susan Whiteford and Andrea Subedi (bone density measurements), Inez Murie (nutrition interviews), Jimmy Gibson (data entry), Dominic Harrington (supervision of serum analyses of vitamin K), and Renata Gorska for assistance in the preparation of the illustrations.

REFERENCES

1. Bonjour J-P, Schurch M-A, Rizzoli R 1996 Nutritional aspects of hip fractures. *Bone* **18**:139S–144S.
2. Reid DM, Macdonald HM 2001 Nutrition and bone: Is there more to it than just calcium and vitamin D? *Q J Med* **94**:53–56.
3. Chapuy MC, Arlot ME, Duboeuf F, Brun J, Crouzet B, Arnaud S, Delmas PD, Meunier PJ 1992 Vitamin D₃ and calcium to prevent hip fractures in elderly women. *N Engl J Med* **327**:1637–1642.
4. Reid IR, Ames RW, Evans MC, Gamble GD, Sharpe SJ 1995 Long-term effects of calcium supplementation on bone loss and fractures in postmenopausal women: A randomized controlled trial. *Am J Med* **98**:331–335.
5. Dawson-Hughes B, Harris SS, Krall EA, Dallal GE 1997 Effect of calcium and vitamin D supplementation on bone density in men and women 65 years of age or older. *N Engl J Med* **337**:670–676.

6. Larsen ER, Mosekilde L, Foldspang A 2004 Vitamin D and calcium supplementation prevents osteoporotic fractures in elderly community dwelling residents: A pragmatic population-based 3-year intervention study. *J Bone Miner Res* **19**:370–378.
7. Lips P, Graafmans WC, Ooms ME, Bezemer PD, Bouter LM 1996 Vitamin D supplementation and fracture incidence in elderly persons: A randomized, placebo-controlled clinical trial. *Ann Intern Med* **124**:400–406.
8. Grant AM, Avenell A, Campbell MK, McDonald AM, MacLennan GS, McPherson GC, Anderson FH, Cooper C, Francis RM, Donaldson C, Gillespie WJ, Robinson CM, Torgerson DJ, Wallace WA, RECORD Trial Group 2005 Oral vitamin D3 and calcium for secondary prevention of low-trauma fractures in elderly people (Randomised Evaluation of Calcium Or vitamin D, RECORD): A randomised placebo-controlled trial. *Lancet* **365**:1621–1628.
9. Porthouse J, Cockayne S, King C, Saxon L, Steele E, Aspray T, Baverstock M, Birks Y, Dumville J, Francis RM, Iglesias C, Puffer S, Sutcliffe A, Watt I, Torgerson DJ 2005 Randomised controlled trial of supplementation with calcium and cholecalciferol (vitamin D3) for prevention of fractures in primary care. *BMJ* **330**:1003–1006.
10. Jackson RD, LaCroix AZ, Gass M, Wallace RB, Robbins J, Lewis CE, Bassford T, Beresford SA, Black HR, Blanchette P, Bonds DE, Brunner RL, Brzyski RG, Caan B, Cauley JA, Chlebowski RT, Cummings SR, Granek I, Hays J, Heiss G, Hendrix SL, Howard BV, Hsia J, Hubbell FA, Johnson KC, Judd H, Kotchen JM, Kuller LH, Langer RD, Lasser NL, Limacher MC, Ludlam S, Manson JE, Margolis KL, McGowan J, Ockene JK, O'Sullivan MJ, Phillips L, Prentice RL, Sarto GE, Stefanick ML, Van Horn L, Wactawski-Wende J, Whitlock E, Anderson GL, Assaf AR, Barad D, Women's Health Initiative Investigators 2006 Calcium plus vitamin D supplementation and the risk of fractures. *N Engl J Med* **354**:669–683.
11. Binkley NC, Suttie JW 1995 Vitamin K nutrition and osteoporosis. *J Nutr* **125**:1812–1821.
12. Vermeer C, Jie K-SG, Knapen MHJ 1995 Role of vitamin K in bone metabolism. *Annu Rev Nutr* **15**:1–22.
13. Vermeer C, Shearer MJ, Zitterman A, Bolton-Smith C, Szulc P, Hodges S, Walter P, Rambeck W, Stöcklin E, Weber P 2004 Beyond deficiency: Potential benefits of increased intakes of vitamin K for bone and vascular health. *Eur J Nutr* **43**:325–335.
14. Hoang QQ, Sicheri F, Howard AJ, Yang DSC 2003 Bone recognition mechanism of porcine osteocalcin from crystal structure. *Nature* **425**:977–980.
15. Boskey AL, Gadaleta S, Gundberg C, Doty SB, Ducey P, Karsenty G 1998 Fourier transform infrared microspectroscopic analysis of bones of osteocalcin-deficient mice provides insight into the function of osteocalcin. *Bone* **23**:187–196.
16. Szulc P, Chapuy MC, Meunier PJ, Delmas PD 1993 Serum undercarboxylated osteocalcin is a marker of the risk of hip fracture in elderly women. *J Clin Invest* **91**:1769–1774.
17. Szulc P, Chapuy MC, Meunier PJ, Delmas PD 1996 Serum undercarboxylated osteocalcin is a marker of the risk of hip fracture: A three year follow up study. *Bone* **18**:487–488.
18. Vergnaud P, Garnero P, Meunier PJ, Breart G, Kamihagi K, Delmas PD 1997 Undercarboxylated osteocalcin measured with a specific immunoassay predicts hip fracture in elderly women: The EPIDOS study. *J Clin Endocrinol Metab* **82**:719–724.
19. Luukinen H, Kakonen SM, Pettersson K, Koski K, Laippala P, Lovgren T, Kivela SL, Vaananen HK 2000 Strong prediction of fractures among older adults by the ratio of carboxylated to total osteocalcin. *J Bone Miner Res* **15**:2473–2478.
20. Szulc P, Arlot M, Chapuy MC, Duboeuf F, Meunier PJ, Delmas PD 1994 Serum undercarboxylated osteocalcin correlates with hip bone mineral density in elderly women. *J Bone Miner Res* **9**:1591–1595.
21. Knapen MHJ, Nieuwenhuijzen Kruseman AC, Wouters RSME, Vermeer C 1998 Correlation of serum osteocalcin fractions with bone mineral density in women during the first 10 years after menopause. *Calcif Tissue Int* **63**:375–379.
22. Sokoll LJ, Booth SL, O'Brien ME, Davidson KW, Tsaion KI, Sadowski JA 1997 Changes in serum osteocalcin, plasma phylloquinone, and urinary γ -carboxyglutamic acid in response to altered intakes of dietary phylloquinone in human subjects. *Am J Clin Nutr* **65**:779–784.
23. Booth SL, O'Brien-Morse ME, Dallal GE, Davidson KW, Gundberg CM 1999 Response of vitamin K status to different intakes and sources of phylloquinone-rich foods: Comparison of younger and older adults. *Am J Clin Nutr* **70**:368–377.
24. Deyl Z, Adam M 1983 Evidence for vitamin D-dependent γ -carboxylation in osteocalcin related proteins. *Biochem Biophys Res Commun* **113**:294–300.
25. Lian JB, Stein GS, Stein JL, van Wijnen AJ 1999 Regulated expression of the bone-specific osteocalcin gene by vitamins and hormones. *Vitam Horm* **55**:443–509.
26. Braam LAJLM, Knapen MHJ, Geusens P, Brouns F, Hamulyak K, Gerichhausen MJW, Vermeer C 2003 Vitamin K1 supplementation retards bone loss in postmenopausal women between 50 and 60 years of age. *Calcif Tissue Int* **73**:21–26.
27. Shiraki M, Shiraki Y, Aoki C, Miura M 2000 Vitamin K₂ (menatetranone) effectively prevents fractures and sustains lumbar bone mineral density in osteoporosis. *J Bone Miner Res* **15**:515–521.
28. Iwamoto J, Takeda T, Ichimura S 2000 Effect of combined administration of vitamin D₃ and vitamin K₂ on bone mineral density of the lumbar spine in postmenopausal women with osteoporosis. *J Orthop Sci* **5**:546–551.
29. Ushiroyama T, Ikeda A, Ueki M 2002 Effect of continuous therapy with vitamin K₂ and vitamin D₃ on bone mineral density and coagulofibrinolysis function in postmenopausal women. *Maturitas* **41**:211–221.
30. Bolton-Smith C, Price RJG, Fenton ST, Harrington DJ, Shearer MJ 1998 The relationship between plasma and dietary phylloquinone (vitamin K₁) in Scottish adults. *Proc Nutr Soc* **57**:148A.
31. Bolton-Smith C, Price RJG, Fenton ST, Harrington DJ, Shearer MJ 2000 Compilation of a provisional UK database for the phylloquinone (vitamin K₁) content of foods. *Br J Nutr* **83**:389–399.
32. Feskanich D, Weber P, Willett WC, Rockett H, Booth SL, Colditz GA 1999 Vitamin K intake and hip fractures in women: A prospective study. *Am J Clin Nutr* **69**:74–79.
33. Booth SL, Tucker KL, Chen H, Hannan MT, Gagnon DR, Cupples LA, Wilson PWF, Ordovas J, Schaefer EJ, Dawson-Hughes B, Kiel DP 2000 Dietary vitamin K intakes are associated with hip fracture but not with bone mineral density in elderly men and women. *Am J Clin Nutr* **71**:1201–1208.
34. Booth SL, Broe KE, Gagnon DR, Tucker KL, Hannan MT, McLean RR, Dawson-Hughes B, Wilson PWF, Cupples LA, Kiel DP 2003 Vitamin K intake and bone mineral density in women and men. *Am J Clin Nutr* **77**:512–516.
35. Booth SL, Broe KE, Peterson JW, Cheng DM, Dawson-Hughes B, Gundberg CM, Cupples LA, Wilson PWF, Kiel DP 2004 Associations between vitamin K biochemical measures and bone mineral density in men and women. *J Clin Endocrinol Metab* **89**:4904–4909.
36. Kalkwarf HJ, Khoury JC, Bean J, Elliot JG 2004 Vitamin K, bone turnover, and bone mass in girls. *Am J Clin Nutr* **80**:1075–1080.
37. Department of Health 1991 Dietary Reference Values for Food Energy and Nutrients for the United Kingdom. Report on Health and Social Subjects no. 41. H. M. Stationery Office, London, UK.
38. New SA, Bolton-Smith C, Grubb DA, Reid DM 1997 Nutritional influences on bone mineral density: A cross-sectional study in pre-menopausal women. *Am J Clin Nutr* **65**:1831–1839.
39. Thane CW, Bates CJ, Shearer MJ, Unadkat N, Harrington DJ, Paul AA, Prentice A, Bolton-Smith C 2002 Plasma phylloquinone (vitamin K₁) concentration and its relationship to intake in a national sample of British elderly people. *Br J Nutr* **87**:615–622.
40. Gundberg CM, Nieman SD, Abrams S, Rosen H 1998 Vitamin

- K status and bone health: An analysis of methods for determination of undercarboxylated osteocalcin. *J Clin Endocrinol Metab* **83**:3258–3266.
41. Prentice A, Parsons TJ, Cole TJ 1994 Uncritical use of bone mineral density in absorptiometry may lead to size-related artifacts in the identification of bone mineral determinants. *Am J Clin Nutr* **60**:837–842.
 42. Bolton-Smith C, Mishra GD, Mole PA, McMurdo MET, Paterson CR 2001 A longitudinal study of the relationship between serum 25-OH vitamin D and parathyroid hormone (PTH) concentrations in older women. *Proc Nutr Soc* **60**:159A.
 43. Harris S, Dawson-Hughes B 1992 Rates of change in bone mineral density of the spine, heel, femoral neck and radius in healthy postmenopausal women. *Bone Miner* **17**:87–95.
 44. Hannan MT, Felson DT, Dawson-Hughes B, Tucker KL, Cupples LA, Wilson PWF, Kiel DP 2000 Risk factors for longitudinal bone loss in elderly men and women: The Framingham Osteoporosis Study. *J Bone Miner Res* **15**:710–720.
 45. Lesser GT 2006 Calcium plus vitamin D and the risk of fractures. *N Engl J Med* **354**:2285–2286. (Letter)
 46. Department of Health 1998 Nutrition and Bone Health: With Particular Reference to Calcium and Vitamin D. Report on Health and Social Subjects no. 49. H. M. Stationery Office, London, UK.
 47. Dawson-Hughes B, Dallal GE, Krall EA, Harris S, Sokoll LJ, Falconer G 1991 Effect of vitamin D supplementation on wintertime and overall bone loss in healthy postmenopausal women. *Ann Intern Med* **115**:505–512.
 48. McMurdo MET, Mole PA, Paterson CR 1997 Controlled trial of weight bearing exercise in relation to bone density and falls. *BMJ* **314**:569.
 49. Caraballo PJ, Gabriel SE, Castro MR, Atkinson EJ, Melton LJ III 1999 Changes in bone density after exposure to oral anti-coagulants: A meta-analysis. *Osteoporos Int* **9**:441–448.
 50. Lips P, Duong T, Oleksik A, Black D, Cummings S, Cox D, Nickelsen T 2001 A global study of vitamin D status and parathyroid function in postmenopausal women with osteoporosis: Baseline data from the multiple outcomes of raloxifene evaluation clinical trial. *J Clin Endocrinol Metab* **86**:1212–1221.
 51. Ooms ME, Roos JC, Bezemer PD, van der Vijgh WJF, Bouter LM, Lips P 1995 Prevention of bone loss by vitamin D supplementation in elderly women: A randomized double-blind trial. *J Clin Endocrinol Metab* **80**:1052–1058.
 52. Thane CW, Paul AA, Bates CJ, Bolton-Smith C, Prentice A, Shearer MJ 2002 Intake and sources of phyloquinone (vitamin K₁): Variation with socio-demographic and lifestyle factors in a national sample of British elderly people. *Br J Nutr* **87**:605–613.
 53. Food and Nutrition Board Institute of Medicine 2001 Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc. National Academy Press, Washington, DC, USA.
 54. Binkley NC, Krueger DC, Engelke JA, Foley AL, Suttie JW 2000 Vitamin K supplementation reduces serum concentrations of under- γ -carboxylated osteocalcin in healthy young and elderly adults. *Am J Clin Nutr* **72**:1523–1528.
 55. Binkley NC, Krueger DC, Kawahara TN, Engelke JA, Chapell RJ, Suttie JW 2002 A high phyloquinone intake is required to achieve maximal osteocalcin γ -carboxylation. *Am J Clin Nutr* **76**:1055–1060.
 56. Schurgers LJ, Geleijnse JM, Grobbee DE, Pols HAP, Hofman A, Witteman JCM, Vermeer C 1999 Nutritional intake of vitamins K₁ (phyloquinone) and K₂ (menaquinone) in The Netherlands. *J Nutr Environ Med* **9**:115–122.
 57. Cockayne S, Adamson J, Lanham-New S, Shearer MJ, Gilbody S, Torgerson DJ 2006 Vitamin K and the prevention of fractures: Systematic review and meta-analysis of randomized controlled trials. *Arch Intern Med* **166**:1256–1261.
 58. Sato Y, Honda Y, Kuno H, Oizumi K 1998 Menatetranone ameliorates osteopenia in disuse-affected limbs of vitamin D- and K-deficient stroke patients. *Bone* **23**:291–296.
 59. Shiomi S, Nishiguchi S, Kubo S, Tamori A, Habu D, Takeda T, Ochi H 2002 Vitamin K₂ (menatetranone) for bone loss in patients with cirrhosis of the liver. *Am J Gastroenterol* **97**:978–981.
 60. Hara K, Akiyama Y, Nakamura T, Murota S, Morita I 1995 The inhibitory effect of vitamin K₂ (menatetranone) on bone resorption may be related to its side chain. *Bone* **16**:179–184.
 61. Ichikawa T, Horie-Inoue K, Ikeda K, Blumberg B, Inoue S 2006 Steroid and xenobiotic receptor SXR mediates vitamin K₂-activated transcription of extracellular matrix-related genes and collagen accumulation in osteoblastic cells. *J Biol Chem* **281**:16927–16934.
 62. Thijssen HHW, Vervoort LMT, Schurgers LJ, Shearer MJ 2006 Menadione is a metabolite of oral vitamin K. *Br J Nutr* **95**:260–266.
 63. Liu G, Peacock M 1998 Age-related changes in serum undercarboxylated osteocalcin and its relationships with bone density, bone quality, and hip fracture. *Calcif Tissue Int* **62**:286–289.
 64. Booth SL, Lichtenstein AH, O'Brien-Morse M, McKeown NM, Wood RJ, Saltzman E, Gundberg CM 2001 Effects of a hydrogenated form of vitamin K on bone formation and resorption. *Am J Clin Nutr* **74**:783–790.

Address reprint requests to:
 Martin J Shearer, PhD
 Vitamin K Research Unit
 Centre for Haemostasis and Thrombosis
 1st Floor North Wing
 St Thomas' Hospital
 Lambeth Palace Road
 London SE1 7EH, UK
 E-mail: martin.shearer@gstt.nhs.uk

Received in original form April 25, 2006; revised form December 28, 2006; accepted January 19, 2007.