

Cancer Epidemiology, Biomarkers & Prevention



No Link between Breast Cancer and Meningioma: Results from a Large Monoinstitutional Retrospective Analysis

Carmen Criscitiello, Davide Disalvatore, Michele Santangelo, et al.

Cancer Epidemiol Biomarkers Prev 2014;23:215-217. Published OnlineFirst October 28, 2013.

Updated version Access the most recent version of this article at:
doi:[10.1158/1055-9965.EPI-13-1041](https://doi.org/10.1158/1055-9965.EPI-13-1041)

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.

Null Results in Brief

No Link between Breast Cancer and Meningioma: Results from a Large Monoinstitutional Retrospective Analysis

Carmen Criscitiello^{1,2}, Davide Disalvatore³, Michele Santangelo⁴, Nicole Rotmensz³, Barbara Bazolli³, Patrick Maisonneuve³, Aron Goldhirsch¹, and Giuseppe Curigliano^{1,2}

Abstract

Background: The etiology of meningioma is largely unknown, although breast cancer has been suggested to play a role.

Methods: A monoinstitutional, retrospective analysis was performed at European Institute of Oncology on 12,330 patients with breast cancer. The cumulative incidence of meningioma was estimated using the Kaplan–Meier method and the log-rank test was used to assess differences between groups.

Results: In total, 33 patients with meningioma were identified from a study population of 12,330, with a 10-year cumulative incidence of meningioma of 0.37%. We did not find a significantly increased risk of meningioma among patients with breast cancer or an association between the hormonal receptor status and the risk of meningioma ($P = 0.65$).

Conclusions: Our results do not support a role of breast cancer or endocrine treatments in meningioma development.

Impact: This analysis adds new information on a debated topic. *Cancer Epidemiol Biomarkers Prev*; 23(1); 215–7. ©2013 AACR.

Introduction

Meningiomas are typically benign tumors arising from the meningeothelial cells of the arachnoid membrane covering the brain and spinal cord. The etiology of meningiomas is largely unknown. There is evidence to suggest that breast cancer may play a role in the development of meningiomas, including the observation that these tumors occur more commonly in women (1, 2). Further evidence for a possible role for hormones in brain tumor etiology includes the presence of progesterone (PgR), estrogen (ER), and androgen receptors in some meningiomas, suggesting a link between breast cancer and meningiomas. Immunohistochemical studies have shown that about 70% of meningiomas express PgR and about 30% express ER, thus corroborating the hypothesis that meningioma is a hormone-sensitive tumor (3, 4). Our study was carried out to explore the possible association between breast cancer, endocrine therapy, and the risk of developing meningioma in a large monoinstitutional population of patients with breast cancer. We focused on endo-

crine-responsive tumors to find a potential link between these tumors and endocrine therapy.

Materials and Methods

Data from 12,330 women with early-stage breast cancer who underwent surgery at the European Institute of Oncology (IEO) in Milan between 1994 and 2011 were retrieved from two institutional databases: the Breast Cancer database, which contains information on patients with breast cancer operated at IEO, and the Tumor Registry. Patients who received neoadjuvant treatment, patients with a previous history of cancer, or patients with metastatic breast cancer at diagnosis were excluded from the analysis. Breast cancer–related data on age, menopausal status, surgery, histopathology (Ki-67 expression, ER/PgR expression, and HER2 status) and systemic treatment were analyzed. Usually, patients were followed up with physical examination every 6 months, annual mammography, breast ultrasound, and blood tests every 6 to 12 months, and additional evaluations when clinically indicated. Whenever possible, the health status of women who did not come back for scheduled follow-up visits for more than 1 year was obtained by phone contact. Institutional Review Board approved this retrospective analysis. The primary endpoint was the cumulative incidence of meningioma (cum-inc ME), calculated from breast surgery to meningioma diagnosis. The cum-inc ME was estimated using the Kaplan–Meier method. The log-rank test was used to assess differences between groups. All analyses were carried out with the SAS software 9.3 (SAS Institute).

Authors' Affiliations: ¹Breast Cancer Program, ²Division of Early Drug Development, ³Division of Epidemiology and Biostatistics, European Institute of Oncology, Milan; and ⁴Department of Advanced Biomedical Sciences, Operative Unit of General Surgery and Transplants, University Naples Federico II, Napoli, Italy

Corresponding Author: Carmen Criscitiello, Division of Early Drug Development, Istituto Europeo di Oncologia, Via Ripamonti 435, 20141, Milano, Italy. Phone: 39-0294-3742-689; Fax: 39-0294-379-224; E-mail: carmen.criscitiello@ieo.it

doi: 10.1158/1055-9965.EPI-13-1041

©2013 American Association for Cancer Research.

Table 1. Association between patients' characteristics and meningioma

Variable	All (% column)	Number of meningioma (% row)	10-year cumulative incidence (95% CI)	P
Number	12,330 (100)	33 (0.3)	0.37 (0.26–0.55)	
Age at surgery				0.24
<35	514 (4.2)	0 (0.0)	0.00 (0.00–0.00)	
35–50	5,005 (40.6)	13 (0.3)	0.38 (0.21–0.70)	
51–65	4,728 (38.3)	12 (0.3)	0.29 (0.16–0.53)	
>65	2,083 (16.9)	8 (0.4)	0.73 (0.32–1.65)	
ER				0.18
ER = 0	1,722 (14.0)	7 (0.4)	0.59 (0.23–1.49)	
ER > 0	10,608 (86.0)	26 (0.2)	0.34 (0.22–0.51)	
PgR				0.60
PgR = 0	3,015 (24.5)	7 (0.2)	0.25 (0.11–0.55)	
PgR > 0	9,315 (75.5)	26 (0.3)	0.42 (0.27–0.64)	
ER/PgR				0.65
Not expressed (Both 0)	1,655 (13.4)	5 (0.3)	0.28 (0.11–0.75)	
Expressed (ER > 0 or PgR > 0)	10,675 (86.6)	28 (0.3)	0.38 (0.25–0.58)	
HER2				0.08
Not expressed	9,557 (77.5)	21 (0.2)	0.36 (0.23–0.56)	
Intense and complete	1,655 (13.4)	8 (0.5)	0.51 (0.24–1.08)	
Ki-67				0.32
<14%	3,416 (27.7)	11 (0.3)	0.44 (0.24–0.83)	
≥14%	8,902 (72.2)	21 (0.2)	0.33 (0.20–0.54)	
Subtype 2013				0.08
LUM A PgR > 20%	2,405 (19.5)	8 (0.3)	0.56 (0.27–1.12)	
LUM B Ki-67 ≥ 14% or PgR ≤ 20%	7,183 (58.3)	15 (0.2)	0.33 (0.19–0.57)	
LUM B HER2 POS	1,086 (8.8)	4 (0.4)	0.38 (0.12–1.19)	
HER2+	630 (5.1)	4 (0.6)	0.71 (0.27–1.88)	
TN	1,024 (8.3)	0 (0.0)	0.00 (0.00–0.00)	
Menopausal status				0.08
Premenopausal	5,769 (46.8)	11 (0.2)	0.28 (0.14–0.55)	
Postmenopausal	6,561 (53.2)	22 (0.3)	0.46 (0.29–0.72)	
Hormone therapy performed				0.53
No	2,061 (16.7)	7 (0.3)	0.35 (0.16–0.79)	
Yes	9,947 (80.7)	26 (0.3)	0.38 (0.25–0.59)	
Hormone therapy				0.054
Tamoxifen	7,037 (75.2)	17 (0.2)	0.31 (0.18–0.52)	
Aromatase inhibitor	2,315 (24.8)	7 (0.3)	0.74 (0.32–1.72)	
Chemotherapy				0.98
No	7,099 (57.6)	18 (0.3)	0.35 (0.21–0.57)	
Yes	4,932 (40.0)	15 (0.3)	0.41 (0.23–0.73)	
CT/HT				0.66
Nil	509 (4.1)	3 (0.6)	0.75 (0.24–2.31)	
ET	6,520 (52.9)	15 (0.2)	0.32 (0.19–0.55)	
CT	1,521 (12.3)	4 (0.3)	0.22 (0.07–0.69)	
ET-CT	3,384 (27.4)	11 (0.3)	0.47 (0.24–0.90)	

Abbreviations: ET, endocrine therapy; CT, chemotherapy; HT, hormone therapy; TN, triple-negative.

Results

From January 1994 to December 2011, we identified a total of 12,330 consecutive patients with early-stage breast cancer. At a median follow-up of 7 years, 33 patients with

meningioma were identified from a study population of 12,330. Table 1 describes selected demographic, clinical, and pathologic characteristics of the study population at the time of breast cancer surgery, systemic treatments

administered, number of meningiomas, and 10-year cumulative incidence (ME). Considering the entire population, the 10-year cumulative incidence (ME) was 0.37% [95% confidence interval (CI), 0.26%–0.55%]. No association was found between ER/PgR and the risk of meningioma. No association was found if we matched analysis coupling ER and PgR ($P = 0.65$) or considering them separately (ER, $P = 0.18$ and PgR, $P = 0.60$). Patients who received aromatase inhibitors had a trend of higher risk to develop meningioma compared with patients who received tamoxifen, with a borderline statistical significance ($P = 0.054$). No other significant association was observed in the analyzed variables.

Discussion

This large, retrospective study evaluated more than 12,000 patients with breast cancer with an average follow-up period of 7 years, and identified a total of 33 individuals with an incident case of meningioma. There was no significant association between meningioma development and breast cancer. Use of aromatase inhibitors conferred a trend of increased risk of meningioma in patients with breast cancer, although no statistical significance was observed. The link between breast cancer and meningioma has been reported in literature, with discordant results. What is behind this association is quite unclear, most likely being a hormonal etiology. Hormone receptors are involved in both tumor types. In meningiomas, PgR is more prevalent and more biologically active than ER. To ascertain a link between hormonal exposure and development of meningioma, we focused on endocrine-responsive breast cancer and on patients receiving endocrine therapy. In our cohort, the 10-year cumulative incidence (ME) was 0.37% (95% CI, 0.26–0.55), similar to that reported by Rao and colleagues (1). Our study has several limitations.

Despite the large cohort, a major limitation of this study is the small number of meningioma cases recorded with exposure to endocrine therapy. However, it has to be acknowledged that the screening for meningioma is not routinely conducted in patients with breast cancer. Also, the endocrine therapy given to ER-positive patients may have influenced the development of meningioma during this period. Another limitation is related to the retrospective analysis. An important strength is that the records of potential cases were manually reviewed and the case status was validated by a data-monitoring system with a confirmation rate of 100%. In summary, this study, in contrast with previous cohort studies, found no association between development of meningioma and breast cancer.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: C. Criscitiello, D. Disalvatore, M. Santangelo, N. Rotmensz, P. Maisonneuve, G. Curigliano

Development of methodology: C. Criscitiello, D. Disalvatore, M. Santangelo, G. Curigliano

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): C. Criscitiello, D. Disalvatore, N. Rotmensz, G. Curigliano

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): C. Criscitiello, D. Disalvatore, M. Santangelo, N. Rotmensz, A. Goldhirsch, G. Curigliano

Writing, review, and/or revision of the manuscript: C. Criscitiello, D. Disalvatore, M. Santangelo, P. Maisonneuve, A. Goldhirsch, G. Curigliano

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): C. Criscitiello, D. Disalvatore, N. Rotmensz, B. Bazolli

Study supervision: C. Criscitiello, M. Santangelo, G. Curigliano

Received October 9, 2013; accepted October 17, 2013; published OnlineFirst October 28, 2013.

References

1. Rao G, Giordano SH, Liu J, McCutcheon IE. The association of breast cancer and meningioma in men and women. *Neurosurgery* 2009;65:483–9.
2. Schoenberg BS, Christine BW, Whisnant JP. Nervous system neoplasms and primary malignancies of other sites. The unique association between meningiomas and breast cancer. *Neurology* 1975;25:705–12.
3. Carroll RS, Zhang J, Black PM. Expression of estrogen receptors alpha and beta in human meningiomas. *J Neurooncol* 1999;42:109–16.
4. Hsu DW, Efird JT, Hedley-Whyte ET. Progesterone and estrogen receptors in meningiomas: prognostic considerations. *J Neurosurg* 1997;86:113–20.