

JAMA. 2021 Jan 5; 325(1): 87-89.

PMCID: PMC7786240

Published online 2020 Dec 7. doi: 10.1001/jama.2020.20991; 10.1001/jama.2020.20991

PMID: [33284323](#)

## Changes in Preterm Birth Phenotypes and Stillbirth at 2 Philadelphia Hospitals During the SARS-CoV-2 Pandemic, March-June 2020

[Sara C. Handley](#), MD, MSCE,<sup>1</sup> [Anne M. Mullin](#), BS,<sup>2</sup> [Michal A. Elovitz](#), MD,<sup>2</sup> [Kristin D. Gerson](#), MD, PhD,<sup>2</sup> [Diana Montoya-Williams](#), MD,<sup>1</sup> [Scott A. Lorch](#), MD, MSCE,<sup>1</sup> and [Heather H. Burris](#), MD, MPH<sup>✉1</sup>

<sup>1</sup>Division of Neonatology, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania

<sup>2</sup>Maternal and Child Health Research Center, University of Pennsylvania Perelman School of Medicine, Philadelphia

✉Corresponding author.

### Article Information

**Corresponding Author:** Heather H. Burris, MD, MPH, Biomedical Research Bldg II/III, Room 1352, 421 Curie Blvd, Philadelphia, PA 19104-6160 ([burrish@email.chop.edu](mailto:burrish@email.chop.edu)).

**Accepted for Publication:** October 6, 2020.

**Published Online:** December 7, 2020. doi:10.1001/jama.2020.20991

**Author Contributions:** Ms Mullin and Dr Burris had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Dr Handley and Ms Mullin contributed equally.

*Concept and design:* Handley, Mullin, Elovitz, Montoya-Williams, Lorch, Burris.

*Acquisition, analysis, or interpretation of data:* All authors.

*Drafting of the manuscript:* Handley, Mullin, Burris.

*Critical revision of the manuscript for important intellectual content:* Mullin, Elovitz, Gerson, Montoya-Williams, Lorch, Burris.

*Statistical analysis:* Handley, Mullin, Burris.

*Administrative, technical, or material support:* Elovitz, Gerson.

*Supervision:* Elovitz, Lorch, Burris.

**Conflict of Interest Disclosures:** Dr Lorch reported receiving grants from the National Institutes of Health outside the submitted work. Dr Burris reported receiving grants from March of Dimes during the conduct of the study. No other disclosures were reported.

**Funding/Support:** This work was supported by the Department of Pediatrics of Children's Hospital of Philadelphia.

**Role of the Funder/Sponsor:** The funder had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Received 2020 Aug 10; Accepted 2020 Oct 6.

[Copyright](#) 2020 American Medical Association. All Rights Reserved.

---

This study uses data from the GeoBirth pregnancy cohort of all births in 2 hospitals in Philadelphia to examine whether rates of preterm birth, spontaneous preterm birth, medically indicated preterm birth, and stillbirth have changed during the SARS-CoV-2 pandemic compared with prepandemic rates.

---

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic has had far-reaching implications, including changes in societal stressors and health care delivery, which may alter preterm birth risk. Previous studies in the US regarding SARS-CoV-2 in pregnancy focused on associations of SARS-CoV-2 infection with cesarean delivery, neonatal transmission, preterm birth, and stillbirth.<sup>1</sup> In a relatively homogeneous Danish population, Hedermann et al<sup>2</sup> reported a decrease in preterm birth during the pandemic among uninfected patients. Given differences in preterm birth across populations,<sup>3</sup> we examined a diverse urban cohort in the US to determine if preterm birth, spontaneous preterm birth, medically indicated preterm birth, and stillbirth rates have changed during the SARS-CoV-2 pandemic.

## Methods

---

GeoBirth is a curated pregnancy cohort of all births in 2 Penn Medicine hospitals in Philadelphia ongoing since 2008 (approximately 9000 births per year), in which each preterm birth (<37 weeks' gestation) is manually classified by 2 independent blinded reviewers, with further adjudication by a third reviewer when there is nonconcordance. Preterm birth phenotypes are categorized as spontaneous preterm birth (eg, preterm labor, spontaneous rupture of membranes) or medically indicated preterm birth (eg, clinician initiated due to a maternal or fetal health condition, such as preeclampsia or intrauterine growth restriction). Stillbirth is defined as intrauterine fetal demise at 20 weeks' gestation or greater. We compared preterm birth, spontaneous preterm birth, medically indicated preterm birth, and stillbirth rates among singleton pregnancies during the pandemic period (March-June 2020) with the same months in 2018 and 2019 (prepandemic) to account for seasonality using a 2-tailed Fisher exact test with a significance threshold of  $P < .05$  using R, version 4.0.2. We used marginal effects models to calculate absolute risk differences between the 2 epochs adjusting for birth month, age, parity, body mass index, race/ethnicity, marital status, smoking, and insurance status. We also performed analyses stratified by race/ethnicity because of persistent preterm birth disparities.<sup>3</sup> This study was approved by the University of Pennsylvania Institutional Review Board with a waiver of informed consent.

## Results

---

There were a total of 8867 singleton, live-born deliveries in March through June of 2018, 2019, and 2020 (42% non-Hispanic Black, 37% non-Hispanic White, and 21% other race/ethnicity); 2992 deliveries occurred during the pandemic period, including 283 preterm births (135 spontaneous and 148 medically indicated) and 15 stillbirths. Prepandemic and pandemic birth outcomes were as follows: 10.5% vs 9.5% of deliveries were preterm births (adjusted difference,  $-1.1\%$  [95% CI,  $-2.4\%$  to  $0.2\%$ ]), 5.7% vs 4.7% were spontaneous preterm births (adjusted difference,  $-0.8\%$  [95% CI,  $-1.8\%$  to  $0.2\%$ ]), 5.4% vs 5.2% were medically indicated preterm births (adjusted difference,  $-0.3\%$  [95% CI,  $-1.4\%$  to  $0.6\%$ ]), and 5.4 per 1000 births vs 5.0 per 1000 births were stillbirths (adjusted difference,  $-0.03$  per 1000 births [95% CI,  $-0.34$  to  $0.29$ ]) (Table). Spontaneous preterm birth among non-Hispanic White patients declined during the pandemic (4.5% vs 2.9%; adjusted difference,  $-1.4\%$  [95% CI,  $-2.8\%$  to  $-0.1\%$ ]); no other racial/ethnic groups had significant changes in outcomes. However, no significant interaction was detected between race/ethnicity and epoch with spontaneous preterm birth ( $P = .09$  for interaction).

In the 2 hospitals, universal SARS-CoV-2 testing began on April 1, 2020, and April 13, 2020. Among 86 patients with test results positive for SARS-CoV-2, the preterm birth rate was 11.6% ( $n = 10$ ; 6 spontaneous and 4 medically indicated preterm births) and there was 1 stillbirth.

## Discussion

---

This study did not detect significant changes in preterm or stillbirth rates during the SARS-CoV-2 pandemic in a racially diverse urban cohort from 2 Philadelphia hospitals. Although these data allow for disaggregation of spontaneous and medically indicated preterm births, no differences in overall rates of these phenotypes were detected.

These findings differ from a Danish report of decreasing preterm birth rates<sup>2</sup> and higher stillbirth rates in a UK hospital<sup>4</sup> during the pandemic. The differences between studies may be due to differences in enforcement of lockdown orders, population heterogeneity, access to health care, or societal stressors.

Study limitations include examination of a single health system, short epochs, limited representation of other races/ethnicities, few stillbirths, and potential for change in delivery hospital choice during the pandemic.

## Notes

---

**Section Editor:** Jody W. Zylke, MD, Deputy Editor.

## References

---

1. Khoury R, Bernstein PS, Debolt C, et al.. Characteristics and outcomes of 241 births to women with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection at five New York City medical centers. *Obstet Gynecol.* 2020;136(2):273-282. doi: 10.1097/AOG.0000000000004025 [PubMed: 32555034] [CrossRef: 10.1097/AOG.0000000000004025]
2. Hedermann G, Hedley PL, Bækvad-Hansen M, et al.. Danish premature birth rates during the COVID-19 lockdown. *Arch Dis Child Fetal Neonatal Ed.* Published online August 11, 2020. doi: 10.1136/archdischild-2020-319990 [PMCID: PMC7421710] [PubMed: 32788391] [CrossRef: 10.1136/archdischild-2020-319990]
3. Martin JA, Hamilton BE, Osterman MJ, Driscoll AK. Births: final data for 2018. *Natl Vital Stat Rep.* 2019;68(13):1-47. [PubMed: 32501202]
4. Khalil A, von Dadelszen P, Draycott T, Ugwumadu A, O'Brien P, Magee L. Change in the incidence of stillbirth and preterm delivery during the COVID-19 pandemic. *JAMA.* 2020;324(7):705-706. doi: 10.1001/jama.2020.12746 [PMCID: PMC7435343] [PubMed: 32648892] [CrossRef: 10.1001/jama.2020.12746]

Table.

**Birth Outcomes by Race/Ethnicity Before (March-June 2018 and 2019) and During (March-June 2020) the Severe Acute Respiratory Syndrome Coronavirus 2 Pandemic in 2 Philadelphia Hospitals<sup>a</sup>**

<sup>a</sup>Race/ethnicity was based on patient self-identification during patient registration.  
<sup>b</sup>Calculated using a 2-tailed Fisher exact test.  
<sup>c</sup>Adjusted for month of birth, age, parity, body mass index, race/ethnicity (except in stratified models), marital status, smoking, and insurance status, calculated using marginal effects models. Presented as percentages for all birth outcomes except for stillbirth.  
<sup>d</sup>Preterm birth calculations exclude stillbirths.  
<sup>e</sup>Spontaneous preterm birth calculations exclude medically indicated preterm births and stillbirths.  
<sup>f</sup>Medically indicated preterm birth calculations exclude spontaneous preterm births and stillbirths.  
<sup>g</sup>Adjusted models did not converge due to small numbers of events.