

## Original Investigation

# Cholecalciferol Treatment to Reduce Blood Pressure in Older Patients With Isolated Systolic Hypertension

## The VitDISH Randomized Controlled Trial

Miles D. Witham, PhD; Rosemary J. G. Price, MSc; Allan D. Struthers, MD; Peter T. Donnan, PhD; Claudia-Martina Messow, PhD; Ian Ford, PhD; Marion E. T. McMurdo, MD

**IMPORTANCE** Observational data link low 25-hydroxyvitamin D levels to both prevalent blood pressure and incident hypertension. No clinical trial has yet examined the effect of vitamin D supplementation in isolated systolic hypertension, the most common pattern of hypertension in older people.

**OBJECTIVE** To test whether high-dose, intermittent cholecalciferol supplementation lowers blood pressure in older patients with isolated systolic hypertension.

**DESIGN** Parallel group, double-blind, placebo-controlled randomized trial.

**SETTING** Primary care clinics and hospital clinics.

**PARTICIPANTS** Patients 70 years and older with isolated systolic hypertension (supine systolic blood pressure >140 mm Hg and supine diastolic blood pressure <90 mm Hg) and baseline 25-hydroxyvitamin D levels less than 30 ng/mL were randomized into the trial from June 1, 2009, through May 31, 2011.

**INTERVENTIONS** A total of 100 000 U of oral cholecalciferol or matching placebo every 3 months for 1 year.

**MAIN OUTCOMES AND MEASURES** Difference in office blood pressure, 24-hour blood pressure, arterial stiffness, endothelial function, cholesterol level, insulin resistance, and b-type natriuretic peptide level during 12 months.

**RESULTS** A total of 159 participants were randomized (mean age, 77 years). Mean baseline office systolic blood pressure was 163/78 mm Hg. Mean baseline 25-hydroxyvitamin D level was 18 ng/mL. 25-Hydroxyvitamin D levels increased in the treatment group compared with the placebo group (+8 ng/mL at 1 year,  $P < .001$ ). No significant treatment effect was seen for mean (95% CI) office blood pressure (−1 [−6 to 4]/−2 [−4 to 1] mm Hg at 3 months and 1 [−2 to 4]/0 [−2 to 2] mm Hg overall treatment effect). No significant treatment effect was evident for any of the secondary outcomes (24-hour blood pressure, arterial stiffness, endothelial function, cholesterol level, glucose level, and walking distance). There was no excess of adverse events in the treatment group, and the total number of falls was nonsignificantly lower in the group receiving vitamin D (36 vs 46,  $P = .24$ ).

**CONCLUSIONS AND RELEVANCE** Vitamin D supplementation did not improve blood pressure or markers of vascular health in older patients with isolated systolic hypertension.

**TRIAL REGISTRATION** isrctn.org Identifier: ISRCTN92186858

*JAMA Intern Med.* doi:10.1001/jamainternmed.2013.9043  
Published online August 12, 2013.

← Invited Commentary

+ Supplemental content at  
jamainternalmedicine.com

**Author Affiliations:** Aging and Health, University of Dundee, Dundee, United Kingdom (Witham, Price, McMurdo); Department of Clinical Pharmacology, University of Dundee, Dundee, United Kingdom (Struthers); Dundee Epidemiology and Biostatistics Unit, Medical Research Institute, University of Dundee, Dundee, United Kingdom (Donnan); Robertson Centre for Biostatistics, University of Glasgow, Glasgow, United Kingdom (Messow, Ford).

**Corresponding Author:** Miles D. Witham, PhD, Ninewells Hospital, Dundee DD1 9SY, United Kingdom (m.witham@dundee.ac.uk).

Isolated systolic hypertension (ISH) is the most common form of hypertension in older people,<sup>1-4</sup> is associated with an increased incidence of both cerebrovascular and cardiovascular events, and is a driver of left ventricular hypertrophy—itself a risk factor for sudden cardiac death and heart failure.<sup>5,6</sup> Antihypertensive therapies improve cardiovascular outcomes in ISH,<sup>7-9</sup> but few studies have enrolled typical older patients with multimorbidity.<sup>3</sup> The recent Hypertension in the Very Elderly trial<sup>10</sup> confirms that the benefits of blood pressure reduction extend to selected very old patients, but despite this evidence, many older patients are undertreated in part because of the treatment constraints imposed by the presence of multimorbidity and polypharmacy and in part because of the adverse effects inherent in many antihypertensive medications, including dizziness and falls. New, simple, safe, and inexpensive ways of treating blood pressure in older people are highly desirable.

Low 25-hydroxyvitamin D (25OHD) levels are strongly associated with both prevalent and incident hypertension<sup>11-13</sup> and cardiovascular disease<sup>14,15</sup> in observational studies, and meta-analyses of previous intervention trials<sup>16</sup> suggest that vitamin D may reduce blood pressure in selected patient groups. Trials to date have not evaluated the effects of vitamin D on ISH. We conducted the VitDISH (Vitamin D in Isolated Systolic Hypertension) randomized controlled trial to test the effect of intermittent, high-dose cholecalciferol on blood pressure and markers of vascular function in older patients with ISH.

## Methods

### Design and Participants

Research ethics approval was obtained from the Fife and Forth Valley National Health Service Research Ethics Committee (08/S0501/90). Clinical trials authorization was obtained from the UK Medicines and Healthcare Regulatory Authority (European Union Drug Regulating Authorities Clinical Trials No. 2008-004534-24). The trial sponsor was the University of Dundee, and the trial was registered at [www.controlled-trials.com](http://www.controlled-trials.com) (ISRCTN92186858). The complete protocol is available from the authors.

The study was a randomized, double-blind, placebo-controlled, parallel-group trial. We studied community-dwelling patients 70 years and older with ISH. Inclusion criteria were as follows: age of 70 years or older, 25OHD level less than 30 ng/mL (to convert to nanomoles per liter, multiply by 2.496), and office systolic blood pressure greater than 140 mm Hg. Full exclusion criteria and detailed trial methods are given in the eMethods in the Supplement. Patients were recruited via 3 routes: from the community via primary care practices, via an article in the local newspaper about the research study, and via secondary care clinics (cardiovascular and medicine for the elderly).

### Intervention

Participants enrolling in the study were allocated to intervention or placebo in a 1:1 ratio. Stratified randomization

was performed using a minimization algorithm, administered by the Robertson Centre for Biostatistics (Glasgow Clinical Trials Unit, University of Glasgow, United Kingdom) using a telephone-based system to conceal study allocation from investigators and participants. Minimization variables used were baseline 25OHD level above or below 20 ng/mL, baseline systolic blood pressure above or below 160 mm Hg, baseline age older or younger than 80 years, and the presence of diabetes mellitus. Identical, masked medication bottles were used; patients were observed ingesting 100 000 U of oral cholecalciferol (Vigantol oil; Merck KgAA) or matching placebo (Mygliol oil; Merck KgAA) after completion of baseline and 3-, 6-, and 9-month assessments.

### Primary Outcome Measures

All outcome measures were performed by researchers who were masked to treatment allocation. Outcomes, including venepuncture for 25OHD levels, were all measured on the same day, with trial medication dosing occurring the next day. The primary outcome measure was prespecified as the between-group difference in office blood pressure at 3 months analyzed using all participants regardless of whether they received their allocated intervention.

### Secondary Outcome Measures

#### 24-Hour Blood Pressure

The 24-hour blood pressure was measured using an ambulatory monitor (Meditech ABPM-04; Meditech Ltd) at 0, 3, 6, 9, and 12 months. The mean of the 24-hour recording was used as the main measure; daytime and nighttime readings were also analyzed separately. Sensitivity analyses using a quality threshold (minimum of 14 daytime and 7 nighttime readings per the British Hypertension Society guidance<sup>17</sup>) were also performed.

#### Soluble Markers of Cardiovascular Risk

Soluble markers were collected at baseline and 3 and 12 months after a minimum 6-hour fast. Samples were collected for measurement of b-type natriuretic peptide, high-sensitivity C-reactive protein, and insulin measurement. Insulin and glucose measurements were used to construct a homeostasis model assessment of insulin resistance<sup>18</sup> using the following formula: (fasting glucose level × fasting insulin level)/405.

#### Endothelial Function

Endothelial function was assessed at 0, 3, and 12 months by measuring flow-mediated dilatation of the brachial artery in response to hyperemia (endothelium-dependent vasodilation) after 5 minutes of forearm cuff occlusion using standard techniques.<sup>19</sup> Mean diameter was measured during baseline acquisition and was compared with the maximum diameter achieved after cuff deflation. Flow-mediated dilatation was then expressed as the percentage change from baseline diameter.

#### Pulse Wave Velocity

Carotid-radial pulse wave velocity and augmentation index were measured using the Sphygmocor applanation tonom-

etry system (AtCor Medical) at 0, 3, and 12 months to give indices of arterial stiffness. Augmentation index is presented normalized to a heart rate of 75/min.

### Other Biochemical Measurements

We measured the levels of glucose, total cholesterol, low-density lipoprotein cholesterol, high-density lipoprotein cholesterol, triglycerides, serum albumin, calcium, parathyroid hormone (PTH), and 25OHD on fasting blood samples. Screening samples were analyzed in small batches before enrollment in the trial to ascertain trial eligibility; follow-up samples were stored and batch analyzed at the end of the trial. Calcium intake was assessed at baseline using the Scottish Collaborative Group Food Frequency Questionnaire.<sup>20</sup>

### Exercise Capacity and Falls

Exercise capacity was determined by 6-minute walks at 0, 6, and 12 months.<sup>21</sup> The walk test was performed on a 25-m flat course, using standardized encouragement every 2 minutes. Fall frequency was prospectively recorded by patients using monthly diaries.

### Statistical Analysis

A 2-sided  $P < .05$  was considered significant for all analyses. For each outcome measure, repeated-measures, mixed-effects analyses were conducted, with estimation and testing of main effects of treatment allocation ( $t$  test) and group  $\times$  time interaction ( $F$  tests). Analyses for outcome measures at individual time points, including the primary outcome measures, were based on analysis of covariance, adjusting for baseline values of the response variable and levels of the stratification variables. Analyses were based on a modified intent-to-treat population (all participants with baseline and follow-up values of the response variable being analyzed). We calculated unadjusted models and models adjusted for baseline values of the outcome measure under study, along with baseline 25OHD level, baseline blood pressure, the presence of diabetes, and age. We also adjusted for use of thiazide diuretics given the known effects of these medications on calcium metabolism and use of statins given their known interaction with vascular health and vitamin D metabolism. Sensitivity analyses were conducted using multiple imputation to account for missing data and excluding patients with changes in antihypertensive medication type or dose for blood pressure analyses. On the basis of blood pressure data from a previous vitamin D intervention trial for hypertension,<sup>22</sup> we powered the study to detect a 7-mm Hg between-group difference in systolic blood pressure change between baseline and 3 months, assuming an SD change of 18 mm Hg. To detect this change with 80% power at an  $\alpha$  level of .05 required 73 patients per group. We originally anticipated a dropout rate of 20% and, thus, aimed to recruit 180 patients into the trial.

## Results

Patients from 13 primary care practices were invited to express interest in the trial; a total of 341 agreed to attend

the screening, and of these, 159 were randomized into the trial from June 1, 2009, through May 31, 2011. The last visit of the last patient occurred at the end of May 2012. Participant flow through the trial is shown in **Figure 1**. The 159 participants randomized had a mean age of 76.8 years, a mean office blood pressure of 163/78 mm Hg, and only minor differences in baseline characteristics between the 2 groups (**Table 1**). Recruitment was terminated at this point because the dropout rate was lower than projected and the trial was on course to obtain the required number of completing participants.

### Effect of Intervention on 25OHD Levels

Mean 25OHD levels increased from a baseline level of 18 ng/mL to 28 ng/mL at 6 months in the intervention group before plateaus. A small (3-ng/mL) increase was seen by 6 months in the placebo group; the overall effect estimate of the intervention during the 12-month study period was equivalent to a 7-ng/mL increase in 25OHD levels in the intervention group. Details are shown in **Figure 2**. Adherence to study medication was high; 301 of 303 planned doses (99%) were ingested in the intervention group, and 285 of 294 planned doses (97%) were ingested in the placebo group. Missing doses were withheld because of elevated calcium level or suspected medication adverse effects; no participant declined to ingest medication at any time point.

### Primary Outcome Measure


Office systolic blood pressure was not statistically different between treatment and placebo groups at 3 months (**Table 2**). Office systolic blood pressure decreased by a mean (SD) of 2.7 (12.5) mm Hg in the treatment group between baseline and 3 months compared with a smaller mean (SD) decrease of 1.4 (14.9) mm Hg in the placebo group. The between-group treatment effect after adjusting for covariates was  $-0.7$  mm Hg (95% CI,  $-5.2$  to  $3.8$  mm Hg;  $P = .76$ ). Season of recruitment (May-October vs November-April, reflecting peak and trough 25OHD levels in population studies) did not change the effect of treatment on blood pressure at 3 months; between-group difference in systolic blood pressure change was  $-0.9$  mm Hg (95% CI,  $-5.3$  to  $3.5$  mm Hg) after adjustment for season. No participant changed antihypertensive therapy between baseline and 3 months.

### Secondary Outcome Measures

Details of blood pressure change in each group over time are given in **Table 2**. No significant between-group differences in blood pressure at any time point for either office blood pressure or daytime 24-hour blood pressure and no significant between-group differences on repeated measures analysis were found. Further analysis that excluded participants with a change in their antihypertensive medication (1 in the placebo group and 3 in the vitamin D group by 12 months) revealed no relevant change in these results. Daytime 24-hour blood pressure results were similar after excluding recordings that had fewer than 14 daytime readings.

**Table 3** lists the changes in other outcome measures at each time point. No significant change in serum calcium

Figure 1. CONSORT Diagram for Participant Flow Through the Trial



BP indicates blood pressure; eGFR, estimated glomerular filtration rate; and 25OHD, 25-hydroxyvitamin D.

levels was noted, and PTH levels decreased by a small (3.8 pg/mL) but significant amount in the intervention group relative to placebo.

### Subgroup Analyses

A series of hypothesis-generating post hoc interaction analyses were conducted. No significant interaction was found between baseline 25OHD level, baseline systolic blood pressure, dietary calcium intake, or PTH level (used as continuous variables) and the treatment effect of vitamin D on systolic blood pressure. A significant interaction was noted between angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB) use and treatment effect, with a greater treatment effect seen in those already taking ACE inhibitors or ARB medication ( $-4$  mm Hg vs  $+8$  mm Hg,  $P = .02$ ). No significant interaction was seen between the effect of treatment on diastolic blood pressure and baseline 25OHD level, ACE inhibitor or ARB use, baseline calcium intake, or baseline PTH level.

### Exercise Capacity and Falls

Exercise capacity as measured by a 6-minute walk did not increase by a clinically significant amount with vitamin D therapy; the between-group difference compared with baseline was 16 m (95% CI, 2 to 31 m;  $P = .03$ ) at 6 months and 12 m (95% CI,  $-4$  to 28 m;  $P = .13$ ) at 12 months. There were nonsignificantly fewer falls in the vitamin D group during the 12-month study period (36 vs 46 falls; hazard ratio, 0.77; 95% CI, 0.50 to 1.20;  $P = .24$ ); the time to first fall in each group was similar ( $P = .94$  by log-rank test) as was the number of participants in each group falling at least once (25 in the intervention group and 26 in the placebo group).

### Safety

Adverse events were similar in both trial arms, with no significant excess of adverse events in any subcategory of events (eTable in the Supplement). One participant in each group had an adjusted serum calcium level greater than 10.6 mg/dL (to convert to millimoles per liter, multiply by 0.25) during the


Table 1. Baseline Characteristics of Randomized Participants

Characteristic	Vitamin D Group (n = 80)	Placebo Group (n = 79)
Age, mean (SD), y	76.9 (4.8)	76.7 (4.5)
Male, No. (%)	40 (50)	42 (53)
BMI	28.5 (5.0)	27.9 (4.5)
Myocardial infarction, No. (%)	5 (6)	5 (6)
Angina, PTCA, or CABG, No. (%)	20 (25)	17 (22)
Peripheral vascular disease, No. (%)	5 (6)	7 (9)
Diabetes mellitus, No. (%)	11 (14)	11 (14)
Stroke or TIA, No. (%)	10 (12)	8 (10)
No. of medications, median (IQR)	4 (3 to 6)	5 (3 to 8)
No. of antihypertensive medications, median (IQR)	2 (1 to 2)	2 (1 to 2.5)
RAAS blockers, No. (%)	41 (51)	50 (63)
β-Blockers, No. (%)	23 (29)	26 (33)
Calcium channel blockers, No. (%)	32 (40)	35 (44)
Diuretics, No. (%)	37 (46)	31 (39)
Statins, No. (%)	41 (51)	46 (58)
25OHD, ng/mL	18 (6)	18 (6)
PTH, pg/mL	5.3 (2.9)	5.3 (1.9)
Serum-adjusted calcium, mg/dL	9.2 (0.28)	9.2 (0.32)
Creatinine, mg/dL	0.93 (0.23)	0.92 (0.25)
Office BP, mm Hg		
Systolic	163 (11)	162 (10)
Diastolic	78 (7)	77 (8)
Daytime BP, mean, mm Hg		
Systolic	136 (11)	133 (11)
Diastolic	71 (9)	67 (12)
Pulse wave velocity, m/s	8.8 (1.2)	8.7 (1.2)
Flow-mediated dilatation of brachial artery, No. (%)	5.1 (2.7)	5.1 (2.7)
Cholesterol, mg/dL		
Total	189 (46)	193 (42)
LDL	108 (39)	108 (39)
HDL	62 (15)	62 (15)
Triglycerides, mg/dL	115 (53)	115 (62)
Glucose, mg/dL	54 (13)	54 (16)
HOMA-IR index	5.5 (4.0)	4.5 (3.7)
BNP, median (IQR), pg/mL	21 (11-68)	44 (21-90)
CRP, median (IQR), mg/L	2.0 (1.0-4.2)	1.5 (0.8-3.4)
6-Minute walk distance, m	398 (94)	402 (105)
Daily mean calcium intake, mg	1127 (407)	1123 (372)

Abbreviations: BMI, body mass index (calculated as weight in kilograms divided by height in meters squared); BNP, b-type natriuretic peptide; BP, blood pressure; CABG, coronary artery bypass graft; CRP, C-reactive protein; HDL, high-density lipoprotein; HOMA-IR, homeostasis model assessment of insulin resistance; IQR, interquartile range; LDL, low-density lipoprotein; 25OHD, 25-hydroxyvitamin D; PTCA, percutaneous transluminal coronary angioplasty; PTH, parathyroid hormone; RAAS, renin-angiotensin-aldosterone system; TIA, transient ischemic attack.

SI conversion factors: To convert 25OHD to nanomoles per liter, multiply by 2.496; PTH to nanograms per liter, multiply by 1; calcium to millimoles per liter, multiply by 0.25; creatinine to micromoles per liter, multiply by 88.4; total cholesterol, LDL, and HDL to millimoles per liter, multiply by 0.0259; triglycerides to millimoles per liter, multiply by 0.0113; glucose to millimoles per liter, multiply by 0.0555; BNP to nanograms per liter, multiply by 1; and CRP to nanomoles per liter, multiply by 9.524.

Figure 2. Effect of Intervention on 25-Hydroxyvitamin D (25OHD) Levels



Error bars indicate SD.

study; neither participant had symptoms or required treatment. Similarly, 3 of 80 participants (4%) in the vitamin D group and 5 of 79 participants (6%) in the placebo group had an increase in creatinine level greater than 20% compared with baseline during the study.

## Discussion

Previous meta-analyses<sup>16,23,24</sup> have suggested that vitamin D supplementation produces a modest reduction in systolic blood pressure between 2 and 4 mm Hg. Selected groups of patients (eg, patients with type 2 diabetes and patients with elevated baseline blood pressure)<sup>25,26</sup> have demonstrated larger improvements in blood pressure with vitamin D supplementation of 8 to 13 mm Hg. Mixed results have been reported for the effects of vitamin D on endothelial function and b-type natriuretic peptide levels, although significant effects were reported in some studies,<sup>22,25-31</sup> and one recent trial did not find any improvement in arterial stiffness with vitamin D supplementation.<sup>32</sup> Previous trials have not found significant effects on lipid profile with vitamin D,<sup>33</sup> but in patients with low 25OHD levels, supplementation may enhance the effect of statins metabolized via the cytochrome 3A4 pathway.<sup>34</sup>

Several explanations are possible for the lack of effect of vitamin D seen in this study. First, vitamin D may have no significant effect on blood pressure. The overall effect size seen by combining small trials to date is modest, and it is possible that a real biological effect on blood pressure is not present.

A second possibility is that vitamin D may not reduce blood pressure in this specific patient group. Vitamin D may not reverse vascular stiffness and calcification during a 1-year period or might only act effectively at an earlier stage in the disease process. Hypertension in older people is characteristically not due to elevated renin levels (in contrast to younger patients); if vitamin D acts predominantly via the renin-angiotensin-aldosterone system as has been previously postulated,<sup>35-37</sup> this could explain the lack of effect seen, especially because 50% to 60% of patients were already taking ACE inhibitors or ARBs.



Table 2. Changes in Office BP at Each Time Point by Randomization Group

Variable	Systolic BP				Diastolic BP			
	Mean (SD)		Treatment Effect (95% CI)		Mean (SD)		Treatment Effect (95% CI)	
	Vitamin D	Placebo	Unadjusted	Adjusted	Vitamin D	Placebo	Unadjusted	Adjusted
Office blood pressure, mm Hg								
Baseline	163 (11)	162 (10)	... <sup>a</sup>	...	78 (7)	77 (8)	...	...
3 mo	161 (15)	161 (16)	-1 (-6 to 4)	-1 (-6 to 3)	77 (8)	78 (9)	-2 (-4 to 1)	-2 (-4 to 1)
6 mo	161 (16)	159 (16)	1 (-3 to 6)	1 (-3 to 6)	78 (9)	77 (9)	1 (-2 to 3)	1 (-2 to 3)
9 mo	164 (16)	161 (16)	3 (-2 to 7)	3 (-2 to 7)	78 (9)	77 (8)	1 (-2 to 3)	1 (-1 to 3)
12 mo	163 (18)	160 (15)	2 (-3 to 6)	2 (-3 to 7)	78 (9)	76 (8)	1 (-1 to 4)	2 (-1 to 4)
Overall treatment effect (95% CI)	...	...	1 (-2 to 4)	1 (-2 to 4)	...	...	0 (-2 to 2)	0 (-2 to 2)
P value for group × time interaction	...	...	.58	.54	...	...	.06	.03
Daytime ambulatory BP, mm Hg								
Baseline	136 (11)	133 (11)	...	...	69 (8)	68 (9)	...	...
3 mo	135 (11)	134 (9)	-1 (-4 to 2)	-1 (-4 to 2)	68 (8)	69 (8)	-1 (-3 to 1)	-1 (-2 to 1)
6 mo	134 (10)	132 (12)	-1 (-4 to 3)	0 (-4 to 3)	68 (8)	68 (9)	0 (-2 to 2)	1 (-1 to 2)
9 mo	136 (12)	132 (10)	1 (-2 to 4)	1 (-2 to 5)	68 (8)	68 (9)	0 (-2 to 2)	0 (-2 to 2)
12 mo	135 (12)	133 (12)	1 (-2 to 4)	1 (-2 to 5)	69 (9)	68 (9)	1 (-1 to 3)	1 (-1 to 3)
Overall treatment effect (95% CI)	...	...	0 (-3 to 2)	0 (-2 to 2)	...	...	0 (-2 to 1)	0 (-1 to 2)
P value for group × time interaction	...	...	.46	.36	...	...	.58	.45

Abbreviation: BP, blood pressure.

<sup>a</sup> Ellipses indicate data not applicable.Table 3. Secondary Outcome Measures: Adjusted Between-Group Treatment Effect at Each Time Point<sup>a</sup>

Outcome	Treatment Effect (95% CI)				
	Month 3	Month 6	Month 9	Month 12	Overall
FMD, %	0.0 (-0.9 to 0.9)	ND	ND	-0.7 (-1.7 to 0.2)	-0.4 (-1.1 to 0.4)
Pulse wave velocity, m/s	-0.1 (-0.5 to 0.3)	ND	ND	-0.2 (-0.6 to 0.2)	-0.2 (-0.5 to 0.2)
Cholesterol, mg/dL					
Total	-3 (-11 to 4)	0 (-8 to 7)	-3 (-11 to 4)	-5 (-12 to 3)	-3 (-9 to 2)
LDL	-2 (-8 to 3)	-1 (-7 to 5)	-3 (-9 to 3)	-5 (-11 to 1)	-3 (-7 to 2)
HDL	0 (-2 to 3)	0 (-3 to 2)	-1 (-3 to 2)	-2 (-4 to 1)	0 (-3 to 2)
Triglycerides, mg/dL	-7 (-19 to 5)	1 (-12 to 13)	-2 (-14 to 12)	-1 (-14 to 12)	-3 (-12 to 7)
Fasting glucose, mg/dL	-1 (-5 to 3)	ND	ND	0 (-3 to 4)	0 (-3 to 3)
BNP, pg/mL	3 (-27 to 32)	ND	ND	35 (5 to 65)	18 (-5 to 41)
HOMA-IR index	-0.4 (-1.3 to 0.4)	ND	ND	0.0 (-1.1 to 1.1)	-0.2 (-1.0 to 0.6)
CRP, mg/L	-0.5 (-3.0 to 2.0)	ND	ND	1.3 (-1.2 to 3.8)	0.4 (-1.7 to 2.5)

Abbreviations: BNP, b-type natriuretic peptide; CRP, C-reactive protein; FMD, flow-mediated dilatation; HDL, high-density lipoprotein; HOMA-IR, homeostasis model assessment of insulin resistance; LDL, low-density lipoprotein; ND, not done.

<sup>a</sup> Adjusted for baseline value of variable, age, 25-hydroxyvitamin D level, the presence of diabetes mellitus, systolic blood pressure, thiazide use, and statin use.

Although office blood pressure readings were all in the hypertensive range in this study, mean daytime blood pressures were markedly lower, with almost 50% of participants having a daytime mean systolic blood pressure less than 135 mm Hg. It is possible that a proportion of patients therefore had white-coat hypertension (ie, elevated blood pressure when measured by health care professionals but lower blood pressure at other times), but the known high prevalence of orthostatic hypotension in older hypertensive patients may also have contributed to this discrepancy; office blood pressures were recorded in the supine position, whereas most daytime ambulatory readings would be recorded in the seated or standing position.

A third possibility is that the dose used was insufficient or incorrect. Intermittent dosing might theoretically have different biological effects to daily dosing, and extrapolation from observational data has suggested that 25OHD levels higher than 30 ng/mL are required for optimal health.<sup>38</sup> The dose used in this study was sufficient to raise mean nadir levels from 18 to 28 ng/mL, and similar intermittent doses producing similar increments in 25OHD levels have caused marked reductions in blood pressure in some previous studies.<sup>22,25</sup>

A fourth possibility is that other permissive factors, for instance, adequate calcium intake, are required for vitamin D to produce a beneficial effect on blood pressure. Patients in this study had a high calcium intake (mean, >1 g/d). It is possible

that vitamin D supplementation has little extra effect in calcium-replete individuals; calcium is itself thought to have a slight antihypertensive effect,<sup>39</sup> although a possible deleterious role in precipitating cardiovascular events is the subject of ongoing controversy.<sup>40</sup> We found no evidence of an interaction between calcium intake and the degree of blood pressure lowering on subgroup analysis to support this hypothesis, however.

Cholecalciferol was well tolerated in this trial, with no increase in adverse events compared with placebo. High-dose cholecalciferol was associated with an increase in falls in a recent trial<sup>41</sup>; our vitamin D group had a reassuringly low number of falls, consistent with previous data.<sup>42</sup> A small improvement was seen in the 6-minute walk distance in keeping with previous trial results; however, the magnitude of the effect seen in this study was less than the minimum clinically important difference of 20 m for the 6-minute walk test.<sup>21</sup>

Our study has several strengths. The projected final evaluable sample size was reached, and the mean age (77 years) was considerably older than that seen in most hypertension trials. Patients had a wide range of comorbidity and concomitant medication use and, thus, the study population reflects that seen in the real world. Despite enrolling older and sometimes frail patients, the dropout rate was only 11% in 12 months. Baseline patient characteristics were well balanced between the 2 groups, and a substantial increase in 25OHD levels was achieved with the intervention.

Limitations of our study include recruitment from a single health board area and patients all being of white ethnicity, limiting generalizability. Recent data on the pharmacokinetics of vitamin D suggest that serum 25OHD levels peak at 7 days before decreasing, with a half-life of a few weeks.<sup>43,44</sup> More frequent doses (ie, monthly or every 2 months) would be more effective at raising and maintaining serum 25OHD levels; this larger dose of vitamin D could conceivably have had an effect. However, 4-month administration of 100 000 U of cholecalciferol was effective at increasing 25OHD levels and reducing fractures in a previous large osteoporosis trial,<sup>45</sup> a finding that underpinned our choice of dose timing for the current trial. Ambulatory blood pressure was somewhat lower than expected, which may have limited our ability to demonstrate reductions in ambulatory blood pressure. The size of our trial means that a small beneficial effect on blood pressure still cannot be excluded, but the clinical relevance of such small improvements, at least at an individual patient level, is questionable.

Our results do not lend support to performing large randomized controlled trials aimed specifically at blood pressure reduction, at least in this patient group. It is still possible, however, that vitamin D supplementation could have beneficial effects on cardiovascular health via non-blood pressure effects,<sup>46</sup> and ongoing large randomized trials are due to report on this in the next few years. In the meantime, our results do not support the use of high-dose, intermittent cholecalciferol to treat ISH in older white patients.

## ARTICLE INFORMATION

**Accepted for Publication:** May 28, 2013.

**Published Online:** August 12, 2013.

doi:10.1001/jamainternmed.2013.9043.

**Author Contributions:** *Study concept and design:* Witham, Price, Struthers, Donnan, Ford, McMurdo. *Acquisition of data:* Witham, Price, Struthers, Donnan, Ford, McMurdo. *Analysis and interpretation of data:* Witham, Struthers, Donnan, Messow, Ford, McMurdo. *Critical revision of the manuscript for important intellectual content:* All authors. *Administrative, technical, and material support:* All authors.

**Conflict of Interest Disclosures:** Dr Witham was funded by a Chief Scientist Office Clinician Scientist Fellowship. We also acknowledge the financial support of National Health Service Research Scotland through National Health Service Tayside in facilitating this research.

**Funding/Support:** This clinical trial was funded by the Chief Scientist Office, Scottish Government, grant CZH/4/470. The trial sponsor was the University of Dundee.

**Role of the Sponsors:** Neither the sponsor nor the funder had influence over the analysis, interpretation of the data, or preparation, review, or approval of the manuscript. Both the funder and the sponsor contributed to the design and conduct of the study, data collection, and data management.

**Additional Contributions:** Lesley McFarlane, BSc (University of Dundee), performed laboratory analyses, the Scottish Primary Care Research Network and Tayside Clinical Research Centre

provided recruitment assistance, and Valona Gouck (University of Dundee) provided secretarial support.

## REFERENCES

- Franklin SS. Hypertension in older people, part 1. *J Clin Hypertens (Greenwich)*. 2006;8(6):444-449.
- Chobanian AV. Clinical practice: isolated systolic hypertension in the elderly. *N Engl J Med*. 2007;357(8):789-796.
- Duprez DA. Systolic hypertension in the elderly: addressing an unmet need. *Am J Med*. 2008;121(3):179-184; e3.
- Primates P, Poulter NR. Hypertension management and control among English adults aged 65 years and older in 2000 and 2001. *J Hypertens*. 2004;22(6):1093-1098.
- Gosse P. Left ventricular hypertrophy as a predictor of cardiovascular risk. *J Hypertens Suppl*. 2005;23(1):S27-S33.
- Liao Y, Cooper RS, McGee DL, Mensah GA, Ghali JK. The relative effects of left ventricular hypertrophy, coronary artery disease, and ventricular dysfunction on survival among black adults. *JAMA*. 1995;273(20):1592-1597.
- SHEP Cooperative Research Group. Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension: final results of the Systolic Hypertension in the Elderly Program (SHEP). *JAMA*. 1991;265(24):3255-3264.
- Staessen JA, Fagard R, Thijs L, et al. The Systolic Hypertension in Europe (Syst-Eur) Trial Investigators. Randomised double-blind comparison of placebo and active treatment for older patients with isolated systolic hypertension. *Lancet*. 1997;350(9080):757-764.
- Liu L, Wang JG, Gong L, Liu G, Staessen JA; Collaborative Group. Comparison of active treatment and placebo in older Chinese patients with isolated systolic hypertension. *J Hypertens*. 1998;16(12, pt 1):1823-1829.
- Beckett NS, Peters R, Fletcher AE, et al; HYVET Study Group. Treatment of hypertension in patients 80 years of age or older. *N Engl J Med*. 2008;358(18):1887-1898.
- Forman JP, Giovannucci E, Holmes MD, et al. Plasma 25-hydroxyvitamin D levels and risk of incident hypertension. *Hypertension*. 2007;49(5):1063-1069.
- Forman JP, Curhan GC, Taylor EN. Plasma 25-hydroxyvitamin D levels and risk of incident hypertension among young women. *Hypertension*. 2008;52(5):828-832.
- Scragg R, Sowers M, Bell C. Serum 25-hydroxyvitamin D, ethnicity, and blood pressure in the Third National Health and Nutrition Examination Survey. *Am J Hypertens*. 2007;20(7):713-719.
- Wang TJ, Pencina MJ, Booth SL, et al. Vitamin D deficiency and risk of cardiovascular disease. *Circulation*. 2008;117(4):503-511.
- Pittas AG, Chung M, Trikalinos T, et al. Systematic review: vitamin D and cardiometabolic outcomes. *Ann Intern Med*. 2010;152(5):307-314.

16. Witham MD, Nadir MA, Struthers AD. Effect of vitamin D on blood pressure: a systematic review and meta-analysis. *J Hypertens*. 2009;27(10):1948-1954.
17. O'Brien E, Coats A, Owens P, et al. Use and interpretation of ambulatory blood pressure monitoring: recommendations of the British Hypertension Society. *BMJ*. 2000;320(7242):1128-1134.
18. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and  $\beta$ -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 1985;28(7):412-419.
19. Corretti MC, Anderson TJ, Benjamin EJ, et al; International Brachial Artery Reactivity Task Force. Guidelines for the ultrasound assessment of endothelial-dependent flow-mediated vasodilation of the brachial artery: a report of the International Brachial Artery Reactivity Task Force. *J Am Coll Cardiol*. 2002;39(2):257-265.
20. Masson LF, McNeill G, Tomany JO, et al. Statistical approaches for assessing the relative validity of a food-frequency questionnaire: use of correlation coefficients and the  $\kappa$  statistic. *Public Health Nutr*. 2003;6(3):313-321.
21. Perera S, Mody SH, Woodman RC, Studenski SA. Meaningful change and responsiveness in common physical performance measures in older adults. *J Am Geriatr Soc*. 2006;54(5):743-749.
22. Witham MD, Dove FJ, Dryburgh M, Sugden JA, Morris AD, Struthers AD. The effect of different doses of vitamin D<sub>3</sub> on markers of vascular health in patients with type 2 diabetes: a randomised controlled trial. *Diabetologia*. 2010;53(10):2112-2119.
23. Pilz S, Tomaschitz A, Ritz E, Pieber TR. Vitamin D status and arterial hypertension: a systematic review. *Nat Rev Cardiol*. 2009;6(10):621-630.
24. Wu SH, Ho SC, Zhong L. Effects of vitamin D supplementation on blood pressure. *South Med J*. 2010;103(8):729-737.
25. Sugden JA, Davies JI, Witham MD, Morris AD, Struthers AD. Vitamin D improves endothelial function in patients with type 2 diabetes mellitus and low vitamin D levels. *Diabet Med*. 2008;25(3):320-325.
26. Witham MD, Dove FJ, Sugden JA, Doney AS, Struthers AD. The effect of vitamin D replacement on markers of vascular health in stroke patients: a randomised controlled trial. *Nutr Metab Cardiovasc Dis*. 2012;22(10):864-870.
27. Tarcin O, Yavuz DG, Ozben B, et al. Effect of vitamin D deficiency and replacement on endothelial function in asymptomatic subjects. *J Clin Endocrinol Metab*. 2009;94(10):4023-4030.
28. Harris RA, Pedersen-White J, Guo DH, et al. Vitamin D<sub>3</sub> supplementation for 16 weeks improves flow-mediated dilation in overweight African-American adults. *Am J Hypertens*. 2011;24(5):557-562.
29. Witham MD, Dove FJ, Khan F, Lang CC, Belch JJ, Struthers AD. Effects of Vitamin D supplementation on markers of vascular function after myocardial infarction: a randomised controlled trial (published online March 27, 2012). *Int J Cardiol*. doi:10.1016/j.ijcard.2012.03.054.
30. Schleithoff SS, Zittermann A, Tenderich G, Berthold HK, Stehle P, Koerfer R. Vitamin D supplementation improves cytokine profiles in patients with congestive heart failure: a double-blind, randomized, placebo-controlled trial. *Am J Clin Nutr*. 2006;83(4):754-759.
31. Witham MD, Crighton LJ, Gillespie ND, Struthers AD, McMurdo ME. The effects of vitamin D supplementation on physical function and quality of life in older patients with heart failure: a randomized controlled trial. *Circ Heart Fail*. 2010;3(2):195-201.
32. Larsen T, Mose FH, Bech JN, Hansen AB, Pedersen EB. Effect of cholecalciferol supplementation during winter months in patients with hypertension: a randomized, placebo-controlled trial. *Am J Hypertens*. 2012;25(11):1215-1222.
33. Wang H, Xia N, Yang Y, Peng DQ. Influence of vitamin D supplementation on plasma lipid profiles: a meta-analysis of randomized controlled trials. *Lipids Health Dis*. 2012;11:42. doi:10.1186/1476-511X-11-42.
34. Schwartz JB. Effects of vitamin D supplementation in atorvastatin-treated patients: a new drug interaction with an unexpected consequence. *Clin Pharmacol Ther*. 2009;85(2):198-203.
35. Li YC, Kong J, Wei M, Chen ZF, Liu SQ, Cao LP. 1,25-Dihydroxyvitamin D<sub>3</sub> is a negative endocrine regulator of the renin-angiotensin system. *J Clin Invest*. 2002;110(2):229-238.
36. Forman JP, Williams JS, Fisher ND. Plasma 25-hydroxyvitamin D and regulation of the renin-angiotensin system in humans. *Hypertension*. 2010;55(5):1283-1288.
37. Tomaschitz A, Pilz S, Ritz E, et al. Independent association between 1,25-dihydroxyvitamin D, 25-hydroxyvitamin D and the renin-angiotensin system: the Ludwigshafen Risk and Cardiovascular Health (LURIC) study. *Clin Chim Acta*. 2010;411(17-18):1354-1360.
38. Bischoff-Ferrari HA, Giovannucci E, Willett WC, Dietrich T, Dawson-Hughes B. Estimation of optimal serum concentrations of 25-hydroxyvitamin D for multiple health outcomes. *Am J Clin Nutr*. 2006;84(1):18-28.
39. Dickinson HO, Nicolson DJ, Cook JV, et al. Calcium supplementation for the management of primary hypertension in adults. *Cochrane Database Syst Rev*. 2006;19(2):CD004639.
40. Bolland MJ, Avenell A, Baron JA, et al. Effect of calcium supplements on risk of myocardial infarction and cardiovascular events: meta-analysis. *BMJ*. 2010;341:c3691.
41. Sanders KM, Stuart AL, Williamson EJ, et al. Annual high-dose oral vitamin D and falls and fractures in older women: a randomized controlled trial. *JAMA*. 2010;303(18):1815-1822.
42. Avenell A, Gillespie WJ, Gillespie LD, O'Connell D. Vitamin D and vitamin D analogues for preventing fractures associated with involutional and post-menopausal osteoporosis. *Cochrane Database Syst Rev*. 2009;15(2):CD000227.
43. Ilahi M, Armas LA, Heaney RP. Pharmacokinetics of a single, large dose of cholecalciferol. *Am J Clin Nutr*. 2008;87(3):688-691.
44. Romagnoli E, Mascia ML, Cipriani C, et al. Short and long-term variations in serum calciotropic hormones after a single very large dose of ergocalciferol (vitamin D<sub>2</sub>) or cholecalciferol (vitamin D<sub>3</sub>) in the elderly. *J Clin Endocrinol Metab*. 2008;93(8):3015-3020.
45. Trivedi DP, Doll R, Khaw KT. Effect of four monthly oral vitamin D<sub>3</sub> (cholecalciferol) supplementation on fractures and mortality in men and women living in the community: randomised double blind controlled trial. *BMJ*. 2003;326(7387):469.
46. Muscogiuri G, Soric GP, Ajjan R, et al. Can vitamin D deficiency cause diabetes and cardiovascular diseases? present evidence and future perspectives. *Nutr Metab Cardiovasc Dis*. 2012;22(2):81-87.