

Investigating the stock status of Indian Ocean Bigeye tuna using the a4a statistical catch-at-age assessment method.

Iago Mosqueira^{†*} Colin P. Millar[†]

October 2013

```
## Warning: replacing previous import 'barchart' when loading 'lattice'
## Warning: replacing previous import 'bwplot' when loading 'lattice'
## Warning: replacing previous import 'densityplot' when loading 'lattice'
## Warning: replacing previous import 'dotplot' when loading 'lattice'
## Warning: replacing previous import 'histogram' when loading 'lattice'
## Warning: replacing previous import 'splom' when loading 'lattice'
## Warning: replacing previous import 'stripplot' when loading 'lattice'
## Warning: replacing previous import 'xyplot' when loading 'lattice'
## Warning: replacing previous import 'expand' when loading 'Matrix'
```

Introduction

A recently developed statistical catch-at-age stock assessment method, designed for evaluating the status of stocks with intermediate data quality, is applied here to Indian Ocean bigeye tuna. The method provides huge flexibility for the rapid evaluation of the effects in parameter estimates and uncertainty of different submodels for fishing mortality-at-age (the F model), survey catchability-at-age (the Q model), stock recruitment (the R model) and observation variance (the V model).

The model does not attempt to substitute more complex stock assessments that make use of all sources of data, like tagging, but to provide a tool for rapid exploration of the influence of different assumptions and data sources. This capability is achieved by combining a powerful optimization engine (ADMB) with an intuitive user interface using linear models as expressed in the R language.

*Contact Author: iago.mosqueira-sanchez@jrc.ec.europa.eu

[†]European Commission, Joint Research Center, IPSC/Maritime Affairs Unit, FISHREG, Via E. Fermi 2749, 21027 Ispra VA, Italy

The a4a stock assessment model

FLR (Kell et al. 2007)

Data

Catch-at-age

The catch-at-age dataset made available by the IOTC Secretariat ¹ was used.




Figure 1: Total catch series for Indian Ocean bigeye, 1950-2012.

CPUE index

Model structure and assumptions

Fixed parameters

Parameter	Ages	Value
M	0-1	0.8
	2-9	0.4

¹http://iotc.org/files/proceedings/2013/wptt/BET_ASPMv2.zip,
27/09/2013

downloaded

Estimated parameters

Model runs

Simulation of catchability fitting

Results

Population trajectories

Reference points

Residual diagnostics

Discussion

Appendix I: a4a model population dynamics

Appendix II: Code of final run

References

Kell, Laurence Thomas, Iago Mosqueira, Philippe Grosjean, Jean-Marc Fromentin, Dorleta Garcia, Richard Hillary, Ernesto Jardim, et al. 2007. “FLR: an open-source framework for the evaluation and development of management strategies.” *ICES Journal of Marine Science* 64 (apr): 640–646. doi:10.1093/icesjms/fsm012.