

# FLsp - a Surplus Production model in FLR

Finlay Scott <finlay.scott@cefas.co.uk>  
Cefas, Lowestoft, UK

June 2011

## 1 Introduction

This package implements the surplus production described in Polacheck REF. At the moment only the model including observation error is implemented. The Pella Tomlinson shape is used (which defaults to Schaeffer) Tested against the three data sets in the paper. Accurate gradients and hessian are returned using automatic differentiation (implemented using ADOLC REF)

## 2 The Model

Following the Polacheck paper Assumptions: Observation error Only  $r$  and  $k$  are estimated  $\sigma$  and  $q$  are approximated as described in the paper The initial biomass is  $k$

The general equation for the biomass through time is:

$$B_{y+1} = B_y + g(B_y) - C_y \quad (1)$$

where  $B$  is the stock biomass at the start of year  $y$ ,  $C$  is the catch during the year and  $g$  is surplus production as a function of biomass.

Here we implement the Pella-Tomlinson form of surplus production:

$$g(B) = \frac{r}{p} B (1 - (B/K)^p) \quad (2)$$

where  $r$  is the intrinsic growth rate parameter and  $K$  is the average biomass level prior to exploitation. By default,  $p$  is set to 1 making the surplus production formulation the same as a Schaefer model.

The biomass is related to an index of abundance:

$$I_y = q B_y \quad (3)$$

Where  $I$  is an index of relative abundance in year  $y$  and  $q$  is the catchability coefficient.

Here we assume that errors are introduced through observation. The population dynamics are assumed to be deterministic and all of the error occurs in the relationship between stock biomass and the index of abundance. It is assumed that the error is multiplicative and log-normal with a constant coefficient of variation. The estimates of the model parameters are ( $B_0$ ,  $r$ ,  $K$  and  $q$ ) are obtained by maximising the likelihood function:

$$L = \prod \exp(\hat{v}_y^2 / (2\hat{\sigma}_v^2)) / (\sqrt{2\pi}\hat{\sigma}_v) \quad (4)$$

where the product is over all years for which CPUE data are available and:

$$\hat{v}_y = \log(C/E)_y - \log(\hat{C}/\hat{E})_y \quad (5)$$

$$\hat{\sigma}_v^2 = \sum \hat{v}_y^2 / n \quad (6)$$

where  $n$  is the number of data points.

The value of  $q$  which maximises the likelihood is given by:

$$\hat{q} = \exp\left(\frac{1}{n} \sum_y \log(I_y / \hat{B}_y)\right) \quad (7)$$

Following Polacheck et al  $B_0$  is set to  $K$ . This means that only two parameters need to be estimated:  $r$  and  $K$ . In *FLsp* the estimation is performed using the *DEoptim* package REF.

### 3 The FLsp class

The *FLsp* class extends the *FLModel* class by including slots to store the catch and index time series. Catch is represented as an *FLQuant* and index is represented as an *FLQuants* object. This allows multiple indices to be used (not yet implemented).

To estimate the parameters  $r$  and  $K$ , an *FLsp* object must be created with catch and index data. The method *fitsp()* is then called. Once the object has been fitted, the biomass trajectory and other variables of interest (e.g.  $\sigma^2$  and  $\hat{q}$  can be calculated).

### 4 Creating and fitting FLsp objects

Here we show how to create and fit a surplus production model using *FLsp*. The data set is New Zealand Rock Lobster, taken from Polcheck REF.

```
> # Load the library
> library(FLsp)
> # Load the New Zealand Rock Lobster data set
> data(nzrl)
> # This is a dataframe with year, catch and cpue
> # Take a look at the top of it
> head(nzrl)
> # Make FLQuant objects of the catch and cpue series
> catch <- FLQuant(nzrl$catch, dimnames=list(year=nzrl$year))
> index <- FLQuant(nzrl$cpue, dimnames=list(year=nzrl$year))
> # Create the FLsp object
> nzrl <- FLsp(catch=catch, index=index)
```

After creating our object we are ready to fit the parameters.

```
> nzrl <- fitsp(nzrl)
```

The published values for this data set are:  $r = 0.0659$ ,  $K = 129000$ ,  $\hat{q} = 2.461\text{e-}5$ ,  $\sigma = 0.207$ ,  $B_{current} = 21150$ . These can be compared to our results by interrogating the *FLsp* object.

```
> # Look at the fitted parameters
```

```
> params(nzrl)
```

```
An object of class "FLPar"
```

```
params
```

```
      r      k
```

```
4.9405e-02 1.4236e+05
```

```
units: NA
```

```
> # qhat
```

```
> qhat(nzrl)
```

```
An object of class "FLQuant"
```

```
, , unit = unique, season = all, area = unique
```

```
      year
```

```
quant 1
```

```
all 2.2126e-05
```

```
units: NA
```

```
> # sigma2
```

```
> sqrt(sigma2(nzrl))
```

```
An object of class "FLQuant"
```

```
, , unit = unique, season = all, area = unique
```

```
      year
```

```
quant 1
```

```
all 0.20695
```

```
units: NA
```

```
> # returns the full biomass timeseries
```

```
> biomass(nzrl)
```

```
An object of class "FLQuant"
```

```
, , unit = unique, season = all, area = unique
```

```
      year
```

```
quant 1945  1946  1947  1948  1949  1950  1951  1952  1953  1954
```

```
all 142356 141547 140733 139893 138653 136959 134544 132075 129222 125651
```

```
      year
```

```
quant 1955  1956  1957  1958  1959  1960  1961  1962  1963  1964
```

```
all 120838 116732 111223 107376 104232 101593 99269 96711 93660 90689
```

```
      year
```

```
quant 1965  1966  1967  1968  1969  1970  1971  1972  1973  1974
```

```
all 87718 84398 80800 77744 74513 71481 68540 65818 64072 62028
```

```
      year
```

```
quant 1975  1976  1977  1978  1979  1980  1981  1982  1983  1984
```

```
all 60115 58843 57238 55692 53949 51554 48989 46518 43734 40846
```

```
      year
```

```
quant 1985  1986  1987  1988  1989  1990
```

```
all 37374 33880 30498 27182 25141 422846
```

It can be seen that there is good agreement between the published results and those generated with *FLsp*. The differences are likely caused by the precision of the printed data set in the Polcheck paper (REF) and the fitting method used.

## 5 Testing FLsp against the other data sets

There are two other data sets available: South Atlantic albacore (*saa*) and Northern Namibian hake (*nnh*). The above process can be repeated and the results checked against the published results.

TABLE OF RESULTS

	New Zealand Rock Lobster		South Atlantic Albacore		Northern Namibian Hake	
Measure	FLsp	Published	FLsp	Published	FLsp	Published
r	0.0494	0.0659	0.3197	0.328	0.3701	0.379
K ('000 t)	142.3564741	129.0	243.3652	239.6	2823.0843	2772.6
$\sigma$	0.207	0.207	0.11	0.111	0.125	0.124
$\hat{q}$ ( $\times 10^4$ )	0	24.61	0.264	26.71	0	4.360

## 6 Plotting results

There is no generic plot for FLsp at the moment. However, it is possible to look at the fitted index and residuals using relatively simple code. For example, to plot the indices with the fitted indices you can use (see Figure 1):

```
> fitted <- cbind(as.data.frame(nzrl@fitted_index),type="fitted")
> index <- cbind(as.data.frame(nzrl@index),type="index")
> index <- rbind(index,fitted)
> print(xyplot(data ~ year | qname, group=type, data=index, type="b",auto.key=TRUE))
```

To look at the residuals and put a loess function through them use (see Figure 2):

```
> residuals <- as.data.frame(nzrl@residuals_index)
> print(xyplot(data ~ year | qname, data = residuals, panel = function(x,
+   y) {
+     panel.xyplot(x, y)
+     panel.loess(x, y, span = 1)
+   }))
```

## 7 Profiling the fit




Figure 1: Indices and fitted indices for New Zealand rock lobster




Figure 2: Residuals for New Zealand rock lobster