

Calculus, Volume 1, 2nd Edition - Tom M.
Apostal

Iain Wong

November 25th, 2021

Contents

1	Introduction	3
1.1	Some Basic Concepts of the Theory of Sets	3
1.1.1	Excercises	3

Chapter 1

Introduction

1.1 Some Basic Concepts of the Theory of Sets

1.1.1 Exercises

1. Use the roster notation to designate the following sets of real numbers.

Proposition 1. $A = \{x|x^2 - 1 = 0\}$ can be designated as $\{-1, 1\}$ in roster notation.

Proof.

$$\begin{aligned} A &= \{x|x^2 - 1 = 0\} \\ &= \{x|(x - 1)(x + 1) = 0\} \\ &\therefore \{-1, 1\} \end{aligned} \tag{1.1}$$

QED

Proposition 2. $B = \{x|(x - 1)^2 = 0\}$ can be designated as $\{1\}$ in roster notation.

Proof.

$$\begin{aligned} B &= \{x|(x - 1)^2 = 0\} \\ &= \{x|x - 1 = \sqrt{0}\} \\ &= \{x|x = 1\} \\ &\therefore \{1\} \end{aligned} \tag{1.2}$$

QED

Proposition 3. $C = \{x|x + 8 = 9\}$ can be designated as $\{1\}$ in roster notation.

Proof.

$$\begin{aligned}
 C &= \{x|x + 8 = 9\} \\
 &= \{x|x = 9 - 8\} \\
 &= \{x|x = 1\} \\
 &\therefore \{1\}
 \end{aligned}
 \tag{1.3}$$

QED

Proposition 4. $D = \{x|x^3 - 2x^2 + x = 2\}$ can be designated as $\{2\}$ in roster notation.

Proof.

$$\begin{aligned}
 D &= \{x|x^3 - 2x^2 + x = 2\} \\
 &= \{x|x^3 - 2x^2 + x - 2 = 0\} \\
 &= \{x|x^2(x - 2) + (x - 2) = 0\} \\
 &= \{x|(x^2 + 1)(x - 2) = 0\} \\
 &\therefore \{2\}
 \end{aligned}
 \tag{1.4}$$

QED

Proposition 5. $E = \{x|(x + 8)^2 = 9^2\}$ can be designated as $\{-17, 1\}$ in roster notation.

Proof.

$$\begin{aligned}
 E &= \{x|(x + 8)^2 = 9^2\} \\
 &= \{x|x + 8 = \pm 9\} \\
 &= \{x|x = \pm 9 - 8\} \\
 &\therefore \{-17, 1\}
 \end{aligned}
 \tag{1.5}$$

QED

Proposition 6. $F = \{x|(x^2 + 16x)^2 = 17^2\}$ can be designated as $\{-17, 1, -8 - \sqrt{47}, -8 + \sqrt{47}\}$ in roster notation.

Proof.

$$\begin{aligned}
 F &= \{x|(x^2 + 16x)^2 = 17^2\} \\
 &= \{x|x^2 + 16x = \pm 17\} \\
 &= \{x|x^2 + 16x \pm 17 = 0\} \\
 &= \{x|x^2 + 16x \pm 17 = 0\}
 \end{aligned} \tag{1.6}$$

Using the quadratic formula:

Definition 1.1.1. Quadratic Equation, analytical method for calculating the roots of a quadratic polynomial.

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}, \text{ where } ax^2 + bx + c = 0 \tag{1.7}$$

Solving when the last term is +17:

$$\begin{aligned}
 x &= \frac{-16 \pm \sqrt{16^2 - 4(1)17}}{2(1)} \\
 &= -8 \pm \frac{\sqrt{188}}{2} \\
 &= -8 \pm \frac{\sqrt{188}}{\sqrt{2^2}} \\
 &= -8 \pm \sqrt{188/4} \\
 &= -8 \pm \sqrt{47} \\
 \therefore &\{-8 - \sqrt{47}, -8 + \sqrt{47}\}
 \end{aligned} \tag{1.8}$$

Solving when the last term is -17:

$$\begin{aligned}
 0 &= x^2 + 16x - 17 \\
 &= (x + 17)(x - 1) \\
 \therefore &\{-17, 1\}
 \end{aligned} \tag{1.9}$$

QED

2. For the sets in Exercise 1, note that $B \subseteq A$. List all the inclusion relations \subseteq that hold among the sets A, B, C, D, E, F .

(a) $B \subseteq A$

(b) $B \subseteq C$

(c) $B \subseteq E$

(d) $B \subseteq F$

(e) $C \subseteq A$

(f) $C \subseteq B$

(g) $C \subseteq E$

(h) $C \subseteq F$

(i) $E \subseteq F$