

# Multiple testing in ANOVA and SEM

Enrico Toffalini, Irene Alfarone

2025-04-17



# Overview

We present two simulated scenarios to illustrate the issue of multiple testing, by looking at false positive rates and power in:

- ▶ ANOVA
- ▶ Structural Equation Modeling (SEM)

# ANOVA

We simulated 5000 ANOVA with 3 binary predictors (A, B, C) and one continuous predictor (D), under  $H_0$  and  $H_1$  ( $y = 1 + 0.20 \cdot C + \text{residual}$ ).

We present here the results of the simulation under  $H_1$ .

# ANOVA

## Anova Table (Type II tests)

Response: y


	Sum Sq	Df	F value	Pr(>F)	
A	1	1	1.1134	0.29135	
B	0	1	0.1302	0.71827	
C	990	1	986.6354	< 2e-16	***
D	0	1	0.4122	0.52084	
A:B	1	1	0.5640	0.45265	
A:C	0	1	0.2892	0.59075	
B:C	1	1	0.5126	0.47402	
A:D	4	1	4.2249	0.03984	*
B:D	0	1	0.4730	0.49161	
C:D	0	1	0.0167	0.89720	
A:B:C	0	1	0.1156	0.73384	
A:B:D	1	1	1.4762	0.22438	
A:C:D	0	1	0.1241	0.72464	
B:C:D	0	1	0.4967	0.48097	

## Results of the simulation


Table 1: Under H1 (true effect of C)

Metric	Value
Power	88.8%
Power (after correction; FDR)	60.6%
At least one significant effect	94.2%

SEM



# SEM



# SEM




Figure 3: Celestial SEM (from Celestial Mediation)



# Results

Table 2: Under H1 (true path  $F3 \sim F1$ )

Metric	Value
Power	94.6%
Power (after correction; FDR)	72.8%
At least one significant effect	97.4%