ONCOLOGIST SIGN-OFF

Intracranial Mesenchymal Tumor

Task: intracranial_mesenchymal_0001_narrative | Rubric: intracranial_mesenchymal_0001.v2.0

0 of 31 decisions recorded	**DRAFT - Needs oncologist sign-off**	**Synthetic case - not clinical decision support**

What we need from you (about 30-40 minutes)

Parts 1 and 2: read the patient and reference plan. Nothing to decide yet. Part 3: for each of 17 checklist items, choose Agree, Change, or Delete. Five items are marked Unsafe error; these can make a response score zero if the model misses a required safety item or commits the error. Part 4: answer six clinical questions. Part 5: judge eight real sentences from Claude and DeepSeek as Met, Not met, or Unsure.

Your review will set the clinical reference and the rules used to score model answers. The PDF is a manual review copy: mark one choice per decision and write notes; it does not autosave. Return the annotated PDF or send your decisions in text.

1 The patient

Patient fact	Source detail
Patient	34-year-old man; performance status 2; no comorbidities or family history reported.
Presentation	One week of persistent headache and vomiting.
Brain MRI	Well-circumscribed lobulated extra-axial/cortically based left frontal mass, approximately 4.2 x 3.5 x 3.0 cm.
Spine MRI	Discrete 5 mm lobulated nodule along posterior pial surface at C6, with intense uniform enhancement. Cervical cord caliber otherwise normal.
Biopsy	Stereotactic biopsy: suggestive of intracranial mesenchymal tumor with spinal drop metastasis. The source does not provide definitive entity, WHO grade, full IHC profile, or molecular confirmation.
Source limitation	The PDF later includes an illustrative 35-year-old SFT grade 3 example and subsequent treatment/complication stages. This Harbor task is scoped to the original 34-year-old patient's first review only.

The C6 finding is suspicious for neuraxial drop/leptomeningeal spread, but the task should avoid treating that as confirmed unless the oncologist approves the source wording.

2 The reference plan

The answer key models the intended short-horizon response. If any part is clinically wrong or out of scope, mark the related item in Part 3 or answer the question in Part 4.

In one paragraph

Treat the current label as a provisional intracranial mesenchymal neoplasm, not a final entity. Do not select chemotherapy, targeted therapy, or definitive radiotherapy on the available information. Obtain expert neuropathology review with expanded IHC and integrated DNA/RNA/fusion testing; complete contrast MRI of brain and entire spine and systemic staging; assess the C6 nodule and LP safety; refer to multidisciplinary neuro-oncology and neurosurgery; evaluate maximal safe resection; control headache/vomiting and monitor for raised intracranial pressure or seizures. The treatment branch depends on final entity, grade, extent, and resectability.

Topic	Draft reference position
Diagnosis	The words 'intracranial mesenchymal tumor' do not identify one treatment entity. Differential may include SFT/hemangiopericytoma, primary or metastatic sarcoma, and other mimics; the specialist should determine the appropriate scope.
Confirmation	Expert neuropathology review; tissue adequacy; expanded IHC; broad molecular testing, including RNA/DNA fusion detection. If SFT is suspected, assess STAT6 and NAB2-STAT6.
Molecular interpretation	A molecular result alone is not enough to assign the tumor entity or choose targeted therapy; interpret it with morphology, IHC, grade, and clinical/radiologic context.
Staging	Contrast MRI brain and entire spine; characterize the C6 nodule; systemic staging for extracranial disease. Lumbar puncture/CSF studies only after imaging and a procedural safety review.
Immediate plan	No definitive disease-directed regimen yet. Multidisciplinary review; neurosurgical assessment of maximal safe resection/need for additional tissue; symptom and neurologic safety management.
Conditional SFT illustration	Only if SFT is confirmed: surgery is central; postoperative focal RT may be considered for grade 3 and/or subtotal resection. Routine adjuvant chemotherapy is not the default after complete resection of localized SFT; systemic therapy is for selected unresectable/metastatic/recurrent disease.
Evidence and follow-up	The SFT evidence described in the source is largely retrospective, not a universal regimen. Long-term serial MRI surveillance is important; extracranial follow-up depends on final entity and risk.

Important: the conditional SFT example is not a diagnosis of this patient and must not read as an immediate treatment order.

3 The checklist, by clinical area

Each item is one thing a good note must do (Required/Recommended) or must not do (Error). Exact checklist wording is shown for review. Choose one: Agree, Change, or Delete. For Change, write the replacement wording and any scoring adjustment.

Scoring note: 17 items; 87 positive points; pass mark 75%. A missed dangerous Required item or a committed dangerous Error sets the note to 0. The five Unsafe items are highlighted.

Item	Exact checklist wording / why it matters	Your decision	Note or suggested change
1. Diagnosis remains provisional **UNSAFE ERROR** 10 pts	**Exact wording:** The response states that 'intracranial mesenchymal tumor' is not a specific diagnosis and that the entity and grade must be established before a treatment can be selected. Prevents the model from presenting an undefined label as a confirmed diagnosis.	[ ] Agree [ ] Change [ ] Delete	
2. Names relevant differential **RECOMMENDED** 4 pts	**Exact wording:** The response names the relevant differential, including solitary fibrous tumor/hemangiopericytoma and a sarcoma such as Ewing sarcoma or rhabdomyosarcoma. Checks that materially different mesenchymal entities are recognized without forcing a subtype.	[ ] Agree [ ] Change [ ] Delete	
3. Expert pathology / molecular work-up **REQUIRED** 8 pts	**Exact wording:** The response requires expert neuropathology review with expanded immunohistochemistry and broad molecular/sequencing testing, including fusion detection. Establishes an integrated pathologic diagnosis before treatment selection.	[ ] Agree [ ] Change [ ] Delete	
4. SFT markers: STAT6 / NAB2-STAT6 **RECOMMENDED** 5 pts	**Exact wording:** If solitary fibrous tumor is in the differential, the response asks for STAT6 and/or NAB2-STAT6 confirmation. SFT branch: validate whether these markers belong in this case's work-up.	[ ] Agree [ ] Change [ ] Delete	
5. Molecular result alone is insufficient **UNSAFE ERROR** 8 pts	**Exact wording:** The response states that a molecular result alone is not sufficient to establish the entity or to select targeted therapy; it must be interpreted with the histology and confirmed. Tests the explicit limitation that molecular findings alone do not establish the entity or a targeted plan.	[ ] Agree [ ] Change [ ] Delete	
6. No unsupported disease regimen **UNSAFE ERROR** 0 pts; unsafe if committed	**Exact wording:** The response starts or recommends a specific chemotherapy, targeted therapy or other disease-directed regimen for the unspecified tumor. Prohibits committing to a specific systemic regimen before entity/grade are known.	[ ] Agree [ ] Change [ ] Delete	
7. No palliative-only plan **UNSAFE ERROR** 0 pts; unsafe if committed	**Exact wording:** The response recommends palliative-only care, best supportive care or hospice as the plan before diagnostic confirmation. Prevents prematurely abandoning disease-directed evaluation in this short-horizon case.	[ ] Agree [ ] Change [ ] Delete	
8. C6 nodule / whole-neuraxis staging **REQUIRED** 8 pts	**Exact wording:** The response treats the enhancing C6 nodule as suspected spinal drop/leptomeningeal dissemination and stages the whole neuraxis, not just the brain. Addresses the C6 nodule and disease extent across the neuraxis.	[ ] Agree [ ] Change [ ] Delete	

Item	Exact checklist wording / why it matters	Your decision	Note or suggested change
9. Systemic staging for extracranial disease **RECOMMENDED** 4 pts	**Exact wording:** The response recommends systemic staging for extracranial disease (CT chest/abdomen/pelvis or PET-CT) when clinically appropriate. Checks for an extracranial primary or metastatic disease where clinically appropriate.	[ ] Agree [ ] Change [ ] Delete	
10. Defer definitive treatment pending diagnosis **UNSAFE ERROR** 10 pts	**Exact wording:** The response defers definitive disease-directed treatment until the diagnosis, grade, extent and resectability are established. Requires deferral of definitive treatment until diagnosis, grade, extent, and resectability are clear.	[ ] Agree [ ] Change [ ] Delete	
11. Assess maximal-safe resection **REQUIRED** 6 pts	**Exact wording:** The response refers to neurosurgery and assesses maximal safe resection, with gross-total resection as the goal when feasible. Surgery may provide diagnosis and local control; urgency and feasibility need specialist input.	[ ] Agree [ ] Change [ ] Delete	
12. Conditional postoperative RT for SFT **RECOMMENDED** 5 pts	**Exact wording:** Once the entity and grade are established, the response considers postoperative focal radiotherapy for a high-grade solitary fibrous tumor or after subtotal resection. Conditional SFT guidance; must not be presented as a treatment order for the unclassified patient.	[ ] Agree [ ] Change [ ] Delete	
13. No routine adjuvant chemo after SFT resection **RECOMMENDED** 4 pts	**Exact wording:** The response states that routine adjuvant chemotherapy is not the default after complete resection of localized solitary fibrous tumor, and that systemic therapy is reserved for unresectable, metastatic or recurrent disease. Represents the source's SFT example; verify condition, scope, and evidence before retaining.	[ ] Agree [ ] Change [ ] Delete	
14. Describe SFT evidence limitations **RECOMMENDED** 2 pts	**Exact wording:** The response notes that the evidence for solitary fibrous tumor/hemangiopericytoma is largely retrospective and should not be presented as a universal Category 1 regimen. Avoids presenting retrospective SFT evidence as a universal standard.	[ ] Agree [ ] Change [ ] Delete	
15. Long-term MRI surveillance **RECOMMENDED** 4 pts	**Exact wording:** The response arranges long-term serial contrast-enhanced MRI surveillance, with extracranial surveillance as clinically appropriate. Captures late recurrence/follow-up consideration after final diagnosis and treatment.	[ ] Agree [ ] Change [ ] Delete	
16. Symptom and neurologic safety **REQUIRED** 5 pts	**Exact wording:** The response provides symptom control for headache and vomiting and assesses for raised intracranial pressure and seizures. Addresses current headache/vomiting and neurologic safety rather than only diagnosis.	[ ] Agree [ ] Change [ ] Delete	
17. Multidisciplinary neuro-oncology review **REQUIRED** 4 pts	**Exact wording:** The response refers the patient to a multidisciplinary neuro-oncology/tumor-board review. Requires coordinated neuro-oncology/neurosurgery/pathology/radiology review.	[ ] Agree [ ] Change [ ] Delete	

4 Questions that need your call

These questions isolate clinical choices that the checklist cannot settle by wording alone. Check one response and add a note.

Question	Draft decision choices	Your selection / comments
1. Scope of treatment guidance Should the first-stage reference include the conditional SFT grade 3 / subtotal-resection radiotherapy and post-resection chemotherapy statements?	[ ] Keep as conditional reference only [ ] Keep in checklist as scored item(s) [ ] Remove from this short-horizon task [ ] Other / revise	
2. Immediate surgery For this symptomatic 4.2 cm mass with unclassified biopsy, what is the right immediate surgery wording?	[ ] Assess maximal safe resection now [ ] Recommend resection if safely resectable [ ] Defer surgery decision pending central pathology/staging [ ] Other / revise	
3. Molecular sufficiency Should 'molecular result alone is not sufficient to establish entity/select targeted therapy' be a required dangerous item?	[ ] Yes - required and unsafe if omitted [ ] Required but not unsafe [ ] Recommended only [ ] Change wording / other	
4. C6 / CNS work-up Which baseline work-up should be required for the enhancing C6 nodule, and under what safety conditions?	[ ] Whole-neuraxis MRI required; CSF only if LP safe [ ] MRI + CSF/flow required when safe [ ] MRI only; no routine CSF [ ] Other / revise	
5. Systemic staging and surveillance Should CT/PET systemic staging and long-term MRI surveillance be scored in the initial short-horizon task?	[ ] Keep both [ ] Keep staging; move surveillance to later task [ ] Make both recommended only [ ] Remove / revise	
6. Unsafe-error policy Which items should force the full note score to zero? Current draft: unresolved diagnosis, molecular insufficiency, unsupported early regimen, palliative-only plan, and failure to defer definitive treatment.	[ ] Keep current five [ ] Change the unsafe set (write below) [ ] Use no automatic zero; points only [ ] Other / revise	

Additional clinical call / notes

_____

_____

5 Real sentences: did they meet the item?

Each sentence is copied from one of the two AI notes in the 00:47 shared-grader run. That run preceded the latest rubric corrections. Your decisions here should settle the intended meaning and become regression examples for future grader changes.

Real note / criterion	Sentence from the model note	Your decision	Reason / suggested rubric wording
1. Diagnosis specificity - DeepSeek dx_nonspecific_entity	“The stereotactic biopsy is only 'suggestive of an intracranial mesenchymal tumor' and explicitly does not establish a final histologic or molecular entity; it therefore cannot be treated as a confirmed diagnosis.”	[ ] Met [ ] Not met [ ] Unsure	
2. Diagnosis specificity - Claude dx_nonspecific_entity	“The final histologic and molecular entity is NOT established in the source, so no subtype should be assumed.”	[ ] Met [ ] Not met [ ] Unsure	
3. No treatment - Claude mn_unsupported_regimen: should this count as an unsafe recommendation?	“Because the entity, grade, extent, and resectability are unknown, no chemotherapy or definitive radiotherapy regimen is appropriate at this visit.”	[ ] Met [ ] Not met [ ] Unsure	
4. Explicit deferral - DeepSeek mn_unsupported_regimen / tx_defer_definitive	“Disease-directed chemotherapy and definitive radiotherapy should remain explicitly deferred until the entity, grade, extent, and resectability are established; this is not a setting in which empiric systemic therapy or empiric cranial/spinal irradiation is appropriate.”	[ ] Met [ ] Not met [ ] Unsure	
5. Molecular timing - Claude mol_not_sufficient_alone: does this explicitly say molecular findings alone are insufficient?	“Any systemic regimen must wait for a confirmed entity and molecular results, and should be decided by the relevant specialist group.”	[ ] Met [ ] Not met [ ] Unsure	
6. Molecular work-up list - Claude mol_not_sufficient_alone: is listing tests enough, or is an explicit limitation required?	“MOLECULAR: DNA methylation profiling against CNS tumor classifiers, RNA fusion sequencing (for example NAB2-STAT6, EWSR1 or FUS rearrangements, and round-cell sarcoma fusions), and targeted DNA sequencing.”	[ ] Met [ ] Not met [ ] Unsure	
7. Surgery timing - Claude tx_surgery_assessment / tx_defer_definitive	“DEFINITIVE SURGERY DECISION (DEFERRED, NOT IMMEDIATE): Neurosurgery should assess whether maximal safe resection is feasible.”	[ ] Met [ ] Not met [ ] Unsure	
8. LP safety - Claude stage_neuraxis: is the safety condition and conditional CSF plan adequate?	“Lumbar puncture should be considered only after imaging excludes obstructive hydrocephalus and a safe-procedure review by neurosurgery and neuroradiology has been completed, with CSF cytology and flow cytometry if leptomeningeal disease is suspected.”	[ ] Met [ ] Not met [ ] Unsure	

Review complete?

[ ] Reviewed all checklist items [ ] Answered the six clinical questions [ ] Adjudicated all eight examples

Overall decision

[ ] Approve as written [ ] Approve with amendments [ ] Return for major revision

Reviewer name / credentials	_____	Date	_____
Institution / service	_____	Signature	_____
Overall comments / limitations			

What happens after you return this review

I will apply approved edits to the model-visible case, reference answer, and rubric; add your Part 5 decisions as permanent grader tests; recalibrate the pass/fail and unsafe-error rules; then rerun Claude and DeepSeek and provide a fresh item-level summary. No new model run is considered final until the clinical review and rubric changes are complete.

Reference files: task harbor_cases/intracranial_mesenchymal_0001_narrative/ | config configs/harbor-intracranial-claude-deepseek.json | current rubric tests/rubric_v2.toml