

Lecture 04

Multivariable linear regression

Recap


- Hypothesis

$$H(x) = Wx + b$$

- Cost function
= **Loss function**

$$cost(W, b) = \frac{1}{m} \sum_{i=1}^m (\overset{\text{prediction}}{H(x^{(i)})} - \overset{\text{true value}}{y^{(i)}})^2$$

- Gradient descent algorithm



convex

Predicting exam score: regression using one input (x)

one-variable
one-feature

x (hours)	y (score)
10	90
9	80
3	50
2	60
11	40

Predicting exam score: regression using three inputs (x_1 , x_2 , x_3)

multi-variable/feature

Test Scores for General Psychology

x_1 (quiz 1)	x_2 (quiz 2)	x_3 (midterm 1)	Y (final)
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142

Hypothesis


$$H(x) = Wx + b$$



$$H(x_1, x_2, x_3) = w_1x_1 + w_2x_2 + w_3x_3 + b$$

Cost function


$$H(x_1, x_2, x_3) = w_1x_1 + w_2x_2 + w_3x_3 + b$$



$$cost(W, b) = \frac{1}{m} \sum_{i=1}^m (H(x_1^{(i)}, x_2^{(i)}, x_3^{(i)}) - y^{(i)})^2$$

Multi-variable

$$H(x_1, x_2, x_3) = w_1x_1 + w_2x_2 + w_3x_3 + b$$




$$H(x_1, x_2, x_3, \dots, \underline{x_n}) = w_1x_1 + w_2x_2 + w_3x_3 + \dots + \underline{w_nx_n} + b$$

Matrix

$$w_1x_1 + w_2x_2 + w_3x_3 + \dots + w_nx_n$$

Matrix multiplication

"Dot Product"

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \times \begin{bmatrix} 7 & 8 \\ 9 & 10 \\ 11 & 12 \end{bmatrix} = \begin{bmatrix} 58 \\ \end{bmatrix}$$


$$1 \times 7 = 7$$

$$2 \times 9 = 18$$

$$3 \times 11 = 33$$

$$7 + 18 + 33 = 58$$

Hypothesis using matrix

$$w_1x_1 + w_2x_2 + w_3x_3 + \dots + w_nx_n$$

$$(x_1 \quad x_2 \quad x_3) \cdot \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = (x_1w_1 + x_2w_2 + x_3w_3)$$

$$H(x) = Wx + b \quad \rightarrow \quad H(X) = XW$$

Hypothesis using matrix

Test Scores for General Psychology

x_1	x_2	x_3	Y
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142

$$H(x) = Wx + b$$



$$H(x_1, x_2, x_3) = x_1w_1 + x_2w_2 + x_3w_3 + b$$

Hypothesis using matrix

Test Scores for General Psychology

x_1	x_2	x_3	Y
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142

$$H(x_1, x_2, x_3) = x_1w_1 + x_2w_2 + x_3w_3 + \cancel{b}$$



$$(x_1 \quad x_2 \quad x_3) \cdot \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = (x_1w_1 + x_2w_2 + x_3w_3)$$

$$H(X) = XW$$

Many x instances

Test Scores for General Psychology

Instance : 5

x_1	x_2	x_3	Y
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142

$$\text{Instance } (x_1 \quad x_2 \quad x_3) \cdot \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = (x_1 w_1 + x_2 w_2 + x_3 w_3)$$

Hypothesis using matrix

x_1	x_2	x_3	Y
73	80	75	152
93	88	93	185
89	91	90	180
96	98	100	196
73	66	70	142

$$w_1x_1 + w_2x_2 + w_3x_3 + \dots + w_nx_n$$

Instance

$$\begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \\ x_{41} & x_{42} & x_{43} \\ x_{51} & x_{52} & x_{53} \end{pmatrix} \cdot \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = \begin{pmatrix} x_{11}w_1 + x_{12}w_2 + x_{13}w_3 \\ x_{21}w_1 + x_{22}w_2 + x_{23}w_3 \\ x_{31}w_1 + x_{32}w_2 + x_{33}w_3 \\ x_{41}w_1 + x_{42}w_2 + x_{43}w_3 \\ x_{51}w_1 + x_{52}w_2 + x_{53}w_3 \end{pmatrix}$$

$$H(X) = XW$$

Hypothesis using matrix

$$\begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \\ x_{41} & x_{42} & x_{43} \\ x_{51} & x_{52} & x_{53} \end{pmatrix} \cdot \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = \begin{pmatrix} x_{11}w_1 + x_{12}w_2 + x_{13}w_3 \\ x_{21}w_1 + x_{22}w_2 + x_{23}w_3 \\ x_{31}w_1 + x_{32}w_2 + x_{33}w_3 \\ x_{41}w_1 + x_{42}w_2 + x_{43}w_3 \\ x_{51}w_1 + x_{52}w_2 + x_{53}w_3 \end{pmatrix}$$

$$\boxed{[5,} \quad 3] \quad [3, \boxed{1}] = \boxed{[5,} \quad \boxed{1]}$$

$$H(X) = XW$$

Hypothesis using matrix

$$\begin{array}{ccc}
 \left[\begin{array}{|c|} \hline X \\ \hline \end{array} \right] & \times & \left[\begin{array}{|c|} \hline W \\ \hline \end{array} \right] = \left[\begin{array}{|c|} \hline H(X) \\ \hline \end{array} \right] \\
 \text{Input} & & \text{Output} \\
 \boxed{[5, 3]} & \boxed{[3, 1]} & \boxed{[5, 1]} \\
 \text{Instance : 5} & & \text{Instance : 5} \\
 X \text{ (variable value) : 3} & & Y \text{ (variable value) : 1}
 \end{array}$$

$H(X) = XW$

Hypothesis using matrix

$$\begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \\ x_{41} & x_{42} & x_{43} \\ x_{51} & x_{52} & x_{53} \end{pmatrix} \cdot \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = \begin{pmatrix} x_{11}w_1 + x_{12}w_2 + x_{13}w_3 \\ x_{21}w_1 + x_{22}w_2 + x_{23}w_3 \\ x_{31}w_1 + x_{32}w_2 + x_{33}w_3 \\ x_{41}w_1 + x_{42}w_2 + x_{43}w_3 \\ x_{51}w_1 + x_{52}w_2 + x_{53}w_3 \end{pmatrix}$$

$[n, 3]$ $[3, 1]$ $[n, 1]$

None $H(X) = XW$ None

Hypothesis using matrix (n output)

$$\begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \\ x_{41} & x_{42} & x_{43} \\ x_{51} & x_{52} & x_{53} \end{pmatrix} \cdot \text{?} = \begin{pmatrix} x_{11}w_{11} + x_{12}w_{21} + x_{13}w_{31} & x_{11}w_{12} + x_{12}w_{22} + x_{13}w_{32} \\ x_{21}w_{11} + x_{22}w_{21} + x_{23}w_{31} & x_{21}w_{12} + x_{22}w_{22} + x_{23}w_{32} \\ x_{31}w_{11} + x_{32}w_{21} + x_{33}w_{31} & x_{31}w_{12} + x_{32}w_{22} + x_{33}w_{32} \\ x_{41}w_{11} + x_{42}w_{21} + x_{43}w_{31} & x_{41}w_{12} + x_{42}w_{22} + x_{43}w_{32} \\ x_{51}w_{11} + x_{52}w_{21} + x_{53}w_{31} & x_{51}w_{12} + x_{52}w_{22} + x_{53}w_{32} \end{pmatrix}$$

$$\begin{matrix} & \text{---} & & \text{---} & & \text{---} \\ \text{[n, 3]} & \text{[3, 2]} & & & & \text{[n, 2]} \end{matrix}$$

Output : 2

$$H(X) = XW$$

Hypothesis using matrix (n output)

$$\begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \\ x_{41} & x_{42} & x_{43} \\ x_{51} & x_{52} & x_{53} \end{pmatrix} \cdot \begin{pmatrix} w_{11} & w_{12} \\ w_{21} & w_{22} \\ w_{31} & w_{32} \end{pmatrix} = \begin{pmatrix} x_{11}w_{11} + x_{12}w_{21} + x_{13}w_{31} & x_{11}w_{12} + x_{12}w_{22} + x_{13}w_{32} \\ x_{21}w_{11} + x_{22}w_{21} + x_{23}w_{31} & x_{21}w_{12} + x_{22}w_{22} + x_{23}w_{32} \\ x_{31}w_{11} + x_{32}w_{21} + x_{33}w_{31} & x_{31}w_{12} + x_{32}w_{22} + x_{33}w_{32} \\ x_{41}w_{11} + x_{42}w_{21} + x_{43}w_{31} & x_{41}w_{12} + x_{42}w_{22} + x_{43}w_{32} \\ x_{51}w_{11} + x_{52}w_{21} + x_{53}w_{31} & x_{51}w_{12} + x_{52}w_{22} + x_{53}w_{32} \end{pmatrix}$$

[n, 3]

[3, 2]

[n, 2]

$$H(X) = XW$$

WX vs XW

- Lecture (theory):

$$H(x) = Wx + b$$

- Implementation (TensorFlow)

$$H(X) = XW$$